

NCBI Bookshelf. A service of the National Library of Medicine, National Institutes of Health.

StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-.

Audiogram Interpretation

Mandy K. Salmon; Jason Brant; Marc H. Hohman; Danielle Leibowitz.

► Author Information and Affiliations

Last Update: March 1, 2023.

Continuing Education Activity

Audiometry is an essential means of assessing hearing thresholds in patients with hearing loss. The audiogram quantifies and visually displays a patient's degree and type of hearing loss (sensorineural, conductive, or mixed). Furthermore, various etiologies of hearing loss may demonstrate characteristic patterns on the audiogram, making this testing modality critical for diagnostic purposes. Audiograms are most useful for objectively identifying deficits in hearing by looking at how sound travels through the auditory system via air or bone conduction. Air conduction measures sound transmission from the outer ear, through the middle ear, and into the inner ear, where signals are sent to the central nervous system (CNS) for auditory processing. In contrast, bone conduction measures transmission only from the inner ear to the CNS. This activity reviews the interpretation of audiograms and their role in identifying patterns of otologic disease.

Objectives:

- Evaluate the concept of pure tone thresholds and how they are measured during audiometry.
- Identify air and bone conduction notation on the audiogram and understand the concept of masking.
- Assess the differences between sensorineural and conductive hearing loss as shown on an audiogram.

[Access free multiple choice questions on this topic.](#)

Introduction

More than 16% of adult Americans experience some degree of hearing loss.[1] Accurately diagnosing and managing hearing loss is essential in patients of all ages, whether for safety and quality of life in adults or for developmental reasons in children. Subtle findings in young patients may warrant referral for audiometry, as children with slight hearing impairment may suffer from speech and language delays. Audiometric threshold data, also known as pure-tone testing, have been used clinically since the 1920s to categorize the degree and type of hearing loss. Audiometric testing utilizes various intensities of sound emitted over a range of frequencies to determine deficits in hearing, and results are plotted on a graph known as an audiogram. The configurations and patterns displayed on the audiogram help healthcare providers understand the etiology and severity of a patient's hearing loss.

Audiograms are most useful for objectively identifying deficits in hearing by looking at how sound travels through the auditory system via air or bone conduction. Air conduction measures

sound transmission from the outer ear, through the middle ear, and into the inner ear, where signals are sent to the central nervous system (CNS) for auditory processing. In contrast, bone conduction measures transmission only from the inner ear to the CNS.[2][3] Despite the ability of this powerful tool to guide clinical practice in managing patients with hearing loss, many healthcare providers have difficulty accurately interpreting an audiogram.[4] This review aims to explore the basic principles of audiometry and discuss the audiogram's interpretation, including examples of common pathologies that can be diagnosed via audiometry.

Procedures

Pure-tone evaluations are performed in sound-treated rooms to eliminate ambient noise while obtaining hearing thresholds for air-conducted and bone-conducted sounds. Air conduction tones are presented through inserted earphones or over-the-ear headphones. Bone conduction thresholds are obtained via an oscillating transducer placed on the mastoid process. Most commonly, tones at the frequency being tested are presented at decreasing intensity levels (in 5 or 10 dB steps) until becoming inaudible and then are again presented at increasing intensity until becoming audible again, a procedure known as the Hughson-Westlake method.[5]

Pure tone thresholds are then plotted on an audiogram, displaying the quietest sound at a given frequency that a patient can discern at least 50% of the time, measured in decibels Hearing Level (dB HL). Hearing Level is a scale based upon normative data for human hearing thresholds, devised such that 0 dB is the loudness at which most people can perceive a sound 50% of the time; individuals that can hear better than average may, therefore, have hearing thresholds that are lower than 0 dB. Some variability across the population is expected, and hearing thresholds up to 25 dB are considered normal. Variability is also expected when testing the same patient due to differences in audiometry equipment, examiner, and patient factors; for this reason, a change in the hearing threshold of 5 dB is considered within the standard error of these measurements.


A typical audiogram measures thresholds at frequencies within a range slightly broader than that of human speech, including 250, 500, 1000, 2000, 3000, 4000, 6000, and 8000 Hz. The primary speech frequencies fall between 500 and 4000 Hz, although normal human hearing ranges from ~20 Hz to ~20,000 Hz.[6] The arithmetic mean of the air conduction thresholds at 500, 1000, and 2000 Hz is calculated for each patient and termed the "pure-tone average" (PTA), which provides a rapid and simple summary of a patient's hearing in the speech frequencies. The PTA, however, does not provide information about high-frequency hearing loss.

If there is a substantial difference in hearing between ears, masking may be utilized. Masking is a frequently employed technique in which noise is introduced into the non-tested ear to prevent a tone being presented to the tested ear from being perceived in the non-tested ear, which might otherwise result in the patient indicating that they had heard the tone despite it being in the wrong ear. In practice, masking is applied for air conduction when there is a significant hearing threshold difference between ears (~40 dB for headphones and ~65 dB for insert earphones), as signals that are loud enough to overcome the interaural attenuation provided by the skull (the difference in loudness perceived between the ears when a sound is presented on 1 side of the head) be transmitted to the contralateral side and may be heard by the non-tested ear if it hears substantially better than the tested ear.[7] Masking is also frequently applied when testing bone conduction because there is essentially no interaural attenuation with bone conduction, making it important to isolate the non-tested ear from the sound source.

After collecting hearing data from the patient, pure tone thresholds at each frequency are displayed on an audiogram to facilitate interpretation. The X-axis represents the frequency of sound in hertz, and the Y-axis represents the (inverted) sound intensity in decibels Hearing Level. The American Speech-Language-Hearing Association (ASHA) sets guidelines for standardizing audiogram symbols.

ASHA Standard Audiogram Symbols

The following table shows the standard audiogram symbols.



Symbol	Frequency (Hz)	Intensity (dB HL)
◊	250	20
◻	500	20
◯	1000	20
◴	2000	20
◵	4000	20
✕	8000	20

Table

Air conduction in the left ear

When color is available, red indicates sounds detected by the right ear, and blue indicates sounds detected by the left ear; the ◊ and ◻ symbols that indicate left and right ear thresholds were chosen to vaguely resemble a patient's ears facing the interpreter of the audiogram.

Speech Audiometry

In addition to pure tone audiometry, speech audiometry is usually conducted as part of a comprehensive audiological evaluation. Two types of speech audiometry are usually performed: speech awareness or detection threshold (SAT or SDT) testing and speech reception threshold (SRT) testing. They provide different information and are both employed in most cases. The SAT is very similar to pure tone testing, except that words replace the tones, and the patient must indicate when a word is heard. SRT involves presenting 2-syllable words with equal emphasis on both syllables, and the patient must repeat the word correctly. This type of word is known as a "spondee," and examples include "racecar," "baseball," and "toothbrush." The SAT is usually within 10-15 dB of the SRT; if not, a cause for the discrepancy should be sought. Likewise, the PTA and SRT should fall within 6-12 dB of each other. Beyond SAT and SRT, additional word recognition or speech discrimination testing may be performed, in which monosyllabic words are presented at 25-40 dB above the SRT, and the patient is asked to repeat the words. A normal score is 80% or greater; if the score decreases as the loudness of the words increases, the phenomenon is known as "rollover," which is indicative of retrocochlear pathologies, such as a vestibular schwannoma.

Indications

Audiologic evaluation may be indicated when a patient perceives hearing loss. It may also be indicated when other otologic symptoms or history are noted, including tinnitus, dizziness, noise exposure, blast injury, or concern for ototoxic exposure. School-aged children and individuals working in loud occupational environments, such as soldiers and pilots, may routinely undergo hearing screenings.[6] If the screening results reveal hearing loss or a significant threshold shift (change in hearing level), they may be referred for a comprehensive evaluation by an audiologist to evaluate their hearing status further. Lastly, audiometry may be used to establish candidacy for hearing aids or even surgical intervention, such as the placement of cochlear implants.

Potential Diagnosis

Audiogram interpretation aims to determine the type and degree of a patient’s hearing loss.
[10] The table below defines the categories of hearing loss based on severity:



Table

Summary of Hearing Loss Categories

The 3 types of hearing loss are sensorineural (SNHL), conductive, and mixed hearing loss.[11] In sensorineural hearing loss, the hearing loss occurs due to damage to the inner ear and CNS higher auditory pathways and tends to be permanent but may be reversible in some situations. On an audiogram, pure SNHL is indicated by overlapping lines representing air conduction and bone conduction without any air-bone gaps >10 dB HL. SNHL may occur with processes such as inner ear (sensory) or vestibulocochlear nerve (neural) pathology, which is seen with presbycusis, noise exposure, infection, Ménière’s disease, aminoglycoside toxicity, and vestibular schwannoma, among many others.[11] See **Graphs**. Audiogram, Presbycusis; Hearing Loss, Ménière Disease; Audiogram, Vestibular Schwannoma/Acoustic Neuroma.

Conductive hearing loss occurs due to abnormal sound transmission through the outer or middle ear without damage to the inner ear. An air-bone gap is noted on the audiogram, which is present when the difference between air conduction and bone conduction thresholds at a specific frequency is greater than 15 dB HL. With a pure CHL, the bone conduction thresholds should be normal, but the air conduction thresholds indicate hearing loss. Conductive hearing loss may occur due to cerumen impaction, otitis media, tympanic membrane perforation, ossicular disruption, and a foreign body in the ear, among other causes.[11] See **Graph**. Audiogram, Otitis Media With Effusion. Finally, mixed hearing loss combines SNHL and CHL, meaning abnormal sound conduction and damage to the inner ear or retrocochlear structures. This type of hearing loss presents on an audiogram as both a decrease in the bone line and an air-bone gap. Mixed hearing loss results from many different pathologies, including otosclerosis, a disease in which both the ossicles (CHL) and the inner ear (SNHL) are affected due to abnormal bone growth. [11] See **Graph**. Audiogram, Otosclerosis.

Normal and Critical Findings

Normal Hearing

A normal audiogram displays air and bone conduction lines with ≤25 dB HL thresholds at each tested frequency in both ears.[12] When hearing loss is encountered, it should be classified by severity (see table above) and frequency, but some characteristic patterns often associated with specific diagnoses are below. According to the National Institute for Occupational Safety and Health, a decrease in hearing acuity from 1 audiogram to the next is considered to represent a "significant threshold shift" when there is a decrease of 15 dB in 1 frequency, and that loss is confirmed by a repeat audiogram 30 days later. This significant threshold shift may indicate noise-induced hearing loss, particularly at or around 4000 Hz. When there is a ≥20 dB asymmetry in 2 consecutive frequencies or a ≥15 dB asymmetry in any 2 frequencies between 2000 and 8000 Hz, particularly when accompanied by unilateral tinnitus, magnetic resonance imaging (MRI) of the cerebellopontine angle should be obtained to rule out the presence of a vestibular schwannoma.[13]

Presbycusis

Presbycusis is the most common type of hearing loss that develops with age. This type of hearing loss occurs due to age-related structural changes in the inner ear, producing a bilateral, symmetric, sensorineural hearing loss that is typically worse at higher frequencies. This pattern is represented by a gradual "downsloping" configuration in both ears on an audiogram.[14]

Ototoxicity

Ototoxicity may cause sensorineural hearing loss by damaging various inner ear structures, such as the cochlea's inner hair cells. Common causes include platinum-based chemotherapeutic agents, aminoglycosides, anti-malarial agents, antineoplastic drugs, loop diuretics, and high dosages of non-steroidal anti-inflammatory drugs. The audiogram pattern for ototoxicity is often similar to presbycusis, with a bilateral sensorineural hearing loss more prominent in the high frequencies.[15]

Ménière Disease

Ménière disease is characterized by episodes of a triad of vertigo, tinnitus, and fluctuating hearing loss in 1 ear.[16] A sensation of aural fullness may also be associated with these symptoms. Ménière's disease most commonly presents as a unilateral sensorineural hearing loss, which is most severe in low frequencies. The audiogram shows an upsloping shape, which may become flat in the late stages, depending on disease progression.[17][18] Over time, 5 to 50% of patients also develop Ménière's Disease in the second ear.[19]

Vestibular Schwannoma

Vestibular schwannoma, or acoustic neuroma, is a benign tumor of the internal auditory canal or the cerebellopontine angle that arises from the Schwann cells of the vestibular nerve. As the tumor grows, it compromises the cochlear or vestibular nerve function, which can, in turn, affect hearing or balance. The hearing loss associated with these tumors is a unilateral or asymmetric high-frequency sensorineural hearing loss on the affected side. These tumors are typically unilateral, except in neurofibromatosis type 2, in which bilateral tumors are diagnostic.[20]

Noise-Induced Hearing Loss

Noise-induced hearing loss is the permanent loss of hearing due to the destruction of stereocilia (hair cells) in the inner ear from repetitive exposure to loud sounds.[21] This type of hearing loss is often studied in populations with high occupational safety risks concerning hearing, such as construction workers, soldiers, and musicians. The pattern of hearing loss in this population may produce a distinct "U" or a "V" shaped "noise notch" on the audiogram, with the poorest hearing in higher frequency ranges (approximately 3000 to 6000 Hz, often centered at approximately 4000 Hz); however, many different patterns exist. These changes may be unilateral or bilateral, depending on the type of noise exposure. Noise-induced hearing loss should be distinguished from the related, yet distinct, entity of acoustic trauma, which is hearing loss caused by a single loud impulse sound, such as an explosion. These patients often present with SNHL or mixed hearing loss and may have tinnitus and imbalance that progresses over many years. Recruitment, or discomfort with loud sounds within a particular frequency range, is also common after acoustic trauma.

Otitis Media with Effusion

Otitis media with effusion is an inflammation of the mucosa of the middle ear that causes fluid accumulation in a typically air-filled space. It may cause conductive hearing loss by affecting the transmission of sound from the air through the structures of the middle ear. This pathology may result in an air-bone gap of >15 dB HL.[22]

Otosclerosis

Otosclerosis occurs due to abnormal growth of the bone surrounding the cochlea, which can lead to fixation of the stapes to the oval window. Due to changes in the middle ear structure, otosclerosis commonly causes conductive hearing loss but may progress to mixed hearing loss in some cases. A classic feature, known as the "Carhart notch," may also be present, represented by a worsening bone conduction hearing of 10 to 15 dB HL at 2000 Hz, sometimes accompanied by an apparent improvement of the air conduction threshold at that same frequency.[23]

Congenital Hearing Loss

Congenital, inherited hearing loss, as may result from Stickler syndrome, Waardenburg syndrome, connexin 26 gene mutations, and others, may present with a "cookie bite" pattern on pure tone audiometry. The "cookie bite" is a "U" shape to the air and bone lines that denotes mid-frequency SNHL centers around 1000 Hz but usually extends from 500 Hz to 2000 Hz before sloping back towards normal. While this pattern is often associated with genetic hearing loss, neither its specificity (18%) nor its positive predictive value (15%) for this diagnosis is high, and a U-shaped audiogram can also result from vestibular schwannoma, autoimmune inner ear disease, carbon monoxide poisoning, and other insults to the cochlea and eighth nerve.[24][25]

Interfering Factors

Several interfering factors may complicate the interpretation of audiograms. Firstly, it is essential to ensure a good fit of the earphones to prevent sound leakage, which may invalidate results by independently preventing reliable testing of the left and right ears.[26] Secondly, if masking is not employed when indicated or if the masking level is insufficient, inaccurate results may be obtained, again because of inadequate isolation of the ears.[27] Thirdly, audiogram validity depends on a patient's ability to indicate accurately and consistently when they hear sounds; accidental or deliberate false positives and negatives compromise results. In functional or non-organic hearing loss cases, patients may report the inability to hear a pure tone even if their auditory system is structurally intact, making audiogram interpretation challenging.[28]

Patient Safety and Education


Overall, audiograms are a relatively safe and non-invasive testing modality. Patients should be informed that pure-tone evaluation does not represent a comprehensive audiological examination, and other tests may be needed to diagnose specific conditions that produce hearing loss.[29] Beyond pure-tone audiometry, testing performed during a comprehensive audiological assessment includes tympanometry, otoscopy, speech audiometry, otoacoustic emissions, and validity checks, such as the Stenger test. These tests are useful for assessing middle ear function, the structural integrity of the outer and middle ear, the ability to interpret speech, cochlear hair cell function, and exaggeration of hearing deficits, respectively.[30][31] Further audiometric testing that may be performed includes auditory brainstem responses and electrocochleography, but these tests are typically only performed for very specific indications, such as to follow up for a failed newborn hearing screen or to diagnose Ménière disease.

Clinical Significance

The audiogram represents a useful clinical test that can aid healthcare providers in determining the need for further diagnostic studies. By interpreting data from the audiogram and the history and physical examination, therapeutic interventions may be offered according to patient presentation and goals. For instance, an audiogram showing an otherwise asymptomatic unilateral significant threshold shift may suggest the presence of a vestibular schwannoma, and an MRI may be warranted. Conversely, bilateral high-frequency SNHL in an older patient may suggest presbycusis, which would not require imaging but may warrant fitting for hearing aids. In this way, the audiogram allows clinicians to quantify and characterize the degree and type of hearing loss, providing more targeted care for patients to improve their quality of life and occupational functioning. Interprofessional team-based care coordination is necessary for these cases, involving family clinicians, ENT specialists, pediatricians, audiologists, and nursing staff, all utilizing open communication and accurate patient records so that every team member can contribute from their specialization leading to improved patient outcomes.


Review Questions

- [Access free multiple choice questions on this topic.](#)
- [Comment on this article.](#)




Figure

Audiogram, Presbycusis. This audiogram of a patient with presbycusis, or age-related hearing loss, is characterized by the typical bilateral, symmetric, high-frequency hearing loss. Contributed by M Salmon, MD




Figure

Hearing Loss, Ménière Disease. A unilateral, up-sloping sensorineural hearing loss is most commonly seen with Ménière disease. Contributed by M Salmon, MD




Figure

Audiogram, Vestibular Schwannoma/Acoustic Neuroma. An audiogram with unilateral sensorineural hearing loss is typically found with vestibular schwannoma/acoustic neuroma. Contributed by M Salmon, MD



Figure

Audiogram, Otitis Media With Effusion. An audiogram showing a right conductive hearing loss in a patient with otitis media with effusion. Contributed by M Salmon, MD



Figure

Audiogram, Otosclerosis. An audiogram showing bilateral mixed hearing loss in a patient with otosclerosis. The typical

dip in bone conduction is also present. Contributed by M Salmon, MD

References

1. Agrawal Y, Platz EA, Niparko JK. Prevalence of hearing loss and differences by demographic characteristics among US adults: data from the National Health and Nutrition Examination Survey, 1999-2004. *Arch Intern Med*. 2008 Jul 28;168(14):1522-30. [PubMed: 18663164]
2. Jerger J. Why the audiogram is upside-down. *Int J Audiol*. 2013 Mar;52(3):146-50. [PubMed: 23316997]
3. CARHART R. An improved method for classifying audiograms. *Laryngoscope*. 1945 Nov;55:640-62. [PubMed: 21007825]
4. Klyn NAM, Letendre C, Shrestha N, Lambert BL, Dhar S. Interpretability of the audiogram by audiologists and physician non-specialists. *Int J Audiol*. 2021 Feb;60(2):133-139. [PubMed: 32815399]
5. Poling GL, Kunnel TJ, Dhar S. Comparing the Accuracy and Speed of Manual and Tracking Methods of Measuring Hearing Thresholds. *Ear Hear*. 2016 Sep-Oct;37(5):e336-40. [PMC free article: PMC4996736] [PubMed: 27232075]
6. Walker JJ, Cleveland LM, Davis JL, Seales JS. Audiometry screening and interpretation. *Am Fam Physician*. 2013 Jan 01;87(1):41-7. [PubMed: 23317024]
7. LIDEN G, NILSSON G, ANDERSON H. Masking in clinical audiometry. *Acta Otolaryngol*. 1959 Mar-Apr;50(2):125-36. [PubMed: 13636845]
8. Guidelines for manual pure-tone threshold audiometry. ASHA. 1978 Apr;20(4):297-301. [PubMed: 656172]
9. Guidelines for audiometric symbols. ASHA. 1988 Dec;30(12):39-42. [PubMed: 3207447]
10. Clark JG. Uses and abuses of hearing loss classification. ASHA. 1981 Jul;23(7):493-500. [PubMed: 7052898]
11. Cunningham LL, Tucci DL. Hearing Loss in Adults. *N Engl J Med*. 2017 Dec 21;377(25):2465-2473. [PMC free article: PMC6457651] [PubMed: 29262274]
12. Vermiglio AJ, Griffin S, Post C, Fang X. An Evaluation of the World Health Organization and American Medical Association Ratings of Hearing Impairment and Simulated Single-Sided Deafness. *J Am Acad Audiol*. 2018 Jul/Aug;29(7):634-647. [PubMed: 29988011]
13. Waterval J, Kania R, Somers T. EAONO Position Statement on Vestibular Schwannoma: Imaging Assessment. What are the Indications for Performing a Screening MRI Scan for a Potential Vestibular Schwannoma? *J Int Adv Otol*. 2018 Apr;14(1):95-99. [PMC free article: PMC6354499] [PubMed: 29764782]
14. Löhler J, Cebulla M, Shehata-Dieler W, Volkenstein S, Völter C, Walther LE. Hearing Impairment in Old Age. *Dtsch Arztebl Int*. 2019 Apr 26;116(17):301-310. [PMC free article: PMC6584833] [PubMed: 31196393]
15. Ganesan P, Schmiedge J, Manchaiah V, Swapna S, Dhandayutham S, Kothandaraman PP. Ototoxicity: A Challenge in Diagnosis and Treatment. *J Audiol Otol*. 2018 Apr;22(2):59-68. [PMC free article: PMC5894487] [PubMed: 29471610]
16. Koenen L, Andaloro C. StatPearls [Internet]. StatPearls Publishing; Treasure Island (FL): Aug 21, 2023. Meniere Disease. [PubMed: 30725640]
17. Sakurai T, Yamane H, Nakai Y. Some aspects of hearing change in Menière's patients. *Acta Otolaryngol Suppl*. 1991;486:92-8. [PubMed: 1842884]
18. Mateijsen DJ, Van Hengel PW, Van Huffelen WM, Wit HP, Albers FW. Pure-tone and speech audiometry in patients with Menière's disease. *Clin Otolaryngol Allied Sci*. 2001

Oct;26(5):379-87. [PubMed: 11678945]

19. Frejo L, Soto-Varela A, Santos-Perez S, Aran I, Batuecas-Caletrio A, Perez-Guillen V, Perez-Garrigues H, Fraile J, Martin-Sanz E, Tapia MC, Trinidad G, García-Arumi AM, González-Aguado R, Espinosa-Sanchez JM, Marques P, Perez P, Benitez J, Lopez-Escamez JA. Clinical Subgroups in Bilateral Meniere Disease. *Front Neurol*. 2016;7:182. [PMC free article: PMC5075646] [PubMed: 27822199]
20. Kim SH, Lee SH, Choi SK, Lim YJ, Na SY, Yeo SG. Audiologic evaluation of vestibular schwannoma and other cerebellopontine angle tumors. *Acta Otolaryngol*. 2016;136(2):149-53. [PubMed: 26479426]
21. Lie A, Skogstad M, Johannessen HA, Tynes T, Mehlum IS, Nordby KC, Engdahl B, Tambs K. Occupational noise exposure and hearing: a systematic review. *Int Arch Occup Environ Health*. 2016 Apr;89(3):351-72. [PMC free article: PMC4786595] [PubMed: 26249711]
22. Cai T, McPherson B. Hearing loss in children with otitis media with effusion: a systematic review. *Int J Audiol*. 2017 Feb;56(2):65-76. [PubMed: 27841699]
23. Danesh AA, Shahnaz N, Hall JW. The Audiology of Otosclerosis. *Otolaryngol Clin North Am*. 2018 Apr;51(2):327-342. [PubMed: 29397946]
24. Ahmadmehrabi S, Li B, Epstein DJ, Ruckenstein MJ, Brant JA. How Does the "Cookie-Bite" Audiogram Shape Perform in Discriminating Genetic Hearing Loss in Adults? *Otolaryngol Head Neck Surg*. 2022 Mar;166(3):537-539. [PubMed: 34058916]
25. Wong L, Bance M. Are all cookie-bite audiograms hereditary hearing loss? *J Otolaryngol*. 2004 Dec;33(6):390-2. [PubMed: 15971657]
26. Bagatto M, Moodie S, Scollie S, Seewald R, Moodie S, Pumford J, Liu KP. Clinical protocols for hearing instrument fitting in the Desired Sensation Level method. *Trends Amplif*. 2005;9(4):199-226. [PMC free article: PMC4111495] [PubMed: 16424946]
27. Seneviratne S, McNeill C, Greenberg SL, Kong J. Naunton's Masking Dilemma Revisited. *Otol Neurotol*. 2019 Jan;40(1):e1-e6. [PubMed: 30531635]
28. Norrix LW, Rubiano V, Muller T. Estimating Nonorganic Hearing Thresholds Using Binaural Auditory Stimuli. *Am J Audiol*. 2017 Dec 12;26(4):486-495. [PubMed: 28915294]
29. Harris PK, Hutchinson KM, Moravec J. The use of tympanometry and pneumatic otoscopy for predicting middle ear disease. *Am J Audiol*. 2005 Jun;14(1):3-13. [PubMed: 16180966]
30. Ooster J, Krueger M, Bach JH, Wagener KC, Kollmeier B, Meyer BT. Speech Audiometry at Home: Automated Listening Tests via Smart Speakers With Normal-Hearing and Hearing-Impaired Listeners. *Trends Hear*. 2020 Jan-Dec;24:2331216520970011. [PMC free article: PMC7720343] [PubMed: 33272109]
31. Kemp DT. Otoacoustic emissions, their origin in cochlear function, and use. *Br Med Bull*. 2002;63:223-41. [PubMed: 12324396]

Disclosure: Mandy Salmon declares no relevant financial relationships with ineligible companies.

Disclosure: Jason Brant declares no relevant financial relationships with ineligible companies.

Disclosure: Marc Hohman declares no relevant financial relationships with ineligible companies.

Disclosure: Danielle Leibowitz declares no relevant financial relationships with ineligible companies.

Copyright © 2025, StatPearls Publishing LLC.

This book is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits others to distribute the work, provided that the article is not altered or used commercially. You are not required to obtain permission to distribute this article, provided that you credit the author and journal.

Bookshelf ID: NBK578179 PMID: 35201707

<https://www.ncbi.nlm.nih.gov/books/NBK578179/>