

SUPPLEMENTAL APPENDIX

Page

PURSUIT-IV Study Group.....2

Phase 2 Analysis.....6

SUPPLEMENTAL TABLES AND FIGURES

Supplemental Table S1. Summary of Baseline Demographics and Disease 7
Characteristics Among All Randomized Patients
in Phase 2

Supplemental Table S2. Efficacy Findings in Phase 2.....8

Supplemental Figure S1. Patient disposition through Week 6 among patients.....10
randomized (A) in Phase 2, (B) while Phase 2 data
were being analyzed, (C) in Phase 3, and (D) all randomized
patients.
The single patient randomized but never treated was assigned to the golimumab (GLM)
2 mg/kg group. W/D, withdrew; LTF, lost to follow-up.

Supplemental Figure S2. Median Serum Golimumab Concentration by Study.....11
Visit (Week) and by Induction Dose and Regimen

SUPPLEMENTAL APPENDIX

The participating investigators for the PURSUIT-IV Study Group were:

Australia

HS Debinski, Melbourne Gastrointestinal Investigation Unit, Cabrini Hospital, Malvern, Victoria; TH Florin, University of Queensland, Department of Medicine, Mater Heath Services' Adult Hospital, South Brisbane, Queensland; DJ Hetzel, Department of Gastroenterology & Hepatology, Q7 North Wing, Royal Adelaide Hospital, North Terrace, Adelaide

Austria

A Kaser, Universitätsklinik für Innere Medizin, Gastroenerologie & Hepatologie, Innsbruck; W Reinisch, Universitätsklinik für Innere Medizin IV, Vienna.

Belgium

GR d'Haens, Imelda General Hospital, Bonheiden; PJ Rutgeerts, University Hospital Gasthuisberg, University of Leuven (Katholieke Universiteit Leuven- KUL), Department of Medicine, Division of Gastroenterology, Leuven

Bulgaria

IA Kotzev, Clinic of Gastroenterology, University Multifunctional Hospital for Active Treatment "St. Marina", Varna; PM Petrov, Gastroenterology Unit at Department of Gastroenterology, Endocrinology, Nephrology and Hematology University, Multifunctional Hospital for Active Treatment, Sofia; GN Vasileva, Third Therapeutical Department – Gastroenterology and Hepatology, Rousse

Canada

FH Anderson, Liver and Intestinal Research Centre, Vancouver, BC; W Depew, Hotel Dieu Hospital, Division of Gastroenterology, Kingston, ON; B Feagan, London Health Science Centre, University Hospital, London, ON; RA Singh, Victoria, BC; TA Sylwestrowicz, Medical Digestive, Saskatoon, SK

France

Y Bouhnik, Hopital Beaujon, Service de Gastro-entérologie, Clichy; JF Colombel, Departement d'Hepato-gastro-entérologie, Hopital Huriez, Lille Cedex; X Hebuterne, Hopital de l'Archet 2, Service de Gastro-entérologie et de nutrition, Nice Cedex 3

SUPPLEMENTAL APPENDIX

Hungary

L Bene, Peterfy S. u. Hospital, Dept. "E" of Internal Medicine, Gastroenterology, Budapest;
Z Dobronte, Vas County Markusovszky Hospital, Dept. of Internal Medicine,
Gastroenterology, Szombathely; A Kiraly, University of Pecs, Faculty of Medicine, 3rd Dept.
of Internal Medicine, Pécs; L Madacsy, Fejer County Szt. Gyorgy Hospital, 1st Dept. of
Internal Medicine, Székesfehérvár; P Orosz, Borsod-Abaúj Zemplén County University and
Teaching Hospital, 2nd Dept. of Internal Medicine, Gastroenterology, Miskolc; I Racz, Petz
Aladar County and Teaching Hospital, 1st Dept. of Medicine and Gastroenterology, Győr; L
Ujszaszy, Miskolci Semmelweis Ignac Egeszegugyi Kozpont es Egyetemi Oktato Korhaz
Nonprofit Kft., 1st Dept. of Medicine, Gastroenterology, I. Belgyogyaszat,
Gasztroenterologia, Miskolc

India

A Chandra, Department of Surgical Gastroenterology, Chatrapati Shahuji Maharaj Medical
University, Lucknow, Uttar Pradesh; M Kalla, S. R. Kalla Memorial Gastro and General
Hospital, Jaipur; NV Pai, Frant Medical Foundation, Ruby Hall Clinic, Pune, Maharashtra;
DN Reddy, Asian Institute of Gastroenterology, Hyderabad, Andhra Pradesh; SC Samal,
Department of Gastroenterology, Apollo Hospitals Education and Research Foundation,
Jubilee Hills, Hyderabad; LV Shimpi, Jehangir Clinical Development Centre Pvt. Ltd.,
Jehangir Hospital Premises, Pune, Maharashtra; A Sood, Dayanand Medical College &
Hospital, Department of Gastroenterology, Ludhiana, Punjab

Israel

Z Fireman, Gastroenterology Department, Hillel-Yaffe Medical Center, Hadera; S Fishman,
Z Halpern (former PI), Tel Aviv Sourasky Medical Center, Institute of Gastroenterology and
Liver Diseases, Tel Aviv; FM Kornikoff, Meir Hospital, Sapir Medical Center, Department
of Gastroenterology and Hepatology, Kfar Sava; E Nussinson, Ha'Emek Medical Center,
Gastroenterology Unit, Afula; D Rachmilewitz, Shaare Zedek Medical Center, Division of
Gastroenterology, Jerusalem

Latvia

A Danilans, P. Stradina Clinical University Hospital, Riga; G Delmans, Daugavpils Regional
Hospital, Daugavpils

SUPPLEMENTAL APPENDIX

Netherlands

G Dijkstra, Department of Gastroenterology and Hepatology, Universitair Medisch Centrum Groningen, Groningen; DW Hommes, Department of Gastroenterology and Hepatology, Leiden University Medical Center, Leiden; CJ van der Woude, Department of Gastroenterology and Hepatology, Erasmus MC Rotterdam, Rotterdam

New Zealand

MM Arnold, Hawkes Bay Hospital, Hastings; S Parry, Department of Gastroenterology and Hepatology, Middlemore Hospital, Otahuhu, Auckland; ID Wallace, Shakespeare Specialist Group, Milford, Auckland

Poland

J Chojnacki, SPZOZ Uniwersytecki Szpital Kliniczny nr 5 im., Gen. Dyw. B. Szareckiego, Uniwersytetu Medycznego w Lodzi, Klinika Gastroenterologii i Chorob Wewnętrznych, Lodz; Z Hebzda, 5 Wojskowy Szpital Kliniczny z Poliklinika, Samodzielny Publiczny Zakład Opieki Zdrowotnej, Krakow; Z Jamrozik-Kruk, Wojewodzki Szpital Specjalistyczny im., Najświętszej Maryi Panny, Oddział Gastroenterologii i Chorob Wewnętrznych, Częstochowa; D Kleczkowski, Endoskopia Sp. z o.o., Sopot; W Laszewicz, Samodzielny SPZOZ Wojewodzki Szpital, Zespólny im. J. Sniadeckiego w Białymstoku, III Oddział Chorob Wewnętrznych i Gastroenterologii, Białystok; M Szura, Specjalistyczne Centrum Diagnostyczno.-Zabiegowe Medicina Sp. z o.o., Krakow

Romania

LS Gheorghe, Institutul Clinic Fundeni, Centrul de Gastroenterologie si Hepatologie Fundeni, Bucuresti

Serbia

G Jankovic, Clinical Center of Serbia, Clinic for Gastroenterology and Hepatology, Belgrade; N Jojic, Clinical Hospital Center Zvezdara, Clinic for Gastroenterology and Hepatology, Belgrade; D Tarabar, Military Medical Academy, Clinic for Gastroenterology, Belgrade

Slovak Republic

I Bunganic, MUCO, Gastroenterologicka ambulancia, Presov; M Gregus, NZZ-KM Management s.r.o., Gastroenterologicke oddelenie, Nitra

SUPPLEMENTAL APPENDIX

Ukraine

A Dorofeyev, Central City Clinical Hospital #3, Department of Gastroenterology, Donetsk State Medical University named after M.Gorkyy, Chair of Internal Medicine #2, Donetsk; AI Golovchenko, Military Medical Center of the Air Force of Ukraine, National Center of Health, Clinic of Gastroenterology, Vynnytsya; V Neyko, Central City Clinical Hospital, Therapeutic Department, Ivano-Frankovsk State Medical University, Chair of Propedeutics of Internal Diseases, Ivano-Frankovsk; Y Zakharash, Hospital of Military Medical division of Security Service of Ukraine, National Medical University named after O.O.Bogomolets, Chair of Faculty Surgery #1, Kyiv

USA

SD Bologna, Center for Digestive Health, Troy, MI; LW Comerford, Medical University of South Carolina, Digestive Disease Center, Charleston, SC; JM Fields, Center for Clinical Studies, Inc., Dearborn, MI; GL Gordon, Center for Digestive and Liver Diseases, Inc., Mexico, MO; JW Hamilton, Dean Clinic, Madison, WI; RN Hansen, Arapahoe Gastroenterology, PC, Littleton, CO; JS Hanson, Charlotte Gastroenterology and Hepatology, P.L.L.C., Charlotte, NC; IE Ibegbu, Kinston, NC; D James, Tulsa, OK; RK Kottoor, University of Florida, Jacksonville, FL; S Lichtiger, New York, NY; RM Matsuyama, Capitol Gastroenterology Consultants Medical Group, Inc., Roseville, CA; PB Miner, Oklahoma Foundation for Digestive Research, Oklahoma City, OK; WM Pandak, McGuire DVAMC, Richmond, VA; M Safdi, Consultants for Clinical Research Inc., Cincinnati, OH; PL Schroeder, Cotton-O'Neil Clinical Research Center, Digestive Health, Topeka, KS; DA Schwartz, Vanderbilt University Medical Center, Nashville, TN; RE Tepper, Long Island Clinical Research Associates, LLP, Great Neck, NY; R Vasudeva, Consultants in Gastroenterology, Columbia, SC; RA Wohlman, Northwest Gastroenterology Associates, Bellevue, WA; ZH Younes, Gastroenterology Center of the Mid South, PC, Germantown, TN

SUPPLEMENTAL APPENDIX

Phase 2 Analysis

In Phase 2, 176 patients were randomized to a single IV infusion at week 0 with golimumab (132 subjects to golimumab [44 to each 1 mg/kg, 2 mg/kg, or 4 mg/kg group]) or placebo (44 patients; Supplemental Figure 1A).

Demographic characteristics were generally comparable across treatment groups, with the exception of CRP which was lower in the 4mg/kg golimumab group (Supplemental Table S1). At baseline, the golimumab and placebo groups had comparable median Mayo scores (8.0 in each treatment group).

The median changes from baseline in the Mayo score at week 6 were lower than anticipated, but greater for golimumab-treated (-2.0 in the 1 mg/kg and 2 mg/kg groups, -1.5 in the 4 mg/kg group) than for placebo-treated patients (-1.0) (Supplemental Table S2). While the proportions of patients in clinical response were greater for the 1- and 2-mg/kg groups, relative to placebo, in general golimumab did not differentiate from placebo with respect to remission and mucosal healing rates at week 6. (Supplemental Table S2).

Golimumab-treated patients in Phase 2 showed greater change in Mayo scores and numerically greater proportions of patients in clinical response at week 6 compared with placebo patients; however, no dose response was observed. The benefit of a single golimumab IV infusion was lower than expected across all efficacy measures (Supplemental Table S2) and lower than that observed in PURSUIT-SC.¹¹

SUPPLEMENTAL APPENDIX

Table S1. Summary of Baseline Demographics and Disease Characteristics; All Randomized Patients in Phase 2

	-----Golimumab-----					
	Placebo (N=44)	1 mg/kg (N=44)	2 mg/kg (N=44)	4 mg/kg (N=44)	All (N=132)	Total (N=176)
Male patients, n (%)	27 (61.4)	30 (68.2)	23 (52.3)	28 (63.6)	81 (61.4)	108 (61.4)
Race, n (%)						
Caucasian	33 (75.0)	33 (75.0)	35 (79.5)	34 (77.3)	102 (77.3)	135 (76.7)
Black	2 (4.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (1.1)
Asian	7 (15.9)	9 (20.5)	9 (20.5)	10 (22.7)	28 (21.2)	35 (19.9)
Other	2 (4.5)	2 (4.5)	0 (0.0)	0 (0.0)	2 (1.5)	4 (2.3)
Age (years)						
Mean $\pm$ SD	41.0 $\pm$ 12.69	41.7 $\pm$ 17.11	41.1 $\pm$ 13.39	40.2 $\pm$ 14.84	41.0 $\pm$ 15.09	41.0 $\pm$ 14.49
Median (IQR)	41.0 (30.5, 49.0)	40.0 (26.0, 52.5)	42.5 (28.5, 52.0)	38.5 (27.0, 49.0)	40.0 (27.0, 51.5)	41.0 (28.0, 50.5)
UC disease duration(yrs)						
Mean $\pm$ SD	6.9 $\pm$ 7.44	5.8 $\pm$ 4.71	6.9 $\pm$ 6.71	6.4 $\pm$ 5.55	6.4 $\pm$ 5.69	6.5 $\pm$ 6.15
Median (IQR)	4.25 (1.98, 9.25)	4.99 (1.88, 8.28)	4.12 (2.69, 8.77)	4.77 (2.18, 9.08)	4.46 (2.26, 8.44)	64.28 (2.15, 8.78)
Extent of disease n (%)						
Limited to left side of colon	25 (56.8)	22 (50.0)	28 (63.6)	25 (56.8)	75 (56.8)	100 (56.8)
Extensive	19 (43.2)	22 (50.0)	16 (36.4)	19 (43.2)	57 (43.2)	76 (43.2)
Mayo score						
Mean $\pm$ SD	8.1 $\pm$ 1.56	8.3 $\pm$ 1.36	8.4 $\pm$ 1.47	8.0 $\pm$ 1.47	8.2 $\pm$ 1.43	8.2 $\pm$ 1.46
Median (IQR)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)
CRP (mg/L), n	42	42	43	40	125	167
Mean $\pm$ SD	11.4 $\pm$ 18.98	12.8 $\pm$ 19.20	13.9 $\pm$ 21.90	7.4 $\pm$ 11.07	11.5 $\pm$ 18.17	11.5 $\pm$ 18.32
Median (IQR)	5.4 (2.7, 11.2)	5.8 (3.0, 14.0)	5.4 (1.6, 14.2)	3.2 (0.7, 8.2)	5.0 (1.3, 12.1)	5.0 (1.5, 12.1)

CRP, C-reactive protein; IQR, interquartile range; SD, standard deviation; UC, ulcerative colitis; yrs, years

SUPPLEMENTAL APPENDIX

Table S2. Efficacy Findings; All randomized patients in Phase 2 (excluding noncompliant site)

	-----Golimumab-----				
Variable	Placebo	1 mg/kg	2 mg/kg	4 mg/kg	Combined
	(N=41)	(N=43)	(N=44)	(N=44)	(N=131)
Mayo score change from baseline at week 6 ^{a, b}					
Baseline					
Mean ± SD	8.0 ± 1.55	8.3 ± 1.37	8.4 ± 1.47	8.0 ± 1.47	8.2 ± 1.44
Median (IQR)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)	8.0 (7.0, 9.0)
week 6 change					
Mean ± SD	-1.3 ± 2.50	-2.0 ± 2.73	-2.3 ± 3.08	-1.8 ± 2.21	-2.0 ± 2.69
Median (IQR)	-1.0 (-3.0, 0.0)	-2.0 (-4.0, 0.0)	-2.0 (-4.5, 0.0)	-1.5 (-3.0, 0.0)	-2.0 (-4.0, 0.0)
P-value		0.241	0.099	0.479	0.152
Clinical response at week 6, n (%) ^{a, b}					
	10 (24.4)	18 (41.9)	18 (40.9)	13 (29.5)	49 (37.4)
P-value		0.090	0.105	0.593	0.126
Clinical remission at week 6, n (%) ^{a, b}					
	4 (9.8)	4 (9.3)	8 (18.2)	3 (6.8)	15 (11.5)
P-value		0.944	0.265	0.622	0.763
Mucosal healing at week 6, n (%) ^{a, b}					
	10 (24.4)	15 (34.9)	13 (29.5)	13 (29.5)	41 (31.3)
P-value		0.293	0.593	0.593	0.398
Partial Mayo score change from baseline					
Baseline					
Mean ± SD	5.6 ± 1.38	5.8 ± 1.17	5.9 ± 1.25	5.6 ± 1.21	5.8 ± 1.21
Median (IQR)	6.0 (5.0, 6.0)	6.0 (5.0, 6.0)	6.0 (5.0, 7.0)	6.0 (5.0, 6.0)	6.0 (5.0, 6.0)

SUPPLEMENTAL APPENDIX

Table S2. Efficacy Findings; All randomized patients in Phase 2 (excluding noncompliant site)

-----Golimumab-----					
Variable	Placebo	1 mg/kg	2 mg/kg	4 mg/kg	Combined
	(N=41)	(N=43)	(N=44)	(N=44)	(N=131)
week 2 change					
Mean ± SD	-0.7 ± 1.48	-1.9 ± 1.90	-1.9 ± 1.97	-1.3 ± 1.29	-1.7 ± 1.75
Median (IQR)	-1.0 (-1.0, 0.0)	-2.0 (-3.0, 0.0)	-2.0 (-3.0, 0.0)	-1.0 (-2.0, -0.5)	-2.0 (-3.0, 0.0)
week 4 change					
Mean ± SD	-0.9 ± 1.68	-2.1 ± 2.14	-1.9 ± 1.97	-1.4 ± 1.51	-1.8 ± 1.90
Median (IQR)	-1.0 (-2.0, 0.0)	-2.0 (-4.0, 0.0)	-1.5 (-3.0, 0.0)	-1.0 (-2.0, 0.0)	-2.0 (-3.0, 0.0)
week 6 change					
Mean ± SD	-0.9 ± 2.02	-1.4 ± 2.24	-1.7 ± 2.39	-1.2 ± 1.73	-1.4 ± 2.13
Median (IQR)	-1.0 (-2.0, 0.0)	-1.0 (-3.0, 0.0)	-1.0 (-3.0, 0.0)	-1.0 (-2.0, 0.0)	-1.0 (-3.0, 0.0)
CRP (mg/L) change from baseline, n ^{a,b}					
Baseline	40	41	43	40	124
Mean ± SD	11.8 ± 19.40	12.4 ± 19.21	13.9 ± 21.87	7.4 ± 11.07	11.3 ± 18.16
Median (IQR)	6.1 (2.5, 11.7)	5.6 (3.0, 13.8)	5.4 (1.6, 14.2)	3.2 (0.7, 8.2)	5.0 (1.3, 12.0)
week 2 change					
Mean ± SD	-3.9 ± 16.83	-7.1 ± 17.02	-5.7 ± 13.22	-3.6 ± 9.64	-5.5 ± 13.62
Median (IQR)	-1.5 (-2.5, 0.2)	-2.7 (-6.5, -0.6)	-1.3 (-6.2, 0.0)	-0.9 (-4.4, -0.1)	-1.9 (-5.9, -0.3)
P-value		0.038	0.218	0.304	0.069
week 4 change					
Mean ± SD	-4.2 ± 15.10	-4.9 ± 20.01	-3.0 ± 16.23	-3.4 ± 10.61	-3.7 ± 16.02
Median (IQR)	-0.2 (-3.5, 1.1)	-1.5 (-5.9, 0.2)	-0.8 (-5.4, 0.0)	-0.6 (-4.2, 0.3)	-0.9 (-5.3, 0.1)
P-value		0.268	0.575	0.713	0.405
week 6 change					
Mean ± SD	-1.8 ± 25.63	-4.8 ± 17.34	-2.2 ± 15.48	-0.1 ± 13.57	-2.4 ± 15.55

SUPPLEMENTAL APPENDIX


Table S2. Efficacy Findings; All randomized patients in Phase 2 (excluding noncompliant site)

Variable	-----Golimumab-----				
	Placebo	1 mg/kg	2 mg/kg	4 mg/kg	Combined
	(N=41)	(N=43)	(N=44)	(N=44)	(N=131)
Median (IQR)	-0.1 (-4.6, 1.8)	-2.6 (-8.8, 1.6)	-0.5 (-6.2, 0.2)	0.1 (-1.6, 3.4)	-0.5 (-3.7, 1.1)
P-value		0.567	0.733	0.162	0.880
IBDQ change from baseline^{a,b}					
Baseline, n	39	43	42	43	128
Mean ± SD	137.2 ± 39.26	128.9 ± 33.17	123.6 ± 32.31	135.0 ± 32.97	129.2 ± 32.90
Median (IQR)	134.0 (113.0, 170.0)	126.0 (104.0, 158.0)	126.5 (102.0, 148.0)	138.0 (115.0, 154.0)	129.0 (108.5, 154.0)
week 6 change					
Mean ± SD	10.3 ± 26.49	24.7 ± 34.70	27.4 ± 33.22	15.1 ± 31.40	22.4 ± 33.29
Median (IQR)	7.0 (-6.0, 25.0)	18.0 (0.0, 43.0)	22.0 (0.0, 46.0)	8.0 (0.0, 31.0)	16.5 (0.0, 38.5)
P-value		0.057	0.010	0.461	0.034
a Patients who had a prohibited change in medication or had an ostomy or colectomy prior to the designated analysis timepoint were considered not to be in clinical response or clinical remission and not to have mucosal healing; for Mayo and partial Mayo scores, IBDQ score, and CRP, the respective baseline value was carried forward from the time of the event onward b Patients who had all 4 mayo subscores missing at the designated analysis timepoint were considered not to be in clinical response or clinical remission and had their last available Mayo subscores carried forward to impute the Missing Mayo score or partial Mayo scores. Patients who had a missing endoscopy subscore at the designated analysis timepoint were considered not to have mucosal healing. Patients who had a missing partial Mayo score, IBDQ score, and CRP at the designated analysis timepoint had the last available value carried forward.					
CRP, C-reactive protein; IBDQ, Inflammatory Bowel Disease Questionnaire; IQR, interquartile range; SD, standard deviation; UC, ulcerative colitis					

SUPPLEMENTAL APPENDIX

Figure S1. Patient disposition through Week 6 among patients randomized (A) in Phase 2, (B) while Phase 2 data were being analyzed, (C) in Phase 3, and (D) all randomized patients.

SUPPLEMENTAL APPENDIX



^a The single patient randomized but never treated was assigned to the golimumab 2 mg/kg group.


^b Excludes patients because of noncompliance at one study site.

^c One patient randomized to golimumab 4 mg/kg only received 0.4 mg/kg and was analyzed for safety with the golimumab 1 mg/kg groups.

LTF = lost to follow-up; W/D = withdrew consent

SUPPLEMENTAL APPENDIX

Figure S2. Median Serum Golimumab Concentration by Study Visit (Week) and by Induction Dose and Regimen



PURSUIT-IV (dashed lines, open symbols); PURSUIT-SC (solid lines, solid symbols)