

SICKLE CELL DISEASE AND THALASSEMIA

How I manage medical complications of β -thalassemia in adults

Ali T. Taher¹ and Maria Domenica Cappellini²

¹Department of Internal Medicine, American University of Beirut Medical Centre, Beirut, Lebanon; and ²Department of Clinical Sciences and Community, University of Milan, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Ca' Granda Foundation Maggiore Policlinico Hospital, Milan, Italy

The complex pathophysiology in β -thalassemia can translate to multiple morbidities that affect every organ system. Improved survival due to advances in management means that patients are exposed to the harmful effects of ineffective erythropoiesis, anemia, and iron overload for a longer duration, and we started seeing new or more frequent complications in adult compared with younger patients. In this article, we highlight particular aspects of managing adult patients with β -thalassemia, using our own experience in treating such patients. We cover both transfusion-dependent and nontransfusion-dependent forms of the disease and tackle specific morbidities of highest interest. (*Blood*. 2018;132(17):1781-1791)

Introduction

The β -thalassemias, a group of inherited hemoglobin disorders, continue to be a concern for health care systems owing to the high burden of disease and its management.¹⁻³ The severity of ineffective erythropoiesis and subsequent anemia depends on several genetic and environmental factors and the disease phenotype was historically labeled as major, intermediate, or minor accordingly.¹ However, in more recent years, we started categorizing patients according to their transfusion requirement, to optimize practical management considerations, although β -thalassemia should always be considered a spectrum of severities with patients able to move from one to another as with management or natural progression of disease.⁴ Transfusion-dependent β -thalassemia (TDT) patients commonly present to our clinics in early childhood with severe anemia that requires lifelong regular transfusion therapy for survival. Nontransfusion-dependent β -thalassemia (NTDT) patients usually present later in childhood or even in adulthood with mild/moderate anemia that only requires occasional or short-course regular transfusions in certain clinical settings. Recent management guidelines have also taken this direction of classification into TDT and NTDT in their recommendations.^{1,5-7}

Over the last few decades, there has been a considerable advance in understanding the disease process of β -thalassemia, and key milestones in optimizing management with transfusion or iron chelation have been achieved. Such advances in supportive management led to a significant improvement in survival in this once fatal disease.^{8,9} For example, mortality rates in western cohorts have declined from 12.7 to 1.65 deaths per 1000 patient-years between the periods 1980 to 1999 and 1999 to 2013 with the leading cause of death moving from iron overload and bone marrow transplant complications to infections and hepatitis C virus complications.^{10,11} However, such advances could not completely abolish the underlying pathophysiology,

which meant that several morbidities continued to manifest at higher incidence with advancing age and chronic exposure to risk factors. Moreover, increased awareness of the disease process prompted clinicians to apply closer and more regular monitoring, which usually leads to higher detection of preclinical and clinical complications especially in adulthood. With this background, we herein share our experience in managing complications in adults with β -thalassemia, especially in their mid-30s and beyond. We limited our coverage to select complications that we most commonly see in our clinics or those persistently reported at higher incidence with advancing age in the literature. It should be noted, however, that such morbidities can still manifest in younger patients, especially in those with severe forms of the disease. Lastly, as clinical trials in this context are limited, we mostly relied on observational data from our own clinics or our expert opinion stemming from direct patient experience.

General management considerations in TDT

In patients with TDT, the culprit of disease process is secondary iron overload from regular transfusion therapy, which can lead to organ damage and failure especially in the heart, liver, and endocrine glands.¹² With advances in magnetic resonance imaging (MRI) that allowed noninvasive estimation of iron levels in key target organs,^{13,14} we realized that significant iron accumulation in these organs can start from early childhood¹⁵⁻¹⁸ and continues to accumulate over time if not optimally treated, leading to the emergence of clinical morbidities.¹⁹ It is hard to assign a specific age of incidence for the various potential complications, as it relies on the specific transfusion and iron chelation practice and patient response. In suboptimally treated patients, we commonly experience an early onset of endocrine disorders in childhood, adolescence, or early adulthood (growth failure, hypogonadism) with an increasing risk as patients age, in

view of cumulative exposure as well as the underlying increased risk seen in the general aging population (eg, hypothyroidism, hypoparathyroidism, diabetes, osteoporosis). Heart failure and arrhythmias can be detected in early adulthood, again with an increasing risk as patients naturally age, whereas we more commonly tend to see arrhythmias in older adults. Early signs of hepatic enzymes that increase secondary to iron overload can be detected at any age, whereas overt hepatic disease such as fibrosis, cirrhosis, or hepatocellular carcinoma are more time-dependent and we more commonly see them in older adults. Management of arrhythmias and hepatic disease in adult patients is covered separately in "Specific morbidities." In a nutshell, early optimization of transfusion and iron-chelation therapy from childhood toward adulthood is essential to ensuring adequate red blood cell supply while still preventing cumulative iron toxicity and subsequent morbidities, especially in later adulthood. More importantly, close and regular monitoring for clinical complications can ensure early detection, specialist consultation, and intervention.⁶ In Figure 1, we highlight key clinical and psychosocial concerns across age groups for patients with TDT. In Figure 2, we highlight standard monitoring recommendations to ensure early detection of abnormalities, although these could be adapted for specific patient populations with varying requirement. Monitoring recommendations for specific morbidities are further discussed in "Specific morbidities."

Although the 3 available iron chelators deferoxamine, deferiprone, and deferasirox have served patients well and have a large body of evidence for efficacy in chelating iron from target organs,^{6,20} we still see a considerable number of patients with high liver and cardiac iron concentration globally,²¹ especially in older adults who were suboptimally treated in childhood due to poor adherence to subcutaneous deferoxamine and imprecise iron measurement tools. Thus, as much as the onus is to prevent iron accumulation by optimal treatment from childhood, adult patients would require adequate chelation to decrease high iron levels to safety thresholds. All 3 iron chelators have established efficacy in significantly reducing iron levels from the liver and heart as evident from trials applying MRI technology.^{6,22-29} The question of which iron chelator is more effective or suitable as first line is futile, as management of patients should be individualized, and different iron chelators may be suitable for different iron overload profiles. Optimal dosing, side-effect monitoring, and adherence are essential for any used iron chelator. In patients with established heart failure, we rely on continuous 24-hour deferoxamine infusion, which was shown to improve cardiac function in earlier studies of TDT patients.³⁰ Similar observations of cardiac function improvement and reversal of heart failure were noted in trials of deferiprone monotherapy and the combination of deferiprone and deferoxamine.³¹⁻³³ The details of management of cardiac complications can be found in the American Heart Association consensus statement on cardiac disease in thalassemia.³⁴ Deferasirox is currently the only chelator to demonstrate stabilization or improvement in hepatic fibrosis.³⁵ Nonclinical trial data have also demonstrated the ability of deferasirox or combination therapy in stabilization or reversal of endocrine and bone disease.^{36,37}

General management considerations in NTDT

Anemia in NTDT can be progressive with age and is directly and independently associated with increased morbidity.³⁸⁻⁴⁰ Ineffective

erythropoiesis also leads to a state of primary iron overload in the absence of transfusion therapy, driven by hepcidin dysregulation.^{41,42} Iron overload is cumulative over time and as patients advance in age,^{38,43,44} with iron overload indices reaching clinically critical thresholds and associated with several morbidities in observational studies including hepatic fibrosis, thrombosis, pulmonary hypertension, endocrinopathies, osteoporosis, and cerebrovascular disease, with notable absence of cardiac iron loading.^{45,46} Ineffective erythropoiesis can also lead to progressive bone marrow expansion and bone changes or mineral density reduction as well as hypercoagulability due to hemolyzed prothrombotic red cell production, the latter leading to vascular events.^{5,47-50} Thus, without intervention, NTDT patients experience increased morbidity as they advance in age with a notable incidence beyond the age of 35 years,^{38,40} with some complications that are less commonly seen in patients with TDT (Figure 3). It should also be noted that quality of life is directly related to age and multiplicity of morbidity in NTDT; hence, attention and appropriate psychological care should be administered.⁵¹ Management of select morbidities that we commonly encounter in adult patients at our clinics will be featured in "Specific morbidities."

In patients with NTDT, no formal clinical trials evaluated the role of regular transfusion therapy as much as occasional transfusions are used in cases of pregnancy or surgery or during infections.⁷ We commonly restrict our use of regular transfusion therapy to specific clinical settings, such as in cases of growth failure or poor sexual development in childhood and adolescence. For adults, consideration of regular transfusion courses is also suggested for prevention or management of certain morbidities (thrombotic events, pulmonary hypertension, leg ulcers, extramedullary hematopoietic pseudotumors) in high-risk patients, backed up by data on transfusion benefit from observational studies of our own patients.^{7,40} The need for transfusion therapy already indicates a more severe phenotype in NTDT that may warrant the need for continuous transfusions if the risk/benefit ratio is favorable.⁷ Careful attention should be paid to the potential risks of iron overload and alloimmunization (especially in splenectomized, pregnant, or previously never-transfused patients).⁷ We almost stopped relying on splenectomy at all owing to the large body of evidence on an increase of a variety of morbidities as well as infections in splenectomized patients.^{7,40} We mainly restrict it to patients with symptomatic splenomegaly or hypersplenism. Iron chelation became standard of care for all patients over 10 years of age with iron overload.⁷ We commonly check all patients older than 10 years for iron overload with serum ferritin or MRI depending on cost and availability. Patients with iron overload (serum ferritin > 800 µg/L or liver iron concentration > 5 mg Fe/g dry weight)^{43,52} are started on iron chelation therapy with deferasirox.⁵³⁻⁵⁵ We follow dosing and iron assessment schedules from management guidelines.⁷

Specific morbidities

In this section, we will feature clinical cases to illustrate the course and management of specific morbidities in patients with TDT or NTDT. These have been selected based on our experience of encountering them most commonly in adults with β -thalassemia, although they may still manifest earlier especially in severe or poorly managed patients. Other clinical complications that may manifest earlier in the natural course of β -thalassemia and that also may increase in incidence with




Figure 1. Management priorities across age groups in TDT. The listing reflects potential incidence of morbidities in respective age groups, but is not exclusive, and individual patients may have different needs. ^aIncludes diabetes mellitus, hypothyroidism, hypoparathyroidism, osteoporosis, hypogonadism.

advancing age (eg, heart failure, endocrine and bone disease, leg ulcers, extramedullary pseudotumors, gallstones, infections) have been featured elsewhere, and management guidelines for those are widely available.^{6,7,56} Lastly, we should also highlight that with advanced age comes an added risk of complications such as cancer, similar to the general population. One epidemiologic study found that the incidence of cancer (3.96 per 1000 person-years) in thalassemia patients was 52% higher than the general population, especially for hematological and abdominal malignancies.⁵⁷ Iron overload and hepatitis C virus infection increase the risk of hepatocellular carcinoma in thalassemia, whereas oxidative stress in the bone marrow is suggested to increase the risk of hematologic malignancies.^{58,59}

Case 1: arrhythmia

A 42-year-old man with TDT was admitted for evaluation of cardiac arrhythmia. He was receiving 2 to 3 units of packed red blood cells every 21 days with a mean pretransfusional hemoglobin of 10.5 g/dL. His comorbidities included glucose intolerance, hypogonadotropic hypogonadism, and subclinical hypothyroidism. He started iron chelation therapy at 4 years of age with subcutaneous deferoxamine. Five years prior to the current presentation, his cardiac T2* MRI showed a moderate myocardial iron overload (14.4 ms), and deferiprone (75 mg/kg per day) was added to his iron chelation regimen in combination with deferoxamine (30 mg/kg per day). His cardiac T2* showed improvement to 24.5 ms 3 years after, and continues to be over 25 ms according to the latest imaging. He started experiencing

several episodes of arrhythmia over the last 2 years (atrial tachycardia with 2:1 atrioventricular block/atrial flutter), which required frequent hospitalizations including the present one. Routine echocardiography has never shown any abnormalities or impairment of left ventricular function except for a mild/moderate atrial dilatation. Subsequently, the patient was sent for catheter ablation. His last 24-hour electrocardiogram (ECG) showed normal sinus rhythm with atrioventricular block (grade 1). He was discharged on bisoprolol 1.25 mg twice daily and amiodarone 200 mg once daily.

Arrhythmia

With advances in cardiac MRI imaging and the availability of oral iron chelators with improved adherence and efficacy in cardiac iron removal, the incidence of heart failure and its associated fatality in TDT patients continue to decline.^{6,10} However, this delay or prevention in heart failure allowed other cardiac manifestations to become more apparent in older adults with thalassemia, such as arrhythmias (especially atrial fibrillation with a prevalence of ~40% in patients over 40 years), cardiac function changes due to restriction, fibrosis, and arterial changes due to loss of vascular compliance.⁶ Symptomatic arrhythmias in thalassemia patients pose a significant clinical risk and are associated with significant myocardial iron overload. They can also occur in patients with otherwise normal iron levels, this being attributed to fibrosis from past iron deposition that was cleared.⁶ Patients should be assessed by history taking for symptoms on every visit, and ECGs should be done annually for all patients,

Iron intake	Record at every transfusion Yearly assessment of iron intake based on transfusion burden		
Serum ferritin	Q 1-3 months		
LIC ^a	Q 2 years if <3 mg/g Q 1 year if 3-15 mg/g Q 6 months if >15 mg/g or rapidly increasing trend in serum ferritin/LIC		
Cardiac T2* ^a	Q 2 years if ≥30 ms Q 1 year if ≥10 to <30 ms Q 6 months if <10 ms		
LVEF ^b	Q 1 year if ≥56% Q 3-6 months if <56% Q 1-3 months if <56% and symptomatic TRV assessment can also be done as needed and simultaneous ECG measurement should be conducted		
Growth	Weight every visit Standing/Sitting Height Q 6 months Bone Age Q 1 year if delayed puberty/growth	Weight every visit	
Sexual development	Tanner staging Q 1 year		Routine assessment for infertility, secondary hypogonadism, impotence
Liver status	Enzymes ^c : Q 3 months Q 1 month if >5 ULN Virology: Q 1 year		
Liver US	Q 1 year Q 6 months if abnormal. TE assessment may also be done if available.		
Endocrine labs ^d	Q 6 months to 1 year Q 3 to 6 months as needed in patients with abnormality		
BMD	Q 2 years Q 1 year as needed in patients with abnormality		
Other	Psychosocial assessment for patient and family		
Age	<10 years	10-18 years	>18 years

Figure 2. General monitoring recommendations across age groups in TDT. ^aBy MRI R2 for liver or MRI T2* for liver and heart with appropriate calibration. LIC and cardiac T2* may be assessed at earlier age (from 6 years) if feasible especially in patients who are on high iron intake. ^bAs assessed by experienced echocardiographer or cardiac MRI. ^cAlanine aminotransferase, aspartate aminotransferase, total and direct bilirubin. ^dThyroid-stimulating hormone; calcium, phosphate, vitamin D, and parathyroid hormone (as indicated); luteinizing hormone, follicle-stimulating hormone, testosterone, estradiol, gonadotropin-releasing hormone (as indicated in cases of abnormal sexual development); fasting blood sugar, oral glucose tolerance test (as indicated). BMD, bone mineral density; ECG, electrocardiogram; LIC, liver iron concentration; LVEF, left-ventricular ejection fraction; Q, every; TE, transient elastography; TRV, tricuspid-regurgitant jet velocity; ULN, upper limit of normal; US, ultrasound.

biannually in asymptomatic patients with abnormality, and monthly or more frequently in symptomatic patients. Assessment can follow the course of echocardiographic or cardiac T2* measurement and should not be ignored. The emerging role of MRI measurement of cardiac fibrosis can also be considered.

In the presence of known or suspected severe cardiac iron overload, we intensify iron chelation therapy when these symptomatic arrhythmias manifest, even if cardiac MRI assessment of myocardial iron status is not feasible, especially as a matter of urgency if the symptoms include syncope or pre-syncope. Arrhythmias in TDT patients can often be controlled or eliminated by aggressive iron chelation, with IV deferoxamine or high-dose deferiprone monotherapy or a combination.⁶⁰ However, this may require some time, and short-term antiarrhythmic therapy is needed in consultation with a cardiologist in view of the constantly changing practice in rhythm control, and the different approaches needed for different arrhythmias, although with low evidence, amiodarone has been used in the limited setting of inpatient and acute care because of its broad spectrum of action and modest compromise of cardiac function.³⁴ It should be noted, however, that amiodarone therapy is associated with side effects especially relevant for patients with thalassemia, particularly for liver and thyroid function that would require close monitoring. Beta-blockers are generally well tolerated, if titrated

slowly, and can be useful in controlling ectopic rhythms.³⁴ Catheter ablation may be considered in patients with chronic symptomatic arrhythmias, but patients should be referred to a cardiac electrophysiology specialist before ablation, given that fibrosis can be challenging to ablate.

It should be noted that arrhythmias may not always be caused by iron overload in β -thalassemia, particularly in younger adults who present with supraventricular arrhythmia. In such cases, other causes such as alcohol or substance abuse or excess of thyroid hormones should be ruled out.

Case 2: liver disease

A 42-year-old man with TDT presented with right upper quadrant pain that has been persistent for 4 months. Initial imaging with contrast-enhanced computed tomography showed foci in his liver that were initially interpreted as extramedullary hematopoiesis. Liver function tests were within normal range as well as albumin level and prothrombin time. His ferritin level was 3100 μ g/L; his liver iron concentration, also measured by MRI, was 17.3 mg Fe/g dry weight. A computed tomography-guided biopsy was recommended and showed the lesions to be compatible with multifocal hepatocellular carcinoma. Hepatitis B and C testing by polymerase chain reaction were negative. The patient received palliative treatment. He developed hepatorenal syndrome and rapidly succumbed to his illness.




Figure 3. Morbidities and risk factors in patients with NTDT. Risk factors and pathophysiologic mechanisms as Venn diagrams within which notable morbidities associated with them are included. Some morbidities are attributed to >1 risk factor. These associations are mostly based on data from observational studies. ^aAs evident on fluorodeoxyglucose positron emission tomography-computed tomography (PET-CT). EMH, extramedullary hematopoietic pseudotumors; GFR, glomerular filtration rate; IE, ineffective erythropoiesis, PHT, pulmonary hypertension.

Liver disease

Liver disease is becoming a leading cause of mortality in TDT, especially owing to the decline in mortality from heart failure associated with advances in cardiac iron monitoring and effective chelation therapy.⁶¹ Liver disease is also responsible for ~10% of the causes of death in NTDT patients.⁶² The liver is the primary site of storage for excess iron; hence, in the absence of effective iron chelation therapy, liver iron concentration can reach clinically critical levels in both TDT and NTDT patients, although at slower rates in NTDT. Primary iron overload from increased intestinal iron absorption in NTDT leads to preferential portal and hepatocyte iron loading whereas hepatic loading in TDT patients also involved the reticuloendothelial system.^{6,7} Irrespectively, chronic hepatic iron deposition promotes liver fibrogenesis and cirrhosis in both patient populations.^{43,52,63-66} A longer duration of hepatic iron exposure is associated with a higher risk of significant fibrosis and cirrhosis, which is why overt morbidity primarily manifests in older adults.^{67,68} In regularly transfused patients, hepatic disease may also be caused or exacerbated by hepatitis C (and less commonly B) virus infection. Although the incidence of viral hepatitis substantially declined after the introduction of regular donor-screening programs in the 1990s and further with the introduction of RNA-polymerase chain reaction testing, adults who may have acquired it earlier continue to have a high prevalence. Hepatitis C virus infection and iron overload are independent risk factors for hepatic fibrosis and cirrhosis, although their coexistence is synergistic in injury.⁶⁶

Both hepatic iron overload and chronic hepatitis C virus (and less so hepatitis B virus) infection can lead to hepatocellular carcinoma. The carcinogenicity of iron is related to its induction of oxidative damage, which results in genotoxicity, and to immunologic dysregulation, which attenuates cancer immune surveillance. Chronic hepatitis C infection leads to necroinflammation, which can prompt progression to cancer.⁶⁹ Several cases of hepatocellular carcinoma in patients with TDT and NTDT from our clinics or those of colleagues have been described in the literature. Most cases occurred in older adults (>45 years) who had cirrhosis and viral hepatitis infection, although malignancy occurred in some patients in the absence of viral hepatitis or cirrhosis while having considerably high iron overload levels.⁶⁹⁻⁷⁴ Additional factors such as obesity or alcohol consumption may also increase steatosis and oxidative stress, which accelerate liver iron uptake and increase risk of liver fibrosis, cirrhosis, and cancer in β -thalassemia patients.⁷⁵

Our approach for the diagnosis and management of liver disease in β -thalassemia is summarized in Figure 4. The hallmark is close monitoring not only for iron overload levels or hepatitis infection status but also for hepatic injury. Optimal management of iron overload is key to preventing liver toxicity. All 3 available iron chelators have data on efficacy in hepatic iron reduction on MRI, although deferasirox has the largest body of evidence in severely iron-loaded patients (liver iron concentration > 15 mg Fe/g dry weight)²⁵⁻²⁷ and has demonstrated ability to reduce necroinflammation and fibrosis in a significant proportion of patients on 3 years' therapy.³⁵ Management of viral hepatitis should be done in consultation with an experienced hepatologist, especially in pregnant women, immunosuppressed patients, and those with cirrhosis.^{6,7} Earlier studies of antiviral therapy in β -thalassemia with peg-interferon and ribavirin showed sustained virological responses in 25% to 64% of patients.⁷⁶ However, ribavirin is known to be associated with worsening anemia and increased transfusion demands, which is a major concern in β -thalassemia patients. Moreover, most clinicians today opt to go for interferon-free regimens. The remarkable revolution in hepatitis C management with direct-acting antiviral drugs offers a new opportunity for β -thalassemia to receive once-daily treatments with no need to check for genetic polymorphisms, and with treatment schedules typically lasting 8 to 12 weeks (eg, glecaprevir-pibrentasvir, ledipasvir-sofosbuvir, or sofosbuvir-velpatasvir for genotype 1). In observational studies and small trials, sustained virological responses at 12 weeks were reported in up to 90% to 100% irrespective of cirrhosis status.⁷⁷⁻⁸⁰ In a recent review, data compiled from 420 thalassemia patients across different studies, with genotypes 1a, 1b, 2, 3, and 4, showed a success rate of at least 93% with direct-acting antiviral drugs.⁸¹ Thus, hepatitis C virus infection in β -thalassemia today is definitely treatable, and much more conveniently than before.

Once manifested, hepatocellular carcinoma would be managed according to standard of care and stage of the disease using chemotherapy, targeted or immune therapy, radiofrequency ablation, transplant, or palliative care. Effective hepatitis C virus clearance in patients with hepatocellular carcinoma improves survival and recurrence rates, but the role of iron chelation in this setting has not been fully evaluated.⁸²

Case 3: thrombotic disease

A 38-year-old splenectomized man with NTDT presented with severe right calf pain. The pain started 4 days prior to presentation,

Iron overload Monitor • LIC^a Q 2 years if <5 NTDT or <3 TDT, Q 1 year if 5-15 NTDT or 3-15 TDT, Q 6 months if >15 mg/g • Serial SF Q 1-3 months TDT and 3-6 months NTDT Prevent • Optimize chelation to iron intake • Ensure adherence to chelation Treat • Optimize chelation to baseline iron levels • Ensure adherence to chelation	Viral hepatitis Monitor • Serology for HCV and HBV^b Q 1 year in transfused patients, positive results confirmed with RNA-PCR, on-treatment monitoring as recommended Prevent • Vaccinate for HBV if to start transfusion • Vaccinate for HAV Treat • DAA (preferred) or peg-interferon plus ribavirin • Consult hepatologist
Hepatic disease Monitor • LFTs: Q 3 months or Q 1 month if >5 ULN • Liver US: Q1 year or Q6 months if abnormal • Transient elastography (measure of hepatic stiffness indicative of fibrosis): Q 1-2 years if available Prevent • Optimize iron chelation in iron overload • Optimize antivirals in viral hepatitis • Lifestyle modifications Treat • Consult hepatologist • Optimize iron chelation	

Figure 4. Approach to monitoring and management of liver disease in adult β -thalassemia patients. ^aBy MRI R2 or T2*, the latter requiring appropriate calibration and preferred if cardiac iron assessment is required at the same time. MRI monitoring can start at the age of 10 years in TDT (or earlier if feasible and deemed necessary) and NTDT. ^bIf not anti-HBsAg⁺. DAA, direct-acting antiviral drug; HAV, hepatitis A virus; HBV, hepatitis B virus; HCV, hepatitis C virus; LFT, liver function test; LIC, liver iron concentration; RNA-PCR, RNA polymerase chain reaction; SF, serum ferritin.

intensified gradually, and was associated with a sensation of hotness around the area. No history of trauma or fever was reported and no personal or family history of thrombosis was documented. A thrombophilia workup was also negative. The pain was unresponsive to analgesics. On physical examination, the patient had diffuse erythema in the right calf which was hot and tender to palpation. No edema was noted. Laboratory workup revealed a total hemoglobin level of 8.4 g/dL, a nucleated red blood cell count of $460 \times 10^6/L$, and a platelet count of $986 \times 10^9/L$. Duplex ultrasonography revealed deep thrombosis of the posterior tibial vein. The patient was started on a treatment dose of low-molecular weight heparin and was started on a course of regular transfusion therapy. Aspirin was also initiated.

Thrombotic disease

A hypercoagulable state has been identified in β -thalassemia patients.^{83,84} It is primarily attributed to abnormalities in platelets and pathological red blood cells, although several additional factors are believed to be involved, including endothelial dysfunction, coagulation system abnormalities, and presence of microparticles, ultimately leading to clinical thrombosis.⁵⁰ The incidence of clinical thrombosis is fourfold higher in NTDT compared with TDT patients, is mostly venous, and is a leading cause of mortality.^{85,86} The frequency of thrombosis in NTDT is significantly higher in patients older than 35 years (28.2%) compared with patients 18 to 35 years (14.9%), or younger than 18 years (4.1%).⁴⁰ Splenectomy, anemia (hemoglobin level < 9 g/dL), and iron overload (serum ferritin > 800 $\mu g/L$ or liver iron concentration > 5 mg Fe/g dry weight) have all been identified as risk factors.^{40,43,52,85,87} Additionally, high nucleated red blood cell ($\geq 300 \times 10^6/L$) and platelet counts ($\geq 500 \times 10^9/L$) were shown to further substantiate the risk in splenectomized

NTDT patients.⁸⁶ Although the incidence of overt strokes is relatively low (~5%),^{85,86} silent strokes have been described in up to 60% of splenectomized adults (mean age, 32 years) with NTDT, with a clear correlation with advancing age.^{88,89} The clinical significance of these white matter lesions or whether they require any intervention, however, is not yet clear.

We treat patients who develop thrombotic disease as per standard local or international guidelines for nonthalassemic patients. Patients who present with unprovoked, spontaneous thrombosis at unusual sites are also worked up for thrombophilia despite the established risk in thalassemia. For primary or secondary prevention, data on risk stratification or prophylaxis in thalassemia are lacking and the approach is mostly individualized. We commonly consider β -thalassemia patients as high risk in medical and surgical settings, especially patients with the aforementioned thalassemia-related (eg, NTDT, splenectomy, low hemoglobin, high platelet or nucleated red blood cell counts) and nonrelated (eg, older age, pregnancy, malignancy, immobility, previous history of thrombosis) risk factors. If prophylaxis is deemed necessary, we commonly use enoxaparin. Newer oral anticoagulants may also be considered, while also acknowledging lack of data in thalassemia, especially if long-term prophylaxis is needed. Blood transfusions may control the hypercoagulability in NTDT patients by improving ineffective erythropoiesis and decreasing the levels of pathological red blood cells with thrombogenic potential.⁹⁰ In observational studies from our clinics, transfusion therapy has been associated with lower rates of thromboembolic events.⁴⁰ It should be noted that a specific transfusion schedule or duration for such patients is not defined. In our practice, we often rely on less frequent regular transfusions to lower the risk of iron overload. These practices should be individualized and based on clinical patient




Figure 5. Approach to diagnosis, prevention, and management of pulmonary hypertension in adult patients with β -thalassemia. ^aPatients with TRV < 2.5 m/s may also be reassessed but at longer intervals (3-5 years). nRBC, nucleated red blood cell count; PHT, pulmonary hypertension.

response. The association between high platelet counts or platelet activation and thrombosis as well as a lower recurrence rate of thrombotic events in splenectomized NTDT patients who took aspirin after their first event, when compared with those who did not, suggests a potential role for aspirin in prevention.^{85,86} We commonly administer aspirin therapy in all splenectomized patients irrespective of platelet count.

Case 4: pulmonary hypertension

A 46-year-old splenectomized woman with NTDT presented with progressive dyspnea of 2 weeks' duration. She had no prior history of cardiac disease. Her laboratory studies revealed a total hemoglobin level of 10.2 g/dL, a platelet count of $850 \times 10^9/L$, and a serum ferritin level of 1520 $\mu g/L$. On continuous-wave Doppler transthoracic echocardiography, she had a peak tricuspid-valve regurgitant jet velocity (TRV) of 3.5 m/s. She was referred to an interventional cardiologist and underwent a right heart cardiac catheterization that revealed a mean pulmonary arterial pressure of 45 mm Hg. She was started on anticoagulant and regular transfusion therapy. She was asked to present back for reevaluation with echocardiography in 6 months.

Pulmonary hypertension

Although the exact mechanisms implicated in the pathogenesis of pulmonary hypertension in β -thalassemia remain unclear, its association with anemia, hemolysis, and hypercoagulability as well as complex interactions of platelets, the coagulation system,

erythrocytes, and endothelial cells along with inflammatory and vascular mediators are suggested.⁹¹⁻⁹³ Pulmonary hypertension in β -thalassemia is pulmonary arterial hypertension, characterized by the presence of precapillary pulmonary hypertension in the absence of left-sided heart disease, lung disease, or chronic thromboembolism.⁹⁴⁻⁹⁶ However, the possibility of pulmonary hypertension occurring secondary to chronic thromboembolic disease cannot be fully excluded with the hypercoagulable state noted in β -thalassemia.^{97,98} In newer classification, it would belong to group 5 pulmonary hypertension associated with chronic hemolytic anemia with unclear/multifactorial mechanism.⁹⁹

Historic studies relying primarily on echocardiographic criteria (TRV exceeding 2.5-2.8 m/s) to document pulmonary hypertension in β -thalassemia provide an overestimate of its prevalence.^{7,40} In a more recent large study of ~1300 patients from Italy, the prevalence of pulmonary hypertension was considerably lower when more strict echocardiographic criteria and confirmatory right heart catheterization were used (5.7% for a TRV > 3.0 m/s, 3.6% for a TRV > 3.2 m/s, and 2.1% on right heart catheterization). Patients with NTDT had a fivefold increased prevalence compared with TDT (4.8% vs 1.1%).¹⁰⁰ Right heart catheterization is needed to confirm the diagnosis. Even in studies relying on catheterization, a significant correlation was noted between prevalence and age, with an exponential rise noted after the age of 45 years.¹⁰⁰ Splenectomy, a history of thrombosis, a platelet count $\geq 500 \times 10^9/L$, nucleated red blood cell counts $\geq 300 \times 10^6/L$, and iron

overload (serum ferritin > 800 µg/L or liver iron concentration > 5 mg Fe/g dry weight) have all been reported as risk factors for pulmonary hypertension in NTDT patients.^{40,43,52,98,100}

When manifest, pulmonary hypertension is associated with functional limitation and can lead to right-sided heart failure.^{97,100-106} Our approach for the diagnosis, prevention, and management of pulmonary hypertension in β-thalassemia is summarized in Figure 5. Although no clinical trials exist, a role for transfusion therapy in preventing the occurrence of pulmonary hypertension is suggested in observational studies, but the specific schedule and duration of use are not defined and should be individualized.^{40,98,107} Similar effects were also noted with hydroxyurea therapy^{40,98,108-111} and iron chelation therapy in observational studies.^{40,98} For management, sildenafil citrate showed promising results in small studies¹¹²⁻¹¹⁴ and trials by improving cardiopulmonary hemodynamics in patients with a TRV > 2.5 m/s.¹¹⁵ Its use, however, should be limited to patients with confirmed pulmonary hypertension by cardiac catheterization.

Future prospects

Several novel therapies are being developed for patients with β-thalassemia. Gene therapy to replace the defective β-globin gene has already shown promising results in reducing or eliminating the transfusion requirement in TDT patients,¹¹⁶ whereas genome editing to reinstate the capacity of fetal hemoglobin production is under way.¹¹⁷ JAK2 inhibitors have also shown some positive results in ameliorating splenomegaly in TDT, although with a modest effect on transfusion requirement.¹¹⁸ The ligand traps sotatercept and luspatercept have shown promising results in phase 2 studies with significant reduction in transfusion requirement for TDT patients and increase in hemoglobin level in NTDT patients; data from randomized clinical trials are awaited.^{119,120} Hepcidin agonists are also being evaluated for their ability to prevent iron overload and improve anemia in β-thalassemia patients.¹²¹ Reductions in transfusion requirement and subsequent prevention of iron overload in TDT and improvement in anemia with prevention of iron overload in NTDT

will surely translate to reduced morbidity risk. Although we expect that such agents, if successful in clinical trials, can transform the disease picture for younger patients and become the mainstay of therapy, older patients who have been exposed to the damaging effects of anemia and iron overload for decades will continue to require conventional therapy for optimal management of their complications risk.

β-thalassemia is a disease of multiple risk factors and multiple morbidities, which logically implies the need for a multidisciplinary management team. This becomes particularly essential for older patients with comorbidities who require the attention of internists and specialists alongside their primary care. Transition from child into adult care facilities becomes more essential for older patients. That said, the ideal treatment strategy will always be an individualized one.

Acknowledgments

The authors thank Khaled Musallam (International Network of Hematology, London, United Kingdom) for constructive review of the manuscript.

Authorship

Contribution: A.T.T. and M.D.C. wrote the manuscript and gave final approval of the manuscript for submission.

Conflict-of-interest disclosure: A.T.T. received honoraria and research grants from Novartis and Celgene. M.D.C. received honoraria from Novartis and Celgene.

Correspondence: Ali T. Taher, Department of Internal Medicine, American University of Beirut Medical Center, P.O. Box 11-0236, Beirut 11072020 Lebanon; e-mail: ataher@aub.edu.lb.

Footnote

Submitted 1 June 2018; accepted 9 September 2018. Prepublished online as *Blood* First Edition paper, 11 September 2018; DOI 10.1182/blood-2018-06-818187.

REFERENCES

- Taher AT, Weatherall DJ, Cappellini MD. Thalassemia. *Lancet*. 2018;391(10116):155-167.
- Weatherall DJ. The definition and epidemiology of non-transfusion-dependent thalassemia. *Blood Rev*. 2012;26(suppl 1):S3-S6.
- Weatherall DJ. The inherited diseases of hemoglobin are an emerging global health burden. *Blood*. 2010;115(22):4331-4336.
- Vitrano A, Calvaruso G, Lai E, et al. The era of comparable life expectancy between thalassemia major and intermedia: is it time to revisit the major-intermedia dichotomy? *Br J Haematol*. 2017;176(1):124-130.
- Musallam KM, Rivella S, Vichinsky E, Rachmilewitz EA. Non-transfusion-dependent thalassemias. *Haematologica*. 2013;98(6):833-844.
- Cappellini MD, Cohen A, Porter J, Taher A, Viprakasit V. Guidelines for the Management of Transfusion Dependent Thalassemia (TDT). 3rd ed. Nicosia, Cyprus: Thalassemia International Federation; 2014.
- Taher A, Musallam K, Cappellini MD. Guidelines for the Management of Non Transfusion Dependent Thalassemia (NTDT). Vol. 2. Nicosia, Cyprus: Thalassemia International Federation; 2017.
- Borgna-Pignatti C, Rugolotto S, De Stefano P, et al. Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica*. 2004;89(10):1187-1193.
- Modell B, Khan M, Darlison M. Survival in beta-thalassemia major in the UK: data from the UK Thalassemia Register. *Lancet*. 2000;355(9220):2051-2052.
- Modell B, Khan M, Darlison M, Westwood MA, Ingram D, Pennell DJ. Improved survival of thalassemia major in the UK and relation to T2* cardiovascular magnetic resonance. *J Cardiovasc Magn Reson*. 2008;10:42.
- Thomas AS, Garbowski M, Ang LA, et al. A decade follow-up of a thalassemia major (TM) cohort monitored by cardiac magnetic resonance imaging (CMR): significant reduction in patients with cardiac iron and in total mortality [abstract]. *Blood*. 2010;116(21). Abstract 1011.
- Taher AT, Saliba AN. Iron overload in thalassemia: different organs at different rates. *Hematology Am Soc Hematol Educ Program*. 2017;2017:265-271.
- Wood JC. Estimating tissue iron burden: current status and future prospects. *Br J Haematol*. 2015;170(1):15-28.
- Wood JC. Guidelines for quantifying iron overload. *Hematology Am Soc Hematol Educ Program*. 2014;2014:210-215.
- Berdoukas V, Nord A, Carson S, et al. Tissue iron evaluation in chronically transfused children shows significant levels of iron loading at a very young age. *Am J Hematol*. 2013;88(11):E283-E285.
- Borgna-Pignatti C, Meloni A, Guerrini G, et al. Myocardial iron overload in

- thalassaemia major. How early to check? *Br J Haematol.* 2014;164(4):579-585.
17. Wood JC, Origa R, Agus A, Matta G, Coates TD, Galanello R. Onset of cardiac iron loading in pediatric patients with thalassemia major. *Haematologica.* 2008;93(6):917-920.
 18. Yang G, Liu R, Peng P, et al. How early can myocardial iron overload occur in beta thalassemia major? *PLoS One.* 2014;9(1): e85379.
 19. Bonifazi F, Conte R, Baiardi P, et al; HTA-THAL Multiregional Registry. Pattern of complications and burden of disease in patients affected by beta thalassemia major. *Curr Med Res Opin.* 2017;33(8): 1525-1533.
 20. Taher AT, Origa R, Perrotta S, et al. New film-coated tablet formulation of deferasirox is well tolerated in patients with thalassemia or lower-risk MDS: results of the randomized, phase II ECLIPSE study. *Am J Hematol.* 2017; 92(5):420-428.
 21. Aydinok Y, Porter JB, Piga A, et al. Prevalence and distribution of iron overload in patients with transfusion-dependent anemias differs across geographic regions: results from the CORDELIA study. *Eur J Haematol.* 2015;95(3):244-253.
 22. Pennell DJ, Porter JB, Piga A, et al; CORDELIA Study Investigators. A 1-year randomized controlled trial of deferasirox vs deferoxamine for myocardial iron removal in β -thalassemia major (CORDELIA). *Blood.* 2014;123(10):1447-1454.
 23. Pennell DJ, Berdoukas V, Karagiorga M, et al. Randomized controlled trial of deferiprone or deferoxamine in beta-thalassemia major patients with asymptomatic myocardial siderosis. *Blood.* 2006;107(9):3738-3744.
 24. Tanner MA, Galanello R, Dessi C, et al. A randomized, placebo-controlled, double-blind trial of the effect of combined therapy with deferoxamine and deferiprone on myocardial iron in thalassemia major using cardiovascular magnetic resonance. *Circulation.* 2007;115(14):1876-1884.
 25. Pennell DJ, Porter JB, Cappellini MD, et al. Deferasirox for up to 3 years leads to continued improvement of myocardial T2* in patients with β -thalassemia major. *Haematologica.* 2012;97(6):842-848.
 26. Cappellini MD, Bejaoui M, Agaoglu L, et al. Iron chelation with deferasirox in adult and pediatric patients with thalassemia major: efficacy and safety during 5 years' follow-up. *Blood.* 2011;110(4):884-893.
 27. Pathare A, Taher A, Daar S. Deferasirox (Exjade) significantly improves cardiac T2* in heavily iron-overloaded patients with beta-thalassemia major. *Ann Hematol.* 2010;89(4): 405-409.
 28. Taher A, Cappellini MD, Vichinsky E, et al. Efficacy and safety of deferasirox doses of >30 mg/kg per d in patients with transfusion-dependent anaemia and iron overload. *Br J Haematol.* 2009;147(5):752-759.
 29. Casale M, Citarella S, Filosa A, et al. Endocrine function and bone disease during long-term chelation therapy with deferasirox in patients with β -thalassemia major. *Am J Hematol.* 2014;89(12):1102-1106.
 30. Davis BA, Porter JB. Long-term outcome of continuous 24-hour deferoxamine infusion via indwelling intravenous catheters in high-risk beta-thalassemia. *Blood.* 2000;95(4): 1229-1236.
 31. Tanner MA, Galanello R, Dessi C, et al. Combined chelation therapy in thalassemia major for the treatment of severe myocardial siderosis with left ventricular dysfunction. *J Cardiovasc Magn Reson.* 2008;10:12.
 32. Filosa A, Vitranò A, Rigano P, et al. Long-term treatment with deferiprone enhances left ventricular ejection function when compared to deferoxamine in patients with thalassemia major. *Blood Cells Mol Dis.* 2013;51(2): 85-88.
 33. Porter JB, Wood J, Olivieri N, et al. Treatment of heart failure in adults with thalassemia major: response in patients randomised to deferoxamine with or without deferiprone. *J Cardiovasc Magn Reson.* 2013;15:38.
 34. Pennell DJ, Udelson JE, Arai AE, et al; American Heart Association Committee on Heart Failure and Transplantation of the Council on Clinical Cardiology and Council on Cardiovascular Radiology and Imaging. Cardiovascular function and treatment in β -thalassemia major: a consensus statement from the American Heart Association [published correction appears in *Circulation.* 2013;128(13):e203]. *Circulation.* 2013; 128(3):281-308.
 35. Deugnier Y, Turlin B, Ropert M, et al. Improvement in liver pathology of patients with beta-thalassemia treated with deferasirox for at least 3 years. *Gastroenterology.* 2011;141(4):1202-1211.
 36. Farmaki K, Tzoumari I, Pappa C, Chouliaras G, Berdoukas V. Normalisation of total body iron load with very intensive combined chelation reverses cardiac and endocrine complications of thalassaemia major. *Br J Haematol.* 2010;148(3):466-475.
 37. Farmaki K, Tzoumari I, Pappa C. Oral chelators in transfusion-dependent thalassemia major patients may prevent or reverse iron overload complications. *Blood Cells Mol Dis.* 2011;47(1):33-40.
 38. Taher AT, Musallam KM, El-Beshlawy A, et al. Age-related complications in treatment-naïve patients with thalassaemia intermedia. *Br J Haematol.* 2010;150(4):486-489.
 39. Taher AT, Musallam KM, Saliba AN, Graziadei G, Cappellini MD. Hemoglobin level and morbidity in non-transfusion-dependent thalassemia [published correction appears in *Blood Cells Mol Dis.* 2015;55(4):419]. *Blood Cells Mol Dis.* 2015;55(2): 108-109.
 40. Taher AT, Musallam KM, Karimi M, et al. Overview on practices in thalassemia intermedia management aiming for lowering complication rates across a region of endemicity: the OPTIMAL CARE study. *Blood.* 2010;115(10):1886-1892.
 41. Camaschella C, Nai A. Ineffective erythropoiesis and regulation of iron status in iron loading anaemias. *Br J Haematol.* 2016; 172(4):512-523.
 42. Rivella S. β -thalassemias: paradigmatic diseases for scientific discoveries and development of innovative therapies. *Haematologica.* 2015;100(4):418-430.
 43. Musallam KM, Cappellini MD, Daar S, et al. Serum ferritin level and morbidity risk in transfusion-independent patients with β -thalassemia intermedia: the ORIENT study. *Haematologica.* 2014;99(11): e218-e221.
 44. Musallam KM, Cappellini MD, Wood JC, et al. Elevated liver iron concentration is a marker of increased morbidity in patients with β thalassemia intermedia. *Haematologica.* 2011;96(11):1605-1612.
 45. Musallam KM, Cappellini MD, Taher AT. Iron overload in β -thalassemia intermedia: an emerging concern. *Curr Opin Hematol.* 2013;20(3):187-192.
 46. Musallam KM, Cappellini MD, Wood JC, Taher AT. Iron overload in non-transfusion-dependent thalassemia: a clinical perspective. *Blood Rev.* 2012;26(suppl 1):S16-S19.
 47. Musallam KM, Taher AT, Duca L, Cesaretti C, Halawi R, Cappellini MD. Levels of growth differentiation factor-15 are high and correlate with clinical severity in transfusion-independent patients with β thalassemia intermedia. *Blood Cells Mol Dis.* 2011;47(4): 232-234.
 48. Musallam KM, Taher AT. Thrombosis in thalassemia: why are we so concerned? *Hemoglobin.* 2011;35(5-6):503-510.
 49. Musallam KM, Taher AT, Rachmilewitz EA. β -thalassemia intermedia: a clinical perspective. *Cold Spring Harb Perspect Med.* 2012;2(7):a013482.
 50. Taher AT, Cappellini MD, Bou-Fakhredin R, Coriu D, Musallam KM. Hypercoagulability and vascular disease. *Hematol Oncol Clin North Am.* 2018;32(2):237-245.
 51. Musallam KM, Khoury B, Abi-Habib R, et al. Health-related quality of life in adults with transfusion-independent thalassaemia intermedia compared to regularly transfused thalassaemia major: new insights. *Eur J Haematol.* 2011;87(1):73-79.
 52. Musallam KM, Cappellini MD, Taher AT. Evaluation of the 5mg/g liver iron concentration threshold and its association with morbidity in patients with β -thalassemia intermedia. *Blood Cells Mol Dis.* 2013;51(1): 35-38.
 53. Taher AT, Cappellini MD, Aydinok Y, et al. Optimising iron chelation therapy with deferasirox for non-transfusion-dependent thalassaemia patients: 1-year results from the THETIS study. *Blood Cells Mol Dis.* 2016;57: 23-29.
 54. Taher AT, Porter JB, Viprakasit V, et al. Deferasirox effectively reduces iron overload in non-transfusion-dependent thalassemia (NTDT) patients: 1-year extension results from the THALASSA study. *Ann Hematol.* 2013;92(11):1485-1493.
 55. Taher AT, Porter J, Viprakasit V, et al. Deferasirox reduces iron overload significantly in nontransfusion-dependent

- thalassemia: 1-year results from a prospective, randomized, double-blind, placebo-controlled study. *Blood*. 2012;120(5):970-977.
56. Musallam KM, Angastiniotis M, Eleftheriou A, Porter JB. Cross-talk between available guidelines for the management of patients with beta-thalassemia major. *Acta Haematol*. 2013;130(2):64-73.
 57. Chung WS, Lin CL, Lin CL, Kao CH. Thalassemia and risk of cancer: a population-based cohort study. *J Epidemiol Community Health*. 2015;69(11):1066-1070.
 58. Halawi R, Cappellini MD, Taher A. A higher prevalence of hematologic malignancies in patients with thalassemia: background and culprits. *Am J Hematol*. 2017;92(5):414-416.
 59. Halawi R, Beydoun H, Cappellini MD, Ferla V, Taher A. Hematologic malignancies in thalassemia: adding new cases to the repertoire. *Am J Hematol*. 2017;92(5):E68-E70.
 60. Anderson LJ, Westwood MA, Holden S, et al. Myocardial iron clearance during reversal of siderotic cardiomyopathy with intravenous desferrioxamine: a prospective study using T2* cardiovascular magnetic resonance. *Br J Haematol*. 2004;127(3):348-355.
 61. Voskaridou E, Ladis V, Kattamis A, et al; Greek Haemoglobinopathies Study Group. A national registry of haemoglobinopathies in Greece: deduced demographics, trends in mortality and affected births. *Ann Hematol*. 2012;91(9):1451-1458.
 62. Matta BN, Musallam KM, Maakaron JE, Koussa S, Taher AT. A killer revealed: 10-year experience with beta-thalassemia intermedia. *Hematology*. 2014;19(4):196-198.
 63. Fargion S, Valenti L, Fracanzani AL. Beyond hereditary hemochromatosis: new insights into the relationship between iron overload and chronic liver diseases. *Dig Liver Dis*. 2011;43(2):89-95.
 64. Musallam KM, Motta I, Salvatori M, et al. Longitudinal changes in serum ferritin levels correlate with measures of hepatic stiffness in transfusion-independent patients with β -thalassemia intermedia. *Blood Cells Mol Dis*. 2012;49(3-4):136-139.
 65. Olivieri NF, Brittenham GM. Iron-chelating therapy and the treatment of thalassemia. *Blood*. 1997;89(3):739-761.
 66. Angelucci E, Muretto P, Nicolucci A, et al. Effects of iron overload and hepatitis C virus positivity in determining progression of liver fibrosis in thalassemia following bone marrow transplantation. *Blood*. 2002;100(1):17-21.
 67. Olynyk JK, St Pierre TG, Britton RS, Brunt EM, Bacon BR. Duration of hepatic iron exposure increases the risk of significant fibrosis in hereditary hemochromatosis: a new role for magnetic resonance imaging. *Am J Gastroenterol*. 2005;100(4):837-841.
 68. Olivieri NF. Progression of iron overload in sickle cell disease. *Semin Hematol*. 2001;38(suppl 1):57-62.
 69. Moukhaider HM, Halawi R, Cappellini MD, Taher AT. Hepatocellular carcinoma as an emerging morbidity in the thalassemia syndromes: a comprehensive review. *Cancer*. 2017;123(5):751-758.
 70. Borgna-Pignatti C, Vergine G, Lombardo T, et al. Hepatocellular carcinoma in the thalassemia syndromes. *Br J Haematol*. 2004;124(1):114-117.
 71. Mancuso A, Sciarrino E, Renda MC, Maggio A. A prospective study of hepatocellular carcinoma incidence in thalassemia. *Hemoglobin*. 2006;30(1):119-124.
 72. Restivo Pantalone G, Renda D, Valenza F, et al. Hepatocellular carcinoma in patients with thalassemia syndromes: clinical characteristics and outcome in a long term single centre experience. *Br J Haematol*. 2010;150(2):245-247.
 73. Maakaron JE, Cappellini MD, Graziadei G, Ayache JB, Taher AT. Hepatocellular carcinoma in hepatitis-negative patients with thalassemia intermedia: a closer look at the role of siderosis. *Ann Hepatol*. 2013;12(1):142-146.
 74. Maakaron JE, Musallam KM, Ayache JB, Jabbour M, Tawil AN, Taher AT. A liver mass in an iron-overloaded thalassemia intermedia patient. *Br J Haematol*. 2013;161(1):1.
 75. Kohgo Y, Ikuta K, Ohtake T, Torimoto Y, Kato J. Iron overload and cofactors with special reference to alcohol, hepatitis C virus infection and steatosis/insulin resistance. *World J Gastroenterol*. 2007;13(35):4699-4706.
 76. Di Marco V, Capra M, Angelucci E, et al; Italian Association for the Study of the Liver. Management of chronic viral hepatitis in patients with thalassemia: recommendations from an international panel. *Blood*. 2010;116(16):2875-2883.
 77. Mehta R, Kabrawala M, Nandwani S, et al. Safety and efficacy of sofosbuvir and daclatasvir for hepatitis C virus infection in patients with β -thalassemia major. *J Clin Exp Hepatol*. 2018;8(1):3-6.
 78. Origa R, Ponti ML, Filosa A, et al; Italy for THALassemia and hepatitis C Advance - Società Italiana Talassemie ed Emoglobinopatie (ITHACA-SITE). Treatment of hepatitis C virus infection with direct-acting antiviral drugs is safe and effective in patients with hemoglobinopathies. *Am J Hematol*. 2017;92(12):1349-1355.
 79. Sinakos E, Kountouras D, Koskinas J, et al. Treatment of chronic hepatitis C with direct-acting antivirals in patients with β -thalassaemia major and advanced liver disease. *Br J Haematol*. 2017;178(1):130-136.
 80. Mangia A, Sarli R, Gamberini R, et al. Randomised clinical trial: sofosbuvir and ledipasvir in patients with transfusion-dependent thalassaemia and HCV genotype 1 or 4 infection. *Aliment Pharmacol Ther*. 2017;46(4):424-431.
 81. Rumi MG, Di Marco V, Colombo M. Management of HCV-related liver disease in hemophilia and thalassemia. *Semin Liver Dis*. 2018;38(2):112-120.
 82. Wirth TC, Manns MP. The impact of the revolution in hepatitis C treatment on hepatocellular carcinoma. *Ann Oncol*. 2016;27(8):1467-1474.
 83. Eldor A, Rachmilewitz EA. The hypercoagulable state in thalassemia. *Blood*. 2002;99(1):36-43.
 84. Eldor A, Durst R, Hy-Am E, et al. A chronic hypercoagulable state in patients with beta-thalassaemia major is already present in childhood. *Br J Haematol*. 1999;107(4):739-746.
 85. Taher A, Isma'eel H, Mehio G, et al. Prevalence of thromboembolic events among 8,860 patients with thalassaemia major and intermedia in the Mediterranean area and Iran. *Thromb Haemost*. 2006;96(4):488-491.
 86. Taher AT, Musallam KM, Karimi M, et al. Splenectomy and thrombosis: the case of thalassemia intermedia. *J Thromb Haemost*. 2010;8(10):2152-2158.
 87. Cappellini MD, Robbiolo L, Bottasso BM, Coppola R, Fiorelli G, Mannucci AP. Venous thromboembolism and hypercoagulability in splenectomized patients with thalassaemia intermedia. *Br J Haematol*. 2000;111(2):467-473.
 88. Musallam KM, Taher AT, Karimi M, Rachmilewitz EA. Cerebral infarction in β -thalassaemia intermedia: breaking the silence. *Thromb Res*. 2012;130(5):695-702.
 89. Taher AT, Musallam KM, Nasreddine W, Hourani R, Inati A, Beydoun A. Asymptomatic brain magnetic resonance imaging abnormalities in splenectomized adults with thalassemia intermedia. *J Thromb Haemost*. 2010;8(1):54-59.
 90. Chen S, Eldor A, Barshstein G, et al. Enhanced aggregability of red blood cells of beta-thalassemia major patients. *Am J Physiol*. 1996;270(6 Pt 2):H1951-H1956.
 91. Farmakis D, Aessopos A. Pulmonary hypertension associated with hemoglobinopathies: prevalent but overlooked. *Circulation*. 2011;123(11):1227-1232.
 92. Morris CR, Vichinsky EP. Pulmonary hypertension in thalassemia. *Ann N Y Acad Sci*. 2010;1202:205-213.
 93. Machado RF, Gladwin MT. Pulmonary hypertension in hemolytic disorders: pulmonary vascular disease: the global perspective. *Chest*. 2010;137(suppl 6):30S-38S.
 94. Galiè N, Hoeper MM, Humbert M, et al; International Society of Heart and Lung Transplantation (ISHLT). Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Respir J*. 2009;34(6):1219-1263.
 95. Simonneau G, Robbins IM, Beghetti M, et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol*. 2009;54(suppl 1):S43-S54.
 96. Galiè N, Hoeper MM, Humbert M, et al; ESC Committee for Practice Guidelines (CPG). Guidelines for the diagnosis and treatment of pulmonary hypertension: the Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS), endorsed by the International Society of Heart and Lung

- Transplantation (ISHLT). *Eur Heart J*. 2009; 30(20):2493-2537.
97. Jootar P, Fucharoen S. Cardiac involvement in beta-thalassemia/hemoglobin E disease: clinical and hemodynamic findings. *Southeast Asian J Trop Med Public Health*. 1990;21(2):269-273.
 98. Karimi M, Musallam KM, Cappellini MD, et al. Risk factors for pulmonary hypertension in patients with β thalassemia intermedia. *Eur J Intern Med*. 2011;22(6):607-610.
 99. Simonneau G, Gatzoulis MA, Adatia I, et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol*. 2013; 62(suppl 25):D34-D41.
 100. Derchi G, Galanello R, Bina P, et al; Webthal Pulmonary Arterial Hypertension Group. Prevalence and risk factors for pulmonary arterial hypertension in a large group of β -thalassemia patients using right heart catheterization: a Webthal study. *Circulation*. 2014;129(3):338-345.
 101. Grisaru D, Rachmilewitz EA, Mosseri M, et al. Cardiopulmonary assessment in beta-thalassemia major. *Chest*. 1990;98(5): 1138-1142.
 102. Aessopos A, Farmakis D, Karagiorga M, et al. Cardiac involvement in thalassemia intermedia: a multicenter study. *Blood*. 2001; 97(11):3411-3416.
 103. Aessopos A, Stamatielos G, Skoumas V, Vassilopoulos G, Mantzourani M, Loukopoulou D. Pulmonary hypertension and right heart failure in patients with beta-thalassemia intermedia. *Chest*. 1995;107(1): 50-53.
 104. Hamdy AM, Zein El-Abdin MY, Abdel-Hafez MA. Right ventricular function in patients with beta thalassemia: relation to serum ferritin level. *Echocardiography*. 2007;24(8): 795-801.
 105. Aessopos A, Kati M, Farmakis D. Heart disease in thalassemia intermedia: a review of the underlying pathophysiology. *Haematologica*. 2007;92(5):658-665.
 106. Kremastinos DT, Farmakis D, Aessopos A, et al. Beta-thalassemia cardiomyopathy: history, present considerations, and future perspectives. *Circ Heart Fail*. 2010;3(3): 451-458.
 107. Atichartakam V, Chuncharunee S, Chandanammattha P, Likittanasombat K, Anyurachai K. Correction of hypercoagulability and amelioration of pulmonary arterial hypertension by chronic blood transfusion in an asplenic hemoglobin E/beta-thalassemia patient. *Blood*. 2004;103(7):2844-2846.
 108. Amoozgar H, Farhani N, Khodadadi N, Karimi M, Cheriki S. Comparative study of pulmonary circulation and myocardial function in patients with β -thalassemia intermedia with and without hydroxyurea, a case-control study. *Eur J Haematol*. 2011; 87(1):61-67.
 109. Karimi M, Borzouee M, Mehrabani A, Cohan N. Echocardiographic finding in beta-thalassemia intermedia and major: absence of pulmonary hypertension following hydroxyurea treatment in beta-thalassemia intermedia. *Eur J Haematol*. 2009;82(3): 213-218.
 110. El-Beshlawy A, Youssry I, El-Saidi S, et al. Pulmonary hypertension in beta-thalassemia major and the role of L-carnitine therapy. *Pediatr Hematol Oncol*. 2008;25(8):734-743.
 111. Musallam KM, Taher AT, Cappellini MD, Sankaran VG. Clinical experience with fetal hemoglobin induction therapy in patients with β -thalassemia. *Blood*. 2013;121(12): 2199-2212.
 112. Littera R, La Nasa G, Derchi G, Cappellini MD, Chang CY, Contu L. Long-term treatment with sildenafil in a thalassemic patient with pulmonary hypertension. *Blood*. 2002; 100(4):1516-1517.
 113. Derchi G, Forni GL, Formisano F, et al. Efficacy and safety of sildenafil in the treatment of severe pulmonary hypertension in patients with hemoglobinopathies. *Haematologica*. 2005;90(4):452-458.
 114. Correale M, De Rosa F, Ieva R, Di Biase M, Brunetti ND. Long-term treatment with high-dose of sildenafil in a thalassemic patient with pulmonary hypertension. *Monaldi Arch Chest Dis*. 2012;78(2):105-106.
 115. Morris CR, Kim HY, Wood J, et al; Thalassemia Clinical Research Network. Sildenafil therapy in thalassemia patients with Doppler-defined risk of pulmonary hypertension. *Haematologica*. 2013;98(9): 1359-1367.
 116. Thompson AA, Walters MC, Kwiatkowski J, et al. Gene therapy in patients with transfusion-dependent β -thalassemia. *N Engl J Med*. 2018;378(16):1479-1493.
 117. Antoniani C, Meneghini V, Lattanzi A, et al. Induction of fetal hemoglobin synthesis by CRISPR/Cas9-mediated editing of the human β -globin locus. *Blood*. 2018;131(17): 1960-1973.
 118. Taher AT, Karakas Z, Cassinerio E, et al. Efficacy and safety of ruxolitinib in regularly transfused patients with thalassemia: results from a phase 2a study. *Blood*. 2018;131(2): 263-265.
 119. Cappellini MD, Porter JB, Viprakasit V, Taher AT. A paradigm shift on beta-thalassaemia treatment: how will we manage this old disease with new therapies? *Blood Rev*. 2018;32(4):300-311.
 120. Guerra A, Musallam KM, Taher AT, Rivella S. Emerging therapies [published correction appears in *Hematol Oncol Clin North Am*. 2018]. *Hematol Oncol Clin North Am*. 2018; 32(2):343-352.
 121. Casu C, Nemeth E, Rivella S. Hepcidin agonists as therapeutic tools. *Blood*. 2018; 131(16):1790-1794.