

Martin A. 'Mac' Cheever

Twelve immunotherapy drugs that could cure cancers

Authors' address

Martin A. 'Mac' Cheever¹

¹Seattle Cancer Care Alliance, Fred Hutchinson/University of Washington Cancer Consortium, Seattle, WA, USA.

Correspondence to:

Martin A. 'Mac' Cheever

Seattle Cancer Care Alliance

Fred Hutchinson/University of Washington Cancer Consortium

825 Eastlake Ave, E., G4-810

PO Box 19023

Seattle, WA 98109-1023, USA

Tel.: +1 206 288 6370

Fax: +1 206 288 6681

e-mail: mcheever@seattlecca.org

Summary: Immune T cells can kill cancer cells. Cancer vaccines function by increasing the number of immune T cells. There are exceedingly strict biologic limits imposed on the immune system to prevent excessive T-cell activation and expansion. The same biological restrictions limit cancer vaccines. Immunotherapeutic agents that circumvent the biological restrictions have been invented and formulated, including (i) dendritic cell activators and growth factors, (ii) vaccine adjuvants, (iii) T-cell stimulators and growth factors, (iv) immune checkpoint inhibitors, and (v) agents to neutralize or inhibit suppressive cells, cytokines, and enzymes. Few of these agents are broadly available for the development of effective multiple component regimens. The major problem facing immunotherapy today is a lack of broad availability of agents already in existence. The National Cancer Institute has developed a well-vetted ranked list of agents with high potential to serve as immunotherapeutic drugs. This review focuses on 12 of the agents, all with proven ability to augment T-cell responses. Alone, each has little chance of making substantial inroads into cancer therapy. In combinations dictated by biology, the agents are overwhelmingly likely to have an impact. Future availability of these agents for development of innovative combination cancer therapy regimens will provide a benchmark for the resolve of the national cancer therapy translational research enterprise.

Keywords: T cell, vaccine, adjuvant, cancer, immunotherapy, translational

Introduction

It is well known that cancer can be cured by immune T cells. Hematologic malignancy can be cured by allogeneic T cells. Probability of relapse after bone marrow transplantation (BMT) for leukemia varies according to type of marrow graft and subsequent development of graft versus host disease (1). 'Mini-transplants' using non-myeloablative regimens with high dose immunosuppression to prevent rejection of the transplanted allogeneic bone marrow can eliminate leukemia and lymphoma exclusively by the donor T-cell-mediated allogeneic graft versus leukemia effect (2).

More common epithelial tumors can also be treated with allogeneic T cells, but less effectively. Mini-transplants for renal cell cancer (3) and breast cancer (4) have induced responses including complete remissions. The approach is limited by inordinate toxicity but establishes the proof of concept that T cells can eliminate epithelial tumors.

Immunological Reviews 2008

Vol. 222: 357–368

Printed in Singapore. All rights reserved

© 2008 The Author

Journal compilation © 2008 Blackwell Munksgaard

Immunological Reviews

0105-2896

A major endeavor for tumor immunotherapists has been trying to figure out how to invoke the therapeutic potential of autologous immune T cells to reproduce the therapeutic efficacy of allogeneic T cells without the inordinate toxicity. It is eminently clear that cancer cells are immunogenic and can be recognized and killed by autologous T cells. Cancer is the result of multiple genetic alterations occurring over many years. The cumulative effect of genetic alterations results in loss of control of cell growth and differentiation. Each change in protein structure, resulting from genetic alterations, represents a potential cancer antigen and target for T-cell attack. Cancer cells commonly possess many tumor-specific and tumor-associated antigens that can be shared or are unique (5). Ongoing and nascent immune responses to autochthonous cancer cells are commonly observed with both circulating and tumor-infiltrating immune T cells with the potential to recognize and kill cancer cells.

It is axiomatic that if large number of immune T cells capable of recognizing and killing cancer cells could be generated *in vivo* with cancer vaccine regimens or *ex vivo* for adoptive transfer many patients could benefit. The prognosis of cancer patients in many instances correlates with the number and character of immune T cells and tumor infiltrating T cells (6, 7). Autologous immune T cells have been used effectively in therapy in small numbers of cancer patients (8). Vaccines that induce immune T-cell responses have proven to be effective, but less so (9).

Immune T-cell number and function are both important components of vaccine efficacy (10). However, the fundamental role of cancer vaccines is to increase the number of cancer antigen-specific T cells *in vivo*. At the simplest level, the therapeutic effect of immune T cells, and thus vaccines, is quantitative and related to the number of T cells present. As the number of immune T cells increases, the effect of the vaccine should increase, all else being equal. The efficacy is related not only to the quantity of T cells but also to the length of time the T cells persist, i.e. in pharmacologic terms, the 'area under the curve.'

Cancer vaccines present antigen to the small number of T cells capable of 'recognizing and responding' to the antigen, which are thereby stimulated to proliferate and increase in number. There are exceedingly strict biologic limits imposed on the immune system to prevent untoward and excessive T-cell expansion. Many of the components and factors that allow T-cell activation and expansion in number have been identified and constructed as immunotherapeutic agents. Similarly, many of the components and factors that disallow activation, expansion, or persistence of immune effector cells

have been identified, and agents that interfere with the disallowment have also been constructed as immunotherapeutic agents (11).

Taken together, the agents needed to greatly augment immune T-cell responses have been discovered. Many new biological agents are available as potential immunotherapeutic drugs to substantially increase the number and therapeutic function of immune T cells *in vivo*.

Agents that could take cancer vaccines from interesting biological phenomena to effective therapeutic agents include:

- T-cell growth factors to increase number and repertoire of naive T cells
- Growth factors to increase the number of dendritic cells (DCs)
- Agonists to activate DCs and other antigen-presenting cells (APCs)
- Adjuvants to allow and augment cancer vaccines
- Agonists to activate and stimulate T cells
- Inhibitors of T-cell checkpoint blockade
- T-cell growth factors to increase the growth and survival of immune T cells
- Agents to inhibit, block, or neutralize cancer cell and immune cell-derived immunosuppressive cytokines

Agents in each category have been discovered, invented, constructed, and proven to have immunologic function in animal models and in human clinical trials. However, the majority of agents needed to improve cancer vaccines are not broadly available for testing in patients, especially for use in innovative multi-component regimens.

Lack of broad availability of immunotherapeutic agents: adjuvant example

The major problem facing immunotherapy today is no longer a lack of knowledge or a lack of agents with proven immunologic effects, rather a lack of broad availability of agents with already proven immunologic function. The example of adjuvants exemplifies that problem. There is one 'Universal Truth' about vaccines: adjuvants are needed to achieve highest levels of immune response. Despite this Universal Truth, few vaccine adjuvants are commonly available for broad testing. The question is 'Why aren't adjuvants available?' The National Cancer Institute (NCI), Food and Drug Administration (FDA), and industry all play a role.

The NCI and other funding agencies have spent literally hundreds of millions, if not billions, for cancer vaccine development. Very little has been spent for essential vaccine components. It is commonly accepted that adjuvants are needed to achieve highest level of immune response, in yet

there is no coordinated effort to assure that the appropriate adjuvants are available, even for ongoing NCI-funded cancer vaccine trials.

By FDA policy and custom, adjuvants are approved only as components of vaccines. Accordingly, the commonest adjuvants used by academics are agents approved for non-adjuvant purposes. Granulocyte-macrophage colony-stimulating factor (GM-CSF) is commonly used in academic vaccine trials, in part because it is readily available. GM-CSF was approved by the FDA as a hematopoietic growth factor, not as a vaccine adjuvant. If GM-CSF had activity only as an adjuvant, it would not be available for testing in cancer vaccines. In a similar vein, imiquimod, a drug approved for therapy of genital warts and superficial basal cell cancer, not as an adjuvant, is beginning to be used as an adjuvant in NCI-funded cancer vaccine trials. The 'Catch 22' is that adjuvants approved for non-adjuvant purposes are broadly available, whereas, adjuvants that function only as adjuvants are not broadly available, regardless of potency. The 'majority of cancer drug development takes place post-approval,' an observation of Robert Capizzi. Adjuvants are not approved as adjuvants and thus not available for cancer drug development. The FDA has given no clear path forward for broad testing and approval of adjuvants that are not effective as monotherapy.

Industry decisions are driven by the 'Invisible hand of the market.' Rational decisions are made based on regulatory and commercial concerns. Companies with great adjuvants have tended to develop them as monotherapy, e.g. CpG and interleukin-12 (IL-12). Agents with adjuvant potential that fail as monotherapy are often 'left on the shelf,' as companies do not see a clear and rational regulatory and commercial path forward.

NCI immunotherapy agent workshop to rank agents with high potential to serve as immunotherapeutic drugs

What is the solution to the lack of availability vaccine adjuvants and other previously invented critical immunotherapeutic agents? The NCI has taken one small step by developing a ranked list of agents with high potential to serve as immunotherapeutic drugs (12). The ranking was exceedingly well vetted, with open and substantial input. The list represents a 'consensus' of what is most needed. There was a range in participant representation from academia, industry, the NCI, and the FDA. Participants were selected from suggestions by the American Association for Cancer Research, American Association of Immunologists, American Society of Oncology, American Society of Hematology, the Cancer Vaccine Consortium,

and the International Society of Biological Therapy of Cancer, and NCI intramural and extramural program heads.

These agents have wide appeal to the immunotherapy community and, by consensus, hold particular promise for use in cancer therapy. The exceedingly broad nature of the consensus behind this list should facilitate NCI discussions on the availability of clinical grade immunotherapeutic drugs for human trials. The agents were not selected specifically with cancer vaccines in mind but are essentially the agents needed to improve cancer vaccines.

The process of developing a ranked list of immunotherapy agents began with the NCI opening a web site to ask for suggestions. The web site was advertised and promoted broadly. A total of 124 agents were suggested. Most had demonstrated immunological or physiological function. This response is substantial evidence of the vigor of the immunology and immunotherapy fields. There was a broad desire for vaccine adjuvants, T-cell growth factors, agents to inhibit immune checkpoint blockade, functional antibodies, cytokines, ligands and receptors to activate or augment T-cell responses and/or to inhibit suppressor circuits, and agents left on the shelf by drug companies.

A workshop was held at the NCI July 12, 2007 to develop a ranked list of agents with high potential for use in cancer therapy (12). The organizing committee selected the top 30 agents winnowed from the 124 suggested agents. The committee focused on agents with the greatest potential for broad usage by multiple investigators in multiple different regimens and excluded specific antigens for vaccines and antigen-specific antibodies regardless of attractiveness or potential utility. The ranked list of 20 agents (Table 1) derives value from the broad and substantial input. The process, participants, and full report including PowerPoint® presentations can be viewed and downloaded from the NCI web site (<http://web.ncifcrf.gov/research/brb/workshops.asp>) (12).

The criteria used for inclusion on ranked list included (i) potential for use in cancer therapy, (ii) perceived need by multiple, independent clinical investigators, (iii) potential use in more than one clinical setting, i.e., against different tumor types or as part of multiple therapy regimens, and (iv) not broadly available for testing in patients. Criteria not used included (i) prior failed attempts to commercialize and (ii) intellectual property.

The final ranked list derived from discussions of each agent. Agents at the top of the list were considered the most desirable based on current evidence. It was well recognized by the participants that many agents with fewer data, including

Table 1. Rankings of agents with high potential for use in treating cancer

Rank*	Agent	Agent category
1	IL-15	T-cell growth factor
2	Anti-PD1 and/or anti-B7-H1 (PD-1L)	*T-cell checkpoint blockade inhibitor
3	IL-12	Vaccine adjuvant
4	Anti-CD40 and/or CD40L	Antigen presenting cell stimulator
5	IL-7	T-cell growth factor
6	CpG	Vaccine adjuvant
7	IMT	Enzyme inhibitor
8	Anti-CD137 (anti-4-1BB)	T-cell stimulator
9	Anti-TGF- β	Signaling inhibitor
10	Anti-IL-10 Receptor or Anti-IL-10	Suppression inhibitor
11	Flt3L	Dendritic cell growth factor/vaccine adjuvant
12	Anti-glucocorticoid-induced TNF receptor (GITR)	T-cell stimulator
13	CCL21 adenovirus	T-cell attracting chemokine
14	MPL	Vaccine adjuvant
15	PolyI:C and/or PolyI:CLC	Vaccine adjuvant
16	Anti-OX40	T-cell stimulator
17	Anti-B7-H4	T-cell checkpoint blockade inhibitor
18	Resiquimod and/or 852A	Vaccine adjuvant
19	LIGHT and/or LIGHT vector	T-cell stimulator
20	Anti-lymphocyte activation gene-3 (LAG-3)	T-cell checkpoint blockade inhibitor

*Anti-CTLA-4, a T-cell checkpoint blockade inhibitor, was considered of exceedingly high value but was not included on the list, as it is being produced by two companies and is likely to be approved by the FDA within the foreseeable future.

agents not currently on the list, will ultimately prove to be more important than those at the top of the list. Although the ranking is well vetted and based on the cumulative knowledge of the broad immunotherapy and cancer research communities, the choice and desirability of individual agents will undoubtedly change with new knowledge. Because the priorities are based on incomplete knowledge, the process should be a dynamic one that can be revised as more data appear. A common suggestion was that a mechanism should be developed to continually update the list.

Possible positive outcomes of having a well-vetted ranked list based on a broad consensus of the immunology and immunotherapy community could include encouragement of (i) Rapid Access to Interventional Development (RAID) applications for the manufacture, (ii) distribution of company-manufactured agents through RAID or the Cancer Therapy Evaluation Program, and (iii) reinvigoration of pharma/biotech efforts to develop them. This rank-setting exercise should also serve as a report card. Future availability

of these agents for broad testing and development will provide a benchmark for the strength and resolve of the national cancer therapy development enterprise.

This review focuses on 12 agents on the list that have proven immunologic efficacy and have been in clinical trials or will shortly be taken into clinical trials (Table 2). If tested creatively and intelligently in multiple combinations and regimens with cancer vaccines, regimens will surely be derived that will be able to benefit many patients with cancer. The major message is that the drugs needed to treat many patients with cancer have already been invented.

T-cell growth factor: IL-15

IL-15 is a T-cell growth factor, similar to IL-2 (13–22). Preclinical data overwhelmingly suggest that IL-15 could have an exceedingly important role in cancer vaccine and adoptive T-cell regimens. In contrast to IL-2, IL-15 inhibits antigen-induced cell death of T cells, whereas IL-2 promotes antigen-induced cell death. IL-15 is made by DCs, macrophages, and stromal cells, not T cells. It acts on CD8⁺ T cells, CD4⁺ T cells, natural killer (NK) cells, and mast cells. It uses a unique IL-15R α chain, but shares the β and γ chains of the IL-2R.

IL-15 is necessary for maintenance of CD8⁺ memory cells and NK cell development. In vaccines, it promotes induction of longer-lived and higher-avidity CD8⁺ T cells that can kill tumor cells very effectively. Greenberg's group (22) showed that IL-15 can reverse T-cell anergy. Palucka's group (14) demonstrated that IL-15 promotes *in vitro* differentiation of

Table 2. Twelve immunotherapy drugs that could cure cancer

Agent category	Agent
T-cell growth factors	IL-15 IL-7
T-cell checkpoint blockade inhibitor	Anti-PD1 and/or anti-B7-H1 (PD-1L)
Antigen-presenting cell stimulator	Anti-CD40 and/or CD40L
Enzyme inhibitor	IMT
T-cell stimulator	Anti-CD137 (anti-4-1BB)
DC growth factor/vaccine adjuvant	Flt3L
Vaccine adjuvants	IL-12 CpG MPL PolyI:C and/or PolyI:CLC Resiquimod and/or 852A

Eight agents on the NCI ranked list (Table 1) are excluded from this list, only as less is known about their effects on T-cell vaccine responses. The agents not included as well as many other might ultimately prove to have greater immunotherapy efficacy than the agents chosen for this review.

monocyte-derived DCs that are potent inducers of CD8⁺ T cells. Unpublished data from Berzofsky's laboratory indicate that IL-15 can overcome lack of CD4⁺ T-cell help in cytotoxic T-lymphocyte (CTL) induction. Unpublished data from Khleif and Pavlakis and Felber show that IL-15-expressing plasmids can induce tumor regression in mice via intratumoral injection or hydrodynamic delivery. No clinical data are available.

The information on IL-15 in this section is taken, largely verbatim, from a presentation by Jay Berzofsky at the NCI Immunotherapy Agent Workshop (12). A full report of the presentation with slides and comments, as well as presentations on the subsequent agents to be discussed in this review, can be downloaded from the NCI web site (<http://web.ncifcrf.gov/research/brb/workshops.asp>).

The contemplated uses for IL-15 include as (i) a vaccine adjuvant to induce longer-lived, higher avidity, more efficacious CD8⁺ T cells, single agent to overcome T-cell energy, (ii) as a T-cell growth factor to be used in conjunction with adoptive T-cell immunotherapy, in place of IL-2, (iii) induction of DC precursors into Langerhans-like DCs for DC-based vaccines, and (iv) intratumoral or systemic cytokine therapy for cancer.

T-cell growth factor: IL-7

IL-7 is a T-cell growth factor responsible for homeostatic expansion of naive cells during lymphopenia (23–31). In mice and primates, IL-7 treatment (24, 26) induces dramatic expansions in peripheral T-cell numbers without obvious toxicity. In mice and primates, following BMT, IL-7 increases the rate and degree of immune reconstitution (27). In humans, IL-7 is required for T-cell development and for naive T-cell survival in the periphery. During lymphopenia, IL-7 accumulates as a result of diminished utilization.

Two-phase I/II clinical trials have been completed in patients with cancer (31, Sportes *et al.*, manuscript submitted). The trials established proof of principle. IL-7 induced dramatic increases in total body CD4⁺ and CD8⁺ T cells with modest increases in NK cells and $\delta\gamma$ T cells with no change in mature B cells or selective increase in T-regulatory cells (Tregs). There appeared to be a preferential expansion of naive T cells with an increased T-cell repertoire diversity, likely related to cycling of recent thymic emigrants. In mice, IL-7 is also a vaccine adjuvant (30) with its most dramatic effects on subdominant responses.

IL-7 and IL-15 are both T-cell growth factors and vaccine adjuvants, but they work at different phases of T-cell activation. Both alter T-cell homeostasis but work on different

populations. IL-7 signals resting T cells and preferentially expands naive T cells, leading to increased repertoire diversity. IL-15 preferentially signals activated T cells and expands effector/memory populations. The information on IL-7 is taken, largely verbatim, from a presentation by Crystal L. Mackall (12).

The contemplated uses of IL-7 are (i) immunorestorative T-cell growth factor for naive T cells, especially in elderly patients and patients with prior lympholytic chemotherapy, and (ii) vaccine adjuvant to enhance the effectiveness of tumor vaccines targeting weak self-antigens.

T-cell checkpoint blockade inhibitor: anti-programmed death-1

Programmed death-1 (PD-1) is a negative regulator of T-cell function (32–38). Blocking PD-1 can increase anti-tumor T-cell immunity. PD-1, a receptor member of the immunoglobulin superfamily, structurally related to cytotoxic T-lymphocyte antigen-4 (CTLA-4) and CD28, is upregulated on activated T and B cells and monocytes. Binding to PD-1 ligand (PD-L1) results in inhibition of T-cell proliferation. PD-L1 is expressed on T and B cells, macrophages, and DCs, as well as parenchymal and tumor cells. In mice, PD-1 is implicated in tolerance induction and appears to 'protect' them from immune attack by CTLs. Blockade of PD-1 or PD-L1 results in long-lasting tumor regression. In humans, abrogation of PD-1 can increase the number of functional cytokine-secreting CTLs, presumably through increased proliferation. PD-L1, expressed on many human tumors, is associated with a reduced T-cell infiltrate and a poorer prognosis. Hatanishi *et al.* (38) showed that ovarian cancer patients who had greater levels of PD-L1 on tumor cells had less T-cell infiltration and lower survival rates than those who expressed little or no PD ligand.

PD-L1 is expressed by esophageal, colon, lung, and ovarian cancer and melanoma by immunohistochemistry. Thus, these tumors might be appropriate candidates for initial testing of anti-PD-1, especially for those tumors with T-cell infiltration. Use of antibody to block PD-L1 is an alternative approach that might also be useful. However, targeting the tumor may be more difficult than targeting T cells, if saturation of all tumor tissue is needed. The information on anti-PD-1 is taken, largely verbatim, from a presentation by Jeffrey Weber (12).

The contemplated uses of anti-PD-1 are (i) combined with vaccines to increase the number of activated T cells, (ii) as monotherapy for cancers that have infiltrating T cells or a nascent or ongoing immune response, and (iii) combined with anti-CTLA-4 therapy.

APC stimulator: CD40 agonists

Anti-CD40 and CD40L are potent agonists for increasing APC function (39–50). CD40, a member of the tumor necrosis factor (TNF) receptor superfamily, is a costimulatory protein found on APCs. Ligation of CD40 on APCs by CD40L expressed by T cells enhances the expression of major histocompatibility complex and costimulatory molecules on APCs, stimulates the production of pro-inflammatory cytokines such as IL-12, and leads to T-cell activation and cell-mediated immunity (44). In mice, agonist CD40 monoclonal antibody (mAb) can mimic the signal of CD40L and substitute for the function of CD4⁺ lymphocytes (39). Agonist CD40 mAb can also overcome T-cell tolerance in tumor-bearing mice, evoke effective CTL responses, and enhance the efficacy of anti-tumor vaccines (42).

In addition to APC activation and augmentation of T-cell immunity, CD40 agonists have other potentially therapeutic effects including direct tumor inhibition, anti-angiogenesis (41), and activation of NK cells (45) and macrophages (40). Engagement of CD40 on tumor cells *in vitro* has been shown to inhibit the growth of solid tumor cells and high-grade B-cell lymphoma lines, a direct cytotoxic effect resulting in tumor regression through apoptosis and necrosis (43).

Two categories of agents have proven efficacy as CD40 agonists, recombinant trimeric ligand, and mAb. Dr Sondel described available unpublished and published data on clinical experience (47–50). One phase I trial enrolled 29 patients with melanoma or other solid tumors. Four subjects had measurable objective responses by RECIST criteria. Most showed upregulation of the CD86 costimulatory molecule. Another phase I trial with recombinant human trimeric CD40 ligand showed two partial responses out of 32 solid tumors or non-Hodgkin's lymphoma (49, R. H. Vonderheide, personal communication regarding unpublished observations, July 2007).

CD40 agonists have great potential for use in combination with other agents. Preclinical models suggest efficacy with chemotherapy, radiation therapy, Toll-like receptor (TLR) agonists, and cytokines. The information on anti-CD40 is taken, largely verbatim, from a presentation by Paul M. Sondel (12).

The contemplated uses for anti-CD40 are (i) for systemic or local activation of APCs as a component of a regimen to induce or augment T-cell responses, (ii) to direct inhibition of CD40⁺ tumors, and (iii) in combination with other cancer therapies.

Inhibitor of an immunosuppressive enzyme inhibitor: 1-methyl tryptophan

1-Methyl tryptophan (1MT) is a potent small molecule inhibitor of the immunosuppressive enzyme indoleamine 2,3-

dioxygenase (IDO) (51–59). IDO is highly expressed in tumor cells and plasmacytoid DCs in tumor-draining lymph nodes. IDO catalyzes tryptophan degradation, leading to insufficient tryptophan and thereby limiting antigen-induced T-cell activation and mediating immunosuppression in cancer. 1MT inhibits IDO, reversing the immunosuppressive effects of IDO. IDO knockout mice are resistant to inflammatory skin carcinogenesis. 1MT limits growth of grafted or transgenic tumors in combination with chemotherapy in models dependent on CD4⁺ T-cell responses. In humans, clinical trials are being planned. The information on 1MT is taken, largely verbatim, from a presentation by George Prendergast (12).

The contemplated uses of 1MT are (i) as a general reversal of a mechanism of tumoral immune suppression alone or in combination with cytotoxic chemotherapy, tumor vaccines, TLR agonists (e.g. CpG), radiotherapy, mAbs, or drugs that target other mechanisms of immune suppression (e.g. OX40, PDL-1), and (ii) for cancer prevention in clinical states of chronic inflammation.

T-cell stimulator: anti-CD137 (anti-4-1BB)

Anti-CD137 is a T-cell stimulator with potent costimulatory, anti-apoptotic, and proliferative activity (60–69). The target, CD137, as a member of the TNF superfamily of receptors and is present on activated T cells, NK cells, and NKT cells. The receptor is not present on tumors. In preclinical studies, anti-CD137 could costimulate T cells that are suboptimally stimulated with anti-CD3, enhance T-cell responses to simian immunodeficiency virus gag DNA vaccines, and treat immunogenic murine tumors as monotherapy or in conjunction with vaccination. Anti-tumor effects have been observed with murine anti-4-1BB in various *in vivo* tumor models. Preclinical toxicity studies in mice suggested a predominance of NKT cells in the liver (e.g. hepatic necrosis, elevated transaminases). In monkey models, occasional mild colitis was observed, possibly related to the high number of activated lymphocytes in intestinal mucosa.

A phase I, first-in-human study is ongoing and is being expanded to a phase II trial. The subjects are advanced cancer patients with a variety of tumors. Toxicities have consisted of faint skin rash, mild neutropenia, and hepatotoxicity (likely to be dose limiting). A phase I trial of anti-4-1BB in combination with paclitaxel and carboplatin is accruing, and another one involving radiotherapy or chemoradiotherapy is being planned. The information on 1MT is taken, largely verbatim, from a presentation by Kim Margolin (12).

The contemplated uses of IMT as part of multiple different antigen-specific strategies involving suboptimal T-cell responses include (i) vaccine and cytokine therapy, (ii) augmentation of nascent anti-tumor responses, and (iii) in combination with checkpoint blockade (e.g. anti-CTLA-4A).

DC growth factor/vaccine adjuvant agent: Flt3 ligand

Flt3 ligand (Flt3L) is a hematopoietic growth factor that induces expansion and differentiation of all DC progenitors, especially interferon-producing killer and plasmacytoid DCs (70–76). Systemic administration of Flt3L increases DC numbers circulating in blood and in secondary lymphoid tissues as well as tumors. Some preclinical models report anti-tumor activity with systemic Flt3L protein alone, though effects are very variable. There are a number of reports demonstrating synergy between Flt3L and DC activators such as CpG and anti-CD40 in generating anti-tumor activity.

Phase I/II studies have examined Flt3L alone, with peptide vaccines, as DC stimulators, and after BMT. Increases in circulating DCs are consistently reported. Increased T-cell responses are commonly reported. CEA-loaded Flt3L-mobilized DC vaccine for colon cancer reported increased CD8⁺ T-cell priming and clinical responses. MAGE-loaded Flt3L-mobilized DC vaccine (with CD40L *in vitro* DC activation for melanoma) reported increased CD8⁺ T-cell responses. Studies were generally too small and variable to make significant statements about clinical efficacy. Immunex terminated trials after lack of obvious efficacy as single agent or with sCD40L. The information on Flt3L is taken, largely verbatim, from a presentation by Drew Pardoll (12).

The contemplated uses of Flt3L are (i) in combination with vaccines, BMT, BMT+vaccine, adoptive T-cell therapy, TLR agonists, and other immune stimulants, (ii) local administration (topical, admixture with vaccine preparations, intratumoral injection) to enhance DCs, (iii) transduction of Flt3L gene into engineered tumor vaccines or *in vivo* delivery (i.e. plasmid vaccination or in viral carriers), and (iv) for DC mobilization for *ex vivo* DC vaccines.

Vaccine adjuvant: IL-12

IL-12 is an exceedingly potent immune adjuvant (77–87). IL-12 promotes interferon- γ release by IL-12R-expressing T and NK cells and induces T-helper 1 cell (Th1) polarization, as well as proliferation of interferon- γ -expressing T cells. IL-12 plays a central role in resistance to mycobacterial and intracellular pathogens (e.g. parasites). It also plays an important part in anti-cancer development and immunity in animal systems.

Phase I and II clinical trials showed very modest efficacy alone with a handful of melanoma and renal cell cancer responses. It did not demonstrate sufficient clinical activity as a stand-alone drug to warrant further development, according to the standard oncology paradigm. It was originally developed as a systemic cytokine, but it proved challenging to administer safely. However, studies in mice have shown IL-12 to be exceedingly effective as a vaccine adjuvant. Owing to systemic toxicity, delivery might best be by local injection including use of viral or plasmid vectors. The information on IL-12 is taken, largely verbatim, from a presentation by Jeffrey Weber (12).

The contemplated uses for IL-12 are (i) as vaccine adjuvant, and (ii) as an adjuvant to cytokine therapy to polarize Th1 responses and augment CD8⁺ T-cell responses in any antigen-specific strategy.

Vaccine adjuvant: CpG

CpG is a potent TLR9 agonist (88–97). Unmethylated CpG is prevalent in bacterial and viral DNA but not in vertebrate genomes. CpGs target TLR9 with activity as a vaccine adjuvant and immune modulator. In humans, TLR9 is expressed on B cells and plasmacytoid DCs. In the mouse, it is expressed on B cells, monocytes, and all DCs. These species-based differences make it a bit difficult to extrapolate preclinical data. Targeting TLR9 with CpG leads to B-cell proliferation and differentiation to plasma cells, immunoglobulin G isotype switching and antibody secretion, maturation of plasmacytoid DCs with type I interferon secretion, activation of NK cells with enhancement of antibody-dependent cell-mediated cytotoxic activity, stimulation of Th1 cell response with CTL activation, but also has opposing effects, e.g. Treg generation, IDO expression, and anti-inflammatory (IL-10) cytokine secretion.

CpG has been studied in phase I and II trials as monotherapy, in combinations, and as a vaccine adjuvant. CpG is a category of agent with many different nucleotide sequences producing similar but not identical results. Two randomized trials of CpG in combination with chemotherapy in patients with non-small cell lung cancer failed to show benefit.

CpG has shown substantial activity as a vaccine adjuvant for peptide, protein, DC, and autologous tumor cell vaccines. It induces strong Th-1-promoting effects and CTL responses, compared with other TLR ligands, and appears to be synergistic with GM-CSF and other adjuvants. As an immune modulator, it has activity in combination with many other types of therapy including anti-CD40, anti-CTLA-4, IL-13 exotoxin, chemotherapy, radiation therapy, and cryoablation.

The information on CpG is taken, largely verbatim, from a presentation by Ellis Reinherz (12).

The contemplated uses of CpG are (i) as a vaccine adjuvant and (ii) in combination with multiple other types of cancer therapeutic interventions.

Vaccine adjuvant: monophosphoryl lipid A

Monophosphoryl lipid A (MPL or MPLA) is an effective and exceedingly safe and non-toxic adjuvant vaccine adjuvant (98–107). It is a highly specific TLR4 agonist. MPL is a component of endotoxin, i.e. lipopolysaccharide (LPS). LPS functions as a vaccine adjuvant but is considered too toxic for clinical use. However, purifying MPL from *Salmonella minnesota* endotoxin yields an excellent, low-toxicity adjuvant capable of activating macrophages and especially DCs. MPL has been shown in animal models to elicit responses to antigens of low immunogenic potential and can be administered by various routes and used in multiple formulations, including combinations with other adjuvants. It is a safe adjuvant with a side effect profile equivalent to that of alum. More than 120 000 doses have been administered to more than 50 000 human subjects. It has been approved as a component of a hepatitis B virus vaccine in the European Union. MPL is being used in combination with QS-21 and CpG in a proposed pivotal trial MAGE-A2 protein for non-small cell lung cancer by GlaxoSmithKline. Synthetic TLR4 agonists are being developed with similar potential as MPL. The information on MPL is taken, largely verbatim, from a presentation by Martin Cheever (12).

The contemplated uses for MPL are (i) as an adjuvant for vaccines as a single adjuvant or combined with other adjuvants with multiple different types of antigens, different formulations, different routes of administration and adjuvant combinations, and (ii) for activation of innate immune responses.

Vaccine adjuvants: polyI:C and polyICLC

PolyI:C and polyICLC are potent TLR3 agonists with vaccine adjuvant potential (108–117). PolyI:C is double-stranded polyinosinic:polycytidylic acid. When stabilized with poly-L-lysine and carboxymethylcellulose, it is known as polyICLC, and is more stable and, in that regard has greater activity. Preclinical studies have demonstrated that these TLR3 agonists activate human DCs, improve antigen presentation, and enhance Th1 polarization and adjuvant vaccines. In humans, they are strong activators of Th1 responses, CD8⁺ T cells, and NK cells. PolyI:C and related compounds are being tested broadly in humans in clinical trials in viral infections. In cancer, an

ongoing phase I/II trial of polyICLC involves patients with malignant gliomas. The agent is also being tested in prostate cancer patients for adjuvant effect with a MUC1 100-mer peptide vaccine. TLR3 agonists have uncertain activity as monotherapy but substantial potential activity as vaccine adjuvants. The information on polyI:C and polyICLC is taken, largely verbatim from a presentation by Anna Karolina Palucka (12).

The contemplated uses for polyI:C and polyICLC are (i) as adjuvant for cancer vaccines for use alone and in combination with other categories of adjuvants and (ii) for prevention or mitigation of viral infections.

Vaccine adjuvants: resiquimod and 852A

The imidazoquinolinamines resiquimod and 852A are TLR7/8 agonists, which induce innate and adaptive immune responses (118–125). Their biology is similar to that of imiquimod, a TLR7 agonist, which is currently FDA approved as a topical medication for genital warts and basal cell skin cancer. Anecdotal reports have indicated that imiquimod is useful for managing some cases of melanoma with cutaneous metastases. Resiquimod induces production of interferon- α , IL-6, IL-8, IL-12, and TNF- α from DCs, monocytes, and macrophages. Activation stimulates the innate immune response and leads to subsequent Th1 cell-mediated immune responses.

Another TLR7 agonist is 852A, which stimulates plasmacytoid DCs and can be administered as an intravenous solution. Scant data are available on 852A, although indications are that it may be more potent than resiquimod. Clinical responses have been reported for carcinoid tumor, melanoma, and breast cancer (124). There is a dearth of publications on 852A. However, mixed TLR 7/8 agonists would be very interesting for local use. Having TLR7 agonists available would add to vaccine adjuvant options. Having topical and systemic formulations could also be useful. The information on resiquimod and 852A is taken, largely verbatim, from a presentation by Louis M. Weiner (12).

The contemplated uses for resiquimod and 852A are (i) as vaccine adjuvant, (ii) as monotherapy for immune activation especially for topical administration, (iii) in combination with chemotherapy agents, and (iv) in combination with antibodies.

Conclusion

The tools needed to raise T-cell levels to extraordinary levels *in vivo* are at hand. The 12 agents described all have proven ability to substantially increase specific T-cell responses. Alone the

agents have very little chance of making substantial inroads into cancer therapy. Together, in innovative and intelligent combinations, the agents are overwhelmingly likely to have an impact.

Imagine sequential treatment of cancer patients with IL-7 to increase the number and repertoire of naive T cells, Flt3 to increase the total body burden of DCs, anti-CD40 to activate the DCs, a polyvalent cancer vaccine with a combination of IL-12, CpG, MPL, polyI:C and resiquimod adjuvants to activate and induce the proliferation of cancer immune T cells, 4-1-BB to further stimulate activation and proliferation of the immune T cells, anti-PD1 to inhibit T-cell checkpoint blockade, IL-15 to maintain proliferation and survival of the immune T cells, and 1MT to inhibit immunosuppression by IDO. The number of immune T cells achievable should make the *in vivo* environment exceedingly inhospitable for cancer cells. If that were not enough, it would be straightforward to expand the tumor immune T cells *ex vivo* and transfer them *in vivo* to achieve even larger numbers.

The above scenario is fraught with difficulties, but development of biologically directed multi-component immunotherapeutic regimens including the use of standard cancer therapies are eminently doable with today's technology.

The scientific discoveries are at hand. Applying the immunotherapy agents already discovered is essentially an engineering feat. The immunology scientific community has done an extraordinary job of developing the principles of T-cell immune responses. Preclinical scientists in academia and industry have combined to develop the immunotherapy agents needed to induce and greatly augment immune T-cell responses. The time is ripe to apply all of the principles of immunology to the therapy of cancer using the immunotherapy agents close at hand. The priorities have been set by the broad immunotherapy community (12).

The list of agents (Table 1) with high potential to serve as immunotherapeutic drugs contained eight additional agents, each with great potential: a signaling inhibitor, anti-transforming growth factor- β ; an immune suppression inhibitor, anti-IL-10 receptor; three T-cell stimulators, anti-glucocorticoid-induced TNF receptor, anti-OX40 and LIGHT; a T-cell attracting chemokine, CCL21; and two T-cell checkpoint blockade inhibitors, anti-B7-H4 and anti-lymphocyte activation gene-3. There is every reason to think that these agents as well as other agents in development will prove to be as effective as the 12 described in this review. The 12 described are just the start.

References

- Horowitz MM, et al. Graft-versus-leukemia reactions after bone marrow transplantation. *Blood* 1990;**5**:555–562.
- Baron F, Storb R. Allogeneic hematopoietic cell transplantation following nonmyeloablative conditioning as treatment for hematologic malignancies and inherited blood disorders. *Mol Ther* 2006;**13**:26–41.
- Lundqvist A, Childs R. Allogeneic hematopoietic cell transplantation as immunotherapy for solid tumors: current status and future directions. *J Immunother* 2005;**28**:281–288.
- Bishop MR, et al. Allogeneic lymphocytes induce tumor regression of advanced metastatic breast cancer. *J Clin Oncol* 2004;**22**:3886–3892.
- Finn OJ. Cancer vaccines: between the idea and the reality. *Nat Rev Immunol* 2003;**3**:630–641.
- Page F, et al. Effector memory T cells, early metastasis, and survival in colorectal cancer. *N Engl J Med* 2005;**353**:2654–2666.
- Zhang L, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med* 2003;**348**:203–213.
- Dudley ME, et al. Adoptive cell transfer therapy following non-myeloablative but lymphodepleting chemotherapy for the treatment of patients with refractory metastatic melanoma. *J Clin Oncol* 2005;**23**:2346–2357.
- Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 2004;**10**:909–915.
- Coulie PG, Connerotte T. Human tumor-specific T lymphocytes: does function matter more than number? *Curr Opin Immunol* 2005;**17**:320–325.
- Pardoll D, Allison J. Cancer immunotherapy: breaking the barriers to harvest the crop. *Nat Med* 2004;**10**:887–892.
- Cheever MA, Creekmore S eds. National Cancer Institute Agent Workshop Proceedings: July 12, 2007. <http://web.ncicrf.gov/research/brb/workshops.asp>
- Burton JD, et al. A lymphokine, provisionally designated interleukin T and produced by a human adult T cell leukemia line, stimulates T cell proliferation and the induction of lymphokine-activated killer cells. *Proc Natl Acad Sci USA* 1994;**91**:4935–4939.
- Mohamadizadeh M, et al. Interleukin 15 skews monocyte differentiation into dendritic cells with features of Langerhans cells. *J Exp Med* 2001;**194**:1013–1020.
- Tan JT, Ernst B, Kieper WC, LeRoy E, Sprent J, Surh CD. Interleukin (IL)-15 and IL-7 jointly regulate homeostatic proliferation of memory phenotype CD8+ cells but are not required for memory phenotype CD4+ cells. *J Exp Med* 2002;**195**:1523–1532.
- Becker TC, et al. Interleukin 15 is required for proliferative renewal of virus-specific memory CD8 T cells. *J Exp Med* 2002;**195**:1541–1548.
- Kobayashi H, et al. Role of trans-cellular IL-15 presentation in the activation of NK cell-mediated killing, which leads to enhanced tumor immunosurveillance. *Blood* 2005;**105**:721–727.
- Oh S, Berzofsky JA, Burke DS, Waldmann TA, Perera L. P. Coadministration of HIV vaccine vectors with vaccinia viruses expressing IL-15 but not IL-2 induces long-lasting cellular immunity. *Proc Natl Acad Sci USA* 2003;**100**:3392–3397.
- Oh S, Perera LP, Burke DS, Waldmann TA, Berzofsky JA. IL-15/IL-15R alpha-mediated avidity maturation of memory CD8+ T cells. *Proc Natl Acad Sci USA* 2004;**101**:15154–15159.
- Burkett PR, Koka R, Chien M, Chai S, Boone DL, Ma A. Coordinate expression and trans presentation of interleukin (IL)-15R alpha

- and IL-15 supports natural killer cell and memory CD8+ T cell homeostasis. *J Exp Med* 2004;**200**:825–834.
21. Klebanoff CA, et al. IL-15 enhances the in vivo antitumor activity of tumor-reactive CD8+ T cells. *Proc Natl Acad Sci USA* 2004;**101**:1969–1974.
22. Teague RM, et al. Interleukin-15 rescues tolerant CD8+ T cells for use in adoptive immunotherapy of established tumors. *Nat Med* 2006;**12**:335–341.
23. Schluns KS, Kieper WC, Jameson SC, Lefrancois L. Interleukin-7 mediates the homeostasis of naive and memory CD8 T cells in vivo. *Nat Immunol* 2000;**1**:426–432.
24. Fry TJ, Mackall CL. Interleukin-7: master regulator of peripheral T cell homeostasis? *Trends Immunol* 2001;**22**:564–571.
25. Geiselhart LA, Humphries CA, Gregorio TA, Mou S, Subleski J, Komschlies KL. IL-7 administration alters the CD4:CD8 ratio, increases T cell numbers, and increases T cell function in the absence of activation. *J Immunol* 2001;**166**:3019–3027.
26. Fry TJ, et al. IL-7 therapy dramatically alters peripheral T cell homeostasis in normal and SIV-infected nonhuman primates. *Blood* 2003;**101**:2294–2299.
27. Storek J, et al. Interleukin-7 improves CD4 T cell reconstitution after autologous CD34 cell transplantation in monkeys. *Blood* 2003;**101**:4209–4218.
28. Fry TJ, Mackall CL. The many faces of IL-7: from lymphopoiesis to peripheral T cell maintenance. *J Immunol* 2005;**174**:6571–6576.
29. Gattinoni L, et al. Removal of homeostatic cytokine sinks by lymphodepletion enhances the efficacy of adoptively transferred tumor-specific CD8+ T cells. *J Exp Med* 2005;**202**:907–912.
30. Melchionda F, Fry TJ, Milliron MJ, McKirdy MA, Tagaya Y, Mackall CL. Adjuvant IL-7 or IL-15 overcomes immunodominance and improves survival of the CD8+memory cell pool. *J Clin Invest* 2005;**115**:1177–1187.
31. Rosenberg SA, et al. IL-7 administration to humans leads to expansion of CD8+ and CD4+ cells but a relative decrease of CD4+ T-regulatory cells. *J Immunother* 2006;**29**:313–319.
32. Dong H, et al. Tumor-associated B7-H1 promotes T cell apoptosis: a potential mechanism of immune evasion [Erratum in: *Nat Med* 2002;**8**:1039]. *Nat Med* 2002;**8**:793–800.
33. Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, Minato N. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci USA* 2002;**99**:12293–12297.
34. Hirano F, et al. Blockade of B7-H1 and PD-1 by monoclonal antibodies potentiates cancer therapeutic immunity. *Cancer Res* 2005;**65**:1089–1096.
35. Konishi J, Yamazaki K, Azuma M, Kinoshita I, Dosaka-Akita H, Nishimura M. B7-H1 expression on non-small cell lung cancer cells and its relationship with tumor-infiltrating lymphocytes and their PD-1 expression. *Clin Cancer Res* 2004;**10**:5094–5100.
36. Liu X, et al. B7DC/PDL2 promotes tumor immunity by a PD-1-independent mechanism. *J Exp Med* 2003;**197**:1721–1730.
37. Barber DL, et al. Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature* 2006;**439**:682–687.
38. Hamanishi J, et al. Programmed cell death 1 ligand 1 and tumor-infiltrating CD8+ T lymphocytes are prognostic factors of human ovarian cancer. *Proc Natl Acad Sci USA* 2007;**104**:3360–3365.
39. Bennett SR, Carbone FR, Karamalis F, Flavell RA, Miller JF, Heath WR. Help for cytotoxic-T cell responses is mediated by CD40 signalling. *Nature* 1998;**393**:478–480.
40. Buhtoiarov IN, Lum H, Berke G, Paulnock DM, Sondel PM, Rakhmilevich AL. CD40 ligation induces antitumor reactivity of murine macrophages via an IFN gamma-dependent mechanism. *J Immunol* 2005;**174**:6013–6022.
41. Chiodoni C, et al. Triggering CD40 on endothelial cells contributes to tumor growth. *J Exp Med* 2006;**203**:2441–2450.
42. French RR, Chan HT, Tutt AL, Glennie MJ. CD40 antibody evokes a cytotoxic T cell response that eradicates lymphoma and bypasses T cell help. *Nat Med* 1999;**5**:548–553.
43. Funakoshi S, et al. Inhibition of human B-cell lymphoma growth by CD40 stimulation. *Blood* 1994;**83**:2787–2794.
44. Grewal IS, Flavell RA. CD40 and CD154 in cell-mediated immunity. *Annu Rev Immunol* 1998;**16**:111–135.
45. Turner JG, et al. Anti-CD40 antibody induces antitumor and anti-metastatic effects: the role of NK cells. *J Immunol* 2001;**166**:89–94.
46. van Kooten C, Banchereau J. CD40-CD40 ligand. *J Leuk Biol* 2000;**67**:2–17.
47. Vonderheide RH. Prospect of targeting the CD40 pathway for cancer therapy. *Clin Cancer Res* 2007;**13**:1083–1088.
48. Vonderheide RH, et al. Clinical activity and immune modulation in cancer patients treated with CP-870,893, a novel CD40 agonist monoclonal antibody. *J Clin Oncol* 2007;**25**:876–883.
49. Vonderheide RH, et al. Phase I study of recombinant human CD40 ligand in cancer patients. *J Clin Oncol* 2001;**19**:3280–3287.
50. Khalil M, Vonderheide RH. Anti-CD40 agonist antibodies: preclinical and clinical experience. *Update Cancer Ther* 2007;**b**:61–65.
51. Friberg M, et al. Indoleamine 2,3-dioxygenase contributes to tumor cell evasion of T cell-mediated rejection. *Int J Cancer* 2002;**101**:151–155.
52. Uyttenhove C, et al. Evidence for a tumoral immune resistance mechanism based on tryptophan degradation by indoleamine 2,3-dioxygenase. *Nat Med* 2003;**9**:1269–1274.
53. Munn DH, et al. Expression of indoleamine 2,3-dioxygenase by plasmacytoid dendritic cells in tumor-draining lymph nodes. *J Clin Invest* 2004;**114**:280–290.
54. Muller AJ, DuHadaway JB, Donover PS, Sutanto-Ward E, Prendergast GC. Inhibition of indoleamine 2,3-dioxygenase, a target of the cancer suppression gene Bin1, potentiates cancer chemotherapy. *Nat Med* 2005;**11**:312–319.
55. Fallarino F, et al. The combined effects of tryptophan starvation and tryptophan catabolites down-regulate T cell receptor zeta-chain and induce a regulatory phenotype in naive T cells. *J Immunol* 2006;**176**:6752–6761.
56. Metz R, DuHadaway JB, Kamasani U, Muller AJ, Prendergast GC. Novel tryptophan catabolic enzyme IDO2 is the preferred biochemical target of the antitumor IDO inhibitory compound D-1MT. *Cancer Res* 2007;**67**:7082–7087.
57. Hou DY, et al. Inhibition of IDO in dendritic cells by stereoisomers of 1-methyl-tryptophan correlates with anti-tumor responses. *Cancer Res* 2007;**67**:792–801.
58. Munn DH, Mellor AL. Indoleamine 2,3-dioxygenase and tumor-induced tolerance. *J Clin Invest* 2007;**117**:1147–1154.
59. Muller AJ, Prendergast GC. Indoleamine 2,3-dioxygenase in immune suppression and cancer. *Curr Cancer Drug Targets* 2007;**7**:31–40.
60. DeBenedette MA, Shahinian A, Mak TW, Watts TH. Costimulation of CD28 – T lymphocytes by 4-1BB ligand. *J Immunol* 1997;**158**:551–559.
61. Hurtado JC, Kim YJ, Kwon BS. Signals through 4-1BB are costimulatory to previously activated splenic T cells and inhibit activation-induced cell death. *J Immunol* 1997;**158**:2600–2609.
62. Melero I, et al. Monoclonal antibodies against the 4-1BB T cell activation molecule eradicate established tumors. *Nat Med* 1997;**3**:682–685.
63. Shuford WW, et al. 4-1BB costimulatory signals preferentially induce CD8+ T cell proliferation and lead to the amplification in

- vivo of cytotoxic T cell responses. *J Exp Med* 1997;**186**:47–55.
64. Takahashi C, Mittler RS, Vella AT. Cutting edge: 4-1BB is a bona fide CD8 T cell survival signal. *J Immunol* 1999;**162**:5037–5040.
 65. Wilcox RA, et al. Provision of antigen and CD137 signaling breaks immunological ignorance, promoting regression of poorly immunogenic tumors. *J Clin Invest* 2002;**109**:651–659.
 66. Wilcox RA, et al. Cutting edge: expression of functional CD137 receptor by dendritic cells. *J Immunol* 2002;**168**:4262–4267.
 67. May KF, Jr, Chen L, Zheng P, Liu Y. Anti-4-1BB monoclonal antibody enhances rejection of large tumor burden by promoting survival but not clonal expansion of tumor-specific CD8+ T cells. *Cancer Res* 2002;**62**:3459–3465.
 68. Lee HW, Park SJ, Choi BK, Kim HH, Nam KO, Kwon BS. 4-1BB promotes the survival of CD8+ T lymphocytes by increasing expression of Bcl-xL and Bfl-1. *J Immunol* 2002;**169**:4882–4888.
 69. Zhu Y, Zhu G, Luo L, Flies AS, Chen L. CD137 stimulation delivers an antigen-independent growth signal for T lymphocytes with memory phenotype. *Blood* 2007;**109**:4882–4889.
 70. Matthews W, Jordan CT, Wiegand GW, Pardoll D, Lemischka IR. A receptor tyrosine kinase specific to hematopoietic stem and progenitor cell-enriched populations. *Cell* 1991;**65**:1143–1152.
 71. Lynch DH, Andreassen A, Maraskovsky E, Whitmore J, Miller RE, Schuh JC. Flt3 ligand induces tumor regression and antitumor immune responses in vivo. *Nat Med* 1997;**3**:625–631.
 72. Marroquin CE, et al. Mobilization of dendritic cell precursors in patients with cancer by flt3 ligand allows the generation of higher yields of cultured dendritic cells. *J Immunother* 2002;**25**:278–288.
 73. Cui Y, et al. Immunotherapy of established tumors using bone marrow transplantation with antigen gene-modified hematopoietic stem cells. *Nat Med* 2003;**9**:952–958.
 74. Furumoto K, Soares L, Engleman EG, Merad M. Induction of potent antitumor immunity by in situ targeting of intratumoral DCs. *J Clin Invest* 2004;**113**:774–783.
 75. Chen W, Chan AS, Dawson AJ, Liang X, Blazar BR, Miller JS. FLT3 ligand administration after hematopoietic cell transplantation increases circulating dendritic cell precursors that can be activated by CpG oligodeoxynucleotides to enhance T cell and natural killer cell function. *Biol Blood Marrow Transplant* 2005;**11**:23–34.
 76. Davis ID, et al. Blood dendritic cells generated with Flt3 ligand and CD40 ligand prime CD8+ T cells efficiently in cancer patients. *J Immunother* 2006;**29**:499–511.
 77. Fallarino F, Uyttenhove C, Boon T, Gajewski TF. Endogenous IL-12 is necessary for rejection of P815 tumor variants in vivo. *J Immunol* 1996;**156**:1095–1100.
 78. Smyth MJ, et al. Differential tumor surveillance by natural killer (NK) and NKT cells. *J Exp Med* 2000;**191**:661–668.
 79. Brunda MJ, et al. Antitumor and antimetastatic activity of interleukin-12 against murine tumors. *J Exp Med* 1993;**178**:1223–1230.
 80. Noguchi Y, Jungbluth A, Richards EC, Old LJ. Effect of interleukin-12 on tumor induction by 3-methylcholanthrene. *Proc Natl Acad Sci USA* 1996;**93**:11798–11801.
 81. Nanni P, et al. Combined allogeneic tumor-cell vaccination and systemic interleukin-12 prevents mammary carcinogenesis in HER-2/neu transgenic mice. *J Exp Med* 2001;**194**:1195–1205.
 82. Colombo MP, Trinchieri G. Interleukin-12 in anti-tumor immunity and immunotherapy. *Cytokine Growth Factor Rev* 2002;**13**:155–168.
 83. Voest EE, et al. Inhibition of angiogenesis in vivo by interleukin-12. *J Natl Cancer Inst* 1995;**87**:581–586.
 84. Cebon J, et al. Two phase I studies of low dose recombinant human IL-12 with Melan-A and influenza peptides in subjects with advanced malignant melanoma. *Cancer Immun* 2003;**3**:7.
 85. Peterson AC, Harlin H, Gajewski TF. Immunization with Melan-A peptide-pulsed peripheral blood mononuclear cells plus recombinant human interleukin-12 induces clinical activity and T cell responses in advanced melanoma. *J Clin Oncol* 2003;**21**:2342–2348.
 86. Hamid O, et al. Alum with interleukin-12 augments immunity to a melanoma peptide vaccine: correlation with time to relapse in patients with resected high-risk disease. *Clin Cancer Res* 2007;**13**:215–222.
 87. Gollob JA, et al. Phase I trial of twice-weekly intravenous interleukin 12 in patients with metastatic renal cell cancer or malignant melanoma: ability to maintain IFN-gamma induction is associated with clinical response. *Clin Cancer Res* 2000;**6**:1678–1692.
 88. Friedberg JW, et al. Combination immunotherapy with a CpG oligonucleotide (1018 ISS) and rituximab in patients with non-Hodgkin lymphoma: increased interferon-alpha/beta-inducible gene expression, without significant toxicity. *Blood* 2005;**105**:489–495.
 89. Speiser DE, et al. Rapid and strong human CD8+ T cell responses to vaccination with peptide, IFA, and CpG oligodeoxynucleotide 7909. *J Clin Invest* 2005;**115**:739–746.
 90. Link BK, et al. Oligodeoxynucleotide CpG 7909 delivered as intravenous infusion demonstrates immunologic modulation in patients with previously treated non-Hodgkin lymphoma. *J Immunother* 2006;**29**:558–568.
 91. Pashenkov M, et al. Phase II trial of a toll-like receptor 9-activating oligonucleotide in patients with metastatic melanoma. *J Clin Oncol* 2006;**24**:5716–5724.
 92. ADIS R&D Profile. CpG 7909: PF 3512676, PF-3512676. *Drugs R D* 2006;**7**:312–316.
 93. Krieg AM. Therapeutic potential of Toll-like receptor 9 activation. *Nat Rev Drug Discov* 2006;**5**:471–484.
 94. Valmori D, et al. Vaccination with NY-ESO-1 protein and CpG in Montanide induces integrated antibody/Th1 responses and CD8 T cells through cross-priming. *Proc Natl Acad Sci USA* 2007;**104**:8947–8952.
 95. Molenkamp BG, et al. Intradermal CpG-B activates both plasmacytoid and myeloid dendritic cells in the sentinel lymph node of melanoma patients. *Clin Cancer Res* 2007;**13**:2961–2969.
 96. Kanzler H, Barrat FJ, Hessel EM, Coffman RL. Therapeutic targeting of innate immunity with Toll-like receptor agonists and antagonists. *Nat Med* 2007;**13**:552–559.
 97. Krieg AM. Development of TLR9 agonists for cancer therapy. *J Clin Invest* 2007;**117**:1184–1194.
 98. Chase JJ, et al. Effect of monophosphoryl lipid A on host resistance to bacterial infection. *Infect Immun* 1986;**53**:711–712.
 99. Childers NK, et al. Adjuvant activity of monophosphoryl lipid A for nasal and oral immunization with soluble or liposome-associated antigen. *Infect Immun* 2000;**68**:5509–5516.
 100. Baldrick P, Richardson D, Elliott G, Wheeler AW. Safety evaluation of monophosphoryl lipid A (MPL): an immunostimulatory adjuvant. *Regul Toxicol Pharmacol* 2002;**35**:398–413.
 101. Sondak VK, et al. Adjuvant immunotherapy of resected, intermediate-thickness, node-negative melanoma with an allogeneic tumor vaccine: overall results of a randomized trial of the Southwest Oncology Group. *J Clin Oncol* 2002;**20**:2058–2066.
 102. Ismaili J, et al. Monophosphoryl lipid A activates both human dendritic cells and T cells. *J Immunol* 2002;**168**:926–932.
 103. Atanackovic D, et al. Vaccine-induced CD4+ T cell responses to MAGE-3 protein in lung cancer patients. *J Immunol* 2004;**172**:3289–3296.
 104. Baldrige JR, et al. Taking a Toll on human disease: toll-like receptor 4 agonists as vaccine

- adjuvants and monotherapeutic agents. *Expert Opin Biol Ther* 2004;**4**:1129–1138.
105. Giannini SL, et al. Enhanced humoral and memory B cellular immunity using HPV16/18 L1 VLP vaccine formulated with the MPL/aluminum salt combination (AS04) compared to aluminium salt only. *Vaccine* 2006;**24**:5937–5949.
106. Nevens F, et al. Immunogenicity and safety of an experimental adjuvanted hepatitis B candidate vaccine in liver transplant patients. *Liver Transplant* 2006;**10**:1489–1495.
107. Mata-Haro V, Cekic C, Martin M, Chilton PM, Casella CR, Mitchell TC. The vaccine adjuvant monophosphoryl lipid A as a TRIF-biased agonist of TLR4. *Science* 2007;**316**:1628–1632.
108. Adams M, et al. Dendritic cell (DC) based therapy for cervical cancer: use of DC pulsed with tumour lysate and matured with a novel synthetic clinically non-toxic double stranded RNA analogue poly [I]: poly [C(12)U] (Ampligen R). *Vaccine* 2003;**21**:787–790.
109. Choe J, Kelker MS, Wilson IA. Crystal structure of human toll-like receptor 3 (TLR3) ectodomain. *Science* 2005;**309**:581–585.
110. Ewel CH, et al. Polyinosinic-polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose in combination with interleukin 2 in patients with cancer: clinical and immunological effects. *Cancer Res* 1992;**52**:3005–3010.
111. Giantonio BJ, et al. Toxicity and response evaluation of the interferon inducer poly ICLC administered at low dose in advanced renal carcinoma and relapsed or refractory lymphoma: a report of two clinical trials of the Eastern Cooperative Oncology Group. *Invest New Drugs* 2001;**19**:89–92.
112. Gitlin L, et al. Essential role of mda-5 in type I IFN responses to polyriboinosinic: polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc Natl Acad Sci USA* 2006;**103**:8459–8464.
113. Kawai T, Akira S. Innate immune recognition of viral infection. *Nat Immunol* 2006;**7**:131–137.
114. Mailliard RB, et al. Alpha-type-1 polarized dendritic cells: a novel immunization tool with optimized CTL-inducing activity. *Cancer Res* 2004;**64**:5934–5937.
115. Uematsu S, Akira S. Toll-like receptors and Type I interferons. *J Biol Chem* 2007;**282**:15319–15323.
116. Yoneyama M, et al. The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol* 2004;**5**:730–737.
117. Marshall-Clarke S, et al. Polyinosinic acid is a ligand for toll-like receptor 3. *J Biol Chem* 2007;**282**:24759–24766.
118. Mark KE, et al. Topical resiquimod 0.01% gel decreases herpes simplex virus type 2 genital shedding: a randomized, controlled trial. *J Inf Dis* 2007;**195**:1324–1331.
119. Wu JJ, Huang DB, Tying SK. Resiquimod: a new immune response modifier with potential as a vaccine adjuvant for Th1 immune responses. *Antiviral Res* 2004;**64**:79–83.
120. Jones T. Resiquimod 3M. *Curr Opin Investig Drugs* 2003;**4**:214–218.
121. Hengge UR, Benninghoff B, Ruzicka T, Goos M. Topical immunomodulators – progress towards treating inflammation, infection, and cancer. *Lancet Infect Dis* 2001;**1**:189–198.
122. Trinchieri G, Sher A. Cooperation of Toll-like receptor signals in innate immune defence. *Nat Rev Immunol* 2007;**7**:179–190.
123. Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol* 2004;**4**:499–511.
124. Dudek AZ, Yunis C, Kumar S, Harrison LI, Hawkinson RW, Miller JS. ASCO Annual Meeting Proceedings. *J Clin Oncol* 2005;**23** (Suppl.):2515.
125. Rechtsteiner G, Warger T, Osterloh P, Schild H, Radsak MP. Cutting edge: priming of CTL by transcutaneous peptide immunization with imiquimod. *J Immunol* 2005;**174**:2476–2480.