



US008163879B2

(12) **United States Patent**
Wong et al.

(10) **Patent No.:** **US 8,163,879 B2**
(45) **Date of Patent:** **Apr. 24, 2012**

(54) **FUSION MOLECULES AND IL-15 VARIANTS**

(75) Inventors: **Hing C. Wong**, Weston, FL (US); **Peter Rhode**, Miami, FL (US); **Xiaoyun Zhu**, Miami, FL (US); **Kai-ping Han**, Miramar, FL (US)

(73) Assignee: **Altor BioScience Corporation**, Miramar, FL (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/700,949**

(22) Filed: **Feb. 5, 2010**

(65) **Prior Publication Data**

US 2010/0278774 A1 Nov. 4, 2010

Related U.S. Application Data

(63) Continuation of application No. 12/151,980, filed on May 9, 2008.

(60) Provisional application No. 60/928,900, filed on May 11, 2007.

(51) **Int. Cl.**

C07K 14/54 (2006.01)

A61K 38/20 (2006.01)

C12N 5/10 (2006.01)

C12N 15/63 (2006.01)

C12P 21/02 (2006.01)

(52) **U.S. Cl.** **530/351**; 424/85.2; 435/69.52; 435/252.3; 435/320.1; 435/325; 435/366; 536/23.5

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2003/0180888 A1 9/2003 Fraser
2004/0156826 A1 8/2004 Dangond et al.
2004/0253587 A1 12/2004 Grabstein et al.
2006/0263857 A1 11/2006 Lefrancois et al.

FOREIGN PATENT DOCUMENTS

EP 1777294 A1 4/2007
WO WO 2005085282 * 2/2005
WO WO-2006063974 A2 6/2006
WO WO-2009117117 A1 9/2009

OTHER PUBLICATIONS

Lazar et al Mol. Cell. Biol., vol. 8, pp. 1247-1252, 1988.*
Wells, Biochemistry, 1990, vol. 29, pp. 8509-8517.*

Pettit et al, The Journal of Biological Chemistry, 1997, vol. 272, No. 4, pp. 2312-2318.*

Waldmann et al, IL-15 Receptors, 2000, pp. 1521-1528.*

Belmont H.J. et al. "Potent antitumor activity of a tumor-specific soluble TCR/IL-2 fusion protein." Clin Immunol. Oct. 2006;121(1):29-39.

Mortier E. et al. "Soluble interleukin-15 receptor alpha (IL-15R alpha)-sushi as a selective and potent agonist of IL-15 action through IL-15R beta/gamma. Hyperagonist IL1-15 x IL-15R alpha fusion proteins." J Biol Chem. Jan. 20, 2006;281(3):1612-9.

Mosquera L. et al. "In vitro and in vivo characterization of a novel antibody-like single-chain TCR human IgG1 fusion protein." J Immunol. Apr. 1, 2005;174(7):4381-8.

Wong LW et al. "Interleukin-15: Interleukin-15 receptor α scaffold for creation of multivalent targeted immune molecules." Protein Eng Des Sel. Dec. 21, 2010; 1-11.

Zhu X et al. "Novel human interleukin-15 agonists." J Immunol. Sep. 15, 2009;183(6):3598-607.

Zhu X. et al. "Visualization of p53(264-272)/HLA-A0201 complexes naturally presented on tumor cell surface by a multimeric soluble single-chain T cell receptor." J Immunol. Mar. 1, 2006;176(5):3223-32.

Extended European Search Report dated Feb. 2, 2011 issued for Application No. PCT/US2008/005918.

Database UniProt [Online], Sequence, retrieved from EBI, Database Accession No. 097687, XP002659759 (1999).

Database UniProt [Online], Sequence, retrieved from EBI, Database Accession No. Q8SPY0, XP002659760 (2002).

Database UniProt [Online], Sequence, retrieved from EBI, Database Accession No. Q6B416, XP002659761 (2004).

Chae et al., "Mutant IL-15 Protein exerting antagonistic effects on IL-15 triggered cell proliferation", Journal of the American Society of Nephrology, 7(9): 1654 (1996).

Bernard et al., "Identification of an interleukin-15alpha receptor-binding site on human interleukin-15", Journal of Biological Chemistry, The American Society of Biological Chemists, Inc., 279(23): 24313-24322 (2004).

European Search Report issued Sep. 22, 2011 and mailed on Oct. 12, 2011, for European Patent Application No. 11 178 091.2.

European Search Report issued Nov. 22, 2011 and mailed on Nov. 30, 2011, for European Patent Application No. 11 178 091.

* cited by examiner

Primary Examiner — Bridget E Bunner

Assistant Examiner — Fozia Hamud

(74) *Attorney, Agent, or Firm* — Edwards Wildman Palmer LLP; Peter F. Corless; Andrew W. Shyjan, Esq.

(57) **ABSTRACT**

The instant invention provides soluble fusion protein complexes and IL-15 variants that have therapeutic and diagnostic use, and methods for making the such proteins. The instant invention additionally provides methods of stimulating or suppressing immune responses in a mammal using the fusion protein complexes and IL-15 variants of the invention.

9 Claims, 43 Drawing Sheets




FIG. 1A




FIG. 1B




FIG. 2B

5' - ACCACCATGGAGACAGACACACTCCTGTTATGGGTACTGCTGCTCTGGGTTCACAGGTTCACCCGGTCAGT
CAGTGACGCAGCCCCGATGCTCGCGTCACTGTCTCTGAAGGAGCCCTCTCTGCAGCTGAGATGCAAGTATTC
CTACTCTGGGACACCTTATCTGTTCTGGTATGTCCAGTACCCGGCAGGGCTGCAGCTGCTCCTCAAG
TACTATTCAGGAGACCCAGTGGTTCAAAGGAGTGAATGGCTTCGAGGCTGAGTTCAGCAAGAGTAACTCTT
CCTTCCACCTGCGGAAAGCCTCTGTGCACTGGAGCGACTCTGCTGTGTACTTCTGTGTTTGAGCGAGGA
TAGCAACTATCAGTTGATCTGGGGCTCTGGGACCAAGCTAATTATAAGCCAGACACTAGTGGTGGCGGT
GGCAGCGCGGTGGTGGTTCCGGTGGCGGGGTCTGGCGGTGGCGGTCTCTCGAGCAATTCAAAAAGTCA
TTCAGACTCCAAGATATCTGGTGAAGGGCAAGGACAAAAGCAAGATGAGGTGTATCCCTGAAAAGGG
ACATCCAGTTGTATTCTGGTATCAACAAAATAAGAACAAATGAGTTTAAATTTTGATTAACTTTCAGAAAT
CAAGAA GTTCTTCAGCAAAATAGACATGACTGAAAAACGATTCTCTGCTGAGTGTCTTCAAACCTCACCTT
GCAGCCTAGAAATTCAGTCCTCTGAGGCAGGAGACTCAGCACTGTACCTCTGTGCCAGCAGTCTGTCAGG
GGCGGCACACAGAAGTTTCTTTGGTAAAGGAACCAAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTT
CCACCCGAGGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGT
GCCTGGCCACAGGCTTCTTCCCTGACCACTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGGTGCACAG
TGGGGTCAGCACGGACCCGAGCCCTCAAGGAGCAGCCCGCCCTCAATGACTCCAGATACTGCCCTGAGC
AGCCGCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCT
ACGGGCTCTCGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGCCGA
GGCCTGGGGTAGAGCAGACGTTAACAACTGGGTGAATGTAATAAGTGATTTGAAAAAAATTTGAAGATCTT
ATTCAATCTATGCATATTGATGCTACTTTATATACGGAAGTGAATGTTACCCCAAGTTGCAAGTAACAG
CAATGAAGTGCTTTCTCTTGGAGTTACAAGTTATTTCACTTGAGTCCGGAGATGCAAGTATTCATGATAC
AGTAGAAAAATCTGATCATCTAGCAAAACAAGTTTGTCTTTCTAATGGGAATGTAAACAGAAATCTGGATGC
AAAGAAATGTGAGGAACTGGAGGAAAAAATAATTAAAGAAATTTTGCAGAGTTTTGTACATATTGTCCAAA
TGTTTCATCAACACTTCTTGATAA -3'

FIG. 2C

```

metdtllllwvlllwpvgstgqsvtqpdarvtvsegaslqlrckysysgtpylfwyvqyprgglqlllkyys
< Signal peptide >< TCR Vα
gdpvvggvngfeae fsksnssfhlrkasvhwdsavvfcvlsedsnyqliwsggtklilikpdtsgggsgg
TCR Vα >< linker
ggsgggggsgggssskvigtprylvkqggqkakmrcipekghpvvfwyqqnknefkflinfqngavlq
linker >< TCR Vβ
gidmtekrfsaecpsnspcsleigsseagdsalylcasslsgggtevfqgktrltvvedlnkvfppevav
TCR Vβ >< TCR Cβ
fepseaeishtqkatlvclatgffpdhvelswvngkevhshqvstdppqlkeqpaldnsryclssrlrvsa
TCR Cβ
tfwgnprnhfrcgqvfyglsendewtqdrakpvtgi vsaaawgradvnnwvnnvisdlkkiedliqsmhida
TCR Cβ > < IL-15
tlytesdvhpckvtamkcflllelqvlslesgdasihdtvenliilannslssngnvtesgckeceeleek
IL-15
nikeflqsfvhi vqm fints
IL-15 >

```

FIG. 3A




FIG. 3B

5' -
ACCACCATGGAGACAGACACACTCCTGTTATGGGTACTGCTGCTGGGTCCAGGTTCCACCGGTCACTCAGTGACGACGCCCGATGC
TCGCGTCACTGTCTCTGAAGGAGCCTCTCTGCAGCTGAGATGCAAGTATTCCTACTCTGGGACACCTTATCTGTTCTGGTATGCCAGT
ACCCGGGCAGGGGCTGCAGCTGCTCCTCAAGTACTATTTCAGGAGACCCAGTGGTTCAGGAGTGAATGGCTTCGAGGCTGAGTTCAGC
AAGAGTAACTCTTCTTCCACCTCGGAAAGCCTCTGTGCACTGGAGCGACTCTGCTGTACTTCTGTGTTTTGAGCGAGGATAGCAA
CTATCAGTTGATCTGGGGCTCTGGGACCAAGCTAATTATAAAGCCAGACACTAGTGGTGGCGGTGGCAGCGCGGCTGGTGGTCCGGTG
GCGGCGGTTCTGGCGGTGGCGGTTCTCGAGCAATTCAAAAGTCATTCAGACTCCAAAGATATCTGGTGAAAGGCAAGGACAAAAAGCA
AAGATGAGGTGTATCCCTGAAAAGGGACATCCAGTTGTATTCTGGTATCAACAAAAATAAGAACAAATGAGTTTTAAATTTTGTAACTT
TCAGAAATCAAGAAGTTCTTCAGCAATAGACATGACTGAAAAACGATTCTCTGCTGAGTGTCTTCAAACTCACCTTCAGCCTAGAAA
TTCAGTCTCTGAGGAGGAGACTCAGCACTGTACCTCTGTGCCAGAGTCTGTCAAGGGGGGGCACAGAAAGTTTCTTTGGTAAAGGA
ACCAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTCCACCCGAGTCGCTGTGTTTGAAGCCATCAGAAGCAGAGATCTCCACAC
CCAAAAGGCCACACTGGTGTGCTGGCCACAGGCTTCTTCCCTGACACGCTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGTGCACA
GTGGGGTCAGCAGGACCCGAGCCCTCAAGGAGCAGCCGCCCTCAATGACTCCAGATCTGCCTGAGCAGCCGCTGAGGGTCTCG
GCCACCTTCTGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGATGACGAGTGGACCCAGGATAG
GGCCAAACCCGTCAACCAGATCGTCAGCGCCGAGGCTGGGTAGAGCAGACGTTAACGAGCCCCAAATCTTCTGACAAAACCTCACACAT
CTCCACCGTCTCCAACGCGTAACTGGGTGAATGTAATAAGTATTTGAAAAAATTGAAGATCTTATTCAATCTATGCATATTGATGCT
ACTTTATATACGGAAGTGATGTTCAACCCAGTTGCAAAAGTAAACAGCAATGAAGTGTCTCTTTGGAGTTACAAGTTATTTCACTTGA
GTCCGGAGATGCAAGTATTATGATACAGTAGAAAAATCTGATCATCTAGCAACAACAGTTTGTCTTCTAATGGGAATGTAACAGAAT
CTGGATGCAAGAATGTGAGGAACTGGAGGAAAAAATATAAGAAATTTTGCAGAGTTTGTGACATATTGTCCAAATGTTTCATCAAC
ACTTCTTGATAA -3'

```

metdtlllwlwlllwpbstgsgsvtqpdarvtvsegaslqlrckysysgtptylfwyvqyprqglqlllkyys
< Signal peptide >< TCR Vα
gdpvvqgvngfgeafsksnssfhlrkasvhwsdsavycvlsedsnyqliwsggtklilikpdtsgggsggg
TCR Vα >< linker
ggsgggsgggsgssnskvigtprylvkgggkakmrceipekghpvfwyqqknnefkflinfqngevlq
linker >< TCR Vβ
qidmtekrfsaecpsnspcsleiqsseaagdsalylcasslsgggtevfqkgtrltvvedlnkvfppevav
TCR Vβ >< TCR Cβ
fepseaeishtqkatlvclatgffpdhvelswvvnqkevhsqvstdbqpakecpalndsryclssrlrvsa
TCR Cβ
tfwgnprnhfrcgvqfyglsendewtqdrakpvtgiysaeawgradvnepkssdkthtspssptnwnvni
TCR Cβ > < linker > < IL-15
sdlkkiedliqsmhidatlytesdvhpckvtamkcflllelqvlesgdasihdtvenliilannslsn
IL-15
gnvtesgcceeeleeknikeflqsfvhiqvmfints
IL-15 >

```

FIG. 3C

FIG. 4A




FIG. 4B

5' - ACCACCATGGAGACAGACACACTCCTGTTATGGGTACTGCTGCTCTGGGTCCAGGTTCCACCGGTCACTCAGTGACGCAGCCCGATGCTCGGTCACCTGCTCTGAAGGAGCCTCTCTGCAGCTGAGATGCAAGTATTCCTACTCTGGGACACCTTATCTGTTCTGGTATGTCCAGTACCGCGGAGGGGCTGCAGCTGCTCCTCAAGTACTATTCAAGGAGACCCAGTGGTTCAAGGAGTGAATGGCTTCGAGGCTGAGTTCAGCAAGAGTAACCTCTTCTCCACCTGCGGAAAGCCTCTGTGCACCTGGAGCGACTCTGCTGTGTAATCTGTGTTTTGAGCGAGGATAGCAACTATCAGTTGATCTGGGCTCTGGGACCAAGCTAATTATAAAGCCAGACACTAGTGGTGGCGGTGGCAGCGGCTGGTGGTCCGGTGGCGGCTCTGGCGGTGGCGGTCCCTCGAGCAATTCAAAAAGTCAATTCAGACTCCAAGATATCTGGTGAAGGGCAAGGACAAAAGCAAAAGATGAGGTGATCCCTGAAAAGGGACATCCAGTTCTTCAGCAAAATAGACATGACTGAAAAACGATTCTCTGCTGAGTGTCCCTTCAAACTCACCTTGCAGCTAGAAATTCAAGTCCCTCTGAGGCAGGAGACTCAGCACTGTACCTCTGTGCCAGCAGTCTGTCAAGGGCGGCACAGAAAGTTTCTTTGGTAAAGGAACCAAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTCCACCCGAGTGGTGGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACAGGCTTCTTCCCTGACCACGTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGTGCACAGTGGGTCAAGCACGGACCCGACGCCCTCAAGGAGCAGCCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCAACCAGATCGTCAGCGCCGAGCCCTGGGTAGAGCAGACGTTAACATCACGTGCCCTCCCCCATGTCCGTGGAACACGCAGACATCTGGTCAAGAGCTACAGCTTGTACTCCAGGGAGCGGTACATTTGTAACCTCTGGTTTCAAGCGTAAAGCCGGCAGTCCAGCCCTGACGGAGTGGTGTGAACAAGGCCACGAATGTGCCCCACTGGACAACCCCCAGTCTCAAATGCAATTAGACCCCGAGCTTCATCTCCAGCTCAAAACAACACAGCGGCCACAACAGCAGCTATTGTCCCGGGCTCCAGCTGATGCCCTTCAAAATCACCTTCCACAGGAACCAACAGAGATAAGCAGTCATGAGTCCCTCCCACGGCACCCCTCTCAGACAAACAGCCAAAGAACTGGGAACTCACAGCATCCGCTCCCAACCAAGCCGCCAGGTGTGTATCCACAGGGCCACAGCGACACCACTTGATAA-3'

FIG. 4C

```

metdtllllwlllvpgstgqsvtqpdarvtvsegaslqlrckysysgtpylfwyvqyprgglqlllkyyys
< Signal peptide >< TCR Vα
gdpvvgvngfeae fsksnssfhlrkasvhwdsavvfcvlsedsnyqliwsgtgkllikpdtsgggsgg
TCR Vα
>< linker
ggsgggsgggssnskviqtprylvkgggkakmrckipekghpvvfwyqgnknefkflinfgngevlq
linker >< TCR Vβ
qidmtekrfsaecpsnspcsleigseagdsalylcasslsgggtvffgkgttrltvvedlnkvfppevay
TCR Vβ
>< TCR Cβ
fepseaei.shtgkat.lvclatgffpdhvel.swwvngkevhsgvstdpqpl.keqpal.ndsryc.lssr.lrvsa
TCR Cβ
tfwgnprnhfrcqvfylsendewtdrakovtdi.vsaewaqradvnitcoppmsvehadiwvksyslys
TCR Cβ
> < IL-15RaDE3
reryicnsqfkrkactssltecvltnkatnvahwtttpslkci.rpaasspsnnntaattaavpqsq.lmpsk
IL-15RaDE3
pstattei.ssheshqtspgttaknwelltasashcpbgvypqghsdt
IL-15RaDE3
>

```




FIG. 5B

5' - ACCACCATGGAGACAGACACACTCCTGTTATGGGTACTGCTGCTCTGGGTCCAGGTTCCACCCGGTCAGT
CAGTGACGCAGCCCGATGCTCGCGTCACTGTCTCTGAAGGAGCCCTCTCTGCAGCTGAGATGCAAGTATTC
CTACTCTGGGACACCTTAICTGTTCTGGTATGTCAGTACCCGGCAGGGCTGCAGCTGCTCCTCAAG
TACTATTCAAGAGACCCAGTGGTTCAAGAGTGAATGGCTTCGAGGCTGAGTTCAGCAAGAGTAACTCTT
CCTTCCACCTGCGGAAAGCCCTCTGTGCACCTGGAGCGACTCTGCTGTGACTTCTGTGTTTGTAGCGAGGA
TAGCAACTATCAGTTGATCTGGGCTCTGGGACCAAGCTAATTATAAGCCAGACACTAGTGGTGGCGGT
GGAGCGCGGTGGTGGTTCGGGTGGCGCGGTCTGGCGGTGGCGGTTCCTCGAGCAATTCAAAAGTCA
TTCAGACTCCAAAGATATCTGGTGAAGGGCAAGGACAAAAGCAAAAGATGAGTGTATCCCTGAAAAGGG
ACATCCAGTTGTATTCTGGTATCAACAAAATAAGAACAAATGAGTTTAAATTTTGTAACTTTCAGAAAT
CAAGAAGTTCTTCAGCAATAAGACATGACTGAAAACGATTCCTGCTGAGTGTCTTCAAACTCACCTT
GCAGCCTAGAAAATTCAGTCCCTCTGAGGCAGGAGACTCAGCACTGTACCTCTGTGCCAGCAGTCTGTCAAG
GGCGGCACAGAAAGTTTCTTTGGTAAAGGAACAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTT
CCACCCGAGGTCGCTGTGTTTGAAGCCATCAGAAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGT
GCCGTGCCACAGGCTTCTTCCCTGACCACGTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGGTGCACAG
TGGGGTCAGCACGGACCCGACGCCCTCAAGGAGCAGCCCGCCCTCAATGACTCCAGATACTGCCTGAGC
AGCCGCCCTGAGGTCCTGGCCACCTTCTTGCCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCT
ACGGGCTCTCGGAGAAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTACCCAGATCGTCAGCGCCGA
GGCTGGGGTAGACGACGTAAACATCACGTGCCCTCCCCCATGTCCGTGGAAACACGCAGACATCTGG
GTCAAGAGCTACAGCTTGTACTCCAGGGAGCGGTACATTTGTAACTCTGGTTTCAAGCGTAAAGCCGGCA
CGTCCAGCCTGACGGAGTGGTGTGAACAAGGCCACGAATGTGCCCCACTGGACAAACCCCAAGTCTCAA
ATGCATTAGA-3'

FIG. 5C

```

metdtllllwlllwpgstgsvtqpdarvtvsegaslqlrckysysgtpylfwyvqyprqglqlllkyys
< Signal peptide >< TCR Vα
gdpvvqgvngfeae fsksnssfhlrkasvhwdsavvfcvlsedsnyqlwsggtklilikpdtsgggsgg
TCR Vα >< linker
ggsgggsgggssnskviqtprylvkgggkakmrcipekghpvvfwyqqknnefkflinfqnqevlq
linker >< TCR Vβ
qidmtekrfsaecpsnspcsleigsseagdsalylcasslsgggtevfkgtrltvvedlnkvfppevav
TCR Vβ >< TCR Cβ
fepseaeishtgkatlvclatgffpdhvelswvngkevhsqvstdpgplkegpalndsryclssrlrvsa
TCR Cβ
tfwqnprrnhfrcqvqfyqlsendewtqdrakpvtqivsaewaqradvnitcpppmsvehadiwvksyslys
TCR Cβ > < IL-15RaSushi
Rervicnsgfkrkaqtssltecvlknatnvahwttpslkcir
IL-15RaSushi >

```




FIG. 6B


5' -
ACCACCATGGAGACAGACACTCCTGTTATGGGTACTGCTGCTCTGGGTCCAGGTTCCACCGGTCACTCAGTGACGACCCCGATGC
TCGCGTCACTGTCTCTGAAGGAGCCTCTCTGCAGCTGAGATGCAAGTATTCCTACTCTGGACACCTTATCTGTTCTGTATGTCCAGT
ACCCGGCAGGGTGCAGCTGCTCCTCAAGTACTATTCAAGAGACCCAGTGGTCAAGGAGTGAATGGCTTCGAGGCTGAGTTCAGC
AAGAGTAACTCTCCTCCACCTCGGAAAGCCTCTGTGCACTGGAGCGACTCTGCTGTACTTCTGTGTTTTCGAGCGAGATAGCAA
CTATCAGTTGATCTGGGCTCTGGGACCAAGCTAATTATAAAGCCAGACACTAGTGGTGGCGGTGCAGCGCGGTGGTGGTCCGGTG
GCGCGGTTCTGGCGGTGGCGGTTCTCTCGAGCAATTCAAAAGTCATTCAAGTCTCAAGATATCTGGTGAAGGGCAAGCAAAAAGCA
AAGATGAGGTGTATCCCTGAAAAGGGACATCCAGTTGTATTCTGGTATCAACAAAATAAGAACAAATGAGTTAAATTTTGTAACTT
TCAGAAATCAAGAACTTCTCAGCAAAATAGACATGACTGAAAAACGATTCTCTGCTGAGTGTCTTCAAACTCACCTTCAGCCTAGAAA
TTTCAGTCTCTGAGGAGGAGACTCAGCACGTACCTCTGTGCCAGCAGTCTGTCAAGGGGGGCACAGAAATTTCTTTGGTAAAGGA
ACCAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTCCACCCGAGGTGGTGTGTTGAGCCATCAGAAAGCAGACATCTCCACAC
CCAAAAGGCCACACTGGTGTGCTGGCCACAGGCTTCTTCCCTGACCACGTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGGTGCACA
GTGGGTTCAGCACGACCCGAGCCCCCTCAAGGAGCAGCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCCTGAGGGTCTCG
GCCACCTTCTGGCAGAAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTCTACGGCTCTCGGAGATGACGAGTGGACCCAGGATAG
GGCCAAACCCGTACCCAGATCGTCAGCGCCGAGGCTGGGTAGAGCAGACGTTAACGAGCCCAATCTCTGACAAAACCTACACAT
CTCCACCGTCTCCAACGGTATCACGTGCCCTCCCCCATGTCCGTGGAACACCGCAGACATCTGGTCAAGAGCTACAGTTGTACTCC
AGGAGCGGTACATTTGTAACTCTGGTTTCAAGCGTAAAGCCGGCAGCTCCAGCCTGACGGAGTGCGTGTTGAACAAGGCCACGAAATGT
CGCCCACTGGACAACCCCGAGTCTCAAAATGCATTAGATGATAA -3'


FIG. 6C

```

metdtlllwlwllwvpstgsgsvtqpdarvtvsegaslqrlckysyqtpylfwyvyprqglqlllkyys
< Signal peptide >< TCR Vα
gdpvvggvgfgeafsksnssfhlrkasvhwdsavvfcvlsedsnyqliwsgtkliikpdtsgggsgg
TCR Vα >< linker
ggsgggsgggssnskvitprylvkgggkakmrcipekghpvvfwyqgnknefkflinfqnqevlq
linker >< TCR Vβ
qidmtekrfsaecpsnspcsleiqsseagdsalylcasslsgggevffgkgtrltvvedlnkvfppevay
TCR Vβ >< TCR Cβ
fepseaeishtqkatlvclatgffpdhvelswvngkevhsqvstdpgplkecpalndsryclssrlrvsa
TCR Cβ
tfwanprnhfrcavqfyqlsendewtdadrakpvtqiysaeawgradvnepkssdkthtsppsptritcbbb
TCR Cβ > < linker > <
msvehadiwksyslvsrvicnsgfkrkagtssltecvlakatnvahwtttpslkcir
IL-15RaSushi >

```












FIG. 9A








FIG. 11A




FIG. 11B




FIG. 11C




FIG. 12A




FIG. 12B





FIG. 14A

NWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSPCKVTAMKCFLLLELQVLSLESGDASI
HDTVENLIILANNLSLSSNGNVTESGCKECEELKEEKNIKEFLQSFVHIVQMFINTS






FIG. 14C-1

c264scTCR-hmt-IL15D8A gene

5' -
 ACCACCATGGAGACAGACACACTCCCTGTTATGGGTAAGTCTGGGTCCAGGTTCCACCGGTCACTCAGTGACGCAGCCCCGATGC
 TCGCGTCACTGTCTCTGAAGGAGCCCTCTCGAGCTGAGATGCAAGTATTCCTACTCTGGGACACCTTATCTGTCTGGTATGTCCAGT
 ACCCGCGGCAGGGGCTGCAGCTGCTCCTCAAGTACTATTCAGGAGACCCAGTGGTCAAGGACTGAATGGCTTCGAGGCTGAGTTCAGC
 AAGATAACTCTTCCCTCCACCTGGGAAAGCCTCTGTGCACCTGGAGCGACTCTGTGTACTTCTGTGTTTGGAGGAGGATAGCAA
 CTATCAGTTGATCTGGGGCTCTGGGACCAAGCTAATAAAGCCAGACACTAGTGGTGGCGGTGGCAGCGCGGTGGTTCGGGTG
 GCGGCGGTTCTGGCGGTGGCGGTTCCTCGAGCAATTCAAAAGTCAATTCAGACTCCAAAGATACTGGTGAAGGGCAAGGACAAAAAGCA
 AAGATGAGGTGTATCCCTGAAAAGGACATCCAGTTGTATTCGTGTATCAACAAAAAAGCAAACTGAGTTTAAATTTTGTAACTT
 TCAGAAACAAGAGTTCTTCAGCAAAATAGACATGACTGAAAAACGATTCTCTGCTGAGTGTCTTCAAACTCACCTTGCAGCCTAGAAA
 TTCAGTCTCTGAGGCAGGAGACTCAGCACTGTACCTCTGTGCCAGCAGTCTGTCAGGGGGGGCACAGAAAGTTTCTTTGGTAAAGGA
 ACCAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTCCACCCGAGGTCGCTGTCTTTGAGCCATCAGAAGCAGAGATCTCCACAC
 CCAAAAGGCCACACTGGTGTGCCTGGCCACAGGCTTCTTCCCTGACCACTGGAGCTGAGCTGGTGGTGAATGGGAAGGAGGTGCACA
 GTGGGTGAGCACGACCCGACGCCCTCAAGGAGCAGCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCGCCCTGAGGGTCTCG
 GCCACCTTCTGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGTACGAGTGCACCCAGGATAG
 GGCCAAACCCGTCAACCAGATCGTCAGCGCCGAGGCCCTGGGTAGAGCAGACGTTAACGAGCCCAATCTTCTGACAAAACTCACACAT
 CTCACCCGTCTCCAACCGTAACTGGGTGAATGTAATAAGT**GCT**TTTGAAGAAAAATGAAGATCTTATCAATCTATGCAATATGATGCT
 ACTTTATATACGGAAGTGATGTTCACCCCGAGTTGCAAAAGTAACAGCAATGAAGTCTTCTCTGGAGTTACAAGTTATTCACCTTGA
 GTCCGGAGATGCAAGTATTCATGATACAGTAGAAAAATCTGATCATCTAGCAAAACACAGTTTGTCTTCTAATGGGAATGTAAACAGAT
 CTGGATGCAAGAAATGTGAGGAACCTGGAGGAAAAAATATAAGAAATTTTTCAGAGATTTTGTACATAATTTTCCAAATGTTCATCAAC
 ACTTCTTGATAA -3'

FIG. 14C-2

c264scTCR-hmt-IL15D8N gene

5' -

ACCACCATGGAGACAGACACACTCCTGTTATGGGTACTGCTGCTCTGGGTCCAGGTTCCACCCGGTCAGTCAGTGACGCAGCCCGATGC
TCGCGTCACGTGCTCTGAAGGAGCCCTCTCTGCAGCTGAGATGCAAGTATTCCTACTCTGGGACACCTTATCTGTTCTGGTATGTCCAGT
ACCCGGCGCAGGGGCTGCAGTGTCTCTCAAGTACTATTCAAGAGACCCAGTGGTTCAGAGAGTGAATGGCTTCGAGGCTGAGTTCAGC
AAGAGTAACTCTTCCCTCCACCTGCGGAAAGCCTCTGTGCACCTGGAGCGACTCTGCTGTGTAATCTCTGTGTTTTCGCGAGGATAGCAA
CTATCAGTTGATCTGGGGCTCTGGGACCAAGCTAATTATAAAGCCAGACACTAGTGGTGGCGGTGGCAGCGGGGTGGTGGTCCCGTG
GGGCGGTTCTGGCGGTGGCGGTTCCCTCGAGCAATTCAAAAGTCATTTCAGACTCCAGATATCTGGTGAAGGGCAAGGACAAAAGCA
AAGATGAGGTGTATCCCTGAAAAGGGACATCCAGTTGTATTCTGTTATCAACAAAAAAGAACAAATGAGTTTAAATTTTTCATTAACTT
TCAGAAATCAAGAAAGTTCTTCAGCAAAATAGACATGACTGAAAAACGATTCTCTGTGAGTGTCTTCAAACCTCACCTTCAGCCCTAGAAA
TTCAGTCCCTCTGAGGCAGGAGACTCAGCACTGTACCTCTGTGCCAGCAGTCTGTACGGGGCGGCACAGAAAGTTTCTTTGGTAAAGGA
ACCAGACTCACAGTTGTAGAGGACCTGAACAAGGTGTTCCCAACCCGAGGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCACAC
CCAAAAGGCCACACTGGTGTGCTGGCCACAGGCTTCTTCCCTGACCACGTGAGCTGGTGGTGAATGGGAAGGAGGTGCACA
GTGGGTGAGCACGACCCGACGCCCTCAAGGAGCAGCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCTGAGGGTCTCG
GCCACCTTCTGGCAGAAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAAATGACGAGTGCACCCAGGATAG
GGCCAAACCCGTCAACCCAGATCGTCAGCGCCGAGGCCCTGGGTAGAGCAGACGTTAACGAGCCCAAAATCTTCTGACAAAACCTCACACAT
CTCCACCGTCTCCAACGCGTAACTGGGTGAATGTAATAAGT**AAT**TTGAAAAAAATGAAGATCTTATTCATCTATGCATATTGATGCT
ACTTTTATATACGGAAAGTGATGTTTCAACCCAGTTGCAAAAGTAACAGCAATGAAGTGCTTCTCTTGGAGTTACAAGTTATTTCACTTGA
GTCCGGAGATGCAAGTATTCATGATACAGTAGAAAATCTGATCATCTAGCAAAACACAGTTTGTCTTCTAAATGGGAATGTAACAGAAAT
CTGGATGCAAAAGAAATCTGAGGAACTGGAGGAAAAAATATTAAGAAATTTTTCAGAGTTTGTACATATTGTCCAAATGTTTCATCAAC
ACTTCTTGATAA -3'

FIG. 14D-1
FIG. 14D-2

FIG. 14D

FIG. 14D-1

c264scTCR-hmt-IL15D8A protein

```

metdtllllwvlllwpvgstgqsvtqpdarvtvsegaslqlrckysysgtpylfwyvqypragqlllkyyys
< Signal peptide >< TCR Vα
gdpvvqgvngfeae fsksnssfhrlrkasvhwdsavycvlsedsnyqliwsgtkllikpdtsgggsgg
TCR Vα >< linker
ggsgggsgggsgssnskvigtprylvkgggkakmrcipekghpvfwyqgnknefkflinfqngvlg
linker >< TCR Vβ
gidmtekrfsaecpsnspcsleigsseagdsalylcasslsgggtevfqkgtrltvvedlnkvfppevav
TCR Vβ >< TCR Cβ
fepseaeishtqkatlvclatgffpdhvelswwvngkevhsgvstdpqpakegpaldnsryclssrlrvsa
TCR Cβ
tfwqnpnrnhfrcaqvqfvglsendewtdqdrakpvtgqvsaawgradvnepkssdkthtsppsptnnwvni
TCR Cβ > < linker > <
salkkiedliqsmhidatlytesdvhpscvktamcfllelqvlesgdasihdtvenliilannslssn
IL-15 D8A
gnvtesgckeceeleeknikeflqsfvhi vqm fints
IL-15 D8A >

```

FIG. 14D-2


c264scTCR-hmt-IL15D8N protein

```

metdtlllwlwlllwpbstgqsvtqpdarvtvsegasqlrckysysgtpylfwyvqyprgqlqlllkyys
< Signal peptide >< TCR Vα
gdpvvgvngfeae fsksnssfhlrkasvhwdsavvyfcvlsedsnyqliwsgtgkllikpdtsgggsgg
TCR Vα >< linker
ggsgggsgggsgssnskviqtprylvkqggqkakmrcipekghpvvfwyqanknnefkflinfqnqevlq
linker >< TCR Vβ
gidmtekrfsaecpsnspcsleiqsseagdsalylcasslsgggtevffgkgtrltvvedlnkvfppevav
TCR Vβ >< TCR Cβ
fepseaei.shtgkatlvcLatgfpdhvel.swwvngckevhsgvstdpgplkegpalndsryclssrlrvsa
TCR Cβ
tfwqnpnrnhfrcavqfyqlsendewtqdrakpvtqivsaeeawgradvnepkssdkthtspspstrnwvni
TCR Cβ > < linker > <
s*llkkiedliqsmhidatlytesdvhpckvtamkcflllelqvlsesgdasihdtvenliilannslsn
IL-15 D8N
gnvtesgckeceeleeknikelfqsfvhivqmfmts
IL-15 D8N >

```

FIG. 15






FIG. 16

FIG. 17A




FIG. 17B




FIG. 17C




FIG. 18A





FIG. 18B







FIG. 19B






FIG. 19C




FIG. 21

Table 1

Mutants	Position	8	61	65	72	108	IL15R β Creceptor binding	IL15R α binding	Proliferation activity
	WT aa	D	D	N	N	Q	+	+	+
1	8	N					-	+	-
2	8	A					-	+	-
3	61		A				-	+	-
4	65			D			-	+	-
5	65			A			-	+	-
6	72				D		3+	+	3+
8	72				R		-	+	-
9	108					A	-	+	-
10	8+65	N		A			-	+	-
11	8+108	A				A	-	+	-
12	8+65	S		R			-	+	-

NWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKVTAMKCFLLLELQVISLES GDASI
HDTVENLIILANNSLSSNGNVTESGCKECEELEEKNIKEFLQSFVHIVQMFINTS

FIG. 22A

AACTGGGTGAATGTAATAAGTGATTTGAAAAAAATGAAGATCTTATTCAATCTATGCA
TATTGATGCTACTTTATATACGGAAAGTGATGTTCACCCCAGTTGCAAAGTAACAGCAA
TGAAGTGCTTTCCTTGGAGTTACAAGTTATTCACTTGAGTCCGGAGATGCAAGTATT
CATGATACAGTAGAAAATCTGATCATCCTAGCAAACAACAGTTTGTCTTCTAATGGGAA
TGTAACAGAAATCTGGATGCAAGAATGTGAGGAACTGGAGGAAAAAATAATTAAAGAAAT
TTTTCAGAGTTTGTACATATTTGTCCAAATGTTTCATCAACACTTCTTGA

FIG. 22B

1

FUSION MOLECULES AND IL-15 VARIANTS**RELATED APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 12/151,980, filed May 9, 2008, which claims the benefit of U.S. Provisional Application Ser. No. 60/928,900, filed May 11, 2007. The entire contents of each of the aforementioned applications is hereby incorporated herein by reference.

GOVERNMENT SUPPORT

Research supporting this application was carried out by the United States of America as represented by the Secretary, Department of Health and Human Services.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Oct. 26, 2010, is named 68295.txt and is 54,760 bytes in size.

BACKGROUND OF THE INVENTION

T Cell Receptors (TCR) are primary effectors of the immune system that have unique advantages as a platform for developing therapeutics. While antibody therapeutics are limited to recognition of pathogens in the blood and extracellular spaces or to protein targets on the cell surface, T cell receptors can recognize antigens displayed with MHC molecules on the surfaces of cells (including antigens derived from intracellular proteins). Depending on the subtype of T cells that recognize displayed antigen and become activated, T cell receptors and T cells harboring T cell receptors can participate in controlling various immune responses. For instance, T cells are involved in regulation of the humoral immune response through induction of differentiation of B cells into antibody producing cells. In addition, activated T cells act to initiate cell-mediated immune responses. Thus, T cell receptors can recognize additional targets not available to antibodies.

T-cells are a subgroup of cells which together with other immune cell types (polymorphonuclear, eosinophils, basophils, mast cells, B-, NK cells) constitute the cellular component of the immune system. Under physiological conditions T-cells function in immune surveillance and in the elimination of foreign antigen. However, under pathological conditions there is compelling evidence that T-cells play a major role in the causation and propagation of disease. In these disorders, breakdown of T-cell immunological tolerance, either central or peripheral is a fundamental process in the causation of autoimmune disease.

The TCR is believed to play an important role in the development and function of the immune system. For example, the TCR has been reported to mediate cell killing, increase B cell proliferation, and impact the development and severity of various disorders including cancer, allergies, viral infections and autoimmune disorders.

It thus would be desirable to provide novel targeting agents based on T cell receptors, as well as methods for producing and using such agents for therapeutic and diagnostic settings. Accordingly, it would be particularly desirable to provide such molecules that would have certain advantages in comparison to prior art complexes based on antibody targeting.

2

Moreover, it is desirable to use the TCR to target various effector molecules to the disease site where they can provide therapeutic benefit without the side effects associated with system non-targeted activity. One such is IL-15, a member of the four alpha-helix bundle family of lymphokines. IL-15 plays a multifaceted role in development and control of the immune system. More specifically, IL-15 influences the function, development, survival, and proliferation of CD8+ T cells, NK cells, killer T cells, B cells, intestinal intraepithelial lymphocytes (IEL) and antigen-presenting cells (APC). It has been demonstrated that both IL-15^{-/-}, and IL-15Ra^{-/-} transgenic mice lack peripheral NK and killer T cell populations, certain IEL subsets, and most memory phenotype CD8+ T cells, suggesting IL-15 plays role in the development, proliferation or/and survival of these cell types. The IL-15 receptor (R) consists of three polypeptides, the type-specific IL-15R alpha ("IL-15Rα" or "IL-15Ra"), the IL-2/IL-15Rbeta ("IL-2Rβ" or "IL-15Rb"), and the common gamma chain ("γC," or "gC" which is shared by multiple cytokine receptors).

IL-15 signaling can occur through the heterotrimeric complex of IL-15Rα, IL-2Rβ and γC; through the heterodimeric complex of IL-2Rβ and γC. A novel mechanism of IL-15 action is that of transpresentation in which IL-15 and IL-15Rα are coordinately expressed by antigen-presenting cells (monocytes and dendritic cells), and IL-15 bound to IL-15Rα is presented in trans to neighboring NK or CD8 T cells expressing only the IL-15RβγC receptor. As a co-stimulatory event occurring at the immunological synapse, IL-15 transpresentation now appears to be a dominant mechanism for IL-15 action in vivo and appears to play a major role in tumor immunosurveillance (Waldmann, T A, 2006, Nature Rev. Immunol. 6:595-601). Soluble IL-2Rα subunits, inducing isoforms containing a deletion of exon3 and the so-called "sushi" domain at the N terminus, have been shown to bear most of the structural elements responsible for cytokine binding. Whereas IL-2Rα alone is a low affinity receptor for IL-2 (Kd ~10 nM), IL-15Rα binds IL-15 with high affinity (Kd ~100 pM). Thus soluble IL-2Rα and IL-15 are able to form stable heterodimeric complexes in solution and these complexes are capable of modulating (i.e. either stimulating or blocking) immune responses via the intermediate or high affinity IL-15R complex (Mortier et al. 2006. J Biol Chem 281: 1612-1619; Stoklasek et al. 2006. J Immunol 177: 6072-6080; Rubinstein et al. 2006. Proc Natl Acad Sci USA 103: 9166-9171).

Given the known effects of IL-15 on the immune system, a number of IL-15-based approaches have been explored to manipulate the immune system for the hosts benefit. While IL-15 administration has been employed to bolster immune responses or augment immune system reconstitution, blockade of IL-15 activity can inhibit autoimmune and other undesirable immune responses (Waldmann, T A, 2006, Nature Rev. Immunol. 6:595-601). In fact, one of the limitations with systemic IL-15 treatment is the possible induction of autoimmune disease. Other limitations include the difficulty in produce this cytokine in standard mammalian cell expression systems as well as its very short half-life in vivo. Therefore, there is a need to generate a suitable immunostimulatory therapeutic form of IL-15 that displays a longer in vivo half-life, increased activity binding to immune cells, or enhanced bioactivity. Additionally there is a need for effective IL-15 antagonists. Ideally it would be desirable that such molecules be selectively targeted to the disease site to avoid unwanted systemic toxicities and provide a more effective therapeutic benefit.

SUMMARY OF THE INVENTION

The instant invention provides a number of IL-15 variants and soluble fusion complexes that have therapeutic use and

3

methods for making such proteins. The instant invention provides methods for killing target cells using the soluble fusion complexes of the invention. The IL-15 variants and soluble complexes described herein have potential therapeutic utility.

Accordingly, in one aspect, the invention provides a soluble fusion protein complex comprising at least two soluble fusion proteins, wherein the first fusion protein comprises a first biologically active polypeptide covalently linked to interleukin-15 (IL-15) or functional fragment thereof, and the second fusion protein comprises a second biologically active polypeptide covalently linked to soluble interleukin-15 receptor alpha (IL-15Ra) polypeptide or functional fragment thereof, wherein IL-15 domain of a first fusion protein binds to the soluble IL-15Ra domain of the second fusion protein to form a soluble fusion protein complex.

In one embodiment, one of the first and second biologically active polypeptides comprises a first soluble T-cell receptor (TCR) or functional fragment thereof. In another embodiment, an other of the biologically active polypeptides comprises the first soluble TCR or functional fragment thereof, thereby creating a multivalent TCR fusion protein complex with increased binding activity. In a further embodiment, the other biologically active polypeptide comprises a second soluble TCR or functional fragment thereof, different than the first soluble TCR.

In another embodiment of the aspect, the TCR is specific for recognition of a particular antigen.

In a further embodiment of the aspect, the TCR is a heterodimer comprising a and b chain TCR.

In still another embodiment of the aspect, the TCR comprises a single chain TCR polypeptide. In a further embodiment, the single chain TCR comprises a TCR V- α chain covalently linked to a TCR V- β chain by a peptide linker sequence. In another further embodiment, the single chain TCR further comprises a soluble TCR C β chain fragment covalently linked to a TCR V- β chain.

In another embodiment, the single chain TCR further comprises a soluble TCR C α chain fragment covalently linked to a TCR V- α chain.

In a further embodiment, one or both of the first and second biologically active polypeptides comprises an antibody or functional fragment thereof.

In still another embodiment, the antibody is specific for recognition of a particular antigen. In a further embodiment, the antibody is a single-chain antibody or single-chain Fv. In another particular embodiment, the single-chain antibody comprises an immunoglobulin light chain variable domain covalently linked to immunoglobulin heavy chain variable domain by polypeptide linker sequence.

In one embodiment of the above described aspects, the first biologically active polypeptide is covalently linked to IL-15 (or functional fragment thereof) by polypeptide linker sequence.

In another embodiment of the above described aspects, the second biologically active polypeptide is covalently linked to IL-15Ra polypeptide (or functional fragment thereof) by polypeptide linker sequence.

In another embodiment, the antigen for the TCR domain comprises peptide antigen presented in an MHC or HLA molecule. In a further embodiment, the peptide antigen is derived from a tumor associated polypeptide or virus encoded polypeptide.

In another embodiment, the antigen for the antibody domain comprises a cell surface receptor or ligand.

In a further embodiment, the antigen comprises a CD antigen, cytokine or chemokine receptor or ligand, growth factor receptor or ligand, tissue factor, cell adhesion molecule,

4

MHC/MHC-like molecules, FC receptor, Toll-like receptor, NK receptor, TCR, BCR, positive/negative co-stimulatory receptor or ligand, death receptor or ligand, tumor associated antigen, or virus encoded antigen.

In another embodiment of the above described aspects, the IL-15Ra polypeptide comprises the extracellular domain of the IL-15 receptor alpha capable for binding IL-15.

In another embodiment of the above described aspects, the IL-15Ra polypeptide comprise either the IL-15a sushi domain (Wei et al. *Journal of Immunology*, 2001, 167: 277-282) or the IL-15a Δ E3 domain (Anderson et al. 1995. *J. Biol. Chem.* 270:29862-29869, Dubois et al. 1999. *J. Biol. Chem.* 274:26978-26984).

In another aspect, the invention provides for an IL-15 variant (also referred to herein as IL-15 mutant) that has a different amino acid sequence that the native (or wild type) IL-15 protein and that binds the IL-15Ra polypeptide and functions as an IL-15 agonist or antagonist. Embodiments of the invention provide an IL-15 variant as a non-fusion protein or as a soluble fusion protein comprising a biologically active polypeptide described above, wherein the IL-15 variant is used in place of the IL-15 domain.

In one embodiment of the above described aspects, the invention features a nucleic acid sequence encoding the first fusion protein of any of the aspects or embodiments as described herein.

In one embodiment of the above described aspects, the invention features a nucleic acid sequence encoding the second fusion protein of any of the aspects or embodiments as described herein.

In one embodiment of the above described aspects, the invention features a nucleic acid sequence encoding the IL-15 variant of any of the aspects or embodiments as described herein.

In a one embodiment, the nucleic acid sequence further comprises a promoter, translation initiation signal, and leader sequence operably linked to the sequence encoding the fusion protein or IL-15 variant. In another embodiment, any of the nucleic acid sequences as described above are contained in a DNA vector.

In another aspect, the invention features a method for making a soluble fusion protein complex of the above-described aspects, the method comprising introducing into a first host cell a DNA vector of the above-described aspects and embodiments that encodes the first fusion protein, culturing the first host cell in media under conditions sufficient to express the first fusion protein in the cell or the media, purifying the first fusion protein from the host cells or media, introducing into a second host cell a DNA vector of the above-described aspects and embodiments encoding the second fusion protein, culturing the second host cell in media under conditions sufficient to express the second fusion protein in the cell or the media, and purifying the second fusion protein from the host cells or media, and mixing the first and second fusion protein under conditions sufficient to allow binding between IL-15 domain of a first fusion protein and the soluble IL-15Ra domain of a second fusion protein to form the soluble fusion protein complex.

In another aspect, the invention features a method for making a soluble fusion protein complex of the above-described aspects, the method comprising introducing into a host cell a DNA vector of the above-described aspects and embodiments, encoding the first fusion protein and a DNA vector as described in the above-described aspects and embodiments, encoding the second fusion protein, culturing the host cell in media under conditions sufficient to express the fusion proteins in the cell or the media and allow association between

5

IL-15 domain of a first fusion protein and the soluble IL-15Ra domain of a second fusion protein to form the soluble fusion protein complex, purifying the soluble fusion protein complex from the host cells or media.

In still another aspect, the invention features a method for making a soluble fusion protein complex of the above-described aspects, the method comprising introducing into a host cell a DNA vector encoding the first and second fusion proteins, culturing the host cell in media under conditions sufficient to express the fusion proteins in the cell or the media and allow association between IL-15 domain of a first fusion protein and the soluble IL-15Ra domain of a second fusion protein to form the soluble fusion protein complex, purifying the soluble fusion protein complex from the host cells or media.

In still other aspects of the above described methods, the DNA vector encoding the IL-15 variant is used in place of the DNA vector encoding the first fusion protein to generate a host cell capable of expressing the IL-15 variant and the IL-15 variant is allowed associate with the IL-15Ra domain of a second fusion protein to form a soluble fusion protein complex.

In another aspect, the invention features a method for making an IL-15 variant of the above-described aspects, the method comprising introducing into a host cell a DNA vector of the above-described aspects and embodiments that encodes an IL-15 variant, culturing the host cell in media under conditions sufficient to express the IL-15 variant in the cell or the media, purifying the an IL-15 variant from the host cells or media.

In another aspect, the invention features a method for killing a target cell, the method comprising contacting a plurality of cells with a soluble fusion protein complex or IL-15 variant of any of the above-described aspects or embodiments, wherein the plurality of cells further comprises immune cells bearing the IL-15R chains recognized by the IL-15 domain of the above-described aspects and the target cells bearing an antigen recognized by at least one of the biologically active polypeptides of the above-described aspects, forming a specific binding complex (bridge) between the antigen on the target cells and the IL-15R chains on the immune cells sufficient to bind and activate the immune cells, and killing the target cells by the bound activated immune cells.

In one embodiment of the method, the target cells are tumor cells or virally infected cells.

In another embodiment of the method, the biologically active polypeptide comprises a TCR.

In still another embodiment of the method, the antigen on the target cells comprises a tumor or virally encoded peptide antigen presented in an MHC or HLA molecule and recognized by the TCR.

In a further embodiment of the method, the immune cells are T-cells, LAK cells or NK cells.

In another aspect, the invention features a method for preventing or treating disease in a patient in which the diseased cells express a disease associated antigen, the method comprising administering to the patient a soluble fusion protein complex or IL-15 variant of any of the above-described aspects or embodiments, comprising a biologically active polypeptide recognizing a disease-associated antigen forming a specific binding complex (bridge) between antigen-expressing diseased cells and IL-15R-expressing immune cells sufficient to localize the immune cells, and damaging or killing the disease cells sufficient to prevent or treat the disease in the patient.

In one aspect, the invention features a method for preventing or treating disease in a patient in which the diseased cells

6

express a disease associated antigen, the method comprising mixing immune cells bearing the IL-15R chains with a soluble fusion protein complex comprising a biologically active polypeptide recognizing a disease-associated antigen, administering to the patient the immune cell-fusion protein complex mixture, forming a specific binding complex (bridge) between antigen-expressing diseased cells and IL-15R-expressing immune cells sufficient to localize the immune cells; and damaging or killing the disease cells sufficient to prevent or treat the disease in the patient.

In another aspect, the invention features a method for preventing or treating an disease in a patient in which the patient's cells express a disease associated antigen, the method comprising administering to the patient a soluble fusion protein complex or IL-15 variant of any of the above-described aspects or embodiments, comprising a biologically active polypeptide recognizing a disease-associated antigen on the patient's cells, localizing soluble fusion protein complex or IL-15 variant on the patient's cells wherein the IL-15 domain of the soluble fusion protein complex or IL-15 variant binds immune cells bearing the IL-15R chains and suppressing the immune response of the immune cells.

In one embodiment of the method, the disease is cancer or viral infection.

In another embodiment of the method, the disease is an immune disorder, autoimmune disease or inflammatory disorder.

In another embodiment of the method, the disease associated antigen is a peptide/MHC complex.

In another embodiment, the invention features a method of stimulating immune responses in a mammal comprising administering to the mammal an effective amount of the soluble fusion protein complex or IL-15 variant of any of the above-described aspects and embodiments.

In another embodiment, the invention features a method of suppressing immune responses in a mammal comprising administering to the mammal an effective amount of the soluble fusion protein complex or IL-15 variant of any of the above-described aspects and embodiments.

DESCRIPTION OF THE DRAWINGS

FIGS. 1(A & B) are schematic drawings. (A) is a schematic depicting an example of a fusion protein complex containing single chain TCR polypeptides. (B) is a schematic depicting representative fusion protein constructs comprising the fusion protein complex (SEQ ID NO: 3).

FIG. 2 (A-C) consists of three panels. (A) depicts a map of pNEF38-c264scTCR/huIL15 expression vector. (B) shows the sequence of c264scTCR/huIL15 fusion gene (SEQ ID NO: 4) and (C) shows the sequence of c264scTCR/huIL15 fusion protein (SEQ ID NO: 5), including the leader sequence.

FIG. 3 (A-C) consists of three panels. (A) depicts a map of pNEF38-c264scTCR-hinge-huIL15 expression vector. (B) shows the sequence of c264scTCR-hinge-huIL15 fusion gene (SEQ ID NO: 6) and (C) shows the sequence of c264scTCR-hinge-huIL15 fusion protein (SEQ ID NO: 7), including the leader sequence.

FIGS. 4 (A-C) consists of three panels. (A) depicts a map of pNEF38-c264scTCR/huIL15RaDE3 expression vector. (B) shows the sequence of c264scTCR/huIL15Ra Δ E3 fusion gene (SEQ ID NO: 8) and (C) shows the sequence of c264scTCR/huIL15Ra Δ E3 fusion protein (SEQ ID NO: 9), including the leader sequence.

FIG. 5 (A-C) consists of three panels. (A) depicts a map of the pNEF38-c264scTCR/huIL15RaSushi expression vector.

7

(B) shows the sequence of c264scTCR/huIL15R α Sushi fusion gene (SEQ ID NO: 10) and (C) shows the sequence of c264scTCR/huIL15R α Sushi fusion protein (SEQ ID NO: 11), including the leader sequence.

FIG. 6 (A-C) consists of three panels. (A) depicts the pNEF38-c264scTCR-hinge-huIL15R α Sushi expression vector. (B) shows the sequence of c264scTCR-hinge-huIL15R α Sushi fusion gene (SEQ ID NO: 12) and (C) shows the sequence of c264scTCR-hinge-huIL15R α Sushi fusion protein (SEQ ID NO: 13), including the leader sequence.

FIG. 7 is a map of pSun-c264scTCRIL15/c264scTCRIL15R α Sushi expression vector.

FIG. 8 is a map of pSun-c264scTCRIL15/c264scTCRIL15R α DE3 expression vector.

FIGS. 9(A & B) set forth characterization of transfected cells expressing TCR/IL15R α fusion protein. (A) is two graphs showing flow cytometric analysis of fusion protein expressing cells. (B) is a graph showing TCR-based ELISA results for fusion protein production.

FIGS. 10(A & B) shows analysis of TCR/IL15 and TCR/IL15R α fusion proteins by reducing SDS PAGE. (A) shows cell culture supernatants containing c264scTCR/huIL15 or c264scTCR/huIL15R α Sushi. (B) shows cell culture supernatants containing c264scTCR/huIL15 or c264scTCR/huIL15R α DE3.

FIGS. 11(A-C) shows analysis of TCR/IL15, TCR/IL15R α and fusion protein complexes by size exclusion chromatography. (A) is a graph showing the SEC chromatography profile of c264scTCR/huIL15. (B) is a graph showing the SEC chromatography profile of c264scTCR/huIL15R α Sushi. (C) is a graph showing the SEC chromatography profile of c264scTCR/huIL15+c264scTCR/huIL15R α Sushi fusion protein complex.

FIGS. 12(A & B) is an analysis of TCR/IL15R α and fusion protein complexes by size exclusion chromatography. (A) is a graph illustrating the SEC chromatography profile of c264scTCR/huIL15R α DE3. (B) is a graph illustrating the SEC chromatography profile of c264scTCR/huIL15+c264scTCR/huIL15R α DE3 fusion protein complex.

FIG. 13 is a graph showing the binding of TCR/IL15, TCR/IL15R α and fusion protein complexes to peptide/MHC complexes displayed on cells, as determined by flow cytometry.

FIG. 14 (A-D) consists of four panels. (A) shows the sequence of mature human IL15 protein (SEQ ID NO:1) and the blue underlined residues are substituted in the IL-15 variants as showed in Table 1. (B) depicts the pNEF38-c264scTCR-hinge-huIL15D8A and pNEF38-c264scTCR-hinge-huIL15D8N expression vectors. (C) shows the sequence of pNEF38-c264scTCR-hinge-huIL15D8A (SEQ ID NO: 14) and pNEF38-c264scTCR-hinge-huIL15D8N (SEQ ID NO: 15) genes and (D) shows the sequence of pNEF38-c264scTCR-hinge-huIL15D8A (SEQ ID NO: 16) and pNEF38-c264scTCR-hinge-huIL15D8N fusion protein (SEQ ID NO: 17), including the leader sequence. The blue underlined nucleotides were changed to generate the indicated IL-15 variants.

FIG. 15 is a graph showing flow cytometric analysis of IL-15R-bearing CTLL2 cells stained with fusion proteins and complexes following by TCR-specific peptide/MHC reagent.

FIG. 16 depicts CTLL-2 cell proliferation assays of the IL15 activity of the TCR/IL15 fusion proteins and TCR/IL15+TCR/IL15R α complexes.

FIG. 17 (A-C) are graphs showing the binding of dimeric fusion proteins complexes of TCR/IL15R α Sushi and TCR/IL15, comprising native and variant forms of IL15, to cognate peptide/MHC complexes displayed on cells loaded with pep-

8

tide, as determined by flow cytometry. Background binding of the dimeric fusion proteins complexes on cells with no loaded peptide is also shown (A) is a graph showing the binding of the dimeric complexes of TCR/IL15R α Sushi and TCR/IL15 wt (native form), or TCR/IL15D8N or TCR/IL15D8A variants to cognate peptide/MHC complexes displayed on cells. (B) is a graph showing the slight background binding of dimeric complexes of TCR/IL15R α Sushi and TCR/IL15 wt (native form) to the cells without loaded peptide. No background binding of dimeric complexes of TCR/IL15R α Sushi and TCR/IL15D8N or TCR/IL15D8A variants was seen to the cells with not loaded. (C) is graph (showing flow cytometric analysis of IL-15R β C-bearing 32D13 cells stained with dimeric complexes of TCR/IL15R α Sushi and TCR/IL15 wt (native form), or TCR/IL15N72D, TCR/IL15D8N or TCR/IL15D8A variants. Enhanced IL-15R β C binding of the complex containing TCR/IL15N72D and decreased IL-15R β C binding of complexes containing TCR/IL15D8N or TCR/IL15D8A was observed.

FIGS. 18 (A&B) are graphs showing binding activities of wide type, antagonist, and agonist TCR/IL15 fusion proteins to cognate peptide/MHC complexes and IL15R α as determined by ELISA. (A) is analysis showing binding activity of fusion proteins to cognate peptide/MHC complexes. (B) is analysis showing binding activity of fusion proteins to IL15R α .

FIG. 19(A-C) are graphs showing the ability of TCR/IL-15 fusion proteins comprising IL-15 variants to inhibit or enhance growth of IL15R-bearing cells, as determined by cell proliferation assay. (A) is graph showing the activity of fusion proteins comprising IL-15 variants to inhibit the proliferation of high affinity IL15R ($\alpha\beta\gamma$ receptor complex) bearing CTLL-2 cells. (B) is graph showing the activity of fusion proteins comprising IL-15 variants to inhibit or enhance the proliferation of low affinity IL15R ($\beta\gamma$ receptor complex) bearing 32D β cells. (C) is graph showing the activity of fusion proteins comprising IL-15 variants to block TCR/IL15wt-stimulated proliferation of high affinity IL15R ($\alpha\beta\gamma$ receptor complex) bearing CTLL-2 cells.

FIG. 20 depicts the effects of in vitro incubation of NK cells with dimeric fusion proteins complexes of TCR/IL15R α Sushi and TCR/IL15 on the survival of xenograft tumor-bearing nude mice. Athymic nude mice were injected with human NSCLC A549-A2 cells to allow establishment of lung metastases. Purified NK cells isolated from spleens of allogeneic donor mice were incubated in vitro with rhIL-2, MART1scTCR-IL2, c264scTCR-IL2 or c264scTCR-IL15/c264scTCR-IL15R α and adoptively transferred into the tumor-bearing mice that had been pretreated with cyclophosphamide (CTX), as indicated in the figure legend. The percent survival following treatment was plotted.

FIG. 21 sets forth Table 1 showing the amino acid replacements in the IL-15 variants and the affects of these changes on IL-15 activity.

FIGS. 22A-B set forth the amino acid sequence of IL-15 (SEQ ID NO:1) and the nucleic acid sequence of IL-15 (SEQ ID NO:2), respectively.

DETAILED DESCRIPTION OF THE INVENTION

It has been established that the IL-15 stably binds to the extracellular domain of the IL-15R α and that the resulting complex is capable of modulating (i.e. either stimulating or blocking) immune responses via the intermediate or high affinity IL-15R complex (Mortier, E. et al., 2006, *J. Biol. Chem.*, 281: 1612-1619; Stoklasek, T. et al., 2006, *J Immunol* 177: 6072-6080; Rubinstein, M. P. et al., 2006, *Proc Natl*

Acad Sci USA 103: 9166-9171, Waldmann, T. A., 2006, *Nat Rev Immunol* 6: 595-601). In addition, it has been demonstrated that single-chain TCR or antibody polypeptides can be fused to cytokines and other immune effector domains and that such bispecific fusion molecules retain functional activity of both fusion domains (Belmont, H. J. et al., 2006, *Clin Immunol* 121: 29-39; Card, K. F. et al., 2004, *Cancer Immunol Immunother* 53: 345-357; Mosquera, L. A. et al., 2005, *J Immunol* 174: 4381-4388; Penichet, M. L., 1997, *Hum Antibodies* 8: 106-118). Further, it has been shown that multivalent forms of the TCR provide enhanced binding to their ligands [(9)] (Zhu, X., H. J., 2006, *J Immunol* 176: 3223-3232).

Definitions

The following definitions are provided for specific terms which are used in the following written description.

As used in the specification and claims, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a cell" includes a plurality of cells, including mixtures thereof. The term "a nucleic acid molecule" includes a plurality of nucleic acid molecules.

As used herein, the term "comprising" is intended to mean that the compositions and methods include the recited elements, but do not exclude other elements. "Consisting essentially of", when used to define compositions and methods, shall mean excluding other elements of any essential significance to the combination. Thus, a composition consisting essentially of the elements as defined herein would not exclude trace contaminants from the isolation and purification method and pharmaceutically acceptable carriers, such as phosphate buffered saline, preservatives, and the like. "Consisting of" shall mean excluding more than trace elements of other ingredients and substantial method steps for administering the compositions of this invention. Embodiments defined by each of these transition terms are within the scope of this invention.

An "antibody" is any immunoglobulin, including antibodies and fragments thereof, that binds a specific epitope. The term encompasses polyclonal, monoclonal, chimeric and single-chain antibodies as well as bispecific antibodies.

The term "antigen" as used herein is meant any substance that causes the immune system to produce antibodies or specific cell-mediated immune responses against it. A disease associated antigen is any substance that is associated with any disease.

The term "biologically active polypeptide" as used herein is meant to refer to an amino acid sequence such as a protein, polypeptide or peptide; a sugar or polysaccharide; a lipid or a glycolipid, glycoprotein, or lipoprotein that can produce the desired effects as discussed herein, including a TCR or antibody fusion protein complex with antigen binding activity.

The term "cell" as used herein is meant to include any prokaryotic, eukaryotic, primary cell or immortalized cell line, any group of such cells as in, a tissue or an organ.

Preferably the cells are of mammalian and particularly of human origin, and can be infected by one or more pathogens. A "host cell" in accord with the invention can be a transfected, transformed, transduced or infected cell of any origin, including prokaryotic, eukaryotic, mammalian, avian, insect, plant or bacteria cells, or it can be a cells of any origin that can be used to propagate a nucleic acid described herein.

The term "conjugate molecule" as it is used herein is meant to refer to a TCR or antibody molecule and an effector molecule usually a chemical or synthesized molecule covalently linked (i.e. fused) by chemical or other suitable method. If desired, the conjugate molecule can be fused at one or several

sites through a peptide linker sequence or a carrier molecule. Alternatively, the peptide linker or carrier may be used to assist in construction of the conjugate molecule. Specifically preferred conjugate molecules are conjugate toxins or detectable labels.

The term "effector molecule" as used herein is meant to refer to an amino acid sequence such as a protein, polypeptide or peptide; a sugar or polysaccharide; a lipid or a glycolipid, glycoprotein, lipoprotein or chemical agent that can produce the desired effects as discussed herein, including an IL-15 domain, IL-15 variant or IL-15 receptor such as IL-15R α , IL-2R β or γ C, or functional fragments thereof.

The terms "fusion molecule" and "fusion protein" are used interchangeably and are meant to refer to a biologically active polypeptide usually a TCR or antibody and an effector molecule usually a protein or peptide sequence covalently linked (i.e. fused) by recombinant, chemical or other suitable method. If desired, the fusion molecule can be fused at one or several sites through a peptide linker sequence. Alternatively, the peptide linker may be used to assist in construction of the fusion molecule. Specifically preferred fusion molecules are fusion proteins. Generally fusion molecule also can be comprised of conjugate molecules.

The term "host cell" is meant to refer to any prokaryotic or eukaryotic cell that contains either a cloning vector or an expression vector. This term also includes those prokaryotic or eukaryotic cells that have been genetically engineered to contain the cloned gene(s) in the chromosome or genome of the host cell.

The term "immune response" as used herein is meant to refer to the process whereby immune cells are stimulated and recruited from the blood to lymphoid as well as non-lymphoid tissues via a multifactorial process that involves distinct adhesive and activation steps. Activation conditions cause the release of cytokines, growth factors, chemokines and other factors, upregulate expression of adhesion and other activation molecules on the immune cells, promote adhesion, morphological changes, and/or extravasation concurrent with chemotaxis through the tissues, increase cell proliferation and cytotoxic activity, stimulate antigen presentation and provide other phenotypic changes including generation of memory cell types. Immune response is also meant to refer to the activity of immune cells to suppress or regulate inflammatory or cytotoxic activity of other immune cells.

As used herein, the terms "polynucleotide" and "nucleic acid molecule" are used interchangeably to refer to polymeric forms of nucleotides of any length. The polynucleotides may contain deoxyribonucleotides, ribonucleotides, and/or their analogs. Nucleotides may have any three-dimensional structure, and may perform any function, known or unknown. The term "polynucleotide" includes, for example, single-, double-stranded and triple helical molecules, a gene or gene fragment, exons, introns, mRNA, tRNA, rRNA, ribozymes, antisense molecules, cDNA, recombinant polynucleotides, branched polynucleotides, aptamers, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A nucleic acid molecule may also comprise modified nucleic acid molecules (e.g., comprising modified bases, sugars, and/or internucleotide linkers).

The term "polypeptide" is meant to refer to any polymer preferably consisting essentially of any of the 20 natural amino acids regardless of its size. Although the term "protein" is often used in reference to relatively large proteins, and "peptide" is often used in reference to small polypeptides, use of these terms in the field often overlaps. The term "polypeptide" refers generally to proteins, polypeptides, and peptides

unless otherwise noted. Peptides useful in accordance with the present invention in general will be generally between about 0.1 to 100 KD or greater up to about 1000 KD, preferably between about 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 30 and 50 KD as judged by standard molecule sizing techniques such as centrifugation or SDS-polyacrylamide gel electrophoresis.

The terms "prevent," "preventing," "prevention," "prophylactic treatment" and the like are meant to refer to reducing the probability of developing a disorder or condition in a subject, who does not have, but is at risk of or susceptible to developing a disorder or condition.

The term "single chain antibody" is meant to refer to an antibody based on a single chain format. Single chain antibodies can consist of the minimal binding subunit of antibodies. Single-chain antibodies can combine only those antigen-binding regions of antibodies on a single stably-folded polypeptide chain. As such, single-chain antibodies are of considerably smaller size than classical immunoglobulins but retain the antigen-specific binding properties of antibodies. Single chain antibodies may be linked to a wide range of ligands, for example effector molecules or drug conjugates.

The term "soluble" as used herein is meant that the fusion molecule and particularly a fusion protein that is not readily sedimented under low G-force centrifugation (e.g. less than about 30,000 revolutions per minute in a standard centrifuge) from an aqueous buffer, e.g., cell media. Further, the fusion molecule is soluble if it remains in aqueous solution at a temperature greater than about 5-37° C. and at or near neutral pH in the presence of low or no concentration of an anionic or non-ionic detergent. Under these conditions, a soluble protein will often have a low sedimentation value e.g., less than about 10 to 50 svedberg units.

Aqueous solutions referenced herein typically have a buffering compound to establish pH, typically within a pH range of about 5-9, and an ionic strength range between about 2 mM and 500 mM. Sometimes a protease inhibitor or mild non-ionic detergent is added. Additionally, a carrier protein may be added if desired such as bovine serum albumin (BSA) to a few mg/ml. Exemplary aqueous buffers include standard phosphate buffered saline, tris-buffered saline, or other well known buffers and cell media formulations.

The term "stimulate" or "stimulating" is meant to refer to increase, to amplify, to augment, to boost an immune response. Stimulation can be a positive alteration. An exemplary increase can be e.g., by 5%, 10%, 25%, 50%, 75%, or even 90-100%. Other exemplary increases include 2-fold, 5-fold, 10-fold, 20-fold, 40-fold, or even 100-fold.

The term "suppress" or "suppressing" is meant to refer to decrease, to attenuate, to diminish, to arrest, or to stabilize an immune response. Suppression may be a negative alteration. An exemplary decrease can be e.g., by 5%, 10%, 25%, 50%, 75%, or even 90-100%. Exemplary decreases include 2-fold, 5-fold, 10-fold, 20-fold, 40-fold, or even 100-fold.

The term "T-cell Receptor" (TCR) is meant to refer to polypeptides of a complex of integral membrane proteins that participates in the activation of T cells in response to the presentation of antigen. T cells recognize a peptide bound to the MHC product through the $\alpha\beta$ heterodimeric T cell receptor (TCR). The TCR repertoire has extensive diversity created by the same gene rearrangement mechanisms used in antibody heavy and light chain genes [Tonegawa, S. (1988) *Bio-sci. Rep.* 8:3-26]. Most of the diversity is generated at the junctions of variable (V) and joining (J) (or diversity, D) regions that encode the complementarity determining region 3 (CDR3) of the α and β chains [Davis and Bjorkman (1988) *Nature* 334:395-402]. However, TCRs do not undergo somatic point mutations as do antibodies and, perhaps not

coincidentally. TCRs also do not undergo the same extent of affinity maturation as antibodies. TCRs as they occur in nature appear to have affinities that range from 10^5 to 10^7 M^{-1} whereas antibodies typically have affinities that range from 10^5 to 10^9 M^{-1} [Davis et al. (1998) *Annu. Rev. Immunol.* 16:523-544; Eisen et al. (1996) *Adv. Protein Chem.* 49:1-56]. While the absence of somatic mutation in TCRs may be associated with lower affinities, it has also been argued that there is not a selective advantage for a TCR to have higher affinity. In fact, the serial-triggering [Valitutti et al. (1995) *Nature* 375:148-151] and kinetic proofreading [Rabinowitz et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:1401-1405] models of T cell activation both suggest that longer off-rates (associated with higher affinity) would be detrimental to the signaling process. It is also possible that higher affinity TCRs might not maintain the peptide specificity required for T cell responses. For example, peptides bound within the MHC groove display limited accessible surface [Bjorkman, P. J. (1997) *Cell* 89:167-170], which may in turn limit the amount of energy that can be generated in the interaction. On the other hand, raising the affinity of a TCR by directing the energy toward the MHC helices would presumably lead to thymic deletion during negative selection [Bevan, M. J. (1997) *Immunity* 7:175-178]. The term "TCR" encompasses polyclonal, monoclonal, chimeric, humanized, heterodimeric and single-chain T-cell receptors or functional fragment thereof, including molecule comprising the TCR V α and V β domains. The term "TCR" also encompasses T-cell receptors disclosed in for example, U.S. Provisional Application Entitled "T CELL RECEPTOR FUSIONS AND CONJUGATES AND METHODS OF USE THEREOF", filed Mar. 19, 2008 and US Patent Publication US 2003 01-44474A1.

The term "vector" is a nucleic acid molecule that is able to replicate autonomously in a host cell and can accept foreign DNA. A vector carries its own origin of replication, one or more unique recognition sites for restriction endonucleases which can be used for the insertion of foreign DNA, and usually selectable markers such as genes coding for antibiotic resistance, and often recognition sequences (e.g. promoter) for the expression of the inserted DNA. Common vectors include plasmid vectors and phage vectors. T-Cell Receptors (TCR)

T-cells are a subgroup of cells which together with other immune cell types (polymorphonuclear, eosinophils, basophils, mast cells, B-, NK cells), constitute the cellular component of the immune system. Under physiological conditions T-cells function in immune surveillance and in the elimination of foreign antigen. However, under pathological conditions there is compelling evidence that T-cells play a major role in the causation and propagation of disease. In these disorders, breakdown of T-cell immunological tolerance, either central or peripheral is a fundamental process in the causation of autoimmune disease.

The TCR is composed of at least seven transmembrane proteins. The disulfide-linked ($\alpha\beta$) heterodimer forms the monotypic antigen recognition unit, while the invariant chains of CD3, consisting of ϵ , γ , δ , and ζ and η chains, are responsible for coupling the ligand binding to signaling pathways that result in T-cell activation and the elaboration of the cellular immune responses. Despite the gene diversity of the TCR chains, two structural features are common to all known subunits. Firstly, they are transmembrane proteins with a single transmembrane spanning domain—presumably α -helical. Secondly, all the TCR chains have the unusual feature of possessing a charged amino acid within the predicted transmembrane domain. The invariant chains have a single negative charge,

13

conserved between the mouse and human, and the variant chains possess one (TCR-beta) or two (TCR-alpha) positive charges. The transmembrane sequence of TCR-alpha is highly conserved in a number of species and thus phylogenetically may serve an important functional role. The octapeptide sequence containing the hydrophilic amino acids arginine and lysine is identical between the species.

A T-cell response is modulated by antigen binding to a TCR. One type of TCR is a membrane bound heterodimer consisting of an α and β chain resembling an immunoglobulin variable (V) and constant (C) region. The TCR α chain includes a covalently linked V- β and C- α chain, whereas the β chain includes a V- β chain covalently linked to a C- β chain. The V- α and V- β chains form a pocket or cleft that can bind a superantigen or antigen in the context of a major histocompatibility complex (MHC) (known in humans as an HLA complex). See generally Davis Ann. Rev. of Immunology 3: 537 (1985); Fundamental Immunology 3rd Ed., W. Paul Ed. Rsen Press LTD. New York (1993).

Fusions Proteins

The soluble fusion protein and conjugate molecule complexes of the invention comprise at least two soluble fusion proteins, where the first fusion protein comprises a first biologically active polypeptide covalently linked to interleukin-15 (IL-15) or functional fragment thereof; and the second fusion protein comprises a second biologically active polypeptide covalently linked to soluble interleukin-15 receptor alpha (IL-15R α) polypeptide or functional fragment thereof, and wherein IL-15 domain of a first fusion protein binds to the soluble IL-15R α domain of the second fusion protein to form a soluble fusion protein complex.

In certain examples, one of the biologically active polypeptides comprises a first soluble TCR or fragment thereof. The other or second biologically active polypeptide comprises the first soluble TCR or functional fragment thereof and thus creates a multivalent TCR fusion protein complex with increased binding activity for cognate ligands compared to the monovalent TCR. Further, the other biologically active polypeptide comprises a second soluble TCR or functional fragment thereof, different than the first soluble TCR. In certain examples, TCRs are produced that have higher affinity, or increased binding affinity for cognate ligands as compared, for example, to the native TCR. If the soluble TCR of the invention as described herein has a higher avidity or affinity for its ligand, then it is useful as a specific probe for cell-surface bound antigen. In certain preferred examples of the invention, the TCR is specific for recognition of a particular antigen.

In exemplary embodiments, TCR is a heterodimer comprising an α chain (herein referred to as cc, alpha or a chain) and a β chain (herein referred to as β , beta or b chain). In other exemplary embodiments, the TCR comprises a single chain TCR polypeptide. The single chain TCR may comprise a TCR V- α chain covalently linked to a TCR V- β chain by a peptide linker sequence. The single chain TCR may further comprise a soluble TCR C β chain fragment covalently linked to a TCR V- β chain. The single chain TCR may further comprise a soluble TCR C α chain fragment covalently linked to a TCR V- α chain.

In a further embodiment, one or both of the first and second biologically active polypeptides comprises an antibody or functional fragment thereof.

As used herein, the term "biologically active polypeptide" or "effector molecule" is meant an amino acid sequence such as a protein, polypeptide or peptide; a sugar or polysaccharide; a lipid or a glycolipid, glycoprotein, or lipoprotein that can produce the desired effects as discussed herein. Effector

14

molecules also include chemical agents. Also contemplated are effector molecule nucleic acids encoding a biologically active or effector protein, polypeptide, or peptide. Thus, suitable molecules include regulatory factors, enzymes, antibodies, or drugs as well as DNA, RNA, and oligonucleotides. The biologically active polypeptides or effector molecule can be naturally-occurring or it can be synthesized from known components, e.g., by recombinant or chemical synthesis and can include heterologous components. A biologically active polypeptides or effector molecule is generally between about 0.1 to 100 KD or greater up to about 1000 KD, preferably between about 0.1, 0.2, 0.5, 1, 2, 5, 10, 20, 30 and 50 KD as judged by standard molecule sizing techniques such as centrifugation or SDS-polyacrylamide gel electrophoresis. Desired effects of the invention include, but are not limited to, for example, forming a TCR fusion protein complex with increased binding activity, killing a target cell, e.g. either to induce cell proliferation or cell death, initiate an immune response, in preventing or treating a disease, or to act as a detection molecule for diagnostic purposes. For such detection, an assay could be used, for example an assay that includes sequential steps of culturing cells to proliferate same, and contacting the cells with a TCR fusion complex of the invention and then evaluating whether the TCR fusion complex inhibits further development of the cells.

Covalently linking the effector molecule to the TCR peptide in accordance with the invention provides a number of significant advantages. TCR fusion complexes of the invention can be produced that contain a single effector molecule, including such a peptide of known structure. Additionally, a wide variety of effector molecules can be produced in similar DNA vectors. That is, a library of different effector molecules can be linked to the TCR molecule for presentation of infected or diseased cells. Further, for therapeutic applications, rather than administration of an TCR molecule to a subject, a DNA expression vector coding for the TCR molecule linked to the effector peptide can be administered for in vivo expression of the TCR fusion complex. Such an approach avoids costly purification steps typically associated with preparation of recombinant proteins and avoids the complexities of antigen uptake and processing associated with conventional approaches.

As noted, components of the fusion proteins disclosed herein, e.g., effector molecule such as cytokines, chemokines, growth factors, protein toxins, immunoglobulin domains or other bioactive molecules and any peptide linkers, can be organized in nearly any fashion provided that the fusion protein has the function for which it was intended. In particular, each component of the fusion protein can be spaced from another component by at least one suitable peptide linker sequence if desired. Additionally, the fusion proteins may include tags, e.g., to facilitate modification, identification and/or purification of the fusion protein. More specific fusion proteins are in the Examples described below.

Linkers

The fusion complexes of the invention preferably also include a flexible linker sequence interposed between the IL-15 or IL-15R α domains and the biologically active polypeptide. The linker sequence should allow effective positioning of the biologically active polypeptide with respect to the IL-15 or IL-15R α domains to allow functional activity of both domains. In embodiments where the biologically active polypeptide is a TCR, the linker sequence positions the TCR molecule binding groove so that the T cell receptor can recognize presenting MHC-peptide complexes and can deliver the effector molecule to a desired site. Successful presentation of the effector molecule can modulate the activity of a

15

cell either to induce or to inhibit T-cell proliferation, or to initiate or inhibit an immune response to a particular site, as determined by the assays disclosed below, including the in vitro assays that includes sequential steps of culturing T cells to proliferate same, and contacting the T cells with a TCR fusion complex of the invention and then evaluating whether the TCR fusion complex inhibits further development of the cells.

In certain embodiments, the soluble fusion protein complex has a linker wherein the first biologically active polypeptide is covalently linked to IL-15 (or functional fragment thereof) by polypeptide linker sequence.

In other certain embodiments, the soluble fusion protein complex as described herein has a linker wherein the second biologically active polypeptide is covalently linked to IL-15R α polypeptide (or functional fragment thereof) by polypeptide linker sequence.

The linker sequence is preferably encoded by a nucleotide sequence resulting in a peptide that can effectively position the binding groove of the TCR molecule for recognition of a presenting antigen. As used herein, the phrase "effective positioning of the biologically active polypeptide with respect to the IL-15 or IL-15R α domains", or other similar phrase, is intended to mean the biologically active polypeptide linked to the IL-15 or IL-15R α domains is positioned so that the IL-15 or IL-15R α domains are capable of interacting with each other to form a protein complex. In certain embodiments, the IL-15 or IL-15R α domains are effectively positioned to allow interactions with immune cells to initiate or inhibit an immune reaction, or to inhibit or stimulate cell development.

Preferably the linker sequence comprises from about 7 to 20 amino acids, more preferably from about 8 to 16 amino acids. The linker sequence is preferably flexible so as not hold the biologically active polypeptide or effector molecule in a single undesired conformation. The linker sequence can be used, e.g., to space the recognition site from the fused molecule. Specifically, the peptide linker sequence can be positioned between the biologically active polypeptide and the effector molecule, e.g., to chemically cross-link same and to provide molecular flexibility. The linker is preferably predominantly comprises amino acids with small side chains, such as glycine, alanine and serine, to provide for flexibility. Preferably about 80 or 90 percent or greater of the linker sequence comprises glycine, alanine or serine residues, particularly glycine and serine residues. For a fusion protein complex that comprise a heterodimer TCR, the linker sequence is suitably linked to the b chain of the TCR molecule, although the linker sequence also could be attached to the a chain of the TCR molecule. Alternatively, linker sequence may be linked to both a and b chains of the TCR molecule. When such a beta peptide chain is expressed along with the a chain, the linked TCR-effector peptide should fold resulting in a functional TCR molecule as generally depicted in FIG. 1. One suitable linker sequence is ASGGGSGGG (i.e., Ala Ser Gly Gly Gly Ser Gly Gly Gly) (SEQ ID NO: 18), preferably linked to the first amino acid of the b domain of the TCR. Different linker sequences could be used including any of a number of flexible linker designs that have been used successfully to join antibody variable regions together, see Whitlow, M. et al., (1991) *Methods: A Companion to Methods in Enzymology* 2:97-105. In some examples, for covalently linking an effector molecule to a TCR b chain molecule, the amino sequence of the linker should be capable of spanning suitable distance from the C-terminal residue of the TCR beta chain to the N-terminal residue of the effector molecule.

16

In general, preparation of the fusion protein complexes of the invention can be accomplished by procedures disclosed herein and by recognized recombinant DNA techniques involving, e.g., polymerase chain amplification reactions (PCR), preparation of plasmid DNA, cleavage of DNA with restriction enzymes, preparation of oligonucleotides, ligation of DNA, isolation of mRNA, introduction of the DNA into a suitable cell, transformation or transfection of a host, culturing of the host. Additionally, the fusion molecules can be isolated and purified using chaotropic agents and well known electrophoretic, centrifugation and chromatographic methods. See generally, Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd ed. (1989); and Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York (1989) for disclosure relating to these methods.

As used herein, biologically active polypeptides or effector molecules of the invention may include factors such as cytokines, chemokines, growth factors, protein toxins, immunoglobulin domains or other bioactive proteins such as enzymes. Also biologically active polypeptides may include conjugates to other compounds such as non-protein toxins, cytotoxic agents, chemotherapeutic agents, detectable labels, radioactive materials and such.

Cytokines of the invention are defined by any factor produced by cells that affect other cells and are responsible for any of a number of multiple effects of cellular immunity. Examples of cytokines include but are not limited to the IL-2 family, interferon (IFN), IL-10, IL-1, IL-17, TGF and TNF cytokine families, and to IL-1 through IL-35, IFN- α , IFN- β , IFN- γ , TGF- β , TNF- α and TNF β .

In an aspect of the invention, the first fusion protein comprises a first biologically active polypeptide covalently linked to interleukin-15 (IL-15) domain or a functional fragment thereof. IL-15 is a cytokine which affects T-cell activation and proliferation. IL-15 activity in affecting immune cell activation and proliferation is similar in some respects to IL-2, although fundamental difference have been well characterized (Waldmann, T A, 2006, *Nature Rev. Immunol.* 6:595-601).

In another aspect of the invention, the first fusion protein comprises an interleukin-15 (IL-15) domain that is an IL-15 variant (also referred to herein as IL-15 mutant). The IL-15 variant preferably comprises a different amino acid sequence that the native (or wild type) IL-15 protein. The IL-15 variant preferably binds the IL-15R α polypeptide and functions as an IL-15 agonist or antagonist. Preferably IL-15 variants with agonist activity have super agonist activity. In some embodiments, the IL-15 variant can function as an IL-15 agonist or antagonist independent of its association with IL-15R α . IL-15 agonists are exemplified by comparable or increased biological activity compared to wild type IL-15. IL-15 antagonists are exemplified by decreased biological activity compared to wild type IL-15 or by the ability to inhibit IL-15-mediated responses. In some examples, the IL-15 variant binds with increased or decreased activity to the IL-15R β RC receptors. In some embodiments, the sequence of the IL-15 variant has at least one amino acid change, e.g. substitution or deletion, compared to the native IL-2 sequence, such changes resulting in IL-15 agonist or antagonist activity. Preferably the amino acid substitutions/deletions are in the domains of IL-15 that interact with IL-15R β and/or γ C. More preferably, the amino acid substitutions/deletions do not affect binding to the IL-15R α polypeptide or the ability to produce the IL-15 variant. Suitable amino acid substitutions/deletions to generate IL-15 variants can be identified based on putative or known IL-15 structures, comparisons of IL-15 with homologous molecules such as IL-2 with known structure, through ratio-

nal or random mutagenesis and functional assays, as provided herein, or other empirical methods. Additionally suitable amino acid substitutions can be conservative or non-conservative changes and insertions of additional amino acids. Preferably IL-15 variants of the invention contain one or more than one amino acid substitutions/deletions at position 8, 61, 65, 72, 92, 101, 108, or 111 of the mature human IL-15 sequence; particularly, D8N ("D8" refers to the amino acid and residue position in the native mature human IL-15 sequence and "N" refers to the substituted amino acid residue at that position in the IL-15 variant), DBA, D61A, N65A, N72R or Q108A substitutions result in IL-15 variants with antagonist activity and N72D substitutions result in IL-15 variants with agonist activity.

While in one aspect of the invention the IL-15 variant is a component of a fusion protein complex, in other aspects the IL-15 variant is a non-fusion protein. Preferably the non-fusion form of the IL-15 variant is a soluble cytokine that functions as an IL-15 agonist or antagonist. In some embodiments, the non-fusion IL-15 variant forms a complex with IL-15Ra whereas in other embodiment it acts independently of IL-15Ra.

Chemokines of the invention, similar to cytokines, are defined as any chemical factor or molecule which when exposed to other cells are responsible for any of a number of multiple effects of cellular immunity. Suitable chemokines may include but are not limited to the CXC, CC, C, and CX₂C chemokine families and to CCL-1 through CCL-28, CXC-1 through CXC-17, XCL-1, XCL-2, CX₃ CL1, MIP-1b, IL-8, MCP-1, and Rantes.

Growth factors include any molecules which when exposed to a particular cell induce proliferation and/or differentiation of the affected cell. Growth factors include proteins and chemical molecules, some of which include: GM-CSF, G-CSF, human growth factor and stem cell growth factor. Additional growth factors may also be suitable for uses described herein.

Toxins or cytotoxic agents include any substance which has a lethal effect or an inhibitory effect on growth when exposed to cells. More specifically, the effector molecule can be a cell toxin of, e.g., plant or bacterial origin such as, e.g., diphtheria toxin (DT), shiga toxin, abrin, cholera toxin, ricin, saporin, pseudomonas exotoxin (PE), pokeweed antiviral protein, or gelonin. Biologically active fragments of such toxins are well known in the art and include, e.g., DT A chain and ricin A chain. Additionally, the toxin can be an agent active at the cell surface such as, e.g., phospholipase enzymes (e.g., phospholipase C).

Further, the effector molecule can be a chemotherapeutic drug such as, e.g., vindesine, vincristine, vinblastin, methotrexate, adriamycin, bleomycin, or cisplatin.

Additionally, the effector molecule can be a detectably-labeled molecule suitable for diagnostic or imaging studies. Such labels include biotin or streptavidin/avidin, a detectable nanoparticles or crystal, an enzyme or catalytically active fragment thereof, a fluorescent label such as green fluorescent protein, FITC, phycoerythrin, cychrome, texas red or quantum dots; a radionuclide e.g., iodine-131, yttrium-90, rhenium-188 or bismuth-212; a phosphorescent or chemiluminescent molecules or a label detectable by PET, ultrasound or MRI such as Gd- or paramagnetic metal ion-based contrast agents. See e.g., Moskaug, et al. J. Biol. Chem. 264, 15709 (1989); Pastan, I. et al. Cell 47, 641, 1986; Pastan et al., Recombinant Toxins as Novel Therapeutic Agents, Ann. Rev. Biochem. 61, 331, (1992); "Chimeric Toxins" Olsnes and Phil, Pharmac. Ther., 25, 355 (1982); published PCT application no. WO 94/29350; published PCT application no. WO 94/04689;

published PCT application no. WO2005046449 and U.S. Pat. No. 5,620,939 for disclosure relating to making and using proteins comprising effectors or tags.

A protein fusion or conjugate complex that includes a covalently linked IL-15 and IL-15Ra domains has several important uses. For example, the protein fusion or conjugate complex comprising a TCR can be employed to deliver the IL-15/IL-15Ra complex to certain cells capable of specifically binding the TCR. Accordingly, the protein fusion or conjugate complex provide means of selectively damaging or killing cells comprising the ligand. Examples of cells or tissue capable of being damaged or killed by the protein fusion or conjugate complexes comprising a TCR include tumors and virally or bacterially infected cells expressing one or more ligands capable of being specifically bound by the TCR. Cells or tissue susceptible to being damaged or killed can be readily assayed by the methods disclosed herein.

The IL-15 and IL-15Ra polypeptides of the invention suitably correspond in amino acid sequence to naturally occurring IL-15 and IL-15Ra molecules, e.g. IL-15 and IL-15Ra molecules of a human, mouse or other rodent, or other mammal.

In some settings it can be useful to make the protein fusion or conjugate complexes of the present invention polyvalent, e.g., to increase the valency of the sc-TCR. In particular, interactions between the IL-15 and IL-15Ra domains of the fusion protein complex provide a means of generating polyvalent complexes. In addition, the polyvalent fusion protein can be made by covalently or non-covalently linking together between one and four proteins (the same or different) by using e.g., standard biotin-streptavidin labeling techniques, or by conjugation to suitable solid supports such as latex beads. Chemically cross-linked proteins (for example cross-linked to dendrimers) are also suitable polyvalent species. For example, the protein can be modified by including sequences encoding tag sequences that can be modified such as the biotinylation BirA tag or amino acid residues with chemically reactive side chains such as Cys or His. Such amino acid tags or chemically reactive amino acids may be positioned in a variety of positions in the fusion protein, preferably distal to the active site of the biologically active polypeptide or effector molecule. For example, the C-terminus of a soluble fusion protein can be covalently linked to a tag or other fused protein which includes such a reactive amino acid(s). Suitable side chains can be included to chemically link two or more fusion proteins to a suitable dendrimer or other nanoparticle to give a multivalent molecule. Dendrimers are synthetic chemical polymers that can have any one of a number of different functional groups of their surface (D. Tomalia, *Aldrichimica Acta*, 26:91:101 (1993)). Exemplary dendrimers for use in accordance with the present invention include e.g. E9 starburst polyamine dendrimer and E9 combust polyamine dendrimer, which can link cystine residues.

Nucleic Acids and Vectors

Nucleic Acids

The invention further provides nucleic acid sequences and particularly DNA sequences that encode the present fusion proteins. Preferably, the DNA sequence is carried by a vector suited for extrachromosomal replication such as a phage, virus, plasmid, phagemid, cosmid, YAC, or episome. In particular, a DNA vector that encodes a desired fusion protein can be used to facilitate preparative methods described herein and to obtain significant quantities of the fusion protein. The DNA sequence can be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. A variety of host-vector systems may be

utilized to express the protein-coding sequence. These include mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage DNA, plasmid DNA or cosmid DNA. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used. See generally Sambrook et al., *supra* and Ausubel et al. *supra*.

Included in the invention are methods for making a soluble fusion protein complex, the method comprising introducing into a host cell a DNA vector as described herein encoding the first and second fusion proteins, culturing the host cell in media under conditions sufficient to express the fusion proteins in the cell or the media and allow association between IL-15 domain of a first fusion protein and the soluble IL-15R α domain of a second fusion protein to form the soluble fusion protein complex, purifying the soluble fusion protein complex from the host cells or media.

In general, a preferred DNA vector according to the invention comprises a nucleotide sequence linked by phosphodiester bonds comprising, in a 5' to 3' direction a first cloning site for introduction of a first nucleotide sequence encoding a TCR chain, operatively linked to a sequence encoding an effector molecule.

The fusion protein components encoded by the DNA vector can be provided in a cassette format. By the term "cassette" is meant that each component can be readily substituted for another component by standard recombinant methods. In particular, a DNA vector configured in a cassette format is particularly desirable when the encoded fusion complex is to be used against pathogens that may have or have capacity to develop serotypes.

To make the vector coding for a TCR fusion complex, the sequence coding for the TCR molecule is linked to a sequence coding for the effector peptide by use of suitable ligases. DNA coding for the presenting peptide can be obtained by isolating DNA from natural sources such as from a suitable cell line or by known synthetic methods, e.g. the phosphate triester method. See, e.g., *Oligonucleotide Synthesis*, IRL Press (M. J. Gait, ed., 1984). Synthetic oligonucleotides also may be prepared using commercially available automated oligonucleotide synthesizers. Once isolated, the gene coding for the TCR molecule can be amplified by the polymerase chain reaction (PCR) or other means known in the art. Suitable PCR primers to amplify the TCR peptide gene may add restriction sites to the PCR product. The PCR product preferably includes splice sites for the effector peptide and leader sequences necessary for proper expression and secretion of the TCR-effector fusion complex. The PCR product also preferably includes a sequence coding for the linker sequence, or a restriction enzyme site for ligation of such a sequence.

The fusion proteins described herein are preferably produced by standard recombinant DNA techniques. For example, once a DNA molecule encoding the TCR protein is isolated, sequence can be ligated to another DNA molecule encoding the effector polypeptide. The nucleotide sequence coding for a TCR molecule may be directly joined to a DNA sequence coding for the effector peptide or, more typically, a DNA sequence coding for the linker sequence as discussed herein may be interposed between the sequence coding for the TCR molecule and the sequence coding for the effector peptide and joined using suitable ligases. The resultant hybrid DNA molecule can be expressed in a suitable host cell to produce the TCR fusion complex. The DNA molecules are ligated to each other in a 5' to 3' orientation such that, after ligation, the translational frame of the encoded polypeptides

is not altered (i.e., the DNA molecules are ligated to each other in-frame). The resulting DNA molecules encode an in-frame fusion protein.

Other nucleotide sequences also can be included in the gene construct. For example, a promoter sequence, which controls expression of the sequence coding for the TCR peptide fused to the effector peptide, or a leader sequence, which directs the TCR fusion complex to the cell surface or the culture medium, can be included in the construct or present in the expression vector into which the construct is inserted. An immunoglobulin or CMV promoter is particularly preferred.

In obtaining variant TCR coding sequences, those of ordinary skill in the art will recognize that TCR-derived proteins may be modified by certain amino acid substitutions, additions, deletions, and post-translational modifications, without loss or reduction of biological activity. In particular, it is well-known that conservative amino acid substitutions, that is, substitution of one amino acid for another amino acid of similar size, charge, polarity and conformation, are unlikely to significantly alter protein function. The 20 standard amino acids that are the constituents of proteins can be broadly categorized into four groups of conservative amino acids as follows: the nonpolar (hydrophobic) group includes alanine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan and valine; the polar (uncharged, neutral) group includes asparagine, cysteine, glutamine, glycine, serine, threonine and tyrosine; the positively charged (basic) group contains arginine, histidine and lysine; and the negatively charged (acidic) group contains aspartic acid and glutamic acid. Substitution in a protein of one amino acid for another within the same group is unlikely to have an adverse effect on the biological activity of the protein.

Homology between nucleotide sequences can be determined by DNA hybridization analysis, wherein the stability of the double-stranded DNA hybrid is dependent on the extent of base pairing that occurs. Conditions of high temperature and/or low salt content reduce the stability of the hybrid, and can be varied to prevent annealing of sequences having less than a selected degree of homology. For instance, for sequences with about 55% G-C content, hybridization and wash conditions of 40-50. degree. C., 6.times.SSC (sodium chloride/sodium citrate buffer) and 0.1% SDS (sodium dodecyl sulfate) indicate about 60-70% homology, hybridization and wash conditions of 50-65. degree. C., 1.times.SSC and 0.1% SDS indicate about 82-97% homology, and hybridization and wash conditions of 52. degree. C., 0.1.times.SSC and 0.1% SDS indicate about 99-100% homology. A wide range of computer programs for comparing nucleotide and amino acid sequences (and measuring the degree of homology) are also available, and a list providing sources of both commercially available and free software is found in Ausubel et al. (1999). Readily available sequence comparison and multiple sequence alignment algorithms are, respectively, the Basic Local Alignment Search Tool (BLAST) (Altschul et al., 1997) and ClustalW programs. BLAST is available on the world wide web at ncbi.nlm.nih.gov and a version of ClustalW is available at 2.ebi.ac.uk.

The components of the fusion protein can be organized in nearly any order provided each is capable of performing its intended function. For example, in one embodiment, the TCR is situated at the C or N terminal end of the effector molecule.

Preferred effector molecules of the invention will have sizes conducive to the function for which those domains are intended. The effector molecules of the invention can be made and fused to the TCR by a variety of methods including well-known chemical cross-linking methods. See e.g., Means, G. E. and Feeney, R. E. (1974) in *Chemical Modifi-*

cation of Proteins, Holden-Day. See also, S. S. Wong (1991) in Chemistry of Protein Conjugation and Cross-Linking, CRC Press. However it is generally preferred to use recombinant manipulations to make the in-frame fusion protein.

As noted, a fusion molecule or a conjugate molecule in accord with the invention can be organized in several ways. In an exemplary configuration, the C-terminus of the TCR is operatively linked to the N-terminus of the effector molecule. That linkage can be achieved by recombinant methods if desired. However, in another configuration, the N-terminus of the TCR is linked to the C-terminus of the effector molecule.

Alternatively, or in addition, one or more additional effector molecules can be inserted into the TCR fusion or conjugate complexes as needed.

Vectors and Expression

A number of strategies can be employed to express protein fusion complexes of the invention. For example, the TCR gene fusion construct described above can be incorporated into a suitable vector by known means such as by use of restriction enzymes to make cuts in the vector for insertion of the construct followed by ligation. The vector containing the gene construct is then introduced into a suitable host for expression of the TCR fusion peptide. See, generally, Sambrook et al., supra. Selection of suitable vectors can be made empirically based on factors relating to the cloning protocol. For example, the vector should be compatible with, and have the proper replicon for the host that is being employed. Further the vector must be able to accommodate the DNA sequence coding for the TCR fusion complex that is to be expressed. Suitable host cells include eukaryotic and prokaryotic cells, preferably those cells that can be easily transformed and exhibit rapid growth in culture medium. Specifically preferred hosts cells include prokaryotes such as *E. coli*, *Bacillus subtilis*, etc. and eukaryotes such as animal cells and yeast strains, e.g., *S. cerevisiae*. Mammalian cells are generally preferred, particularly J558, NSO, SP2-O or CHO. Other suitable hosts include, e.g., insect cells such as Sf9. Conventional culturing conditions are employed. See Sambrook, supra. Stable transformed or transfected cell lines can then be selected. Cells expressing a TCR fusion complex of the invention can be determined by known procedures. For example, expression of a TCR fusion complex linked to an immunoglobulin can be determined by an ELISA specific for the linked immunoglobulin and/or by immunoblotting. Other methods for detecting expression of fusion proteins comprising TCRs linked to IL-15 or IL-15Ra domains are disclosed in the Examples.

As mentioned generally above, a host cell can be used for preparative purposes to propagate nucleic acid encoding a desired fusion protein. Thus a host cell can include a prokaryotic or eukaryotic cell in which production of the fusion protein is specifically intended. Thus host cells specifically include yeast, fly, worm, plant, frog, mammalian cells and organs that are capable of propagating nucleic acid encoding the fusion. Non-limiting examples of mammalian cell lines which can be used include CHO dhfr-cells (Urlaub and Chasm, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)), 293 cells (Graham et al., J Gen. Virol., 36:59 (1977)) or myeloma cells like SP2 or NSO (Galfre and Milstein, Meth. Enzymol., 73(B):3 (1981)).

Host cells capable of propagating nucleic acid encoding a desired fusion protein encompass non-mammalian eukaryotic cells as well, including insect (e.g., *Sp. frugiperda*), yeast (e.g., *S. cerevisiae*, *S. pombe*, *P. pastoris*, *K. lactis*, *H. polymorpha*; as generally reviewed by Fleer, R., Current Opinion

in Biotechnology, 3(5):486496 (1992)), fungal and plant cells. Also contemplated are certain prokaryotes such as *E. coli* and *Bacillus*.

Nucleic acid encoding a desired fusion protein can be introduced into a host cell by standard techniques for transfecting cells. The term "transfecting" or "transfection" is intended to encompass all conventional techniques for introducing nucleic acid into host cells, including calcium phosphate co-precipitation, DEAE-dextran-mediated transfection, lipofection, electroporation, microinjection, viral transduction and/or integration. Suitable methods for transfecting host cells can be found in Sambrook et al. supra, and other laboratory textbooks.

Various promoters (transcriptional initiation regulatory region) may be used according to the invention. The selection of the appropriate promoter is dependent upon the proposed expression host. Promoters from heterologous sources may be used as long as they are functional in the chosen host.

Promoter selection is also dependent upon the desired efficiency and level of peptide or protein production. Inducible promoters such as tac are often employed in order to dramatically increase the level of protein expression in *E. coli*. Over-expression of proteins may be harmful to the host cells. Consequently, host cell growth may be limited. The use of inducible promoter systems allows the host cells to be cultivated to acceptable densities prior to induction of gene expression, thereby facilitating higher product yields.

Various signal sequences may be used according to the invention. A signal sequence which is homologous to the TCR coding sequence may be used. Alternatively, a signal sequence which has been selected or designed for efficient secretion and processing in the expression host may also be used. For example, suitable signal sequence/host cell pairs include the *B. subtilis* sacB signal sequence for secretion in *B. subtilis*, and the *Saccharomyces cerevisiae* alpha-mating factor or *P. pastoris* acid phosphatase phoI signal sequences for *P. pastoris* secretion. The signal sequence may be joined directly through the sequence encoding the signal peptidase cleavage site to the protein coding sequence, or through a short nucleotide bridge consisting of usually fewer than ten codons, where the bridge ensures correct reading frame of the downstream TCR sequence.

Elements for enhancing transcription and translation have been identified for eukaryotic protein expression systems. For example, positioning the cauliflower mosaic virus (CaMV) promoter 1000 bp on either side of a heterologous promoter may elevate transcriptional levels by 10- to 400-fold in plant cells. The expression construct should also include the appropriate translational initiation sequences. Modification of the expression construct to include a Kozak consensus sequence for proper translational initiation may increase the level of translation by 10 fold.

A selective marker is often employed, which may be part of the expression construct or separate from it (e.g., carried by the expression vector), so that the marker may integrate at a site different from the gene of interest. Examples include markers that confer resistance to antibiotics (e.g., bla confers resistance to ampicillin for *E. coli* host cells, nptII confers kanamycin resistance to a wide variety of prokaryotic and eukaryotic cells) or that permit the host to grow on minimal medium (e.g., HIS4 enables *P. pastoris* or His.sup.-*S. cerevisiae* to grow in the absence of histidine). The selectable marker has its own transcriptional and translational initiation and termination regulatory regions to allow for independent expression of the marker. If antibiotic resistance is employed as a marker, the concentration of the antibiotic for selection

will vary depending upon the antibiotic, generally ranging from 10 to 600. μ g of the antibiotic/mL of medium.

The expression construct is assembled by employing known recombinant DNA techniques (Sambrook et al., 1989; Ausubel et al., 1999). Restriction enzyme digestion and ligation are the basic steps employed to join two fragments of DNA. The ends of the DNA fragment may require modification prior to ligation, and this may be accomplished by filling in overhangs, deleting terminal portions of the fragment(s) with nucleases (e.g., ExoIII), site directed mutagenesis, or by adding new base pairs by PCR. Polylinkers and adaptors may be employed to facilitate joining of selected fragments. The expression construct is typically assembled in stages employing rounds of restriction, ligation, and transformation of *E. coli*. Numerous cloning vectors suitable for construction of the expression construct are known in the art (λ .ZAP and pBLUESCRIPT SK-1, Stratagene, LaJolla, Calif., pET, Novagen Inc., Madison, Wis.—cited in Ausubel et al., 1999) and the particular choice is not critical to the invention. The selection of cloning vector will be influenced by the gene transfer system selected for introduction of the expression construct into the host cell. At the end of each stage, the resulting construct may be analyzed by restriction, DNA sequence, hybridization and PCR analyses.

The expression construct may be transformed into the host as the cloning vector construct, either linear or circular, or may be removed from the cloning vector and used as is or introduced onto a delivery vector. The delivery vector facilitates the introduction and maintenance of the expression construct in the selected host cell type. The expression construct is introduced into the host cells by any of a number of known gene transfer systems (e.g., natural competence, chemically mediated transformation, protoplast transformation, electroporation, biolistic transformation, transfection, or conjugation) (Ausubel et al., 1999; Sambrook et al., 1989). The gene transfer system selected depends upon the host cells and vector systems used.

For instance, the expression construct can be introduced into *S. cerevisiae* cells by protoplast transformation or electroporation. Electroporation of *S. cerevisiae* is readily accomplished, and yields transformation efficiencies comparable to spheroplast transformation.

The present invention further provides a production process for isolating a fusion protein of interest. In the process, a host cell (e.g., a yeast, fungus, insect, bacterial or animal cell), into which has been introduced a nucleic acid encoding the protein of the interest operatively linked to a regulatory sequence, is grown at production scale in a culture medium in the presence of the fusion protein to stimulate transcription of the nucleotides sequence encoding the fusion protein of interest. Subsequently, the fusion protein of interest is isolated from harvested host cells or from the culture medium. Standard protein purification techniques can be used to isolate the protein of interest from the medium or from the harvested cells. In particular, the purification techniques can be used to express and purify a desired fusion protein on a large-scale (i.e. in at least milligram quantities) from a variety of implementations including roller bottles, spinner flasks, tissue culture plates, bioreactor, or a fermentor.

An expressed protein fusion complex can be isolated and purified by known methods. Typically the culture medium is centrifuged and then the supernatant is purified by affinity or immunoaffinity chromatography, e.g. Protein-A or Protein-G affinity chromatography or an immunoaffinity protocol comprising use of monoclonal antibodies that bind the expressed fusion complex such as a linked TCR or immunoglobulin region thereof. The fusion proteins of the present invention

can be separated and purified by appropriate combination of known techniques. These methods include, for example, methods utilizing solubility such as salt precipitation and solvent precipitation, methods utilizing the difference in molecular weight such as dialysis, ultra-filtration, gel-filtration, and SDS-polyacrylamide gel electrophoresis, methods utilizing a difference in electrical charge such as ion-exchange column chromatography, methods utilizing specific affinity such as affinity chromatography, methods utilizing a difference in hydrophobicity such as reverse-phase high performance liquid chromatography and methods utilizing a difference in isoelectric point, such as isoelectric focusing electrophoresis, metal affinity columns such as Ni-NTA. See generally Sambrook et al. and Ausubel et al. supra for disclosure relating to these methods.

It is preferred that the fusion proteins of the present invention be substantially pure. That is, the fusion proteins have been isolated from cell constituents that naturally accompany it so that the fusion proteins are present preferably in at least 80% or 90% to 95% homogeneity (w/w). Fusion proteins having at least 98 to 99% homogeneity (w/w) are most preferred for many pharmaceutical, clinical and research applications. Once substantially purified the fusion protein should be substantially free of contaminants for therapeutic applications. Once purified partially or to substantial purity, the soluble fusion proteins can be used therapeutically, or in performing in vitro or in vivo assays as disclosed herein. Substantial purity can be determined by a variety of standard techniques such as chromatography and gel electrophoresis.

Truncated TCR fusion complexes of the invention contain a TCR molecule that is sufficiently truncated so the TCR fusion complex can be secreted into culture medium after expression. Thus, a truncated TCR fusion complex will not include regions rich in hydrophobic residues, typically the transmembrane and cytoplasmic domains of the TCR molecule. Thus, for example, for a preferred truncated DR1 TCR molecule of the invention, preferably from about residues 199 to 237 of the b chain and from about residues 193 to 230 of the a chain of the TCR molecule are not included in the truncated TCR fusion complex.

The present TCR fusion and conjugate complexes are suitable for in vitro or in vivo use with a variety of cells that are infected or that may become infected by one or more diseases.

Methods

Therapeutic

Included in the invention are methods for preventing or treating disease in a patient in which the diseased cells express a disease associated antigen, the method comprising administering to the patient a soluble fusion protein complex comprising a biologically active polypeptide recognizing a disease-associated antigen, forming a specific binding complex (bridge) between antigen-expressing diseased cells and IL-15R-expressing immune cells sufficient to localize the immune cells, and damaging or killing the disease cells sufficient to prevent or treat the disease in the patient.

Included are methods for preventing or treating disease in a patient in which the diseased cells express a disease associated antigen, the method comprising mixing immune cells bearing the IL-15R chains with a soluble fusion protein complex comprising a biologically active polypeptide recognizing a disease-associated antigen, for example a peptide/MHC complex, administering to the patient the immune cell-fusion protein complex mixture, forming a specific binding complex (bridge) between antigen-expressing diseased cells and IL-15R-expressing immune cells sufficient to localize the immune cells, and damaging or killing the disease cells sufficient to prevent or treat the disease in the patient.

25

Also included in the invention are methods for killing a target cell, the method comprising contacting a plurality of cells with a soluble fusion protein complex, where the plurality of cells further comprises immune cells bearing the IL-15R chains recognized by the IL-15 domain of claim 1 and the target cells bearing an antigen recognized by at least one of the biologically active polypeptides as described herein, forming a specific binding complex (bridge) between the antigen on the target cells and the IL-15R chains on the immune cells sufficient to bind and activate the immune cells; and killing the target cells by the bound activated immune cells.

Also included in the inventions are methods to increase in vivo half life of IL-15 and/or enhance its ability to stability bind immune cells (e.g. increase cell surface residency time) through generation of a soluble fusion protein complex. For example, evaluation of the pharmacokinetic parameters and cell surface residency time of the fusion protein complex are conducted and compared to IL-15, as described herein. Fusion protein complexes with an increased serum half life or cell surface residency time are preferable as based on their improved therapeutic utility.

Examples of diseases that can be treated include, but are not limited to, neoplasia, including cancer, or viral infection. By "neoplasia" is meant any disease that is caused by or results in inappropriately high levels of cell division, inappropriately low levels of apoptosis, or both. For example, cancer is an example of a neoplasia. Examples of cancers include, without limitation, leukemias (e.g., acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia), polycythemia vera, lymphoma (Hodgkin's disease, non-Hodgkin's disease), Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors such as sarcomas and carcinomas (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, uterine cancer, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma). Lymphoproliferative disorders are also considered to be proliferative diseases.

Also included are methods of stimulating immune responses in a mammal comprising administering to the mammal an effective amount of the soluble fusion protein complex or IL-15 variant as described herein. Also included are methods of suppressing immune responses in a mammal comprising administering to the mammal an effective amount of the soluble fusion protein complex or IL-15 variant as described herein. In the case of immune suppression, a fusion protein complex or IL-15 variant comprising IL-15 antago-

26

nists or IL-15 domains that lack the ability to bind the IL-15 $\beta\gamma_c$ complex may be particularly advantageous.

As an illustration of the use of the fusion protein complex therapeutics, a cultured cell can be infected by a pathogen of a single serotype. The infected cell is then contacted by a specified fusion protein complex in vitro. As discussed previously, the fusion protein complex is configured so that the toxic domain is presented to the infected cell by the association of the TCR. After providing for introduction of the bioactive molecule to the cell (generally less than about 30 minutes), the cells are allowed to cause a desired effect for a time period of about up to about 2 to 24 hours, typically about 18 hours. After this time, the cells are washed in a suitable buffer or cell medium and then evaluated for viability. The time allotted for cell killing or injury by the fusion protein complex will vary with the particular effector molecule chosen. However viability can often be assessed after about 2 to 6 hours up to about 24 hours. As will be explained in more detail below, cell viability can be readily measured and quantified by monitoring uptake of certain well-known dyes (e.g., trypan blue) or fluors.

Cells incubated with the fusion protein complex of the present invention can be assayed for viability by standard methods. In one exemplary approach, cell viability can be readily assayed by measuring DNA replication following or during incubation. For example, a preferred assay involves cell uptake of one or more detectably labeled nucleosides such as radiolabelled thymidine. The uptake can be conveniently measured by several conventional approaches including trichloroacetic acid (TCA) precipitation followed by scintillation counting. Other cell viability methods include well-known trypan blue exclusion techniques or WST-1-based proliferation assays.

The TCR molecules of the fusion complexes of the invention suitably correspond in amino acid sequence to naturally occurring TCR molecules, e.g. TCR molecules of a human, mouse or other rodent, or other mammal.

Accordingly, one treatment method of the invention for inhibition of an autoimmune or inflammatory response would include a fusion protein complex which comprises a T cell receptor or antibody with binding specificity to a disease associated antigen. Preferably, a "truncated" soluble TCR complex is administered, i.e. the TCR complex does not contain a transmembrane portion. The fusion protein complex also comprises an IL-15 variants that functions as an IL-15 antagonist to suppress the unwanted immune response. Following administration, the TCR or antibody domain targets the fusion protein complex to the disease site where the IL-15 antagonist suppresses the autoimmune or inflammatory response. Such fusion protein complex may particularly useful for treatment of allergies, autoimmune diseases such as multiple sclerosis, insulin-dependent diabetes mellitus and rheumatoid arthritis or transplant rejection. Similar non-targeted approaches could be carried out using antagonist IL-15 variants as non-fusion proteins.

Another treatment method of the invention for induction of an immune response provides for the administration of an effective amount of a protein fusion complexes of the invention in the presence of any costimulatory effector molecule such as a cytokine to thereby induce a desired immune response at the location of the presented antigen which binds the biologically active polypeptide.

Different therapies of the invention also may be used in combination as well as with other known therapeutic agents such as anti-inflammatory drugs to provide a more effective treatment of a disorder. For example, immunosuppressive protein fusion complexes or IL-15 variants can be used in

combination with anti-inflammatory agents such as corticosteroids and nonsteroidal drugs for the treatment of autoimmune disorders and allergies.

Compounds of the invention will be especially useful to a human patient who has or is suspected of having a malignant disease, disorder or condition. Compounds of the invention will be particularly useful in targeting particular tumor antigens in human patients. Specific examples of diseases which may be treated in accordance with the invention include cancers, e.g. breast, prostate, etc, viral infections, e.g. HCV, HIV, etc. as well as other specific disorders of conditions mentioned herein.

Without wishing to be bound by theory, it is believed the multiple and distinct covalently linked compounds of this invention (i.e. at least IL-15 in combination with at least one TCR) can significantly enhance efficacy of the IL-15, e.g., by increasing targeting of IL-15 to target antigen in subject individuals.

Moreover, by virtue of the covalent linkage, the conjugates of the invention present the IL-15 and the TCR to the subject cell essentially simultaneously, an effect that may not be readily achieved by administering the same compounds in a drug "cocktail" formulation without covalently linking the compounds.

It also has been reported that treatment with treatment with one drug can in turn sensitize a patient to another drug. Accordingly, the essentially simultaneous presentation to the subject cell of IL-15 and TCR via a conjugate of the invention may enhance drug activity, e.g., by providing synergistic results and/or by enhancing production an immune response.

Diagnostic

High affinity or multivalent TCR proteins specific for a particular pMHC ligand are useful in diagnosing animals, including humans believed to be suffering from a disease associated with the particular pMHC. The fusion protein complexes of the present invention are useful for detecting essentially any antigen, including but not limited to, those associated with a neoplastic condition, an abnormal protein, an autoimmune disease or an infection or infestation with a bacterium, a fungus, a virus, a protozoan, a yeast, a nematode or other parasite.

In one such method for detecting a tumor or virally infected cell or tissue in a subject, wherein the cell or tissue comprises a tumor or virus-associated peptide antigen presented on the cells or tissue in the context of an MHC complex, comprises: a) administering to the subject a soluble fusion protein complex of the invention comprising a soluble TCR under conditions that form a specific binding complex between the presented peptide antigen and the TCR; and b) detecting the specific binding complex as being indicative of a tumor or virally infected cell or tissue comprising the presented tumor or viral-associated peptide antigen.

The fusion protein complexes can also be used in the diagnosis of certain genetic disorders in which there is an abnormal protein produced. Exemplary applications for fusion protein complexes are in the diagnosis and treatment of autoimmune diseases in which there is a known pMHC. Type I diabetes is relatively well characterized with respect to the autoantigens which attract immune destruction. Multiple sclerosis, celiac disease, inflammatory bowel disease, Crohn's disease and rheumatoid arthritis are additional candidate diseases for such application.

The fusion protein complexes of the present invention comprising IL-15 variant polypeptides may be particularly useful in these applications. For example, for a fusion protein complex comprising TCR molecules, interactions between the IL-15 variant domain and the IL-15Ra polypeptide gen-

erate multivalent TCR molecules with enhanced antigen binding activity, as disclosed herein. Moreover, the IL-15 variant contains amino acid changes that potentially eliminate binding to cells bearing IL-15R $\beta\gamma$ C receptors, thereby reducing non-specific or non-targeted binding to immune cells. As a results, improved detection of TCR-specific antigens can be achieved with such fusion protein complexes. Additionally fusion protein complexes of the invention can be further multimerized via peptide tags sequences or conjugation to detectable labels, as disclosed herein.

Dosage and Administration

Administration of compounds of the invention may be made by a variety of suitable routes including oral, topical (including transdermal, buccal or sublingal), nasal and parenteral (including intraperitoneal, subcutaneous, intravenous, intradermal or intramuscular injection) with oral or parenteral being generally preferred. It also will be appreciated that the preferred method of administration and dosage amount may vary with, for example, the condition and age of the recipient.

Compounds of the invention may be used in therapy alone or in conjunction with other medicaments such those with recognized pharmacological activity to treat the desired indications. Exemplary medicaments include recognized therapeutics such as surgery, radiation, chemotherapy and other forms of immunotherapy (e.g. vaccines, antibody based therapies). The compounds of this invention can be administered before, during or after such therapies as needed.

While one or more compounds of the invention may be administered alone, they also may be present as part of a pharmaceutical composition in mixture with conventional excipient, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, oral or other desired administration and which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof. Pharmaceutical compositions of the invention in general comprise one or more fusion protein complex or IL-15 variant of the invention or DNA constructs coding for such compounds together with one or more acceptable carriers. The carriers must be "acceptable" in the sense of being compatible with other ingredients of the formulation and not deleterious to the recipient thereof. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds.

For parenteral application, particularly suitable are solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Ampules are convenient unit dosages.

For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

Therapeutic compounds of the invention also may be incorporated into liposomes. The incorporation can be carried out according to known liposome preparation procedures, e.g. sonication and extrusion. Suitable conventional methods of liposome preparation are also disclosed in e.g. A. D. Bangham et al., *J. Mol. Biol.*, 23:238-252 (1965); F. Olson et al., *Biochim. Biophys. Acta*, 557:9-23 (1979); F. Szoka et al., *Proc. Nat. Acad. Sci.*, 75:4194-4198 (1978); S. Kim et al., *Biochim. Biophys. Acta*, 728:339-348 (1983); and Mayer et al., *Biochim. Biophys. Acta*, 858:161-168 (1986).

The invention also provides methods for invoking an immune response in a mammal such as a human, including vaccinating a mammal such as a human against an infectious agent or a targeted disorder such as cancer.

These methods comprise administering to a mammal an effective amount of a DNA sequence that comprises a DNA vector that codes for a fusion protein complex or IL-15 variant of the invention. Preparation of expression vectors of fusion protein complexes and IL-15 variants is described above and in the Examples which follow. Methods for administration of plasmid DNA, uptake of that DNA by cells of the administered subject and expression of protein has been reported. See Ulmer, J. B., et al., *Science* (1993) 259: 1745-1749.

DNA vectors that encode fusion protein complexes and IL-15 variants of the invention are suitably administered to a mammal including a human preferably by intramuscle injection. Administration of cDNA to skeletal muscle of a mammal with subsequent uptake of administered expression vector by the muscle cells and expression of protein encoded by the DNA has been described by Ulmer et al. and represents an exemplary protocol [Ulmer, J. B., et al., *Science* 259: 1745-1749]. The optimal dose for a given therapeutic application can be determined by conventional means.

In addition to treatment of human disorders, fusion protein complexes and IL-15 variants of the invention and DNA constructs of the invention that encode such molecules will have significant use for veterinary applications, e.g., treatment of disorders of livestock such as cattle, sheep, etc. and pets such as dog and cats.

It will be appreciated that actual preferred amounts of a given fusion protein complex and IL-15 variant of the invention or DNA construct coding for same used in a given therapy will vary according to the particular active compound or compounds being utilized, the particular compositions formulated, the mode of application, the particular site of administration, the patient's weight, general health, sex, etc., the particular indication being treated, etc. and other such factors that are recognized by those skilled in the art including the attendant physician or veterinarian. Optimal administration rates for a given protocol of administration can be readily determined by those skilled in the art using conventional dosage determination tests conducted e.g. with regard to the foregoing guidelines and the assays disclosed herein.

EXAMPLES

It should be appreciated that the invention should not be construed to be limited to the examples that are now described; rather, the invention should be construed to include any and all applications provided herein and all equivalent variations within the skill of the ordinary artisan.

Example 1

Design of a Fusion Protein Complex Comprising scTCR/huIL15 and scTCR/huIL15R α Fusion Proteins

It has been established that the IL-15 stably binds to the extracellular domain of the IL-15R α and that the resulting

complex is capable of modulating (i.e. either stimulating or blocking) immune responses via the intermediate or high affinity IL-15R complex (Mortier, E. et al., 2006, *J. Biol. Chem.*, 281: 1612-1619; Stoklasek, T. et al., 2006, *J. Immunol* 177: 6072-6080; Rubinstein, M. P. et al., 2006, *Proc Natl Acad Sci USA* 103: 9166-9171, Waldmann, T. A., 2006, *Nat Rev Immunol* 6: 595-601). In addition, it has been demonstrated that single-chain TCR or antibody polypeptides can be fused to cytokines and other immune effector domains and that such bispecific fusion molecules retain functional activity of both fusion domains (Belmont, H. J. et al., 2006, *Clin Immunol* 121: 29-39; Card, K. F. et al., 2004, *Cancer Immunol Immunother* 53: 345-357; Mosquera, L. A. et al., 2005, *J Immunol* 174: 4381-4388; Penichet, M. L., 1997, *Hum Antibodies* 8: 106-118). Further, it has been shown that multivalent forms of the TCR provide enhanced binding to their ligands [(9)] (Zhu, X., H. J., 2006, *J Immunol* 176: 3223-3232). Therefore a feature of the invention provides for a fusion protein complex comprising at least one fusion protein wherein a first TCR polypeptide is fused to IL-15 and at least one fusion wherein a second TCR polypeptide is fused to the extracellular domain of IL-15R α , such that the two fusion proteins form a complex through binding interactions between the IL-15 and IL-15R α domains. In such a fusion protein complex, the TCR polypeptides can be the same or different and in either single-chain or heterodimeric format.

An example of a fusion protein complex containing single-chain TCR polypeptides is shown schematically in FIG. 1A. In this fusion protein complex, the multivalent TCR domains provide increased binding avidity/affinity for their ligands. Exemplary ligands include, but are not limited to, peptide/MHC complexes. The IL-15/IL-15R α domains provide immunomodulatory activity. Representative fusion protein constructs comprising the fusion protein complex are schematically shown in FIG. 1B. In these constructs the TCR polypeptide is a single-chain TCR (264scTCR) comprised of TCR-V α and TCR-V β -C β domains linked by a peptide linker sequence ((G₄S)₄) (SEQ ID NO: 3). The scTCR polypeptide is fused to either the IL-15 or IL-15R α domains, directly or via a peptide linker sequence. Proceeding the scTCR polypeptide is a signal peptide (or leader peptide) sequence that permits soluble expression. The signal peptide is subsequently cleaved during protein transport to generate the mature fusion protein. In other examples of the fusion protein complex, an antibody domain can substitute a TCR domain depicted in FIGS. 1A and 1B. Such an antibody can be in a single-chain or heteromultimeric format. For any of the fusion protein complexes described above, sequences can be human or non-human, for example, but not limited to mouse. These sequences can be employed for part or all of the fusion protein domains. In addition, the arrangement of the domains can vary so long as the fusion proteins remain soluble and functional.

Example 2

Construction of the c264scTCR/huIL15 Gene Fusion in an Expression Vector

Isolation and characterization of TCR genes for the p53 (aa264-272)-specific TCR were described previously (Belmont, H. J. et al., 2006, *Clin Immunol* 121: 29-39; Card, K. F. et al., 2004, *Cancer Immunol Immunother* 53: 345-357; Mosquera, L. A. et al., 2005, *J Immunol* 174: 4381-4388). To obtain the human IL15 and IL15R α genes, human PBMC were isolated from 200 mL of blood of a donor (Lot#2238789, Community Blood Bank, Miami, Fla.) with

31

HISTOPAGUE-1077 (Sigma). The cells (1.5×10^7) were activated by 30 ng/ml of PMA (Sigma), 200 ng/ml of ionomycin, and 220 ng/ml of recombinant human IL-2 in IMDM containing 10% FBS in a CO₂ incubator for 10 days. The activated cells (1×10^7 per mL) were frozen at -70 C for the further applications. To purify the total RNA from the activated PBMC, RNEASY PLUS MINI (Qiagen) was used according to the manufacturer's protocol. Human IL15 gene containing the coding region and a portion of 5' and 3' flanking regions was amplified from the total RNA with the front primer

5' -CACCTTGCCATAGCCAGCTCTTC-3' (SEQ ID NO: 19)

and the back primer

5' -GTCTAAGCAGCAGAGTGATGTTTG-3' (SEQ ID NO: 20)

by SUPERSRIPT III One-Step RT-PCR Platinum Tag HiFi (Invitrogen) according to the following conditions: for RT; 55 C 30 min; 94 C, 2 min; for amplifying cDNA; 94 C, 30s; 53 C, 30s; 68 C, 1 min; $\times 40$ cycles; 68 C, 5 min. The 600 bp human IL15 PCR-cDNA product was separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The gene of the mature human IL15 protein was amplified from the 600 bp human IL15 cDNA with the front primer

(SEQ ID NO: 21)
5' -TGTTAACAACCTGGGTGAATGTAATAAGTG-3'

and the back primer

(SEQ ID NO: 22)
5' -ACGCGTTTATCAAGAAGTGTGATGAACATTGGAC-3'

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 1 min; 63 C, 1 min; 72 C 1 min; $\times 35$ cycles; 72 C, 10 min. The mature human IL15 protein gene was gel-purified and cloned into the shuttle vector, pcDNA3.1 Directional TOPO Expression Vector (Invitrogen), with the TOPO reaction according to the manufacturer's protocol. The clone containing the mature human IL15 protein gene insert was identified based on the diagnostic PCR with the front primer

5' -TGTTAACAACCTGGGTGAATGTAATAAGTG-3' (SEQ ID NO: 21)

and the back primer

(SEQ ID NO: 22)
5' -ACGCGTTTATCAAGAAGTGTGATGAACATTGGAC-3'

by RedTag (Sigma) under the following condition: 94 C, 1 min; 63 C, 1 min; 72 C, 1 min; $\times 35$ cycles; 72 C, 10 min. The sequence of the correct clone was verified by DNA sequencing with GenomeLab Dye Termination Cycle Sequencing with a QUICK START KIT (Beckman Coulter) according to the manufacturer's protocol. The mature human IL15 protein gene was removed from the shuttle vector by digestion with HpaI and MluI and ligated into an expression vector pNEF38-c264scTCR which had been digested with HpaI and MluI. The pNEF38-c264scTCR expression vector contains the gene fragment encoding an immunoglobulin light chain leader (or secretory signal) sequence linked to the p53 (aa264-272) peptide-specific soluble chimeric single-chain

32

TCR protein (c264scTCR) (Belmont, H. J. et al., 2006, *Clin Immunol* 121: 29-39). The vector also contains 5' regulatory/enhancer regions, transcription regulatory and promoter regions, translational regulatory/initiation/termination sequences including a Kozak consensus sequence and poly-A termination region, and 3' regulatory regions with putative matrix attachment regulatory elements. The vector also contains DNA sequences allowing selective growth in mammalian cells (SV40 promoter/neoR gene/poly-A) and bacterial cells (ori/amp gene). Cloning of the DNA fragment encoding mature human IL15 protein into the pNEF38-c264scTCR vector resulted in a c264scTCR/huIL15 fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V- α -peptide linker -264 TCR V- β -human TCR C- β -human IL-15. The resulting vector (pNEF38-c264scTCR/huIL15), shown in FIG. 2A, was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The sequences of the c264scTCR/huIL15 fusion gene and protein (including the leader sequence) are shown at FIG. 2B and FIG. 2C, respectively.

Example 3

Construction of the c264scTCR/huIL15 Gene Fusion Containing a Mutated Human IgG1 Hinge Region in an Expression Vector

Construction of the pNEF38-c264scTCR/huIL15 vector was described at Example 2. A mutated hinge region from human IgG1 H chain where three cysteine residues were substituted with three serine residues was used to link c264scTCR and huIL15. The hinge region was mutated and amplified from 264scTCR/IgG1 gene described previously (Mosquera, L. A. et al., 2005, *J Immunol* 174: 4381-4388) with the front primer

5' -TGTTGGGTTAACGAGCCCAATCTTCTG-3' (SEQ ID NO: 23)

and the back primer

5' -ATTATTACGCGTTGGAGACGGTGGAGATG-3' (SEQ ID NO: 24)

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 30 sec; 65 C, 30 sec; 70 C 1 min; $\times 35$ cycles; 72 C, 10 min. The 70 bp mutated human IgG1 hinge PCR-cDNA product was separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The mutated hinge region gene was digested with HpaI and MluI and ligated into pNEF38-c264scTCR which had been digested with HpaI and MluI. The clone containing the mutated hinge region gene insert was identified based on the diagnostic PCR with the front primer

(SEQ ID NO: 25)
5' -TGAGTGATCGATACCACCATGGAGACAGACAC-3'

and the back primer

5' -ATTATTACGCGTTGGAGACGGTGGAGATG-3' (SEQ ID NO: 24)

by RedTag (Sigma) under the following condition: 94 C, 30 sec; 64 C, 30 sec; 70 C 1 min; $\times 35$ cycles; 72 C, 10 min. The

33

huIL15 was amplified from pNEF38-c264scTCR/huIL15 vector described at Example 2 with the front primer

5' - TGGTGGACGCGTAACTGGGTGAATG - 3' (SEQ ID NO: 26)

and the back primer

(SEQ ID NO: 27)
5' - TGGTGGTCTAGAATTATCAAGAAGTGTGATG - 3'

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 30 sec; 65 C, 30 sec; 70 C 1 min; $\times 35$ cycles; 72 C, 10 min. The 380 bp huIL15 PCR-cDNA product was separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The huIL15 gene was digested with MluI and XbaI and ligated into pNEF38-c264scTCR containing mutated hinge gene which had been digested with MluI and XbaI. The clone containing the huIL15 gene insert was identified based on the diagnostic PCR with the front primer

(SEQ ID NO: 25)
5' - TGAGTGATCGATACCACCATGGAGACAGACAC - 3'

and the back primer

(SEQ ID NO: 27)
5' - TGGTGGTCTAGAATTATCAAGAAGTGTGATG - 3'

by RedTag (Sigma) under the following condition: 94 C, 30 sec; 64 C, 2 min; 70 C 2 min; $\times 35$ cycles; 72 C, 10 min. The sequence of the correct clone was verified by DNA sequencing with GenomeLab Dye Termination Cycle Sequencing with a QUICK START KIT (Beckman Coulter) according to the manufacturer's protocol. The pNEF38-c264scTCR expression vector is described above at Example 2. Cloning of the DNA fragment encoding mutated human IgG1 hinge region and mature human IL15 protein into the pNEF38-c264scTCR vector resulted in a c264scTCR-hmt-huIL15 fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V- α -peptide linker -264 TCR V- β -human TCR C- β -mutated human IgG1 hinge-human IL-15. The resulting vector (pNEF38-c264scTCR-hmt-huIL15), shown in FIG. 3A, was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The sequences of the c264scTCR-hmt-huIL15 fusion gene and protein (including the leader sequence) are shown at FIG. 3B and FIG. 3C, respectively.

Example 4

Construction of the c264scTCR/huIL15R α AE3 Gene Fusion in an Expression Vector

The total RNA of PBMC was prepared as described above. Human IL15R α gene containing coding region and a portion of 5' and 3' flanking regions was amplified from the total RNA of the PBMC with the front primer

5' - AGTCCAGCGGTGCTCCTGTGG - 3' (SEQ ID NO: 28)

and the back primer

(SEQ ID NO: 29)
5' - TGACGCGTTTAAGTGGTGTGCTGTGCCCTG - 3'

34

by SUPERScript III One-Step RT-PCR Platinum Tag HiFi (Invitrogen) according to the following condition: for RT; 55 C, 30 min; 94 C, 2 min; for amplifying cDNA; 94 C, 1 min; 66 C, 1 min; 72 C, 1 min; $\times 35$ cycles; 72 C, 5 min. The 970 bp human IL15 R α PCR cDNA product was separated by electrophoresis on a 1% agarose gel and isolated. The cDNA was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The human IL15 R α extracellular domain gene was amplified from the 970 bp human IL15 R α cDNA with the front primer

5' - TGGTTAACATCAGTGCCTCCCCCATG - 3' (SEQ ID NO: 30)

and the back primer

(SEQ ID NO: 29)
5' - TGACGCGTTTAAGTGGTGTGCTGTGCCCTG - 3'

by PfuULTRA (Stratagene) under following PCR conditions: 94 C, 1 min; 72 C 2 min; $\times 35$ cycles; 72 C, 10 min. The human IL15 R α extracellular domain gene was gel-purified and cloned into the shuttle vector, pcDNA3.1 Directional TOPO Expression Vector (Invitrogen), by TOPO reaction according to the manufacturer's protocol. The clone containing the correct human IL15 R α extracellular domain gene insert was chosen based on diagnostic PCR and reconfirmed by DNA sequencing with the GenomeLab Dye Termination Cycle Sequencing with a Quick Start Kit according to the manufacturer's protocol. The gene was determined to be human IL15 R α AE3 extracellular domain gene. The human IL15 R α AE3 extracellular domain gene was removed from the shuttle vector by digestion with HpaI and MluI and ligated into pNEF38-c264scTCR which had been digested with HpaI and MluI. Cloning of the DNA fragment encoding the human IL15 R α AE3 extracellular domain into the pNEF38-c264scTCR vector resulted in a c264scTCR/huIL15R α fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V- α -peptide linker -264 TCR V- β -human TCR C- β -IL15 R α AE3 extracellular domain. The resulting vector (pNEF38-c264scTCR/huIL15R α AE3), shown in FIG. 4A, containing the correct IL15 R α AE3 extracellular domain gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The sequences of the c264scTCR/huIL15 R α AE3 gene and protein are shown at FIG. 4B and FIG. 4C, respectively.

Example 5

Construction of c264scTCR/huIL15R α Sushi Gene Fusion in an Expression Vector

The total RNA of PBMC was prepared as described above. Human IL15R α Sushi gene was amplified from the 970 bp human IL15 R α cDNA (see Example 3) with the front primer

5' - TGGTTAACATCAGTGCCTCCCCCATG - 3' (SEQ ID NO: 30)

and the back primer

(SEQ ID NO: 31)
5' - TTGTTGACGCGTTTATCTAATGCATTGAGACTGG - 3'

by PfuULTRA (Stratagene) under following PCR conditions: 94 C, 1 min; 66 C, 1 min; 70 C, 1 min; $\times 35$ cycles; 72 C, 10 min. The PCR product of human IL15R α I Sushi gene was gel-purified and digested with HpaI and MluI. The gene was ligated into pNEF38-c264scTCR which had been digested

35

with HpaI and MluI. Cloning of the DNA fragment encoding the human IL15R α Sushi domain into the pNEF38-c264scTCR vector resulted in a c264scTCR/huIL15R α fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V α -peptide linker -264 TCR V β -human TCR C β -human IL15R α Sushi. The resulting vector, shown in FIG. 5A, containing the correct human IL15R α Sushi gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The sequences of the c264scTCR/huIL15 R α Sushi gene and protein are shown at FIG. 5B and FIG. 5C, respectively.

Example 6

Construction of c264scTCR/huIL15R α Sushi Gene Fusion Containing a Mutated Human IgG1 Hinge Region in an Expression Vector

Construction of the pNEF38-c264scTCR/huIL15R α Sushi vector was described above. A mutated hinge region from human IgG1 H chain where three cysteine residues were replaced by three serine residues was used to link c264scTCR and huIL15R α Sushi. The hinge region was mutated, amplified, ligated, and verified as above. The huIL15R α Sushi was amplified from pNEF38-c264scTCR/huIL15R α Sushi vector described above with the front primer

5' -TAATAAACGCGTATCACGTGCCCTC-3' (SEQ ID NO: 32)

and the back primer

(SEQ ID NO: 33)
5' -TGGTGGTCTAGATTATCATCTAATGCATTG-3'

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 30 sec; 65 C, 30 sec; 70 C 1 min; $\times$ 35 cycles; 72 C, 10 min. The 250 bp huIL15R α Sushi PCR-cDNA product was separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The huIL15R α Sushi gene was digested with MluI and XbaI and ligated into pNEF38-c264scTCR containing mutated hinge gene which had been digested with MluI and XbaI. The clone containing the huIL15 gene insert was identified based on the diagnostic PCR with the front primer

5' -TGGTGGGTTAACGAGCCCAATCTTCTG-3' (SEQ ID NO: 23)

and the back primer

(SEQ ID NO: 33)
5' -TGGTGGTCTAGATTATCATCTAATGCATTG-3'

by RedTag (Sigma) under the following condition: 94 C, 30 sec; 65 C, 1 min; 70 C 1 min; $\times$ 35 cycles; 72 C, 10 min. The sequence of the correct clone was verified by DNA sequencing with GenomeLab Dye Termination Cycle Sequencing with a QUICK START KIT (Beckman Coulter) according to the manufacturer's protocol. The pNEF38-c264scTCR expression vector is described above. Cloning of the DNA fragment encoding mutated human IgG1 hinge region and human IL15R α Sushi protein into the pNEF38-c264scTCR vector resulted in a c264scTCR-hmt-huIL15R α Sushi fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V α -peptide linker -264 TCR

36

V β -human TCR C β -mutated human IgG1 hinge-human IL15R α Sushi. The resulting vector (pNEF38-c264scTCR-hmt-huIL15R α Sushi), shown in FIG. 6A, was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The sequences of the c264scTCR-hmt-huIL15R α Sushi fusion gene and protein (including the leader sequence) are shown at FIG. 6B and FIG. 6C, respectively.

Example 7

Construction of the c264scTCR/huIL15R α Sushi and c264scTCR/huIL15 Genes in a Single Expression Vector

To achieved expression of two fusion proteins of the invention in a single host cell, the genes encoding c264scTCR/huIL15R α Sushi and c264scTCR/huIL15 were cloned into a single expression vector. The c264scTCR/huIL15R α Sushi gene was amplified from the template described in Example 5 by PfuUltra (Stratagene) with the front primer 5'-TGAGTGTCCGGAACCACCATGGAGACAGACAC-3' (SEQ ID NO: 34) and the back primer 5'-TTGTTGGCGGCCGTATCATCTAATGCATTGAG-3' (SEQ ID NO: 35) under the following condition: 94 C, 1 min; 68 C, 1 min; 72 C, 2 min; $\times$ 35 cycles; 72 C, 10 min. The PCR product of c264scTCR/huIL15R α Sushi gene was gel-purified, digested with BspEI and NotI and ligated into the pSUN34R1 expression vector which had been digested with BspEI and NotI. The pSUN34R1 expression vector contains two sites for cloning genes-of-interest as well as 5' regulatory/enhancer regions, transcription regulatory and promoter regions, translational regulatory/initiation/termination sequences including a Kozak consensus sequence and poly-A termination region, and intron and 3' regions with regulatory elements. This vector also contains DNA sequences allowing selective growth in mammalian cells (SV40 promoter/neoR gene/poly-A) and bacteria (ori/amp gene). The vector containing the correct c264scTCR/IL15R α Sushi gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The c264scTCR/huIL15 gene was amplified from the template described in Example 2 by PfuUltra (Stratagene) with the front primer

(SEQ ID NO: 25)
5' -TGAGTGATCGATACCACCATGGAGACAGACAC-3'

and the back primer

(SEQ ID NO: 36)
5' -TGAGTGTTCGAATTATCAAGAAGTGTGTGATGAAC-3'

under the following condition: 94 C, 1 min; 65 C, 1 min; 72 C, 2 min; $\times$ 35 cycles; 72 C, 10 min. The PCR product of c264scTCR/huIL15 gene was gel-purified, digested with ClaI and Csp45I and ligated into pSUN34R1-c264scTCR/huIL15R α Sushi expression vector which had been digested with ClaI and Csp45I. The resulting vector (pSun-c264scTCRIL15/c264scTCRIL15R α Sushi), shown in FIG. 7, containing the correct c264scTCR/huIL15 gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. This vector contains both c264scTCR/huIL15R α Sushi and c264scTCR/huIL15 genes.

37

Example 8

Construction of c264scTCR/huIL15R α ΔE3 and
c264scTCR/huIL15 Genes in a Single Expression
Vector

The c264scTCR/huIL15R α ΔE3 fusion gene was amplified from the template described in Example 4 by PfuUltra (Stratagene) with the front primer

(SEQ ID NO: 34)
5' - TGAGTGTCCGGAACCCATGGAGACAGACAC - 3'

and the back primer

5' - TTGTTGGCGGCCGCTTATCAAGTGGTGTGCTG - 3'
(SEQ ID NO: 37)

under the following condition: 94 C, 1 min; 68 C, 1 min; 72 C, 2 min; $\times 35$ cycles; 72 C, 10 min. The PCR product of c264scTCR/huIL15R α ΔE3 gene was gel-purified, digested with BspEI and NotI and ligated to the expression vector pSUN34R1 which had been digested with BspEI and NotI. The vector containing the correct c264scTCR/huIL15R α ΔE3 gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. The c264scTCR/huIL15 gene was amplified and cloned into the expression vector as described on Example 7. The resulting vector (pSun-c264scTCRIL15/c264scTCRIL15R α ΔE3), shown in FIG. 8, containing the correct c264scTCR/huIL15 gene insert was identified based on the diagnostic PCR and reconfirmed by DNA sequencing. This vector contains both c264scTCR/huIL15R α ΔE3 and c264scTCR/huIL15 genes.

Example 9

Generation of Transfected Host Cell Lines Producing
Fusion Proteins

The expression vectors can be introduced into a variety of host cell lines by several different transformation, transfection or transduction methods. In one such method, CHO-K1 cells (5×10^4) were seeded in a 6-well plate and cultured overnight in a CO₂ incubator. The cells were transfected with 5 μ g of expression vector containing the TCR/IL15 and/or TCR/IL15R α fusion genes using 10 μ L of Mirus TransIT-LT1 reagent (Mirus) according to the manufacturer's protocol. The cells were selected with 4 mg/mL of G418 (Invitrogen) one day after the transfection. The G418 resistant cells were expanded and TCR fusion protein expressing cells were enriched by 3-5 rounds of MACS selection as described below. The cells were detached in 10 mM EDTA and washed once with IMDM containing 10% FBS. Cells were resuspended (10^7 cells in 100 μ L) and incubated with 5 μ g of R-Phycoerythrin (PE) conjugated p53 (aa264-272)/HLA-A2 tetramer reagent for 15 min at 4 C. The cells were washed once and incubated with anti-PE antibody conjugated magnetic beads (Miltenyi Biotec) for 15 min at 4 C. The cells were loaded to a magnetic column (in a magnetic field) and the unbound cells were removed with wash buffer (PBS containing 0.5% BSA). The column-bound cells were eluted with IMDM containing 10% FBS after the column had been removed from the magnetic field. This procedure allows enrichment of fusion protein-expressing cells based on the transient display of the soluble fusion protein on the cell surface during the production/secretion process. The cell sur-

38

face association of the fusion proteins was monitored after each enrichment. Levels of cell surface-bound fusion proteins determined by flow cytometry were compared to levels of soluble fusion proteins present in the cell culture media as determined by ELISA. An example of the comparison is shown in FIGS. 9A and 9B. In this example, CHO-K1 cells transfected with pNEF38-c264scTCR/huIL15R α Sushi were enriched by MACS for one to five times and were then seeded (1×10^6 cells/well) on a 6-well plate. After 24 hours, cells were then detached with 10 mM EDTA, washed once with IMDM+10% FBS, and stained (at 2×10^5 cells/100 μ L of IMDM+10% FBS) with 0.6 μ g of PE-conjugated p53 (aa264-272)/HLA-A2 tetramer or same amount of control PE-conjugated CMVpp65 (aa495-503)/HLA-A2 tetramer for 30 min at 4 C. Cells were washed once and analyzed for levels of cell surface associated soluble fusion protein by flow cytometry, as shown in FIG. 9A. The level of soluble fusion protein secreted into the cell culture medium was also determined by TCR-specific ELISA with a capture antibody, anti-human TCR C β antibody (BF1), and a detection antibody, biotinylated anti-human TCR C β antibody (W4F) described previously (5), as shown in FIG. 9B. The results indicate that the magnetic bead-based enrichment process yielded transfectants that produced increased levels of soluble fusion protein. The enriched transfectant cells were then subcloned three times by the limiting dilution and production cell lines were screened based on the level of soluble fusion protein secreted into the culture media (determined by ELISA described above). Production cell lines were expanded and grown in IMDM+10% FBS or serum-free media under conditions (i.e. flasks, spinners, fermenters, bags, bottles) suitable to generate the soluble fusion protein.

In some cases, host cells were co-transfected with different expression vectors to generate transfectants capable of expressing multiple fusion proteins. Transfectants expressing one fusion protein could also be re-transfected with a one or more expression vectors to generate transfectants expressing multiple fusion proteins. Cells were also transfected with an expression vector containing more than one fusion protein genes, as exemplified in Examples 7 and 8, to generate a transfectant expressing multiple fusion proteins. The resulting cells could be used to produce the multi-component fusion protein complexes of the invention as soluble molecules in the cell culture medium.

High levels of fusion protein or fusion protein complex production can also be achieved through cell transfection and selection methods described in U.S. Ser. No. 09/204,979.

Example 10

Purification of the TCR/IL15 and TCR/IL15R α
Fusion Proteins or Fusion Protein Complexes

Soluble fusion proteins or fusion protein complexes of the invention can be purified from the host cells or cell culture media using a variety of methods, including by selective partitioning or solubility in solvents or by separation (i.e. via chromatography) based on charge, hydrophobicity, hydrophilicity, size, and/or selective or semi-selective binding to a ligand. Soluble fusion proteins or fusion protein complexes can be generated from insoluble materials through use of the appropriate protein folding conditions. In one example, c264scTCR/IL15 fusion protein was purified from cell culture media by affinity chromatography using an antibody (BF1) recognizing the human TCR-C β domain. Typically, a column containing BF1-conjugated Sepharose was first equilibrated with 20mM Tris-HCl pH 8.0 (loading buffer) and

39

then loaded at 2 ml/min with pH adjusted cell culture media containing c264scTCR/IL15 fusion protein. The column was then washed with 5 column volumes of the loading buffer to remove unbound proteins, and the c264scTCR/IL15 fusion protein was eluted with 4 column volumes of 0.5M Na-citrate, pH 4. After collection, the eluate was adjusted to pH 8.0 by 2M Tris-HCl pH 8.0. The purified protein was buffer exchanged into PBS and filtered using 0.22 µm filter. The BF1 column was stripped with 50 mM Glycine-HCl pH 3.0, and stored in 20% ethanol at 4 C for further use. The fusion protein could be further purified by ion exchange and/or size exclusion chromatography. Cell culture supernatants containing c264scTCR/IL15, c264scTCR/IL15RαSushi and c264scTCR/IL15RαΔE3 fusion proteins were purified by the above methods and samples of the purified fusion proteins were analyzed by electrophoresis on SDS polyacrylamide gels under reducing conditions and followed by staining with Coomassie brilliant blue. Examples of such gels are shown in FIG. 10. The major protein bands correspond to the correct molecular weights expected based on fusion protein sequences.

Example 11

Generation of a Fusion Protein Complex of the TCR/IL15 and TCR/IL15Rα Fusion Proteins

IL15 specifically binds to the extracellular IL15Rα domain with high affinity (4). Thus a complex of fusion proteins bearing the IL-15 and IL15Rα domains can be formed under a variety of conditions, including within the expression cell or extracellularly with unpurified or purified fusion proteins. In one example, equal molar amounts of purified fusion proteins can be mixed under the appropriate conditions (i.e. 10 min at room temperature) to form a fusion protein complex. Complex formation can be monitored in using a variety of techniques including direct binding assays, competitive binding assays, immunoprecipitation, surface plasma resonance, or analyses based on complex size, activity or other properties. For example, as shown in FIG. 11, size exclusion chromatography can monitor the formation of complexes comprising c264scTCR/huIL15 and c264scTCR/huIL15RαSushi fusion proteins based on molecular weight. In this study, about 100 µg of c264scTCR/huIL15 (0.5 mg/ml) was loaded on a Superdex 200 HR 10/30 column for the analysis. The calculated molecular weight for c264scTCR/huIL15 is about 57 kD. Based on SEC profile (FIG. 11A), the estimated molecular weight is about 98 kD, suggesting that this fusion protein is likely a monomer. Similarly, about 60 µg of c264scTCR/huIL15RαSushi fusion protein (0.3 mg/ml) was loaded on the Superdex column. The calculated molecular weight for c264scTCR/huIL15RαSushi is about 52 kD. Based on SEC profile (FIG. 11B), estimated molecular weight of the fusion protein is about 81 kD, again suggesting this fusion protein is a monomer. Previous SEC analysis of other TCR-based fusion proteins showed similar differences between the calculated monomeric molecular weight and the estimated molecular weight of the glycosylated fusion protein. When the c264scTCR/huIL15 and c264scTCR/huIL15RαSushi fusion proteins were mixed in equal molar amounts and about 126 µg of the mixed proteins (0.63 mg/ml) were loaded on the column, the profile shown in FIG. 11C was obtained. Molecular weights of two major peaks were estimated: one at about 170 kD, which is a heterodimer of the two fusion proteins and another one at about 91 kD, which is likely a mix of monomeric forms of the fusion proteins. Thus, the appearance of the 170 kD species in the mixed c264scTCR/huIL15+

40

c264scTCR/huIL15RαSushi fusion protein preparation is evidence that the fusion protein complex of the invention can be generated.

Analysis of the fusion protein complex comprising c264scTCR/huIL15 and c264scTCR/huIL15RαΔE3 fusion proteins was also conducted. About 100 µg of c264scTCR/huIL15RαΔE3 fusion protein (0.5 mg/ml) was loaded on the Superdex column. The calculated molecular weight for c264scTCR/huIL15RαΔE3 is about 60 kD. Based on SEC profile (FIG. 12A), estimated molecular weight of the protein is about 173 kD, suggesting this protein exists as a homodimer. When the c264scTCR/huIL15 and c264scTCR/huIL15RαΔE3 fusion proteins were mixed in equal molar amounts and about 118 µg of the mixed proteins (0.59 mg/ml) were loaded on the column, the profile shown in FIG. 12B was obtained. Molecular weights of two major peaks were estimated: one is >210 kD, which is likely a tetramer composed of two heterodimers and the other is about 93 kD, likely to be c264scTCR/huIL15 monomer. Thus, the appearance of the 170 kD species in the mixed c264scTCR/huIL15+c264scTCR/huIL15RαΔE3 fusion protein preparation is evidence that the fusion protein complex of the invention can be generated.

Example 12

Fusion Protein Complex of the TCR/IL15 and TCR/IL15Rα Fusion Proteins Exhibits Enhanced Binding for Peptide/MHC Complexes

The fusion protein complexes generated as described above were characterized for their ability to bind the TCR-specific antigen, p53 (aa264-272)/HLA-A2.1. To generate cells presenting this antigen, HLA-A2.1-positive T2 cells were loaded with p53 (aa264-272) peptide at 26 C overnight and then stored at 5×10⁹ cells/mL in a liquid nitrogen. T2 cells that were not incubated with peptide serve as controls. The p53 peptide-loaded or control T2 cells were thawed and resuspended in 1 mL of IMDM+10% FBS. The cells (5×10⁵/100 µL) were then stained for 30 min at RT with 0.5 µg of following fusion proteins: c264scTCR/huIL15, c264scTCR/huIL15RαSushi, c264scTCR/huIL15+c264scTCR/huIL15RαSushi complex. Cells were washed once with washing buffer (PBS containing 0.5% BSA and 0.05% sodium azide) and stained with 0.1 µg of biotinylated mouse monoclonal anti-human TCR Cβ antibody (BF1) in 100 µL of washing buffer for 30 min at RT. Cells were washed once and stained with 0.5 µg of R-Phycoerythrin conjugated streptavidin in 100 µL of washing buffer for 30 min at RT. Cells were washed and resuspended for analysis by flow cytometry. As shown in FIG. 13, each of the fusion proteins was capable of specifically staining p53 peptide-loaded cells. In addition, the c264scTCR/huIL15+c264scTCR/huIL15RαSushi fusion protein complex displayed enhanced specific binding via the multivalent c264scTCR domains to the p53 (aa264-272)/HLA-A2.1 complexes displayed on the T2 cells. In particular, the dimeric fusion protein complex showed better staining of the p53 peptide-loaded T2 cells than the monomeric c264scTCR/huIL15 or c264scTCR/huIL15RαSushi fusion proteins. These data suggest that the multimeric fusion protein complex will provide better antigen recognition properties than monomeric form of the fusion proteins.

Example 13

Generation of huIL-15 Mutant Genes and Construction of c264scTCR-hmt-huIL15 Mutant Gene Expression Vectors

As described above, c264scTCR/huIL15+c264scTCR/huIL15RαSushi polypeptides are able to form a complex

41

through interactions of the IL-15 and IL-15R α domains and the multivalent fusion protein complex has enhanced binding for peptide/MHC complexes. Such a fusion protein complex has advantages as an antigen-specific or targeted research, diagnostic and therapeutic agent based on the enhanced binding activity. The ability of the IL-15/IL-15R α domains of the fusion protein to bind cells expressing IL-15 receptors is also an desirable feature as indicated herein. However, there are applications where it is advantageous to increase or decrease the ability of the IL-15/IL-15R α domains to interact with and/or effect the responses of cells expressing IL15 receptors. For example, it may be desirable to reduce this interaction in the applications (i.e. research and diagnostic uses) where the primary goal is to use the fusion protein complex for specific detection of peptide/MHC complexes. In therapeutic applications, it may also be desirable to generated fusion protein complexes that contain IL-15 domains capable of increasing or decreasing IL-15-mediated responses. To address this issue, mutational analysis was carried out to identify residues in IL-15 that effect its binding to IL-2/15R $\beta\gamma_C$ complex without effecting its interactions to IL-15R α . The resulting mutations may create IL-15 variants including antagonists or agonists. In addition to use in the fusion proteins of the invention, the resulting IL-15 antagonists and agonists also may have utility as soluble cytokines (i.e., non-fusion proteins) or as a complex with IL-15R α domains, for research, diagnostic or therapeutic applications. For example, IL-15 antagonists may be useful in suppressing unwanted immune responses whereas IL-15 agonists may be used to stimulate immune responses in therapeutic strategies to treat various diseases.

Based on a comparison between the amino acid sequence and structure of IL-15 with IL-2, several amino acids were identified that could potentially effect interactions between IL-15 and IL-15R α , IL-15R β and/or γ_C . As showed in Table 1 and FIG. 14A, IL-15 variants were created where the potential binding sites of the mature human IL-15 protein to IL-15R $\beta\gamma_C$ receptors, amino acids at positions 8, 61, 65, 72, and 108 (numbering based on mature native human IL-15 sequence), were each substituted or combined with two or more other substitutions. The aspartic acid at position 8 was substituted with alanine or asparagine. The aspartic acid at position 61 was substituted with alanine. The asparagine at position 65 was substituted with alanine or aspartic acid. The asparagine at position 72 was substituted with arginine or aspartic acid. The glutamine at position 108 was substituted with alanine. Both Asp at position 8 and Gln at position 108 were each substituted with an alanine. Both Asp at position 8 and Asn at position 65 were each substituted with an asparagine or alanine. Both Asp at position 8 and Asn at position 65 were each substituted with a serine or arginine. To generate IL-15 mutants, the overlapping PCR was used.

For example, to generate Asp at position 8 with the substitution of alanine or asparagines residues, pNEF38-c264scTCR/huIL15 vector was used as template to amplify two overlapping c-DNA fragments with the front primer for fragment 1

5' -TGGTGGACGCGTAACTGGGTGAATG-3' (SEQ ID NO: 26)

and with the back primer for fragment 1

5' -AGATCTTCAATTTTTCAMKHACTTATTACATTCACCCAG-3' (SEQ ID NO: 38)

42

and with the front primer for fragment 2

(SEQ ID NO: 39)
5' -ACTGGGTGAATGAATAAGTDMKTTGAAAAAATGAAGATC-3'

and with the back primer for fragment 2

(SEQ ID NO: 40)
5' -TGGTGGTCTAGATTATCAAGAAGTGTGATG-3'

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 1 min; 66 C, 1.5 min; 72 C 1.5 min; $\times 35$ cycles; 72 C, 10 min. The fragments 1 and 2 PCR-cDNA products were separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The fragments 1 and 2 PCR-cDNA products were fused together with PfuUltra (Stratagene) under following PCR conditions: 94 C, 1 min; 66 C, 1.5 min; 72 C 1.5 min; $\times 10$ cycles. The overlapping PCR-cDNA fragment was amplified with the front primer

5' -TGGTGGACGCGTAACTGGGTGAATG-3' (SEQ ID NO: 26)

and with the back primer

(SEQ ID NO: 40)
5' -TGGTGGTCTAGATTATCAAGAAGTGTGATG-3'

by PfuUltra (Stratagene) under following PCR conditions: 94 C, 1 min; 64 C, 1.5 min; 69 C 1.5 min; $\times 30$ cycles; 72 C, 10 min. The huIL-15 mutant PCR-cDNA products were separated by electrophoresis on a 1% agarose gel and isolated. The cDNA product was purified from agarose with a Qiaquick Gel Extraction Kit (Qiagen). The huIL15 mutant gene was digested with MluI and XbaI and ligated into pNEF38-c264scTCR-hmt which had been digested with MluI and XbaI. The clone containing the huIL15 gene at position 8 with a substitution of alanine or asparagine was verified by DNA sequencing with GenomeLab Dye Termination Cycle Sequencing with a QUICK START KIT (Beckman Coulter) according to the manufacturer's protocol. The pNEF38-c264scTCR expression vector is described above. Cloning of the DNA fragment encoding mutated human IL-15 protein into the pNEF38-c264scTCR vector resulted in a c264scTCR-hmt-huIL15D8A or c264scTCR-hmt-huIL15D8N fusion gene comprising the following sequence: 3'-immunoglobulin light chain leader -264 TCR V- α -peptide linker -264 TCR V- β -human TCR C- β -mutated human IgG1 hinge-human IL15D8A or -human IL15D8N. The resulting vector (pNEF38-c264scTCR-hmt-huIL15D8A or pNEF38-c264scTCR-hmt-huIL15D8N), shown in FIG. 14B, was confirmed by DNA sequencing. The sequences of the pNEF38-c264scTCR-hmt-huIL15D8A or pNEF38-c264scTCR-hmt-huIL15D8N fusion gene and protein (including the leader sequence) are shown at FIG. 14C and FIG. 14D, respectively.

Other mutations were introduced in a similar fashion and expression vector constructed as described above. The expression vectors were introduced into CHO.K1 cells to generated stable transfectants as described in Example 9. Production and purification of the TCR/IL15 fusion proteins and fusion protein complexes comprising IL-15 variants was carried out using similar methods as described in Examples 10 and 11. Generation of IL-15 variants as soluble cytokines can be carried out through a variety of methods known in the art, including production in prokaryotic and eukaryotic expression systems (see for example, WO9527722; Hsu et al. 2005 J. Immunol. 175:7226).

43

Example 14

Functional Characterization of the TCR/IL15 and TCR/IL15R α Fusion Proteins and Fusion Protein Complexes

Functional binding of the fusion protein TCR domain was demonstrated based on the ELISA and cell staining methods using p53 (aa264-272)/HLA-A2.1 reagents described in Example 9 and the antigen presenting cell staining methods described in Example 12. The ability of the fusion protein IL15 and IL15R α domains to interact was demonstrated as described in Example 11. Further the functional activity of the IL15 and IL15R α domains can be assessed through a variety of methods including binding to IL-2/15R $\beta\gamma_C$ receptors or modulation of the activity of immune cells bearing IL-15 receptors. In one example, CTLL-2 cells, which express the heterotrimeric IL-15R ($\alpha\beta\gamma_C$ chains), were incubated with 0.5 μ g of the individual fusion proteins: c264scTCR/huIL15, c264scTCR/huIL15R α Sushi, or c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex for 30 min at RT. Cells were washed once with washing buffer (PBS containing 0.5% BSA and 0.05% sodium azide) and stained with 0.5 μ g of R-Phycoerythrin (PE) conjugated p53 (aa264-272)/HLA-A2 tetramer for 30 min at RT. Cells were washed and resuspended for analysis by flow cytometry. As shown in FIG. 15, association of the c264scTCR/huIL15 fusion protein and c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex via their huIL15 domains with the IL-15 receptors on CTLL-2 cells can be detected with PE-conjugated p53 (aa264-272)/HLA-A2 tetramer recognizing the c264scTCR domain of the bound fusion protein. These results indicate that both the IL-15 and TCR domains of the fusion protein/fusion protein complexes are capable of functionally interacting with their cognate ligands.

In addition, CTLL-2 cells are dependent upon cytokines for growth and can respond to recombinant human IL-15. A cell-based WST-1 proliferation assay using CTLL-2 cells was developed to assess the IL-15 biological activity of fusion proteins and fusion protein complexes. WST-1 (Roche) is a reagent that can be converted into formazan by dehydrogenase enzymes found in metabolically active cells. In the WST-1 assay, the quantity of formazan in the culture media measured by the amount of 440-450 nm absorbance is directly proportional to the number of living cells in culture. CTLL-2 cells (2×10^4 /200 μ L) were incubated with the indicated concentrations of fusion proteins (0-28 ng/mL): c264scTCR/huIL15, c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex or c264scTCR/huIL15+c264scTCR/huIL15R α AE3 complex for 3 days in 96-well plates at 37 C in a CO₂ incubator. Cells were incubated with 10 μ L of WST-1 for 4 hours before harvesting 100 μ L of culture medium for 440-450 nm absorbance measurement with a microtiter plate reader. As shown in FIG. 16, c264scTCR/huIL15 fusion protein can support the proliferation of CTLL-2 cells at a concentration as low as 1.8 ng/mL (31.25 pM), suggesting activation of CTLL-2 cells with c264scTCR/huIL15 fusion protein via the high affinity IL15 receptor. Interestingly, fusion protein complexes also supported CTLL-2 cell proliferation but to a lesser degree suggesting that c264scTCR/huIL15 stimulatory activity was inhibited following complex formation with c264scTCR/huIL15R α Sushi or c264scTCR/huIL15R α AE3 (by one fold or four fold, respectively). This suggests the binding of c264scTCR/huIL15 to the high affinity IL15 receptor is inhibited by c264scTCR/huIL15R α Sushi or c264scTCR/huIL15R α AE3 fusion proteins. These results provide evi-

44

dence that the fusion proteins and fusion protein complexes can activate or suppress responses of immune cells under different conditions.

Similar assays were performed with cell lines expressing only the intermediate affinity IL-15 $\beta\gamma_C$ receptors, such as 32D β cell lines (see below). In some cases, it is possible that the biological activity of IL15 in stimulating proliferation of the IL-15R-bearing cells will be enhanced when it is in a complex with the IL15R α domain (Mortier, E. et al., 2006, *J. Biol. Chem.*, 281: 1612-1619; Stoklasek, T. et al., 2006, *J. Immunol* 177: 6072-6080; Rubinstein, M. P. et al., 2006, *Proc Natl Acad Sci USA* 103: 9166-9171). Stimulation of cell proliferation by the c264scTCR/huIL15+c264scTCR/huIL15R α Sushi or c264scTCR/huIL15+c264scTCR/huIL15R α AE3 complexes will be assessed and may provide additional evidence that the fusion protein complexes can stimulate or activate immune responses of immune cells.

Example 15

Dimeric Fusion Protein Complexes of the TCR/IL15R α Sushi and TCR/IL15 Variants Exhibit TCR-Specific Binding to Peptide/MHC Complex but Less Binding to the IL-15R $\beta\gamma_C$ Receptors

The fusion protein complexes comprising IL-15 variants as described above were characterized for their ability to bind the TCR-specific antigen, p53 (aa264-272)/HLA-A2.1. To generate cells presenting p53 (aa264-272)/HLA-A2.1, HLA-A2.1-positive T2 cells (2×10^6 /mL) were loaded with 20 μ M p53 (aa264-272) peptide at 37 C in the presence of 1 $\times$ PLE (Altor Bioscience) for 2-3 hrs. T2 cells that were not incubated with peptide and 32D β cells expressing IL-2/15R $\beta\gamma_C$ serve as controls. The p53 peptide-loaded T2 cells, control T2 cells, or 32D β cells (2×10^5 /100 μ L) were then incubated for 30 min at 4 C with 320 nM of following dimeric fusion protein complexes: 1) c264scTCR/huIL15+c264scTCR/huIL15R α Sushi, 2) c264scTCR/huIL15D8A+c264scTCR/huIL15R α Sushi, and 3) c264scTCR/huIL15D8N+c264scTCR/huIL15R α Sushi. These complexes were generated by incubating 160 nM of purified c264scTCR/huIL15 fusion protein and 160 nM of purified c264scTCR/huIL15R α Sushi fusion protein at 4 C for 3 hours: Following staining, cells were washed once with washing buffer (PBS containing 0.5% BSA and 0.05% sodium azide) and stained with 0.5 μ g of biotinylated mouse monoclonal anti-human TCR C β antibody (BF1) in 100 μ L of washing buffer for 30 min at 4 C. Cells were washed once and stained with 0.5 μ g of R-Phycoerythrin conjugated streptavidin in 100 μ L of washing buffer for 30 min at 4 C. Cells were washed and resuspended for analysis by flow cytometry. As shown in FIG. 17A, the c264scTCR/huIL15D8A+c264scTCR/huIL15R α Sushi complex and c264scTCR/huIL15D8N+c264scTCR/huIL15R α Sushi complex exhibited equivalent activity as the c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex for specifically staining p53 peptide-loaded T2 cells. These results indicate that the multivalent scTCR domains are fully functional in each of these fusion complexes. However, as shown in FIG. 17B and FIG. 17C, the mutant c264scTCR/huIL15 fusion protein complexes showed less background staining on control T2 cells (FIG. 17B) and IL-15R $\beta\gamma_C$ -positive 32D β cells (FIG. 17C) than the wide type c264scTCR/IL15 fusion protein complex. Thus these fusion protein complexes comprising IL-15 variants (D8A and D8N) do not show binding activity to the IL-15R $\beta\gamma_C$ receptors present on the 32D β cells. Similar studies of IL-15R $\beta\gamma_C$ receptor binding were carried out with other

45

fusion proteins comprising IL-15 variants and are summarized in Table 1. The results indicate that fusion proteins and fusion protein complexes of the invention comprising IL-15 variants retain activity to recognize peptide/MHC complexes and exhibit decreased or increased binding activity for IL-15R $\beta\gamma_c$ receptors.

To confirm the above fusion proteins functional TCR and IL-15 domains, peptide/MHC and IL-15R α binding activity was measured by ELISA analysis. The 96-well microtiter plates were precoated with 20 nM BF1, an anti-TCR C β antibody, or 20 nM TCR/IL15R α Sushi in carbonated buffer pH 9.1 (sodium bicarbonate 35 mM, Na₂ CO₃, 17.5 mM, NaCl 50 mM) over 3 hours at 4 C. Plates were washed with washing buffer (Imidazole 40 mM, NaCl 150 mM) for 4 times and blocked with 1% BSA-PBS for 10 minutes. The indicated fusion proteins at a concentration of 0.03-4 nM were added to the plates and incubated at RT for 30 minutes. The plates were washed four times. The BF1 captured fusion proteins were incubated with 1 μ g/mL of HRP-conjugated p53/HLA-A2.1 tetramer for 45 minutes at RT and the TCR/IL15R α Sushi captured fusion proteins were incubated with 50 ng/mL of biotinylate mouse anti-human IL-15 for 30 minutes at RT. After washing for 4 times, the plate incubated with biotinylate mouse anti-human IL-15 was incubated with 0.25 μ g/mL of HRP-streptavidin for 15 minutes. The plates were washed 4 times and incubated with peroxidase substrate ABT for 1-10 minutes and developed for 405 nm absorbance measurement with a microtiter plate reader. As shown at FIG. 18A and FIG. 18B, the fusion proteins shared similar TCR-specific binding activity for p53/HLA-A2 tetramer and equivalent IL-15 binding activity for IL15R α Sushi. Similar studies of IL-15R α binding were carried out with other fusion proteins comprising IL-15 variants and are summarized in Table 1. The results indicate that fusion proteins and fusion protein complexes of the invention comprising IL-15 variants retain activity to recognize peptide/MHC complexes and IL-15R α receptors.

Example 16

Functional Characterization of the TCR/IL15 Mutant Fusion Proteins and Fusion Protein Complexes

As indicated above, fusion proteins comprising an IL-15 antagonist or agonist may be a useful as a targeted agents for inhibiting or stimulating IL-15-mediated responses (i.e., T cell or NK cell activity) at the disease site. To determine the IL-15 bioactivity of these fusion proteins to effect immune responses, cell proliferation studies were carried out with CTLL-2 cells expressing the high affinity IL-15R ($\alpha\beta\gamma_c$ chains) and with 32D β cells expressing the intermediate IL-15R ($\beta\gamma_c$ chains). The cells (2 $\times$ 10⁴/200 μ L) were incubated with 0.4-40 nM of the above described TCR/IL15 fusion proteins for 3 days in 96-well plates at 37 C in a CO₂ incubator. Cells were incubated with 10 μ L of WST-1 for 4 hours before harvesting 150 μ L of culture medium for 440 nm absorbance measurement with a microtiter plate reader. As shown in FIG. 19A and FIG. 19B, the c264scTCR/huIL15 fusion protein comprising the wild type IL-15 domain can support the proliferation of CTLL-2 and 32D β cells at a concentration as low as 40 pM or 1 nM respectively. Interestingly, the fusion protein comprising an IL15 variant with an asparagine to aspartic acid substitution at position 72 with an amino acid (c264scTCR/huIL15N72D) was much more active than the fusion protein comprising the wild type IL-15 domain at supporting the proliferation of 32D β cell line, showing biological activity at a concentration as low as 80 pM (FIG. 19B). In this respect the fusion protein comprising

46

IL-15 variant (huIL15N72D) showed super agonist activity. In a complex with c264scTCR/IL15R α Sushi at one to one ratio, the c264scTCR/huIL15N72D had similar binding ability as c264scTCR/huIL15wt to p53/HLA-A2.1 complex on T2 cells (FIG. 17A) but exhibited increased binding ability to IL-15R $\beta\gamma_c$ receptors on 32D β cells (FIG. 17C). In contrast, the fusion proteins comprising IL-15 variants with substitutions at position 8 (c264scTCR/huIL15D8N or c264scTCR/huIL15D8A), position 65 (c264scTCR/huIL15N65A), position 108 (c264scTCR/huIL15Q108A), or a different substitution at position 72 (c264scTCR/huIL15N72R) were less active in supporting proliferation of both CTLL-2 and 32D β cells compared to c264scTCR/huIL15wt fusion protein (FIG. 19A and FIG. 19B). Similar studies of IL-15-dependent proliferative activity were carried out with other fusion proteins comprising IL-15 variants and are summarized in Table 1. The data support the hypothesis that mutations at positions 8, 61, 65, 72 and 108 of the IL-15 protein can result in IL-15 antagonists with decreased binding to IL-15R and little or no activity to stimulate immune responses. The results with the position 72 substitutions are unexpected given that one mutant (c264scTCR/huIL15N72R) acted as an IL-15 antagonist whereas a different mutant (c264scTCR/huIL15N72D) showed increased binding to IL-15 R and enhanced activity at stimulate immune responses.

In a typical circumstance, IL-15 is trans-presented by IL15R α on a dendritic cell surface to IL-15R $\beta\gamma_c$ receptors on memory T, NKT, or NK cell to support cell survival and stimulate immune responses. An antagonist should block the trans-presentation of IL-15 by binding IL15R α . To evaluate if the antagonist fusion proteins can compete with c264scTCR/huIL15wt to block its activity to support CTLL-2 cell growth, 4 $\times$ 10⁴ CTLL-2 cells were incubated with 0.5 nM of c264scTCR/huIL15wt in the presence or absence of 50 nM (100-fold molar excess) of various c264scTCR/huIL15 mutant fusion proteins at 37 C in a CO₂ incubator for 24 hours. Cells were incubated with 10 μ L of WST-1 for 4 hours before harvesting 150 μ L of culture medium for 440 nm absorbance measurement with a microtiter plate reader. As shown in FIG. 19C, the ability of c264scTCR/huIL15wt to support proliferation of CTLL-2 cells was totally blocked in the presence of 100-fold more c264scTCR/huIL15D8N, c264scTCR/huIL15D8A, c264scTCR/huIL15D8A/Q108A, c264scTCR/huIL15Q108A, or c264scTCR/huIL15D8N/N65A, and was reduced 62% in the presence of c264scTCR/huIL15N72R fusion protein. It suggested that these fusion proteins were the antagonists to c264scTCR/IL15 fusion protein. This data indicates that c264scTCR/huIL15 mutant fusion proteins were functional antagonist of the IL-15 activity, as expected based on the ability of these proteins to bind IL-15R α but not IL-15R $\beta\gamma_c$ receptors.

Similar studies will be carried out with the other TCR/IL15 fusion proteins and IL-15 variants described herein to demonstrate IL-15 antagonist and agonist activity. As summarized in Table 1, the substitutions at positions 8, 61, 65, 72, and 108 of IL-15 show the ability to affect the binding of IL-15 to IL-15R ($\beta\gamma_c$ chains). Other substitutions at positions 92, 101, and 111 of IL-15 will also be assessed as potential binding sites for IL-15R interaction. In addition, combinations of changes including substitutions at all or several of these residues may create the effective antagonists or agonists of IL-15. Including the molecules described above, IL-15 variants to be assessed include those with changes at position 8 to alanine, asparagine, serine, lysine, threonine, or tyrosine; position 61 to alanine, asparagine, serine, lysine, threonine, or tyrosine; position 65 to alanine, aspartic acid, or arginine;

47

position 72 to alanine, aspartic acid, or arginine; and positions 92, 101, 108, or 111 to alanine or serine.

Example 17

Cell-Cell Conjugation and Immune Cell Retargeting
by the TCR/IL15 and TCR/IL15R α Fusion Proteins
and Fusion Protein Complexes

To demonstrate that the fusion proteins or fusion protein complexes can bridge IL-15 receptor-bearing cells with peptide/MHC bearing target cells, T2 cells will be loaded with either p53 (aa264-272) peptide or control CMVpp65 (aa495-503) peptide and then labeled with dihydroethidium. CTLL-2 cells will be labeled with calcein AM and the two labeled cell populations will be mixed and incubated in the presence or absence of the fusion proteins or fusion protein complexes. In the absence of the fusion protein complexes or when the T2 cells were loaded with control peptide, the cells are anticipated to remain as two distinct populations as assessed by flow cytometry. However, when the T2 cells are loaded with p53 (aa264-272) and incubated with the CTLL-2 cells in the presence of fusion proteins or complexes, the appearance of a double staining population of cells would be indicative of conjugation of T2 cells to CTLL-2 cells via the fusion proteins or fusion protein complexes.

Similarly, studies can be conducted to demonstrate that the fusion protein complexes can bridge IL-15 receptor-bearing immune cells with peptide/MHC bearing target cells and direct immune cytotoxicity against the target cells. For example, T2 cells will be loaded with either p53 (aa264-272) peptide or control CMVpp65 (aa495-503) peptide and then labeled with calcein AM. Immune effector cells bearing IL-15 receptors (i.e. activated NK cells or T cells) will be mixed at different ratios and incubated under appropriate conditions (i.e. 37 C for 2-4 hours) in the presence or absence of the fusion protein complex. Cytotoxicity will be assessed based on release of calcein from the T2 target cells into the culture media by standard methods. The specific release of calcein-AM will be measured or compared to the non-specific control of spontaneous released calcein-AM. In the absence of the fusion protein complex or when the T2 cells were loaded with control peptide, low levels of target cell cytotoxicity are expected. However, when the T2 cells are loaded with p53 (aa264-272) and incubated with the immune effector cells in the presence of fusion protein complex, specific lysis of the T2 cells would be an indication that the immune effector cells are retargeted against the p53 peptide-presenting cells via the fusion protein complex. Similar studies will be conducted with tumor cell lines presenting p53 (aa264-272)/HLA-A2.1 complexes as target cells.

Example 18

In Vivo Demonstration of Anti-Tumor Effects of
IL-15 Variant Agonists, TCR/IL15 Fusion Proteins
and Fusion Protein Complexes

To determine if the fusion protein complexes or IL-15 variant agonists have anti-tumor activity in vivo, an experimental xenograft tumor model will be used. Human tumor cell lines expressing p53 (aa264-272)/HLA-A2.1 complexes, such as A375 melanoma, MDA-MB-231 mammary adenocarcinoma, PANC1 pancreatic carcinoma, have been employed in similar animal efficacy studies using other TCR-based fusion proteins (5-7). For example, A375 human melanoma cells will be injected subcutaneously into the flank of

48

nude mice and tumors will be allowed to establish for three days. Tumor bearing mice will be injected intravenously with c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex or an IL-15 variant agonist (dose range—0.1 to 2 mg/kg), or the dose volume equivalent of PBS daily for four or more days. During the study, tumor size will be measured and the tumor volumes will be calculated. All mice treated with PBS are expected to develop tumors. Suppression of tumor growth or complete tumor regression in some or all the mice treated with the fusion protein complex or IL-15 variant agonist would be an indication of an antitumor effect of the treatment. Alternative dosing schedules including multi-cycle dosing may also demonstrate the antitumor efficacy of the fusion protein or IL-15 variant agonist. Tumor cell lines lacking p53 (aa264-272)/HLA-A2.1 complexes (such as HT-29 or AsPC-1 (5, 9)) can be used as controls for antigen specific recognition by the c264scTCR-domain of the fusion protein complex. Alternative fusion protein complexes comprising other TCR domains (i.e. specific to CMVpp65 (aa495-503) peptide (9) could be used as non-tumor targeting controls.

In addition, adoptive cell transfer studies will be carried out in xenograft tumor bearing mice. For example, immune cells bearing the IL-15 receptor, such as naïve or activated (or memory) splenocytes, NK cell or T cells, will be isolated from mice and incubated with c264scTCR/huIL15+c264scTCR/huIL15R α Sushi complex or an agonist IL-15 variant under conditions permitting binding to the cells. In some cases the fusion protein complex or an agonist IL-15 variant will be used to activate the immune cells. The IL-15 variant activated cells or fusion protein complex-coated cells will then be transferred into nude mice bearing A375 tumors. Controls will include transfer of the untreated immune cells, the fusion protein complex alone and PBS. Tumor growth will be monitored and all mice treated with PBS are expected to develop tumors. Suppression of tumor growth or complete tumor regression in some or all the mice treated with the IL-15 variant activated cells or fusion protein complex-coated cells would be an indication of an antitumor effect of the treatment. Alternatively, the IL-15 variant or fusion protein complex and immune cells will be administered at the same time, or separately at the same or different times. The immune cells may be autologous or allogeneic in relation with the tumor-bearing host. The number of cells transferred and dosing schedule will be varied to assess and optimize antitumor efficacy. As described above, other tumor lines or fusion protein complexes will be employed to determine the role of antigen targeting in any observed antitumor activity.

Example 19

In Vitro Treatment of Immune Cells with
TCR/IL15:TCR/IL15R α Fusion Protein Complexes
Followed by Adoptive Cell Transfer Provide
Improved Survival in Xenograft Tumor Animal
Model

To demonstrate the anti-tumor efficacy of enriched allogeneic mouse NK cells preincubated with TCR/IL15:TCR/IL15R α fusion protein complexes on tumor growth, the following study was carried out using human NSCLC A549A2 tumor cells in an experimental metastasis model in nude mice.

Athymic nude mice (n=4 per group, female, 5-6 week old) were intravenously

(IV) injected through the lateral tail vein with the human NSCLC tumor cell line A549-A2 at 5×10^6 cells/mouse. The

49

A549-A2 cell line represents a transfectant of the p53-positive A549 parental line carrying a vector expressing human HLA-A2.1 cDNA.

Spleens from A2 mice (B6 background) were collected and NK cells were isolated using a NK cell isolation kit from Miltenyi Biotech, Inc. according to the manufacturer's instruction. Briefly, a single cell suspension of splenocytes was prepared by homogenizing the spleens through a metal screen (60 mesh) in HBSS. Red blood cells were lysed in ACK red blood lysing buffer. Cells will be incubated with biotin-antibody cocktail (10 μ L for 10⁷ cells) for 10 min at 4-8 C. The number of leukocytes was determined and 30 μ L of buffer (PBS pH 7.2, 0.5% BSA and 2 mM EDTA) and 20 μ L of anti-biotin MicroBeads per 10⁷ cells was added and the mixture was incubated at 4-8 C for 15 min. The cells were washed in 2 mL buffer and centrifuge at 300 $\times$ g for 10 min. The cells were resuspended in 500 μ L of buffer for loading to the MACS column. The flow through was collected and the purity of the NK cells was determined using FACS analysis.

In order to activate the cells, NK cells (5 $\times$ 10⁶) were cultured at 37 C overnight in the presence or absence of c264scTCR/IL15: c264scTCR/IL15R α fusion protein complex, TCR-IL2 fusion protein or rhIL-2 in T25 flasks in 10 ml RPMI1640 supplemented with 10% FBS. c264scTCR/IL15: c264scTCR/IL15R α fusion protein complex and TCR-IL2 fusion protein was added at a concentration of 0.8 μ g/mL and rhIL-2 was added at 0.2 μ g/mL. After overnight incubation, cells were harvested and preincubated in 0.5 mg/mL c264scTCR/IL15: c264scTCR/IL15R α fusion protein complex or TCR-IL2 fusion protein or 0.125 mg/mL rhIL-2 in 100 μ L on ice for 30 min. After wash in PBS (1 mL), cells were resuspended in PBS at 10 $\times$ 10⁶/mL for adoptive transfer.

On day 1, mice were injected i.v. via the tail vein with A549A2 tumor cells (5 $\times$ 10⁶) to establish pulmonary tumors. Fourteen days post tumor cell injection, mice were randomized and divided into 5 groups (n=4). Mice were treated with cyclophosphamide (CTX) via intraperitoneal injection at a dose of 200 mg/kg on days 14 and 21. NK cells (1 $\times$ 10⁶/mouse) preincubated with different fusion proteins or rhIL-2 were injected i.v. on days 16 and 22, and mice receiving PBS served as controls. A summary of the treatment schedule is as follows:

Group	CTX		NK cells	
	Dose (mg/kg)	Injection (ip)	Dose ($\times$ 10 ⁶)	Injection (iv)
CTX	200	Days 14, 21	0	Days 16, 22
CTX+NK/rhIL2	200	Days 14, 21	1	Days 16, 22
CTX+NK/MART-1scTCR-IL2	200	Days 14, 21	1	Days 16, 22
CTX+NK/c264scTCR-IL2	200	Days 14, 21	1	Days 16, 22
CTX+NK/c264scTCR/IL15: c264scTCR/IL15R α fusion protein complex	200	Days 14, 21	1	Days 16, 22

Survival of tumor-bearing mice was monitored every day. Mice that became moribund were sacrificed and counted as dead. Mice surviving longer than 100 days post-tumor injection were considered as cured.

Median survivals for mice in the CTX, CTX+NK/rhIL-2, CTX+NK/MART1scTCR-IL2, CTX+NK/c264scTCR-IL2 and c264scTCR/IL15: c264scTCR/IL15R α fusion protein complex treatment groups are 52, 67.5, 64.5, 85.5, and 80 days, respectively (FIG. 20). Thus, adoptive transfer of

50

c264scTCR/IL15: c264scTCR/IL15R α -activated NK cells resulted a longer median survival time than observed in tumor-bearing animals treated with chemotherapy alone or with NK cells activated by the non-targeted MARTscTCR-IL2 or rhIL-2. The results from this pilot experiment indicate that activation and targeting mouse NK cells with c264scTCR/IL15: c264scTCR/IL15R α may provide enhanced antitumor activity.

Example 20

Enhanced Binding of TCR/IL15:TCR/IL15R α Fusion Protein Complexes to IL-15R Bearing Immune Cells as Evidenced by an Extended Cell Surface Residency Time

The cell surface residency time of the fusion protein complexes on the IL-15R-bearing cells may influence the ability of the fusion protein to target or bridge effector cells with the TCR-specific tumor cells. To investigate this, binding of the scTCR/IL-15 fusion proteins, TCR/IL15:TCR/IL15R α fusion protein complexes and recombinant IL-15 to IL-15 α β γ C receptor-bearing CTLL-2 cells and IL-15R β γ C receptor-bearing CTLL-2 cells 32D β cells will be directly compared by flow cytometry. Following incubation with the various proteins, cells will be washed and incubated in media at 37 $^{\circ}$ C. for up to 180 min and the level of proteins remaining on the cell surface was detected with PE-labeled anti-IL-15 mAb. Comparisons between the initial time zero staining and subsequent times will allow determination of the cell surface residency time for each proteins binding to IL-15R. Increased cell surface residency time of the scTCR/IL-15 fusion proteins or TCR/IL15:TCR/IL15R α fusion protein complexes compared to IL-15 would be an indication of enhanced and more stable receptor binding activity.

Example 21

Increased In Vivo Half Life of TCR/IL15 Fusion Proteins and TCR/IL15:TCR/IL15R α Fusion Protein Complexes Compared to IL-15 in Mice

The pharmacokinetic parameters of c264scTCR/IL-15, c264scTCR/IL15: c264scTCR/IL15R α complex, recombinant IL-15 or soluble IL-15:IL-15R α complex will be evaluated in the HLA-A2.1/Kb-transgenic mouse strain. The presence of the HLA-A2.1 domain, for which c264scTCR/IL-2 is restricted, may influence the pharmacokinetics of this fusion protein and should give a more relevant "humanized" view of the pharmacokinetics than other mouse strains. Mice will be injected intravenous with molar equivalent amounts of c264scTCR/IL-15, c264scTCR/IL15: c264scTCR/IL15R α complex, recombinant IL-15 or soluble IL-15:IL-15R α complex and blood will be collected at various time points from 5 minutes to two weeks post injection. Serum concentrations of the fusion proteins will be evaluated using ELISA formats, disclosed above. Concentrations of IL-15 were detected with a standard IL-15-specific ELISA.

The in vivo pharmacokinetic parameters of c264scTCR/IL-15, c264scTCR/IL15: c264scTCR/IL15R α complex, recombinant IL-15 or soluble IL-15:IL-15R α complex will be determined using curve fitting software (e.g., WinNonlin). Elevated C_{max} values, increased serum half life or decreased clearance for the c264scTCR/IL-15 or c264scTCR/IL15: c264scTCR/IL15R α complex compared to recombinant IL-15 or soluble IL-15:IL-15R α complex would be an indication that generation of the TCR-IL15 fusion or TCR/IL15:

51

TCR/IL-15Ra complex provides more favorable pharmacokinetics that is observed with IL-15 alone.

Example 22

In Vivo Demonstration of Immunosuppressive Effects of IL-15 Variant Antagonists, TCR/IL15 Fusion Proteins and Fusion Protein Complexes

To determine if the fusion protein complexes or IL-15 variant antagonists have immuno-suppressive activity in vivo, an experimental autoimmune arthritis model will be used. It has been demonstrated that autoimmune arthritis can be induced following administration of type II collagen (CII) in HLA-DR4-transgenic mice (Rosloniec et al. 1998. J Immunol. 160:2573-8). Additionally, CII-specific T-cells involved in the pathology of this disease have been characterized. The TCR genes from these T-cells will be used to construct the appropriate expression vectors and host cell lines to generate CIIscTCR/IL15 comprising IL-15 variant antagonists and CIIscTCR/IL15RaSushi fusion proteins as described in previous examples. Following induction of arthritis by CII administration, the HLA-DR4-transgenic mice will be injected intravenously with CIIscTCR/IL15-antagonist+CIIscTCR/IL15RaSushi complex or an IL-15 variant antagonist (dose range—0.1 to 2 mg/kg), or the dose volume equivalent of PBS daily for four or more days. During the study, the mouse paw joints will be evaluated and scored for the degree of inflammation on a scale of 0 to 4. All mice treated with PBS are expected to develop arthritis. Suppression of arthritis (e.g. decreased incidence or clinical score) in some or all the mice treated with the fusion protein complex or IL-15 variant would be an indication of an immunosuppressive effect of the treatment. Alternative dosing schedules including multi-cycle dosing may also demonstrate the immunosuppressive efficacy of the fusion protein or IL-15 variant. Alternative fusion protein complexes comprising other TCR domains (i.e. specific to p53 peptide) could be used as non-disease targeting controls to demonstrate specificity of the targeted TCR fusion proteins ability to direct immunosuppressive activity to the disease site.

Incorporation by Reference
The contents of all references, patents, pending patent applications and published patents, cited throughout this application are hereby expressly incorporated by reference.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many

52

equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

REFERENCES

1. Mortier, E., A. Quemener, P. Vusio, I. Lorenzen, Y. Boublik, J. Grotzinger, A. Plet, and Y. Jacques. 2006. Soluble interleukin-15 receptor alpha (IL-15R alpha)-sushi as a selective and potent agonist of IL-15 action through IL-15R beta/gamma. Hyperagonist IL-15xIL-15R alpha fusion proteins. J Biol Chem 281: 1612-1619.
2. Stoklasek, T. A., K. S. Schluns, and L. Lefrancois. 2006. Combined IL-15/IL-15Ra immunotherapy maximizes IL-15 activity in vivo. J Immunol 177: 6072-6080.
3. Rubinstein, M. P., M. Kovar, J. F. Purton, J. H. Cho, O. Boyman, C. D. Surh, and J. Sprent. 2006. Converting IL-15 to a superagonist by binding to soluble IL-15R{alpha}. Proc Natl Acad Sci USA 103: 9166-9171.
4. Waldmann, T. A. 2006. The biology of interleukin-2 and interleukin-15: implications for cancer therapy and vaccine design. Nat Rev Immunol 6: 595-601.
5. Belmont, H. J., S. Price-Schiavi, B. Liu, K. F. Card, H. I. Lee, K. P. Han, J. Wen, S. Tang, X. Zhu, J. Merrill, P. A. Chavillaz, J. L. Wong, P. R. Rhode, and H. C. Wong. 2006. Potent antitumor activity of a tumor-specific soluble TCR/IL-2 fusion protein. Clin Immunol 121: 29-39.
6. Card, K. F., S. A. Price-Schiavi, B. Liu, E. Thomson, E. Nieves, H. Belmont, J. Builes, J. A. Jiao, J. Hernandez, J. Weidanz, L. Sherman, J. L. Francis, A. Amirkhosravi, and H. C. Wong. 2004. A soluble single-chain T-cell receptor IL-2 fusion protein retains MHC-restricted peptide specificity and IL-2 bioactivity. Cancer Immunol Immunother 53: 345-357.
7. Mosquera, L. A., K. F. Card, S. A. Price-Schiavi, H. J. Belmont, B. Liu, J. Builes, X. Zhu, P. A. Chavillaz, H. I. Lee, J. A. Jiao, J. L. Francis, A. Amirkhosravi, R. L. Wong, and H. C. Wong. 2005. In Vitro and In Vivo Characterization of a Novel Antibody-Like Single-Chain TCR Human IgG1 Fusion Protein. J Immunol 174: 4381-4388.
8. Penichet, M. L., E. T. Harvill, and S. L. Morrison. 1997. Antibody-IL-2 fusion proteins: a novel strategy for immune protection. Hum Antibodies 8: 106-118.
9. Zhu, X., H. J. Belmont, S. Price-Schiavi, B. Liu, H. I. Lee, M. Fernandez, R. L. Wong, J. Builes, P. R. Rhode, and H. C. Wong. 2006. Visualization of p53264-272/HLA-A*0201 Complexes Naturally Presented on Tumor Cell Surface by a Multimeric Soluble Single-Chain T Cell Receptor. J Immunol 176: 3223-3232.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 40

<210> SEQ ID NO 1

<211> LENGTH: 114

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu Asp Leu Ile
1 5 10 15

Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val His
20 25 30

-continued

Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu Gln
 35 40 45

Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val Glu
 50 55 60

Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn Val
 65 70 75 80

Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn Ile
 85 90 95

Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile Asn
 100 105 110

Thr Ser

<210> SEQ ID NO 2
 <211> LENGTH: 345
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

aactgggtga atgtaataag tgatttgaaa aaaattgaag atcttattca atctatgcat      60
attgatgcta ctttatatac ggaaagtgat gttcacocca gttgcaaagt aacagcaatg      120
aagtgtcttc tcttggaagt acaagttatt tcaactgagt ccggagatgc aagtattcat      180
gatacagtag aaaatctgat catcctagca aacaacagtt tgtcttctaa tgggaatgta      240
acagaatctg gatgcaaaga atgtgaggaa ctggaggaaa aaaatattaa agaatttttg      300
cagagttttg tacatattgt ccaaatgttc atcaacactt cttga                      345

```

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 3

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
 1 5 10 15

Gly Gly Gly Ser
 20

<210> SEQ ID NO 4
 <211> LENGTH: 1563
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic construct

<400> SEQUENCE: 4

```

accaccatgg agacagacac actcctgtta tgggtactgc tgctctgggt tccaggttcc      60
accggtcagt cagtgaacga gcccgatgct cgcgtcactg tctctgaagg agcctctctg      120
cagctgagat gcaagtattc ctactctggg acaccttata tgttctgta tgtccagtag      180
ccgcggcagg ggctgcagct gctcctcaag tactattcag gagaccagt ggttcaagga      240
gtgaatggct tcgaggctga gttagcaag agtaactctt ccttccacct gcggaagacc      300
tctgtgcact ggagcgactc tgctgtgtac ttctgtgttt tgagcgagga tagcaactat      360
cagttgatct ggggctctgg gaccaagcta attataaagc cagacactag tgggtggcgg      420

```

-continued

```

ggcagcggcg gtgggtgggtc cgggtggcggc ggttctggcg gtggcggttc ctgagcaat 480
tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaagatg 540
aggtgtatcc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
gagtttaaat ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
gaaaaacgat tctctgctga gtgtccttca aactcacctt gcagcctaga aattcagtc 720
tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
gaagttttct ttggtaaagg aaccagactc acagtgttag aggacctgaa caaggtgttc 840
ccacccgagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
acactggtgt gcctggccac aggtttcttc cctgaccacg tggagctgag ctggtgggtg 960
aatgggaagg aggtgcacag tggggtcagc acggaccgcg agccctcaa ggagcagccc 1020
gccctcaatg actccagata ctgcctgagc agccgcctga ggtctctggc cacttcttg 1080
cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggctctc ggagaatgac 1140
gagtggaacc aggatagggc caaacccgtc acccagatcg tcagcgcgga ggctgggg 1200
agagcagacg ttaacaactg ggtgaatgta ataagtgatt tgaaaaaaat tgaagatctt 1260
attoaatcta tgcataattg tgctacttta tatacggaaa gtgatgttca cccagttgc 1320
aaagtaacag caatgaagtg ctttctcttg gagttacaag ttatttctact tgagtcggga 1380
gatgcaagta ttcatgatac agtagaaaat ctgatcatcc tagcaacaaa cagtttgtct 1440
tctaattgga atgtaacaga atctggatgc aaagaatgtg aggaactgga ggaaaaaat 1500
attaaagaat ttttgacag tttgtacat attgtccaaa tggtcatcaa cacttcttga 1560
taa 1563

```

```

<210> SEQ ID NO 5
<211> LENGTH: 517
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

```

```

<400> SEQUENCE: 5

```

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Trp Val Pro
 1           5           10           15
Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
          20           25           30
Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
          35           40           45
Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
          50           55           60
Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
          65           70           75           80
Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
          85           90           95
Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
          100          105          110
Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
          115          120          125
Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
          130          135          140
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Ser Asn Ser Lys
          145          150          155          160

```

-continued

Val Ile Gln Thr Pro Arg Tyr Leu Val Lys Gly Gln Gly Gln Lys Ala
 165 170 175
 Lys Met Arg Cys Ile Pro Glu Lys Gly His Pro Val Val Phe Trp Tyr
 180 185 190
 Gln Gln Asn Lys Asn Asn Glu Phe Lys Phe Leu Ile Asn Phe Gln Asn
 195 200 205
 Gln Glu Val Leu Gln Gln Ile Asp Met Thr Glu Lys Arg Phe Ser Ala
 210 215 220
 Glu Cys Pro Ser Asn Ser Pro Cys Ser Leu Glu Ile Gln Ser Ser Glu
 225 230 235 240
 Ala Gly Asp Ser Ala Leu Tyr Leu Cys Ala Ser Ser Leu Ser Gly Gly
 245 250 255
 Gly Thr Glu Val Phe Phe Gly Lys Gly Thr Arg Leu Thr Val Val Glu
 260 265 270
 Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser
 275 280 285
 Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala
 290 295 300
 Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly
 305 310 315 320
 Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln Pro Leu Lys Glu
 325 330 335
 Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu Arg
 340 345 350
 Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln
 355 360 365
 Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp Arg
 370 375 380
 Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg Ala
 385 390 395 400
 Asp Val Asn Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu
 405 410 415
 Asp Leu Ile Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser
 420 425 430
 Asp Val His Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu
 435 440 445
 Glu Leu Gln Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp
 450 455 460
 Thr Val Glu Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn
 465 470 475 480
 Gly Asn Val Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu
 485 490 495
 Lys Asn Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met
 500 505 510
 Phe Ile Asn Thr Ser
 515

<210> SEQ ID NO 6
 <211> LENGTH: 1614
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 construct
 <400> SEQUENCE: 6

-continued

```

accacatg agacagac actcctgtta tgggtactgc tgctctgggt tccaggttcc 60
accggtcagt cagtgcagca gcccgatgct cgcgtcactg tctctgaagg agcctctctg 120
cagctgagat gcaagtattc ctactctggg acaccttate tgttctggta tgtccagtac 180
ccgcggcagg ggtgcagct gctcctcaag tactattcag gagaccagt ggttcaagga 240
gtgaatggct tgcaggctga gttcagcaag agtaactctt ccttccacct gcggaaagcc 300
tctgtgcaact ggagcgactc tgctgtgtac ttctgtgttt tgagcgagga tagcaactat 360
cagttgatct ggggctctgg gaccaagcta attataaagc cagacactag tgggtggcgg 420
ggcagcgagg gtgggtgttc cgggtggcggc ggttctggcg gtggcggttc ctgcagcaat 480
tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaagatg 540
aggtgtatcc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
gagtttaaat ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
gaaaaacgat tctctgtgta gtgtccttca aactcacctt gcagcctaga aattcagttc 720
tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
gaagttttct ttggtaaagg aaccagactc acagtgttag aggacctgaa caaggtgttc 840
ccacccgagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
acactgggtg gctgggccac aggtcttctc cctgaccacg tggagctgag ctgggtgggtg 960
aatgggaagg aggtgcacag tggggtcagc acggacccgc agccctcaa ggagcagccc 1020
gccctcaatg actccagata ctgcctgagc agccgcctga gggctctcggc cacttctctg 1080
cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggtctctc ggagaatgac 1140
gagtggaacc aggatagggc caaacccgtc acccagatcg tcagcgccga ggccctgggg 1200
agagcagacg ttaacgagcc caaatcttct gacaaaaactc acacatctcc accgtctcca 1260
acgcgtaact ggggtgaatg aataagtgat ttgaaaaaaa ttgaagatct tattcaatct 1320
atgcataatg atgctacttt atatacgga agtgatgttc accccagtgt caaagtaaca 1380
gcaatgaagt gctttctctt ggagttacaa gttatttcac ttgagtcagg agatgcaagt 1440
attcatgata cagttagaaa tctgatcctc ctacaaaca acagtttctc ttctaattgg 1500
aatgtaacag aatctggatg caaagaatgt gaggaactgg aggaaaaaaa tattaagaa 1560
tttttcgaga gttttgtaca tattgtccaa atgttcacac acacttcttg ataa 1614

```

```

<210> SEQ ID NO 7
<211> LENGTH: 534
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

```

```

<400> SEQUENCE: 7

```

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Trp Val Pro
 1             5             10             15
Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
          20             25             30
Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
          35             40             45
Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
          50             55             60
Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
          65             70             75             80

```

Gly	Phe	Glu	Ala	Glu	Phe	Ser	Lys	Ser	Asn	Ser	Ser	Phe	His	Leu	Arg	
				85					90					95		
Lys	Ala	Ser	Val	His	Trp	Ser	Asp	Ser	Ala	Val	Tyr	Phe	Cys	Val	Leu	
				100					105					110		
Ser	Glu	Asp	Ser	Asn	Tyr	Gln	Leu	Ile	Trp	Gly	Ser	Gly	Thr	Lys	Leu	
				115					120					125		
Ile	Ile	Lys	Pro	Asp	Thr	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	
				130					135					140		
Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Ser	Ser	Asn	Ser	Lys	
				145					150					155		
Val	Ile	Gln	Thr	Pro	Arg	Tyr	Leu	Val	Lys	Gly	Gln	Gly	Gln	Lys	Ala	
				165					170					175		
Lys	Met	Arg	Cys	Ile	Pro	Glu	Lys	Gly	His	Pro	Val	Val	Phe	Trp	Tyr	
				180					185					190		
Gln	Gln	Asn	Lys	Asn	Asn	Glu	Phe	Lys	Phe	Leu	Ile	Asn	Phe	Gln	Asn	
				195					200					205		
Gln	Glu	Val	Leu	Gln	Gln	Ile	Asp	Met	Thr	Glu	Lys	Arg	Phe	Ser	Ala	
				210					215					220		
Glu	Cys	Pro	Ser	Asn	Ser	Pro	Cys	Ser	Leu	Glu	Ile	Gln	Ser	Ser	Glu	
				225					230					235		
Ala	Gly	Asp	Ser	Ala	Leu	Tyr	Leu	Cys	Ala	Ser	Ser	Leu	Ser	Gly	Gly	
				245					250					255		
Gly	Thr	Glu	Val	Phe	Phe	Gly	Lys	Gly	Thr	Arg	Leu	Thr	Val	Val	Glu	
				260					265					270		
Asp	Leu	Asn	Lys	Val	Phe	Pro	Pro	Glu	Val	Ala	Val	Phe	Glu	Pro	Ser	
				275					280					285		
Glu	Ala	Glu	Ile	Ser	His	Thr	Gln	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ala	
				290					295					300		
Thr	Gly	Phe	Phe	Pro	Asp	His	Val	Glu	Leu	Ser	Trp	Trp	Val	Asn	Gly	
				305					310					315		
Lys	Glu	Val	His	Ser	Gly	Val	Ser	Thr	Asp	Pro	Gln	Pro	Leu	Lys	Glu	
				325					330					335		
Gln	Pro	Ala	Leu	Asn	Asp	Ser	Arg	Tyr	Cys	Leu	Ser	Ser	Arg	Leu	Arg	
				340					345					350		
Val	Ser	Ala	Thr	Phe	Trp	Gln	Asn	Pro	Arg	Asn	His	Phe	Arg	Cys	Gln	
				355					360					365		
Val	Gln	Phe	Tyr	Gly	Leu	Ser	Glu	Asn	Asp	Glu	Trp	Thr	Gln	Asp	Arg	
				370					375					380		
Ala	Lys	Pro	Val	Thr	Gln	Ile	Val	Ser	Ala	Glu	Ala	Trp	Gly	Arg	Ala	
				385					390					395		
Asp	Val	Asn	Glu	Pro	Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	
				405					410					415		
Ser	Pro	Thr	Arg	Asn	Trp	Val	Asn	Val	Ile	Ser	Asp	Leu	Lys	Lys	Ile	
				420					425					430		
Glu	Asp	Leu	Ile	Gln	Ser	Met	His	Ile	Asp	Ala	Thr	Leu	Tyr	Thr	Glu	
				435					440					445		
Ser	Asp	Val	His	Pro	Ser	Cys	Lys	Val	Thr	Ala	Met	Lys	Cys	Phe	Leu	
				450					455					460		
Leu	Glu	Leu	Gln	Val	Ile	Ser	Leu	Glu	Ser	Gly	Asp	Ala	Ser	Ile	His	
				465					470					475		
Asp	Thr	Val	Glu	Asn	Leu	Ile	Ile	Leu	Ala	Asn	Asn	Ser	Leu	Ser	Ser	

-continued

500	505	510	
Glu Lys Asn Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln			
515	520	525	

Met Phe Ile Asn Thr Ser
530

<210> SEQ ID NO 8
 <211> LENGTH: 1647
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic construct

<400> SEQUENCE: 8

accacatg	agacagacac	actcctgtta	tgggtactgc	tgtcttgggt	tccaggttcc	60
accggtcagt	cagtgcacga	gcccgatgct	cgcgctactg	tctctgaagg	agcctctctg	120
cagctgagat	gcaagtattc	ctactctggg	acaccttate	tgttctggta	tgtccagtac	180
ccgcggcagg	ggctgcagct	gctcctcaag	tactattcag	gagaccagct	ggttcaagga	240
gtgaatggct	tcgaggctga	gttcagcaag	agtaactctt	ccttccacct	gcggaagacc	300
tctgtgcact	ggagcgactc	tgtgtgtac	ttctgtgttt	tgagcgagga	tagcaactat	360
cagttgatct	ggggctctgg	gaccaagcta	attataaagc	cagacactag	tgggtggcgg	420
ggcagcggcg	gtgggtggtc	cggtggcggc	ggttctggcg	gtggcggttc	ctcgagcaat	480
tcaaaagtca	ttcagactcc	aagatatctg	gtgaaagggc	aaggacaaaa	agcaaagatg	540
aggtgtatcc	ctgaaaaggg	acatccagtt	gtattctggt	atcaacaaaa	taagaacaat	600
gagtttaaat	ttttgattaa	ctttcagaat	caagaagttc	ttcagcaaat	agacatgact	660
gaaaaacgat	tctctgtcga	gtgtccttca	aactcacctt	gcagcctaga	aattcagttc	720
tctgaggcag	gagactcagc	actgtacctc	tgtgccagca	gtctgtcagg	gggcggcaca	780
gaagttttct	ttggtaaagg	aaccagactc	acagttgtag	aggacctgaa	caaggtgttc	840
ccacccgagg	tcgctgtgtt	tgagccatca	gaagcagaga	tctccacac	ccaaaaggcc	900
acactgggtg	gcctggccac	aggcttcttc	cctgaccacg	tggagctgag	ctgggtgggtg	960
aatgggaagg	agggtgcacg	tggggtcagc	acggacccgc	agcccccaa	ggagcagccc	1020
gcccctcaatg	actccagata	ctgcctgagc	agccgcctga	gggtctcggc	caccttcttg	1080
cagaaccccc	gcaaccactt	ccgctgtcaa	gtccagttct	acgggctctc	ggagaatgac	1140
gagtggaccc	aggatagggc	caaacccgtc	acccagatcg	tcagcgccga	ggcctggggt	1200
agagcagacg	taaacatcac	gtgccctccc	cccatgtccg	tggaaacacg	agacatctgg	1260
gtcaagagct	acagcttgta	ctccagggag	cggtacattt	gtaactctgg	tttcaagcgt	1320
aaagccggca	cgtccagcct	gacggagtgc	gtgttgaaca	aggccacgaa	tgtcgccccc	1380
tggacaaccc	ccagcttcaa	atgcattaga	cccgcagctt	catctcccag	ctcaaacac	1440
acagcggcca	caacagcagc	tattgtcccg	ggctcccagc	tgatgccttc	aaaatcacct	1500
tccacaggaa	ccacagagat	aagcagtcac	gagtcctccc	acggcacccc	ctctcagaca	1560
acagccaaga	actgggaact	cacagcatcc	gcctcccacc	agccgcaggg	tgtgtatcca	1620
cagggccaca	gcgacaccac	ttgataa				1647

<210> SEQ ID NO 9
 <211> LENGTH: 545
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

<400> SEQUENCE: 9

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1             5             10             15

Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
 20             25             30

Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
 35             40             45

Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
 50             55             60

Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
 65             70             75             80

Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
 85             90             95

Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
100            105            110

Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
115            120            125

Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
130            135            140

Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Ser Asn Ser Lys
145            150            155            160

Val Ile Gln Thr Pro Arg Tyr Leu Val Lys Gly Gln Gly Gln Lys Ala
165            170            175

Lys Met Arg Cys Ile Pro Glu Lys Gly His Pro Val Val Phe Trp Tyr
180            185            190

Gln Gln Asn Lys Asn Asn Glu Phe Lys Phe Leu Ile Asn Phe Gln Asn
195            200            205

Gln Glu Val Leu Gln Gln Ile Asp Met Thr Glu Lys Arg Phe Ser Ala
210            215            220

Glu Cys Pro Ser Asn Ser Pro Cys Ser Leu Glu Ile Gln Ser Ser Glu
225            230            235            240

Ala Gly Asp Ser Ala Leu Tyr Leu Cys Ala Ser Ser Leu Ser Gly Gly
245            250            255

Gly Thr Glu Val Phe Phe Gly Lys Gly Thr Arg Leu Thr Val Val Glu
260            265            270

Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser
275            280            285

Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala
290            295            300

Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly
305            310            315            320

Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln Pro Leu Lys Glu
325            330            335

Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu Arg
340            345            350

Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln
355            360            365

Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp Arg
370            375            380

Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg Ala
385            390            395            400

```

-continued

Asp Val Asn Ile Thr Cys Pro Pro Pro Met Ser Val Glu His Ala Asp
 405 410 415
 Ile Trp Val Lys Ser Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys
 420 425 430
 Asn Ser Gly Phe Lys Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys
 435 440 445
 Val Leu Asn Lys Ala Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu
 450 455 460
 Lys Cys Ile Arg Pro Ala Ala Ser Ser Pro Ser Ser Asn Asn Thr Ala
 465 470 475 480
 Ala Thr Thr Ala Ala Ile Val Pro Gly Ser Gln Leu Met Pro Ser Lys
 485 490 495
 Ser Pro Ser Thr Gly Thr Thr Glu Ile Ser Ser His Glu Ser Ser His
 500 505 510
 Gly Thr Pro Ser Gln Thr Thr Ala Lys Asn Trp Glu Leu Thr Ala Ser
 515 520 525
 Ala Ser His Gln Pro Pro Gly Val Tyr Pro Gln Gly His Ser Asp Thr
 530 535 540
 Thr
 545

<210> SEQ ID NO 10
 <211> LENGTH: 1410
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 construct
 <400> SEQUENCE: 10

accaccatgg agacagacac actcctgtta tgggtactgc tgcctctgggt tccaggttcc 60
 accggtcagt cagtgcagca gcccgatgct cgcgtcactg tctctgaagg agcctctctg 120
 cagctgagat gcaagtattc ctactctggg acaccttacc tgttctggtta tgtccagtac 180
 ccgcggcagg ggctgcagct gctcctcaag tactattcag gagaccaggt ggttcaagga 240
 gtgaatggct tcgaggctga gttcagcaag agtaactctt ccttccacct gcggaaaagcc 300
 tctgtgcact ggagcgactc tgctgtgtac ttctgtgttt tgagcgagga tagcaactat 360
 cagttgatct ggggctctgg gaccaagcta attataaagg cagacactag tgggtggcgg 420
 ggcagcggcg gtggtggttc cgggtggcggc ggttctggcg gtggcggttc ctcgagcaat 480
 tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaagatg 540
 aggtgtatcc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
 gagtttaaat ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
 gaaaaacgat tctctgtgta gtgtccttca aactcacctt gcagcctaga aattcagttc 720
 tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
 gaagttttct ttggtaaagg aaccagactc acagtgttag aggacctgaa caagtggttc 840
 ccaccggagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
 aactgggtgt gcctggccac aggtctcttc cctgaccacg tggagctgag ctggtgggtg 960
 aatgggaagg aggtgcacag tgggtcagc acggacccgc agccctcaa ggagcagccc 1020
 gccctcaatg actccagata ctgcctgagc agccgcctga gggctctggc caccttcttg 1080
 cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggctctc ggagaatgac 1140

-continued

```

gagtggaccc aggatagggc caaacccgtc acccagatcg tcagcgccga ggccctgggggt 1200
agagcagacg ttaacatcac gtgccctccc cccatgtccg tggaacacgc agacatctgg 1260
gtcaagagct acagcttgta ctccaggag cggtacattt gtaactctgg ttccaagcgt 1320
aaagccggca cgtccagcct gacggagtgc gtgttgaaca aggccacgaa tgtcgcccac 1380
tggacaaccc ccagttctca atgcattaga 1410

```

```

<210> SEQ ID NO 11
<211> LENGTH: 468
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

```

```

<400> SEQUENCE: 11

```

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1           5           10           15
Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
          20           25           30
Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
          35           40           45
Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
          50           55           60
Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
          65           70           75           80
Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
          85           90           95
Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
          100          105          110
Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
          115          120          125
Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
          130          135          140
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Ser Asn Ser Lys
          145          150          155          160
Val Ile Gln Thr Pro Arg Tyr Leu Val Lys Gly Gln Gly Gln Lys Ala
          165          170          175
Lys Met Arg Cys Ile Pro Glu Lys Gly His Pro Val Val Phe Trp Tyr
          180          185          190
Gln Gln Asn Lys Asn Asn Glu Phe Lys Phe Leu Ile Asn Phe Gln Asn
          195          200          205
Gln Glu Val Leu Gln Gln Ile Asp Met Thr Glu Lys Arg Phe Ser Ala
          210          215          220
Glu Cys Pro Ser Asn Ser Pro Cys Ser Leu Glu Ile Gln Ser Ser Glu
          225          230          235          240
Ala Gly Asp Ser Ala Leu Tyr Leu Cys Ala Ser Ser Leu Ser Gly Gly
          245          250          255
Gly Thr Glu Val Phe Phe Gly Lys Gly Thr Arg Leu Thr Val Val Glu
          260          265          270
Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser
          275          280          285
Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala
          290          295          300
Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly
          305          310          315          320

```

-continued

Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln Pro Leu Lys Glu
 325 330 335
 Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu Arg
 340 345 350
 Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln
 355 360 365
 Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp Arg
 370 375 380
 Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg Ala
 385 390 395 400
 Asp Val Asn Ile Thr Cys Pro Pro Pro Met Ser Val Glu His Ala Asp
 405 410 415
 Ile Trp Val Lys Ser Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys
 420 425 430
 Asn Ser Gly Phe Lys Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys
 435 440 445
 Val Leu Asn Lys Ala Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu
 450 455 460
 Lys Cys Ile Arg
 465

<210> SEQ ID NO 12
 <211> LENGTH: 1467
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 construct
 <400> SEQUENCE: 12

accaccatgg agacagacac actcctgtta tgggtactgc tgcctctgggt tccaggttcc 60
 accggtcagt cagtgcagca gcccgatgct cgcgtcactg tctctgaagg agcctctctg 120
 cagctgagat gcaagtattc ctactctggg acaccttata tgttctgtga tgtccagtac 180
 ccgcggcagg ggctgcagct gctcctcaag tactattcag gagaccaggt ggttcaagga 240
 gtgaatggct tcgaggctga gttcagcaag agtaactctt ccttccacct gcggaaagcc 300
 tctgtgcact ggagcgactc tgctgtgtac ttctgtgttt tgagcgagga tagcaactat 360
 cagttgatct ggggctctgg gaccaagcta attataaagg cagacactag tgggtggcgg 420
 ggcagcggcg gtgtgtgttc cgggtggcgc ggttctggcg gtggcggttc ctcgagcaat 480
 tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaagatg 540
 aggtgtatcc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
 gagttaaata ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
 gaaaaacgat tctctgtga gtgtccttca aactcacctt gcagcctaga aattcagttc 720
 tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
 gaagttttct ttggtaaagg aaccagactc acagtgttag aggacctgaa caaggtgttc 840
 ccaccgagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
 aactgggtgt gcctggccac aggtttcttc cctgaccacg tggagctgag ctggtgggtg 960
 aatgggaagg aggtgcacag tgggtcagc acggacccgc agccctcaa ggagcagccc 1020
 gccctcaatg actccagata ctgcctgagc agccgcctga gggctctggc caccttcttg 1080
 cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggctctc ggagaatgac 1140

-continued

```

gagtggaccc aggatagggc caaacccgtc acccagatcg tcagcgccga ggccctggggt 1200
agagcagacg ttaacgagcc caaatcttct gacaaaactc acacatctcc accgtctcca 1260
acgcgtatca cgtgccctcc ccccatgtcc gtggaacacg cagacatctg ggtcaagagc 1320
tacagcttgt actccagga gcggtacatt tgtaactctg gtttcaagcg taaagccggc 1380
acgtccagcc tgacggagtg cgtgttgaac aaggccaaga atgtcgccca ctggacaacc 1440
cccagctcca aatgcattag atgataa 1467

```

```

<210> SEQ ID NO 13
<211> LENGTH: 485
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

```

```

<400> SEQUENCE: 13

```

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1          5          10          15
Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
          20          25          30
Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
          35          40          45
Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
          50          55          60
Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
          65          70          75          80
Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
          85          90          95
Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
          100         105         110
Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
          115         120         125
Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
          130         135         140
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Ser Asn Ser Lys
          145         150         155         160
Val Ile Gln Thr Pro Arg Tyr Leu Val Lys Gly Gln Gly Gln Lys Ala
          165         170         175
Lys Met Arg Cys Ile Pro Glu Lys Gly His Pro Val Val Phe Trp Tyr
          180         185         190
Gln Gln Asn Lys Asn Asn Glu Phe Lys Phe Leu Ile Asn Phe Gln Asn
          195         200         205
Gln Glu Val Leu Gln Gln Ile Asp Met Thr Glu Lys Arg Phe Ser Ala
          210         215         220
Glu Cys Pro Ser Asn Ser Pro Cys Ser Leu Glu Ile Gln Ser Ser Glu
          225         230         235         240
Ala Gly Asp Ser Ala Leu Tyr Leu Cys Ala Ser Ser Leu Ser Gly Gly
          245         250         255
Gly Thr Glu Val Phe Phe Gly Lys Gly Thr Arg Leu Thr Val Val Glu
          260         265         270
Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser
          275         280         285
Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala
          290         295         300

```

-continued

Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly
 305 310 315 320
 Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln Pro Leu Lys Glu
 325 330 335
 Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu Arg
 340 345 350
 Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln
 355 360 365
 Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp Arg
 370 375 380
 Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg Ala
 385 390 395 400
 Asp Val Asn Glu Pro Lys Ser Ser Asp Lys Thr His Thr Ser Pro Pro
 405 410 415
 Ser Pro Thr Arg Ile Thr Cys Pro Pro Pro Met Ser Val Glu His Ala
 420 425 430
 Asp Ile Trp Val Lys Ser Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile
 435 440 445
 Cys Asn Ser Gly Phe Lys Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu
 450 455 460
 Cys Val Leu Asn Lys Ala Thr Asn Val Ala His Trp Thr Thr Pro Ser
 465 470 475 480
 Leu Lys Cys Ile Arg
 485

<210> SEQ ID NO 14
 <211> LENGTH: 1614
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 construct

<400> SEQUENCE: 14

accaccatgg agacagacac actcctgtta tgggtactgc tgctctgggt tccaggttcc 60
 accggtcagt cagtgaacga gcccgatgct cgcgtcactg tctctgaagg agcctctctg 120
 cagctgagat gcaagtattc ctactctggg acaccttata tgttctggtta tgtccagtag 180
 ccgcggcagg ggctgcagct gctcctcaag tactattcag gagaccaggt ggttcaagga 240
 gtgaatggct tcgaggctga gttcagcaag agtaactctt ccttccacct gcggaaagcc 300
 tctgtgcact ggagcgactc tgctgtgtac ttctgtgttt tgagcgaggga tagcaactat 360
 cagttgatct ggggctctgg gaccaagcta attataaagc cagacactag tgggtggcgg 420
 ggcagcggcg gtggtgggtc cgggtggcggc ggttctggcg gtggcggttc ctcgagcaat 480
 tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaaagatg 540
 aggtgtatcc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
 gagtttaaat ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
 gaaaaacgat tctctgtgta gtgtccttca aactcacctt gcagcctaga aattcagttc 720
 tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
 gaagttttct ttggtaaagg aaccagactc acagtgttag aggacctgaa caaggtgttc 840
 ccacccgagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
 aactgggtgt gcctggccac aggtcttctc cctgaccacg tggagctgag ctggtgggtg 960
 aatgggaagg aggtgcacag tggggtcagc acggaccgcc agccctcaa ggagcagccc 1020

-continued

```

gccctcaatg actccagata ctgcctgagc agccgcctga gggctctcggc caccttctgg 1080
cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggctctc ggagaatgac 1140
gagtggaacc aggatagggc caaacccgtc acccagatcg tcagcgccga ggcctggggg 1200
agagcagacg ttaacgagcc caaatcttct gacaaaactc acacatctcc accgtctcca 1260
acgcgtaact ggggtaagt aataagtgtc ttgaaaaaaa ttgaagatct tattcaatct 1320
atgcataatg atgctacttt atatacgga agtgatgttc accccagttg caaagtaaca 1380
gcaatgaagt gctttctctt ggagttacaa gttatttcac ttgagtcggg agatgcaagt 1440
attcatgata cagtagaaaa tctgatcacc ctagcaaaaa acagtttggt ttctaattgg 1500
aatgtaacag aatctggatg caaagaatgt gaggaactgg agggaaaaaa tattaagaa 1560
tttttcgaga gttttgtaca tattgtccaa atgttcacca acacttcttg ataa 1614

```

```

<210> SEQ ID NO 15
<211> LENGTH: 1614
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
                        construct

```

```

<400> SEQUENCE: 15

```

```

accacatg agacagac actcctgtta tgggtactgc tgctctgggt tccaggttcc 60
accggtcagt cagtgcgc gcccgatgct cgcgtcactg tctctgaagg agcctctctg 120
cagctgagat gcaagtattc ctactctggg acaccttacc tgttctggta tgtccagtac 180
ccgcggcagg ggcgtcagct gctcctcaag tactattcag gagaccagt ggttcaagga 240
gtgaatggct tcgaggctga gttcagcaag agtaactctt ccttccacct gcggaaagcc 300
tctgtgcact ggagcgactc tgctgtgtac ttctgtgttt tgagcgagga tagcaactat 360
cagttgatct ggggctctgg gaccaagcta attataaagc cagacactag tgggtggcgg 420
ggcagcggcg gtgggtggtc cgggtggcggc ggttctggcg gtggcggttc ctogagcaat 480
tcaaaagtca ttcagactcc aagatatctg gtgaaagggc aaggacaaaa agcaaatgat 540
agggtgtacc ctgaaaaggg acatccagtt gtattctggt atcaacaaaa taagaacaat 600
gagtttaaat ttttgattaa ctttcagaat caagaagttc ttcagcaaat agacatgact 660
gaaaaacgat tctctgtgta gtgtccttca aactcacctt gcagcctaga aattcagttc 720
tctgaggcag gagactcagc actgtacctc tgtgccagca gtctgtcagg gggcggcaca 780
gaagttttct ttggtaaagg aaccagactc acagttgtag aggacctgaa caaggtgttc 840
ccaccggagg tcgctgtgtt tgagccatca gaagcagaga tctcccacac ccaaaaggcc 900
acactgggtg gcctggccac aggtctcttc cctgaccacg tggagctgag ctgggtgggtg 960
aatgggaagg aggtgcacag tggggtcagc acggacccgc agccctcaa ggagcagccc 1020
gccctcaatg actccagata ctgcctgagc agccgcctga gggctctcggc caccttctgg 1080
cagaaccccc gcaaccactt ccgctgtcaa gtccagttct acgggctctc ggagaatgac 1140
gagtggaacc aggatagggc caaacccgtc acccagatcg tcagcgccga ggcctggggg 1200
agagcagacg ttaacgagcc caaatcttct gacaaaactc acacatctcc accgtctcca 1260
acgcgtaact ggggtaagt aataagtaat ttgaaaaaaa ttgaagatct tattcaatct 1320
atgcataatg atgctacttt atatacgga agtgatgttc accccagttg caaagtaaca 1380
gcaatgaagt gctttctctt ggagttacaa gttatttcac ttgagtcggg agatgcaagt 1440

```

-continued

```

atcattgata cagtagaaaa tctgatcatc ctagcaaaca acagtttgtc ttctaattggg 1500
aatgtaacag aatctggatg caaagaatgt gaggaactgg aggaaaaaaa tattaagaa 1560
tttttcgaga gttttgtaca tattgtccaa atgttcacac acatttcttg ataa 1614

```

```

<210> SEQ ID NO 16
<211> LENGTH: 534
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
construct

```

```

<400> SEQUENCE: 16

```

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1          5          10          15
Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
 20          25          30
Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
 35          40          45
Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
 50          55          60
Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
 65          70          75          80
Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
 85          90          95
Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
100          105          110
Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
115          120          125
Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
130          135          140
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Asn Ser Lys
145          150          155          160
Val Ile Gln Thr Pro Arg Tyr Leu Val Lys Gly Gln Gly Gln Lys Ala
165          170          175
Lys Met Arg Cys Ile Pro Glu Lys Gly His Pro Val Val Phe Trp Tyr
180          185          190
Gln Gln Asn Lys Asn Asn Glu Phe Lys Phe Leu Ile Asn Phe Gln Asn
195          200          205
Gln Glu Val Leu Gln Gln Ile Asp Met Thr Glu Lys Arg Phe Ser Ala
210          215          220
Glu Cys Pro Ser Asn Ser Pro Cys Ser Leu Glu Ile Gln Ser Ser Glu
225          230          235          240
Ala Gly Asp Ser Ala Leu Tyr Leu Cys Ala Ser Ser Leu Ser Gly Gly
245          250          255
Gly Thr Glu Val Phe Phe Gly Lys Gly Thr Arg Leu Thr Val Val Glu
260          265          270
Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser
275          280          285
Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala
290          295          300
Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly
305          310          315          320
Lys Glu Val His Ser Gly Val Ser Thr Asp Pro Gln Pro Leu Lys Glu
325          330          335

```

-continued

Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu Arg
 340 345 350
 Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln
 355 360 365
 Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp Arg
 370 375 380
 Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg Ala
 385 390 395 400
 Asp Val Asn Glu Pro Lys Ser Ser Asp Lys Thr His Thr Ser Pro Pro
 405 410 415
 Ser Pro Thr Arg Asn Trp Val Asn Val Ile Ser Ala Leu Lys Lys Ile
 420 425 430
 Glu Asp Leu Ile Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu
 435 440 445
 Ser Asp Val His Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu
 450 455 460
 Leu Glu Leu Gln Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His
 465 470 475 480
 Asp Thr Val Glu Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser
 485 490 495
 Asn Gly Asn Val Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu
 500 505 510
 Glu Lys Asn Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln
 515 520 525
 Met Phe Ile Asn Thr Ser
 530

<210> SEQ ID NO 17
 <211> LENGTH: 534
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 construct

<400> SEQUENCE: 17

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1 5 10 15
 Gly Ser Thr Gly Gln Ser Val Thr Gln Pro Asp Ala Arg Val Thr Val
 20 25 30
 Ser Glu Gly Ala Ser Leu Gln Leu Arg Cys Lys Tyr Ser Tyr Ser Gly
 35 40 45
 Thr Pro Tyr Leu Phe Trp Tyr Val Gln Tyr Pro Arg Gln Gly Leu Gln
 50 55 60
 Leu Leu Leu Lys Tyr Tyr Ser Gly Asp Pro Val Val Gln Gly Val Asn
 65 70 75 80
 Gly Phe Glu Ala Glu Phe Ser Lys Ser Asn Ser Ser Phe His Leu Arg
 85 90 95
 Lys Ala Ser Val His Trp Ser Asp Ser Ala Val Tyr Phe Cys Val Leu
 100 105 110
 Ser Glu Asp Ser Asn Tyr Gln Leu Ile Trp Gly Ser Gly Thr Lys Leu
 115 120 125
 Ile Ile Lys Pro Asp Thr Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 130 135 140
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ser Ser Asn Ser Lys
 145 150 155 160

-continued

Val	Ile	Gln	Thr	Pro	Arg	Tyr	Leu	Val	Lys	Gly	Gln	Gly	Gln	Lys	Ala	
				165					170					175		
Lys	Met	Arg	Cys	Ile	Pro	Glu	Lys	Gly	His	Pro	Val	Val	Phe	Trp	Tyr	
	180						185						190			
Gln	Gln	Asn	Lys	Asn	Asn	Glu	Phe	Lys	Phe	Leu	Ile	Asn	Phe	Gln	Asn	
	195					200					205					
Gln	Glu	Val	Leu	Gln	Gln	Ile	Asp	Met	Thr	Glu	Lys	Arg	Phe	Ser	Ala	
	210				215						220					
Glu	Cys	Pro	Ser	Asn	Ser	Pro	Cys	Ser	Leu	Glu	Ile	Gln	Ser	Ser	Glu	
225				230					235						240	
Ala	Gly	Asp	Ser	Ala	Leu	Tyr	Leu	Cys	Ala	Ser	Ser	Leu	Ser	Gly	Gly	
	245							250						255		
Gly	Thr	Glu	Val	Phe	Phe	Gly	Lys	Gly	Thr	Arg	Leu	Thr	Val	Val	Glu	
	260					265							270			
Asp	Leu	Asn	Lys	Val	Phe	Pro	Pro	Glu	Val	Ala	Val	Phe	Glu	Pro	Ser	
	275					280						285				
Glu	Ala	Glu	Ile	Ser	His	Thr	Gln	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ala	
	290				295						300					
Thr	Gly	Phe	Phe	Pro	Asp	His	Val	Glu	Leu	Ser	Trp	Trp	Val	Asn	Gly	
305				310					315					320		
Lys	Glu	Val	His	Ser	Gly	Val	Ser	Thr	Asp	Pro	Gln	Pro	Leu	Lys	Glu	
	325							330						335		
Gln	Pro	Ala	Leu	Asn	Asp	Ser	Arg	Tyr	Cys	Leu	Ser	Ser	Arg	Leu	Arg	
	340						345						350			
Val	Ser	Ala	Thr	Phe	Trp	Gln	Asn	Pro	Arg	Asn	His	Phe	Arg	Cys	Gln	
	355					360					365					
Val	Gln	Phe	Tyr	Gly	Leu	Ser	Glu	Asn	Asp	Glu	Trp	Thr	Gln	Asp	Arg	
	370				375						380					
Ala	Lys	Pro	Val	Thr	Gln	Ile	Val	Ser	Ala	Glu	Ala	Trp	Gly	Arg	Ala	
385				390					395					400		
Asp	Val	Asn	Glu	Pro	Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	
	405							410					415			
Ser	Pro	Thr	Arg	Asn	Trp	Val	Asn	Val	Ile	Ser	Asn	Leu	Lys	Lys	Ile	
	420					425						430				
Glu	Asp	Leu	Ile	Gln	Ser	Met	His	Ile	Asp	Ala	Thr	Leu	Tyr	Thr	Glu	
	435				440						445					
Ser	Asp	Val	His	Pro	Ser	Cys	Lys	Val	Thr	Ala	Met	Lys	Cys	Phe	Leu	
	450				455					460						
Leu	Glu	Leu	Gln	Val	Ile	Ser	Leu	Glu	Ser	Gly	Asp	Ala	Ser	Ile	His	
465			470						475					480		
Asp	Thr	Val	Glu	Asn	Leu	Ile	Ile	Leu	Ala	Asn	Asn	Ser	Leu	Ser	Ser	
	485							490					495			
Asn	Gly	Asn	Val	Thr	Glu	Ser	Gly	Cys	Lys	Glu	Cys	Glu	Glu	Leu	Glu	
	500						505					510				
Glu	Lys	Asn	Ile	Lys	Glu	Phe	Leu	Gln	Ser	Phe	Val	His	Ile	Val	Gln	
	515				520							525				
Met	Phe	Ile	Asn	Thr	Ser											
	530															

<210> SEQ ID NO 18

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

-continued

<400> SEQUENCE: 18

Ala Ser Gly Gly Gly Gly Ser Gly Gly Gly
1 5 10

<210> SEQ ID NO 19

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 19

caccttgcca tagccagctc ttc

23

<210> SEQ ID NO 20

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 20

gtctaagcag cagagtgatg ttg

24

<210> SEQ ID NO 21

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 21

tgggtaacaa ctgggtgaat gtaataagt

30

<210> SEQ ID NO 22

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 22

acgcgtttat caagaagtgt tgatgaacat ttggac

36

<210> SEQ ID NO 23

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 23

tgggtgggta acgagcccaa atcttctg

28

<210> SEQ ID NO 24

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

```

<400> SEQUENCE: 24

attattacgc gttggagacg gtggagatg                               29

<210> SEQ ID NO 25
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 25

tgagtgatcg ataccaccat ggagacagac ac                             32

<210> SEQ ID NO 26
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 26

tgggtggacgc gtaactgggt gaatg                                   25

<210> SEQ ID NO 27
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 27

tgggtgtcta gaattatcaa gaagtgtga tg                             32

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 28

agtcacgcgg tgcctgtgg                                           20

<210> SEQ ID NO 29
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 29

tgacgcgttt aagtggtgtc gctgtgcct g                             31

<210> SEQ ID NO 30
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 30

```

-continued

tggttaacat cacgtgcct cccccatg 29

<210> SEQ ID NO 31
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 31

ttgttgacgc gtttatctaa tgcatttgag actgg 35

<210> SEQ ID NO 32
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 32

taataaacgc gtatcacgtg ccctc 25

<210> SEQ ID NO 33
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 33

tggtggtcta gattatcatc taatgcattt g 31

<210> SEQ ID NO 34
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 34

tgagtgtccg gaaccacat ggagacagac ac 32

<210> SEQ ID NO 35
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 35

ttgttgccg cgccttatca tctaagcat ttgag 35

<210> SEQ ID NO 36
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 36

tgagtgttcg aattatcaag aagtgttgat gaac 34

-continued

```

<210> SEQ ID NO 37
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 37

ttgttggcgg ccgcttatca agtgggtgctg ctg                               33

<210> SEQ ID NO 38
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 38

agatcttcaa tttttttcaa mkhacttatt acattcaccc ag                       42

<210> SEQ ID NO 39
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 39

actgggtgaa tgtaataagt dmkttgaaaa aaattgaaga tc                       42

<210> SEQ ID NO 40
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 40

tggtggtcta gattatcaag aagtgttgat g                                   31

```

What is claimed is:

1. An isolated human interleukin 15 (hIL-15) variant comprising an amino acid substitution at position 61, 72, 92, or 111 of a mature human IL-15 sequence of SEQ ID NO:1, wherein the hIL-15 variant has increased or decreased binding activity for the interleukin 15 receptor beta/gamma chain (IL-15R $\beta\gamma$ C) receptors compared to native hIL-15 polypeptide.

2. The isolated hIL-15 variant of claim 1, wherein the hIL-15 variant functions as an IL-15 agonist or antagonist.

3. The isolated hIL-15 variant of claim 1, where the amino acid substitution is in the domain of IL-15 that interacts with IL-15R $\beta\gamma$ C, or both IL-15R β and γ C.

4. The isolated hIL-15 variant of claim 1, wherein the amino acid substitution is of D to A at position 61 or N to R at position 72 of the mature human IL-15 sequence.

5. The isolated hIL-15 variant of claim 4, wherein the amino acid substitution results in an hIL-15 variant that has

45

IL-15 antagonist activity or decreased binding activity for the IL-15R $\beta\gamma$ C receptors compared to the native hIL-15 polypeptide.

50

6. The isolated hIL-15 variant of claim 1, wherein the amino acid substitution is of N to D at position 72 of the mature human IL-15 sequence.

55

7. The isolated hIL-15 variant of claim 6, wherein the amino acid substitution results in an IL-15 variant that has IL-15 agonist activity or increased binding activity for the IL-15R $\beta\gamma$ C receptors compared to the native hIL-15 polypeptide.

60

8. An isolated human interleukin 15 (hIL-15) variant, comprising an amino acid substitution at position 72 of a mature human IL-15 sequence.

9. The isolated hIL-15 variant of claim 8, wherein the amino acid substitution is selected from the group consisting of alanine, aspartic acid, and arginine.

* * * * *