


Volume 56 No. 1 1999

CODEN PTCEDJ ISSN 0167-6857

# **Plant Cell, Tissue and Organ Culture**

**An International Journal on  
the Cell Biology of Higher Plants**



**Kluwer Academic Publishers**



# Plant Cell, Tissue and Organ Culture

An International Journal on the Cell Biology of Higher Plants

## Editorial Board Chair

**Daniel C.W. Brown**, Agriculture and Agri-Food Canada, Southern Crop Protection & Food Research Centre, 1391 Sandford Street, London, Ontario, N5V 4T3, Canada. tel: +1 519 457 1470; fax: +1 519 457 3997; email: brown dc@em.agr.ca

## Science Editors

**Jan M. Bonga**, Dept. of Natural Resources, Canadian Forest Service, P.O. Box 4000, Fredericton, New Brunswick, E3B 5P7, Canada; email: jbonga@fcmr.forestry.ca

**Daniel Cohen**, The Horticulture and Food Research Institute of New Zealand Ltd, Mt Albert Research Centre, Private Bag 92-169, Auckland, New Zealand; email: dcohen@marccri.marc.cri.nz

**Didier Courtois**, Centre Recherche Nestlé, 101 Avenue Gustave Eiffel, BP 9715, 37097 Notre Dame d'Oé, France; email: didier.courtois@frtour01.nestrd.ch

**Richard E. Litz**, University of Florida, IFAS, Tropical Research and Education Center, 18905 SW 280 St., Homestead, FL 33031, USA; email: rel@gnv.ifas.ufl.edu

**Wayne A. Parrott**, Dept. of Crop and Soil Science, The University of Georgia, Athens, GA 30602, USA; email: wparrott@uga.cc.uga.edu

**John E. Preece**, Dept. Plant & Soil Sciences, Southern Illinois University, Carbondale, IL 62901, USA; email: jpreece@siu.edu

**Alan H. Scragg**, Dept. of Biological Sciences, University of the West of England, Coldharbour Lane, Frenchay Campus, Bristol BS16 1QY, UK; email: a-scragg@wpg.uwe.ac.uk

## Editorial Board

**Aitken-Christie, J.**, Carter Holt Harvey Forests Ltd., Rotorua, New Zealand

**Allan, E.J.**, Aberdeen University, UK

**Arcioni, S.**, Istituto di Ricerche sul Miglioramento Genetico delle Piante Foraggere del CNR, Perugia, Italy

**Austin Burns, J.**, Monsanto Company, St. Louis, USA

**Becwar, M. R.**, Westvaco Corporation, Summerville, USA

**Benson, E.E.**, University of Abertay, Dundee, Scotland, UK

**Berlin, J.**, Gesellschaft für Biotechnologische Forschung mbH, Braunschweig, Germany

**Brettell, R.I.S.**, CSIRO, Canberra, Australia

**Cassells, A.C.**, University College, Cork, Ireland

**Chávez Avila, V.M.**, Jardín Botánico del Instituto de Biología, UNAM, Ciudad Universitaria, Mexico

**Chibbar, R.N.**, National Research Council of Canada, Saskatoon, Canada

**Clemente, T.E.**, University of Nebraska, Lincoln, USA

**Collin, H.A.**, University of Liverpool, UK

**Compton, M.E.**, University of Wisconsin, Platteville, USA

**Debergh, P.**, University of Gent, Gent, Belgium

**De Klerk, G.J.**, Centre for Plant Tissue Culture Research, Lisse, The Netherlands

**DiCosmo, F.**, University of Toronto, Ontario, Canada

**Doran, P.M.**, University of New South Wales, Sydney, Australia

**Douglas, G.**, Agriculture and Food Development Authority, Dublin, Ireland

**Drew, R.A.**, Griffith University, Nathan, Australia

**Economou, A.S.**, Aristotle University, Thessaloniki, Greece

**Grosser, J.W.**, University of Florida, Lake Alfred, USA

**Harms, C.T.**, Novartis Crop Protection, Protection AG, Basel, Switzerland

**Hunter, C.S.**, University of the West of England, Bristol, UK

**Kane, M.E.**, University of Florida, Gainesville, USA

**Komamine, A.**, Japan Women's University, Tokyo, Japan

**Korban, S.S.**, University of Illinois, Urbana, USA

**Kozai, T.**, Chiba University, Japan

**Kumar, P.P.**, National University of Singapore, Singapore

**Lindsey, K.**, University of Durham, UK

**Marks, T.R.**, Horticultural Research International, East Malling, UK

**McComb, J.**, Murdoch University, W. Australia

**Merkle, S.A.**, University of Georgia, Athens, USA

**Morris, P.**, Institute for Grassland and Environmental Research, Aberystwyth, UK

**Navarro, L.**, IVIA, Valencia, Spain

**Pliego Alfaro, F.**, University of Málaga, Málaga, Spain

**Power, J.B.**, University of Nottingham, Nottingham, UK

**Saxena, P.K.**, University of Guelph, Ontario, Canada

**Smith, M.A.L.**, University of Illinois, Urbana, USA

**Smith, M.K.**, Queensland Horticulture Institute, Nambour, Australia

**Songstad, D.D.**, Monsanto, St. Louis, USA

**Thorpe, T.A.**, University of Calgary, Canada

**Verpoorte, R.**, Leiden University, The Netherlands

**Vieitez, A.M.**, CSIC, Santiago de Compostela, Spain

**Wetzstein, H.Y.**, University of Georgia, Athens, USA

**Wilde, H.D.**, Westvaco, Summerville, USA

# Plant Cell, Tissue and Organ Culture

An International Journal on  
the Cell Biology of Higher Plants

Volume 56 / 1999

STEENBOCK  
MEMORIAL LIBRARY

OCT 15 1999

U.W.-MADISON



Kluwer Academic Publishers

Dordrecht / Boston / London



# Plant Cell, Tissue and Organ Culture

An International Journal on the Cell Biology of Higher Plants

## Editorial Board Chair

**Daniel C.W. Brown**, Agriculture and Agri-Food Canada, Southern Crop Protection & Food Research Centre, 1391 Sandford Street, London, Ontario, Canada N5V 4T3. tel: +1 519 457 1470; fax: +1 519 457 3997; email: brown dc@em.agr.ca

## Science Editors

**Jan M. Bonga**, Dept. of Natural Resources, Canadian Forest Service, P.O. Box 4000, Fredericton, New Brunswick E3B 5P7, Canada; email: jbonga@fcmr.forestry.ca

**Daniel Cohen**, The Horticulture and Food Research Institute of New Zealand Ltd, Mt Albert Research Centre, Private Bag 92 169, Auckland, New Zealand; email: dcohen@marccri.marc.cri.nz

**Didier Courtois**, Centre Recherche Nestlé, 101 avenue Gustave Eiffel, BP 9715, 37097 Notre Dame d'Oé, France; email: didier.courtois@frtour01.nestrd.ch

**Richard E. Litz**, University of Florida, IFAS, Tropical Research and Education Center, 18905 SW 280 St., Homestead, FL 33031, USA; email: rel@gnv.ifas.ufl.edu

**Wayne E. Parrott**, Dept. of Crop and Soil Science, The University of Georgia, Athens, GA 30602-7272, USA; email: wparrott@uga.cc.uga.edu

**John E. Preece**, Dept. Plant & Soil Sciences, Southern Illinois University, Carbondale, IL 62901, USA; email: jpreece@siu.edu

**Alan H. Scragg**, Dept. of Biological Sciences, University of the West of England, Coldharbour Lane, Frenchay Campus, Bristol BS 16 1QY, UK; email: a-scragg@wpg.uwe.ac.uk

## Editorial Board

**Aitken-Christie, J.**, Carter Holt Harvey Forests Ltd., Rotorua, New Zealand

**Allan, E.J.**, Aberdeen University, Scotland

**Arcioni, S.**, Istituto di Ricerche sul Miglioramento Genetico delle Piante Foraggere del CNR, Perugia, Italy

**Austin Burns, J.**, Monsanto Company, St. Louis, USA

**Becwar, M.R.**, Westvaco Corporation, Summerville, USA

**Benson, E.E.**, University of Abertay, Dundee, Scotland, UK

**Berlin, J.**, Gesellschaft für Biotechnologische Forschung mbH, Braunschweig, Germany

**Brettell, R.I.S.**, CSIRO, Canberra, Australia

**Cassells, A.C.**, University College, Cork, Ireland

**Chávez Avila, V.M.**, Jardín Botánico del Instituto de Biología, UNAM, Ciudad Universitaria, Mexico

**Chibbar, R.N.**, National Research Council of Canada, Saskatoon, Canada

**Clemente, T.E.**, University of Nebraska, Lincoln, USA

**Collin, H.**, University of Liverpool, UK

**Compton, M.E.**, University of Wisconsin, Platteville, USA

**Debergh, P.**, University of Gent, Gent, Belgium

**De Klerk, G.J.**, Centre for Plant Tissue Culture Research, Lisse, The Netherlands

**DiCosmo, F.**, University of Toronto, Ontario, Canada

**Doran, P.M.**, University of New South Wales, Sydney, Australia

**Douglas, G.**, Agriculture and Food Development Authority, Dublin, Ireland

**Drew, R.A.**, Griffith University, Nathan, Australia

**Economou, A.S.**, Aristotle University, Thessaloniki, Greece

**Grosser, J.W.**, University of Florida, Lake Alfred, USA

**Harms, C.T.**, Novartis Crop Protection AG, Basel, Switzerland

**Hunter, C.S.**, University of the West of England, Bristol, UK

**Kane, M.E.**, University of Florida, Gainesville, USA

**Komamine, A.**, Japan Women's University, Tokyo, Japan

**Korban, S.S.**, University of Illinois, Urbana, USA

**Kozai, T.**, Chiba University, Japan

**Kumar, P.P.**, National University of Singapore, Singapore

**Lindsey, K.**, University of Durham, UK

**Marks, T.R.**, East Malling Horticulture Research International, UK

**McComb, J.**, Murdoch University, W. Australia

**Merkle, S.A.**, University of Georgia, Athens, USA

**Morris, P.**, Institute for Grassland and Environment Research, Aberystwyth, UK

**Navarro, L.**, IVIA, Valencia, Spain

**Pliego Alfaro, F.**, University of Málaga, Málaga, Spain

**Power, J.B.**, University of Nottingham, Nottingham, UK

**Saxena, P.K.**, University of Guelph, Ontario, Canada

**Smith, M.A.L.**, University of Illinois, Urbana, USA

**Smith, M.K.**, Queensland Horticulture Institute, Nambour, Australia

**Songstad, D.D.**, Pioneer Hi-Bred International, Johnston, USA

**Thorpe, T.A.**, University of Calgary, Calgary, Canada

**Verpoorte, R.**, Leiden University, Leiden, The Netherlands

**Vieitez, A.M.**, CSIC, Santiago de Compostela, Spain

**Wetzstein, H.**, University of Georgia, Athens, USA

**Wilde, H.D.**, Westvaco, Summerville, USA

## Journals Editorial Office

Manuscripts submitted for publication and communications concerning editorial matters should be sent to:

**Journals Editorial Office**, *Plant Cell, Tissue and Organ Culture*, P.O. Box 990, 3300 AZ, Dordrecht, The Netherlands  
tel.: +31 78 6392 392 fax: +31 78 6392 555 e-mail: editdept@wkap.nl

ISSN 0167-6857

All Rights Reserved

©1999 Kluwer Academic Publishers

No part of the material protected by this copyright notice may be reproduced or utilised in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without written permission from the copyright owner.

Printed in the Netherlands



# Plant Cell, Tissue and Organ Culture

An International Journal on the Cell Biology of Higher Plants

CONTENTS Volume 56 No. 1 1999

## Somatic embryogenesis in Canary Island date palm

L.T.L. Huong, M. Baiocco, B.P. Huy, B. Mezzetti, R. Santilocchi, P. Rosati

1-7

## Protoplast and leaf explant culture of *Lycopersicon cheesmanii* and salt tolerance of protoplast-derived calli

N. Dorion, N. Wies, A. Burteaux, C. Bigot

9-16

## Plant regeneration from seedling explants of *Eucalyptus grandis* × *E. urophylla*

L.P. Barrueto Cid, A.C.M.G. Machado, S.B.R.C. Carvalheira, A.C.M. Brasileiro

17-23

## A neural network based pattern recognition system for somatic embryos of Douglas fir

C. Zhang, R. Timmis, W.-S. Hu

25-35

## The use of glufosinate as a selective agent in *Agrobacterium*-mediated transformation of soybean

Z. Zhang, A. Xing, P. Staswick, T.E. Clemente

37-46

## Rapid plant regeneration from callus cultures of *Plumbago zeylanica*

G.R. Rout, C. Saxena, S. Samantaray, P. Das

47-51

## Research notes

### BA enhances the germination of oil palm somatic embryos derived from embryogenic suspension cultures

F. Aberlenc-Bertossi, M. Noirot, Y. Duval

53-57

### *In vitro* propagation of *Syzygium travancoricum* Gamble – an endangered tree species

A. Anand, C. Srinivasa Rao, P. Balakrishna

59-63

### Assessment of Al<sup>3+</sup> availability in callus culture media for screening tolerant genotypes of *Cynodon dactylon*

S. Ramgareeb, M.P. Watt, C. Marsh, J.A. Cooke

65-68

*Plant Cell, Tissue and Organ Culture* is an international monthly journal that publishes original results of fundamental studies on the behavior of plant cells, tissues and organs in vitro. Topics include growth and developmental, anatomical, histological and ultrastructural aspects, physiology and biochemistry, genetics and breeding, phytopathological aspects and biotechnology. Methodological and technical aspects of the culture of plant cells, tissues and organs will be considered for publication on the basis of novelty and potential contributions to studies of the in vitro biology of these systems.

*In the USA:* This journal is registered at the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923.

Authorization does not extend to other kinds of copying, such as that for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

*In the rest of the world:* Permission to photocopy must be obtained from the copyright owner. Please apply to Kluwer Academic Publishers, P.O. Box 17, 3300 AA Dordrecht, The Netherlands.

Subscription price per year (1999): Subscription price NLG 2480.00/USD 1240.00, including postage and handling.

US Mailing Agent: Mercury Airfreight International Ltd., 365 Blair Road, Avenel, NJ 07001.

Published by Kluwer Academic Publishers, Spuiboulevard 50, P.O. Box 17, 3300 AA Dordrecht, The Netherlands, and 101 Philip Drive, Norwell, MA 02061, USA.

Postmaster: Please send all address corrections to: *Plant Cell, Tissue and Organ Culture*  
c/o Mercury Airfreight International Ltd., 365 Blair Road, Avenel, NJ 07001, USA.

Corteva Exhibit 2079 - Page 6 of 16  
Inari v. Corteva - PGR2023-00022



## The use of glufosinate as a selective agent in *Agrobacterium*-mediated transformation of soybean

Zhanyuan Zhang<sup>1</sup>, Aiqiu Xing<sup>2</sup>, Paul Staswick<sup>1</sup> & Thomas E. Clemente<sup>1,2,\*</sup>

<sup>1</sup>Department of Agronomy and <sup>2</sup>Center for Biotechnology, University of Nebraska-Lincoln, Plant Transformation Core Research Facility, E303 Beadle Center, Lincoln, Nebraska, 68588-0665; (\*requests for offprints; e-mail: tclemente1@unl.edu)

Received 19 May 1998; accepted in revised form 10 March 1999

**Key words:** *Agrobacterium tumefaciens*, *bar*, *Glycine max*, phosphinothricin

### Abstract

The soybean transformation procedure using the *Agrobacterium*-cotyledonary node transformation system and the *bar* gene as the selectable marker coupled with glufosinate as a selective agent is described. Soybean cotyledonary explants were derived from 5 day old seedlings and co-cultivated with *Agrobacterium tumefaciens* for 3 days. Explants were cultured on Gamborg's B5 medium supplemented with 1.67 mg l<sup>-1</sup> BAP and glufosinate at levels of 3.3 mg l<sup>-1</sup> or 5.0 mg l<sup>-1</sup> for 4 weeks. After 4 weeks explants were subcultured to medium containing MS major and minor salts and B5 vitamins (MS/B5) supplemented with 1.0 mg l<sup>-1</sup> zeatin-riboside, 0.5 mg l<sup>-1</sup> GA<sub>3</sub> and 0.1 mg l<sup>-1</sup> IAA amended with 1.7 mg l<sup>-1</sup> or 2.0 mg l<sup>-1</sup> glufosinate. Elongated shoots were rooted on a MS/B5 rooting medium supplemented with 0.5 mg l<sup>-1</sup> NAA without further glufosinate selection. Plantlets were transplanted to soil and grown to maturity and set seed in the greenhouse. Primary transformants and their progeny were characterized by Southern blot analysis and a leaf paint assay.

**Abbreviations:** AS – acetosyringone; BAP – 6-benzyl-aminopurine; IAA – indole-3-acetic acid; NAA – naphthalene acetic acid

### Introduction

Introduction of agronomically important traits into soybean (*Glycine max* [L.] Merr.) has been accomplished via plant transformation (Trick et al., 1997). Those traits include herbicide tolerance (Padgett et al., 1995), amino acid modification (Falco et al., 1995), virus resistance (Di et al., 1996) and insect resistance (Stewart et al., 1996). Development of more efficient procedures for *Agrobacterium*-mediated soybean transformation could make it possible for routine production of transgenic soybean lines with simple inserts that are inherited as a single functional locus to facilitate breeding.

*Agrobacterium*-mediated transformation of soybean has been very difficult. Only a few studies on this system have been reported to date. Plant regeneration of soybean via the cotyledonary node explant was first

reported by Cheng et al. (1980). An *Agrobacterium*-mediated transformation system exploiting this explant was first reduced to practice by Hinchey et al. (1988). Di et al. (1996) used this system to introduced the coat protein of bean pod mottle virus into soybean. Kanamycin is the only selective agent that has been published for use in the system.

The herbicide resistance gene, *bar* (Thompson et al., 1987), has been widely used as a selectable marker in plant transformation systems (D'Halluin et al., 1992). The *bar* gene encodes for phosphinothricin acetyltransferase (PAT) which detoxifies glufosinate, the active ingredient in the herbicide Liberty® (AgrEvo USA). To our knowledge, no work has been reported to date evaluating this selectable marker in a soybean transformation system. Due to the rapid translocation mechanism of glufosinate in soybean tissue through both the xylem and phloem (Shelp et al.,

1992), we thought that glufosinate selection would provide efficient recovery of transformed tissue with the *Agrobacterium*-cotyledonary system. We report here the use of the *bar* gene as a selectable marker for *Agrobacterium*-mediated transformation of soybean.

## Materials and methods

### Plant material

Soybean seeds, genotype A3237 (Asgrow Seed Company; Des Moines, Iowa), were surface sterilized by an overnight exposure to chlorine gas (Di et al., 1996). Sterilized seeds were germinated, in  $100 \times 20$  mm Petri dishes, on Gamborg's B5 basal medium (B5) (Gamborg et al., 1968) supplemented with 2% sucrose, pH 5.8. The plates were stacked 5 high and placed in plastic bags in which 4, approx. 3 inch, slits were made with scissors. Seeds were germinated for 5 days in a growth room, at 24 °C, 18/6 light regime, under a light intensity of approximately  $150 \mu\text{mol s}^{-1} \text{m}^{-2}$ . The light source was a mixture of cool white (Sylvania model # F72T12/CW/VHO) and Gro-lux® bulbs (Sylvania model # F72T12/GRO/VHO).

### Plant transformation vectors

The binary vector, pPTN101 (Figure 1A), is a derivative of pGPTV-*bar* (Becker et al., 1992), carrying the GUS cassette from pE7131-GUS (Mitsuhashi et al., 1996). The binary vector pPTN125 (Figure 1B) is a derivative of pGPTV-*bar* (Becker et al., 1992) carrying a GUS cassette under the control of the soybean vegetative storage protein B (*vspB*) promoter (Rhee and Staswick, 1992). An intermediate vector carrying the *bar* gene under the control of the 35S promoter was generated by isolating the *bar* open reading frame (ORF) from pDM302 (Cao et al., 1992) as a *Sma* I fragment and subcloning it into the plant expression cassette pRTL2 (Carrington, 1990). The resultant vector pPTN117 (not shown) contained the *bar* ORF in a translation fusion with the N-terminus of the Tobacco Etch Virus (TEV) polyprotein coupled with the TEV translation enhancer element. A third binary was assembled that harbored a 35S *bar* cassette, a derivative of the *bar* cassette within pPTN117, lacking the N-terminus of the TEV polyprotein, hence, the *bar* ORF is directly ligated to the TEV translational enhancer element (Carrington, 1990). This latter *bar* cassette was generated by introducing a *Nco* I site at the start of translation of the *bar* ORF, by

PCR. Primers used were referred to as BAR-5: 5' CCCGGGATCTACCATGGCTAGCCCAGCCCAGA-ACGAC-3' and BAR-3: 5'-TCATCAGATCTCGGT-GACGG-3'. This manipulation introduced an Ala residue between the Met and Ser residues of the native *bar* protein. The PCR product was subcloned into pPTN117 as a *Nco* I/ *Kpn* I fragment replacing the 5' region of the *bar* ORF within the vector with the engineered fragment. The resultant vector is referred to as pPTN129 (not shown). The 35S *bar* cassette within pPTN129 was sequenced at the University of Nebraska-Lincoln DNA Sequencing Core Research Facility to confirm the integrity of the PCR reaction. The 35S *bar* cassette from pPTN129 was subcloned as a *Hind* III fragment into the binary vector pPZP202 (Hajdukiewicz et al., 1994) that harbored a 35S GUS cassette, the resultant vector is referred to as pPTN140 (Figure 1C).

### *Agrobacterium tumefaciens* strains

The binary vectors were mobilized into *Agrobacterium tumefaciens* strains EHA105 (Hood et al., 1993) or EHA101 (Hood et al., 1986) by triparental mating (Ditta et al., 1980). Binary vectors pPTN101 and pPTN125 were introduced into EHA105, and transconjugants were selected on 25 mg l<sup>-1</sup> chloramphenicol [EHA105 chromosomal drug marker] and 50 mg l<sup>-1</sup> kanamycin [Binary vector drug marker]. Binary vector pPTN140 was introduced into EHA101, and transconjugants were selected on 25 mg l<sup>-1</sup> chloramphenicol [EHA101 chromosomal drug marker], 50 mg l<sup>-1</sup> kanamycin [Ti plasmid pEHA101 drug marker], 100 mg l<sup>-1</sup> spectinomycin [binary vector drug marker] and 100 mg l<sup>-1</sup> streptomycin [binary vector drug marker]. Integrity of the binary vectors in the respective *Agrobacterium* strains was confirmed by plasmid rescue.

### *Agrobacterium* culturing conditions

*Agrobacterium* cultures were grown in YEP medium (10 g l<sup>-1</sup> peptone, 5 g l<sup>-1</sup> yeast extract and 5 g l<sup>-1</sup> NaCl, pH 7.0) amended with the appropriate antibiotics to an OD<sub>650</sub> = 0.6 to 0.8 at 27 °C. The bacterial cultures were centrifuged at 3,500 rpm for 10 min., and the pellets resuspended to a final OD<sub>650</sub> = 0.6 to 0.8 in 1/10 Gamborg's B5 medium (Gamborg et al., 1968) amended with 1.67 mg l<sup>-1</sup> BA, 0.25 mg l<sup>-1</sup> GA<sub>3</sub>, 200 μM acetosyringone (AS) and 3% sucrose. The medium was buffered with 20 mM MES, pH 5.4.




**Figure 1C**

**Figure 1.** Binary vectors utilized in the experiments. See Materials and methods section for description of the binary vectors. Abbreviations: RB – right border, LB – left border, T35s and pAg7 are polyadenylation signals, Pnos – nopaline synthase promoter, aadA – bacterial drug resistance marker for spectinomycin and streptomycin.

All growth regulators, vitamin components and AS were filter sterilized post autoclaving.

#### Plant transformation

Cotyledonary explants were prepared from the 5-day-old soybean seedlings by making a horizontal slice through the hypocotyl region, approximately 3–5 mm below the cotyledon. A subsequent vertical slice was made between the cotyledons, and the embryonic axis was removed. This manipulation generated 2 cotyledonary node explants. Approximately 7–12 vertical

slices were made on the adaxial surface of the explant about the area encompassing 3 mm above the cotyledon/hypocotyl junction and 1 mm below the cotyledon/hypocotyl junction. Explant manipulations were conducted with a No. 15 scalpel blade.

Explants were immersed in the *Agrobacterium* inoculum for 30 min and then co-cultured on 100 × 15 mm Petri plates containing the *Agrobacterium* resuspension medium solidified with 0.5% purified agar (BBL Cat # 11853). The co-cultivation plates were overlaid with a piece of Whatman #1 filter paper (Mullins et al., 1990; Janssen and Gardner,

1993; Zhang et al., 1997). The explants (5 per plate) were cultured adaxial side down on the co-cultivation plates, that were overlaid with filter paper, for 3 days at 24 °C, under an 18/6 hour light regime with an approximate light intensity of 80  $\mu\text{mol s}^{-1} \text{m}^{-2}$  (F17T8/750 cool white bulbs, Litetronics®). The co-cultivation plates were wrapped with Parafilm®.

Following the co-cultivation period explants were briefly washed in B5 medium supplemented with 1.67 mg l<sup>-1</sup> BAP, 3% sucrose, 500 mg l<sup>-1</sup> ticarcillin and 100 mg l<sup>-1</sup> cefotaxime. The medium was buffered with 3 mM MES, pH 5.6. Growth regulator, vitamins and antibiotics were filter sterilized post autoclaving. Following the washing step, explants were cultured (5 per plate) in 100 × 20 mm Petri plates, adaxial side up with the hypocotyl imbedded in the medium, containing the washing medium solidified with 0.8% purified agar (BBL Cat # 11853) amended with either 3.3 or 5.0 mg l<sup>-1</sup> glufosinate (AgrEvo USA). This medium is referred to as shoot initiation medium (SI). Plates were wrapped with 3M pressure sensitive tape (Scotch™, 3M, USA) and cultured under the environmental conditions used during the seed germination step.

After 2 weeks of culture, the hypocotyl region was excised from each of the explants, and the remaining explant, cotyledon with differentiating node, was subsequently subcultured onto fresh SI medium. Following an additional 2 weeks of culture on SI medium, the cotyledons were removed from the differentiating node. The differentiating node was subcultured to shoot elongation medium (SE) composed of Murashige and Skoog (MS) (1962) basal salts, B5 vitamins, 1 mg l<sup>-1</sup> zeatin-riboside, 0.5 mg l<sup>-1</sup> GA<sub>3</sub> and 0.1 mg l<sup>-1</sup> IAA, 50 mg l<sup>-1</sup> glutamine, 50 mg l<sup>-1</sup> asparagine, 3% sucrose and 3 mM MES, pH 5.6. The SE medium was amended with either 1.7 or 2.0 mg l<sup>-1</sup> glufosinate. The explants were subcultured biweekly to fresh SI medium until shoots reached a length greater than 3 cm. The elongated shoots were rooted on Murashige and Skoog salts with B5 vitamins, 1% sucrose, 0.5 mg l<sup>-1</sup> NAA without further selection in either Magenta boxes® or Sundae cups (Industrial Soap Company, St. Louis MO).

#### *Southern blot analysis*

Total DNA was extracted from soybean leaf tissue as described by Dellaporta et al. (1983). Ten or 20  $\mu\text{g}$  of DNA were digested with a restriction enzyme for which only one recognition sequence is present

with in the T-DNA element [either *Eco* RI or *Hind* III]. The digested genomic DNA was separated by agarose gel electrophoresis prior to transfer to Zeta-Probe® GT nylon membrane (BIO-RAD catalog # 162-0196). The DNA was fixed to the membrane by UV cross linking. Hybridization and washing conditions for Southern blot analysis (Southern, 1975) followed the Zeta-Probe® GT manufacturer's instructions. A DNA fragment containing the GUS ORF was used to generate a <sup>32</sup>P-labeled probe. The probe was prepared by random primer synthesis incorporating <sup>32</sup>P-dCTP utilizing Prime-It® II kit (Stratagene, catalog # 300385). The synthesized probe was purified by passing the reaction product over NucTrap® columns (Stratagene, catalog # 400701).

#### *GUS assays*

Fluorometric GUS assays (Jefferson et al., 1987) were conducted by grinding leaf tissue in lysis buffer composed of 50 mM NaPO<sub>4</sub> (pH 7.0), 1 mM EDTA, 0.1% Triton-X100, 10 mM  $\beta$ -mercaptoethanol and 0.1% sarkosyl. Protein samples were quantified using a Bio-Rad protein assay kit (Catalog # 500-0002). GUS fluorometric reactions were conducted using 1 mM 4-methyl umbeliferone as the substrate. Reactions were carried out at 37 °C, and the reactions were stopped by the addition of 0.2 M Na<sub>2</sub>CO<sub>3</sub> at 30 min intervals. Fluorescence was monitored using a DyNA Quant 200 fluorometer (Hoefer Scientific).

Histochemical GUS assays (Jefferson et al., 1987) were conducted by immersing soybean tissue in a substrate solution composed of 0.1 M NaHPO<sub>4</sub> buffer (pH 7.0), 0.5 mM K<sub>3</sub>[Fe(CN)<sub>6</sub>], 0.5 mM K<sub>4</sub>[Fe(CN)<sub>6</sub>], 10 mM EDTA and 800 mg l<sup>-1</sup> X-Gluc. Methanol and Triton-X100 were subsequently added to final concentrations of 20% and 0.06%, respectively. Soybean tissue was allowed to incubate in the X-Gluc substrate for 3 to 8 hours at 37 °C. The tissue was subsequently cleared in 70% ethanol prior to visualization.

*Leaf painting.* Progeny were screened for tolerance to the herbicide Liberty® by application of a 0.1% solution of the herbicide with a cotton swab to the upper surface of the first unifoliate leaf 3 weeks post germination. A 200 g l<sup>-1</sup> stock of Liberty® was diluted in water for the leaf painting assay. Leaf tissue was scored for herbicide tolerance 5 days post application.

Table 1. Glufosinate kill response.

| Selection pressure                      | Co-cultivation          | Total number of explants | Regenerating explants | % regeneration |
|-----------------------------------------|-------------------------|--------------------------|-----------------------|----------------|
| <i>Experiment 1</i>                     |                         |                          |                       |                |
| 3.3 mg l <sup>-1</sup> (4 weeks)        | No <i>Agrobacterium</i> | 50                       | 2                     | 4              |
| 3.3 mg l <sup>-1</sup> (4 weeks)        | <i>Agrobacterium</i>    | 40                       | 34                    | 85             |
| <i>Experiment 2</i>                     |                         |                          |                       |                |
| 3.3 mg l <sup>-1</sup> (4 weeks)        | No <i>Agrobacterium</i> | 50                       | 0                     | 0              |
| 3.3 mg l <sup>-1</sup> (4 weeks)        | <i>Agrobacterium</i>    | 50                       | 40                    | 80             |
| <i>Experiment 3</i>                     |                         |                          |                       |                |
| 5.0 to 3.3 mg l <sup>-1</sup> (4 weeks) | No <i>Agrobacterium</i> | 50                       | 0                     | 0              |
| 5.0 to 3.3 mg l <sup>-1</sup> (4 weeks) | <i>Agrobacterium</i>    | 50                       | 42                    | 84             |

Selection pressure column refers to the level of glufosinate added to the shoot induction medium. In Experiment 3, shoot initiation medium contained 5 mg l<sup>-1</sup> glufosinate during the first 2 weeks and the following 2 weeks the selection level was decreased to 3.3 mg l<sup>-1</sup>. Co-cultivation column indicates if the explants were exposed to *Agrobacterium* (EHA105 carrying pPTN101) or merely mock-inoculated. Regenerating explants and % regeneration columns, refer to the total number of explants in which shoots were initiated and percent regeneration based on total explants put into culture, respectively.

Table 2. Summary of 5 independent transformation attempts under the higher glufosinate selection regime.

| Experiment number | N   | Total no. in G.H. | I.E. in G.H. | GUS+ in G.H. | %GUS+ efficiency |
|-------------------|-----|-------------------|--------------|--------------|------------------|
| 246               | 100 | 7                 | 4            | 3            | 3.0              |
| 247               | 100 | 3                 | 3            | 0            | 0                |
| 248               | 200 | 2                 | 2            | 0            | 0                |
| 249               | 150 | 3                 | 3            | 1            | 0.7              |
| 252               | 200 | 9                 | 3            | 1            | 0.5              |

N refers to the total number of explants put into culture. Total No. in G.H. indicates the total number of soybean plants that were established in the greenhouse following selection. 'I.E. in G.H.' indicates the number of independent lines (plants arising from independent explants) represented. 'GUS+ in G.H.' represents the total number of independent soybean transformants co-expressing GUS. %GUS+ Efficiency indicates the efficiency of recovering independent soybean transformants co-expressing GUS.

## Results and discussion

### Glufosinate response

Kill curve experiments were used to determine the dose response of soybean cotyledonary explants to glufosinate. Our objective was to ascertain glufosinate levels that would allow for differential growth between transformed and non-transformed tissues. Glufosinate levels ranging from 2 mg l<sup>-1</sup> up to 33 mg l<sup>-1</sup> were tested. Explants were prepared for these studies as described in M & M, Plant Transformation section. A total of 10 plates were evaluated per glufosinate

Table 3. GUS activity in primary soybean transformants carrying T-DNA from pPTN101.

| Plant ID  | GUS Activity (nM 4-MU/min./μg total protein) |
|-----------|----------------------------------------------|
| 207-5     | 91.00±41.44                                  |
| 210-16    | 36.15±29.49                                  |
| 211-2B    | 43.75±9.55                                   |
| 207-5-2   | 12.80*                                       |
| 210-11    | 0.00±0.00                                    |
| 209-15    | 24.50±3.25                                   |
| 209-15B   | 31.10±3.25                                   |
| Wild Type | 0.01±0.00                                    |

Plant ID column refers to the designation of the soybean putative transformant. Wild type is a A3237 non-transformant. GUS activity is the mean of 2 independent assays ± S.D.

\* indicates only 1 tissue sample was assayed.

level per experiment (i.e. 50 explants per treatment). Explants were scored after 2 and 4 weeks on selection. Explants cultured on glufosinate levels greater than 10 mg l<sup>-1</sup> failed to regenerate (Data not shown). We decided to focus our efforts on selection levels 3.3 or 5.0 mg l<sup>-1</sup> during shoot initiation. A different response was observed depending on whether the explants were inoculated with *Agrobacterium* and subsequently co-cultivated for 3 days or merely mock inoculated and co-cultivated for 3 days (Table 1). A mock inoculation was conducted by placing the cotyledonary explants in the bacterial resuspension medium [see M & M, *Agrobacterium* culturing con-




Figure 2. Primary transformants carry the T-DNA of pPTN140. Panel 1A: Histochemical GUS expression in mature leaflet of transformant (left) and control (right). Panel 1B: Histochemical GUS expression in developing pericarp (right) and control pericarp (left). Panel 1C: Progeny screen for herbicide tolerance. Black arrow pointing to herbicide susceptible response in T<sub>1</sub> plants and red arrows pointing to the herbicide tolerant response in T<sub>1</sub> plants following leaf painting with a 0.1% glufosinate solution.

ditions] without the *Agrobacterium*. The explants that were exposed to *Agrobacterium* regenerated on 3.3 or 5.0 mg l<sup>-1</sup> glufosinate, while those explants that were mock inoculated, were either significantly reduced in regeneration, or regeneration was impeded on the levels of glufosinate tested, across 3 independent experiments (Table 1). This observation suggests that is important to incorporate all the steps of a transformation system when evaluating alternative selection agents and/or conducting experiments to optimize this soybean transformation system.

#### Selection regimes

Two selection regimes were employed for the recovery of whole plants: (1) 3.3 mg l<sup>-1</sup> glufosinate during the shoot initiation culture period and stepping down to 1.7 mg l<sup>-1</sup> during shoot elongation, and; (2) 5.0 mg l<sup>-1</sup> during the shoot initiation culture period and stepping down to 2.0 mg l<sup>-1</sup> during shoot elongation. After 4 weeks on SI medium under 3.3 mg l<sup>-1</sup> glufosinate, 70–80% of the explants survived and were transferred to SE medium. Under the more stringent

## Soybean Primary Transformants Derived from pPTN101




Figure 3. Southern blot analysis of soybean transformants carrying the T-DNA of pPTN101. Total genomic DNA was digested with *Hind* III. The membrane was probed with the GUS ORF. Lane 1: Molecular marker, Lanes 2-6 soybean plants carrying the T-DNA of pPTN101. Lane 7: 10  $\mu$ g of A3237 wild type DNA. Lanes 8 and 9: are 12.5 pg and 25 pg, respectively, of pPTN101 digested with *Hind* III.

selection regime, 70–80% of the explants regenerated after 2 weeks on SI medium. However, after the second subculture period, only 40–50% of the explants survived and were subcultured to SE medium.

GUS-positive soybean plants were recovered at frequencies ranging from 0.0% to 0.5% and 0.0% to 3.0%, across independent transformation experiments under the lower and higher stringent selection regimes, respectively. Data from 5 transformation attempts utilizing EHA101 (pPTN140) employing the higher selection regime are given in Table 2. Histochemical GUS expression and herbicide tolerance in transformed progeny carrying pPTN140 are shown in Figure 2. GUS expression of primary transform-

ants derived from the lower selection scheme carrying pPTN101 is quantified in Table 3. The events listed in Table 3 were apparent chimeras (see below), with only one being a germline transformation event, 207-5-2.

### Molecular characterization of transgenic plants

Southern blot analyses on primary transformants carrying pPTN101 and pPTN125 are shown in Figures 3 and 4, respectively. The Southern data confirms the integration of the transgenes in the soybean genome.

Ten plants, derived from the lower selection regime, (arising from independent explants) were characterized by Southern blot analysis. From the 10

## Soybean Primary Transformants Derived from pPTN125




Figure 4. Southern blot analysis of soybean transformants carrying the T-DNA of pPTN125. Total genomic DNA was digested with *EcoR* I. The membrane was probed with the GUS ORF. Lane 1: Molecular marker, Lanes 2–5: soybean plants carrying the T-DNA of pPTN125. Lane 6: 20  $\mu$ g of A3237 wild type DNA. Lanes 7 and 8: 18 pg and 54 pg, respectively, of pPTN125 digested with *EcoR* I.

Table 4. Segregation analysis on primary transformants derived from glufosinate selection.

| Event  | T <sub>0</sub><br>GUS | T <sub>1</sub><br>GUS <sup>+</sup> /HT | T <sub>1</sub><br>GUS <sup>-</sup> /HT | T <sub>1</sub><br>GUS <sup>-</sup> /HS |
|--------|-----------------------|----------------------------------------|----------------------------------------|----------------------------------------|
| 246-2C | +                     | 4                                      | 0                                      | 3                                      |
| 252-3A | +                     | 4                                      | 0                                      | 4                                      |
| 252-3B | +                     | 7                                      | 0                                      | 1                                      |
| 255-1B | +                     | 0                                      | 8                                      | 0                                      |
| 230-1  | +                     | 5                                      | 1                                      | 2                                      |
| Z149   | +                     | 3                                      | 0                                      | 5                                      |

Event column refers to the identification number of the soybean transformants. T<sub>0</sub> GUS column indicates the primary transformants as being positive for GUS expression. T<sub>1</sub> GUS<sup>+</sup>/HT refers to the number of progeny for which the phenotype was GUS positive and herbicide tolerant. T<sub>1</sub> GUS<sup>-</sup>/HT column is the number of progeny for which the phenotype was GUS negative and herbicide tolerant. T<sub>1</sub> GUS<sup>-</sup>/HS indicates the number of progeny for which the phenotype was GUS negative and herbicide susceptible.

Southern positive plants, 3 transmitted the transgenes on to the progeny, while the others were non-germ-line events. Segregation analysis for two of these events, E230 and Z149, is shown in Table 4. Molecular analysis of the progeny from these events is shown in Figure 5. Herbicide tolerance and GUS expression correlated with the presence of the approximately 10-kb and 8-kb GUS-hybridizing fragments (Figure 5), in progeny derived from events E230 and Z149, respectively. An exception to the correlation between the molecular and phenotypic segregation analysis was observed in event E230. Individual number 4 (T<sub>1</sub>-4, Figure 5A) carried the 10-kb hybridizing fragment, yet the observed phenotype was GUS-negative and herbicide tolerant.

Ten plants, derived from the higher selection regime, (arising from independent explants) were also characterized by Southern blot analysis. Progeny ana-




**Figure 5.** (A) A Southern blot analysis on a primary transformant E230 (same event as in Figure 4 and Table 4) carrying pPTN125 and 8 T<sub>1</sub> progeny. The parent, lane 2, E230, had less DNA loaded (~5 µg) as compared to the progeny, designated T<sub>1</sub>, and control lanes, ck, (2.0 µg). The DNA samples were digested with *Eco* RI (cuts once within the T-DNA) and probed with the GUS ORF. The 14.9-kb linearized binary vector (18 pg) is shown with arrow on right side. The approximate 10-kb hybridizing fragment (arrow left side) correlated with GUS expression and herbicide tolerance, except for T<sub>1</sub> number 4, which we can not explain. (B) Southern blot analysis on a primary transformant Z149 carrying pPTN101 and 8 R<sub>1</sub> progeny. Lane 1: parent, Z149. Lanes 2–9: T<sub>1</sub> plants derived from Z149. Lane 10, control, A3237, designated ck. Last lane contains 18 pg of pPTN101. The DNA samples were digested with *Eco* RI and probed with the GUS ORF. The 16.7-kb linearized binary vector is shown on the right side. The approximate 8.0-kb hybridizing fragment (arrow left side) correlated with GUS expression and herbicide tolerance.

lysis on four of the primary transformants using the leaf painting technique was performed. The segregation data are given in Table 4 for events 246-2C, 252-3A, 252-3B, and 255-1B. The data demonstrate that these plants transmitted the transgenes to the next generation.

Generally a single plant is recovered from one explant. However, we have taken to soil multiple plants derived from the same explant. In Figure 3, events 207-5 and 207-5a were derived from the same explant and appear to carry the same integration event. Similarly, event 137 had two plants recovered from a single explant (137-1 and 137-2; Figure 4) and both of these appear to be derived from the same integration event. In one case, a GUS-negative plant, 211-2 (Figure 3) recovered under the lower stringency selection regime, appears to be an escape, while another plant derived from the same explant was positive for GUS expression, both histochemically and fluorometrically (211-2B, Table 2). For this reason our transformation frequencies do not include multiple plants derived

from the same explant. Hence, we assume that plants derived from the same explants are clonal events, though this may not necessarily be the case.

## Conclusion

*Agrobacterium*-mediated transformation of soybean using the cotyledonary node axillary meristem as the target for gene transfer was first achieved using kanamycin as a selective agent. Here we have demonstrated the utility of the *bar* gene, employing glufosinate as the selective agent, in the system. Germ-line transformation events were recovered at frequencies up to 3% using a selection regime of 5 mg l<sup>-1</sup> glufosinate during the shoot initiation stage and 2 mg l<sup>-1</sup> during shoot elongation. The time frame from explant in culture to collecting T<sub>1</sub> seed from a primary transformant ranges from 7 to 10 months under the conditions outlined in this study.

## Acknowledgements

All work was conducted at the University of Nebraska's Plant Transformation Core Research Facility. Funding was provided by the North Central Soybean Board, NFS-EPSCoR program, Center for Biotechnology at University of Nebraska-Lincoln and Nebraska's Research Initiative. Gratitude to extended to Ray Wu (Cornell University), Jim Carrington (Washington State University), Elke Kemper (Max-Planck-Institut für Züchtungsforschung), Pal Maliga (Waksman Institute, Rutgers) and Masashi Ugaki (National Institute of Agobiological Resources, Japan) for plasmids pDM302, pRTL2, pGPTV-bar, pPZP binary family and pE7131-GUS, respectively. We would also like to thank Dale Kinney and Mark Alexander (AgrEvo USA) for kindly providing Liberty® and glufosinate. A very special thanks to R.A. Zimmerman for keeping things in perspective. Note: The growth regulator regime outlined within this study was developed at Monsanto Company and is described in the manuscript, 'Progeny analysis of glyphosate selected transgenic soybeans derived from *Agrobacterium*-mediated transformation.' The manuscript has been submitted for publication.

## References

- Becker D, Kemper E, Schell J & Masterson R (1992) New plant binary vectors with selectable markers located proximal to the left T-DNA border. *Plant Mol. Biol.* 20: 1195–1197
- Cao J, Duan X, McElroy D & Wu R (1992) Regeneration of herbicide resistant rice plants following microprojectile-mediated transformation of suspension culture cells. *Plant Cell Rep.* 11: 586–591
- Carrington J & Freed DD (1990) Cap-independent enhancement of translation by a plant potyvirus 5' nontranslated region. *J. Virol.* 64: 1590–1597
- Cheng TY, Saka H & Voqui-Dinh TH (1980) Plant regeneration from soybean cotyledonary node segments in culture. *Plant Science Letters.* 19: 91–99
- Dellaporta SL, Wood J & Hicks JB (1983) A plant DNA miniprep: Version II. *Plant Mol. Biol. Repr.* 1: 19–21
- D'Halluin K, DeBlock M, Denecke J, Janssens J, Leemans J & Reynaerts A (1992) The *bar* gene as a selectable and screenable marker in plant engineering. *Methods Enzymol.* 216: 415–426
- Di R, Purcell V, Collins GB & Ghabrial SA (1996) Production of transgenic soybean lines expressing the bean pod mottle virus coat protein precursor gene. *Plant Cell Rep.* 15: 746–750
- Ditta G, Stanfield S, Corbin D & Helinski D (1980) Broad host range DNA cloning system for gram-negative bacteria: Construction of a gene bank of *Rhizobium meliloti*. *Proc. Natl. Acad. Sci. USA.* 77: 7347–7351
- Falco SC, Guida T, Locke M, Mauvais J, Sanders C, Ward RT & Webber P (1995) Transgenic canola and soybean seeds with increased lysine. *Bio/Technology.* 13: 577–582
- Gamborg OL, Miller RA & Ojima K (1968) Nutrient requirements of suspension cultures and soybean root cells. *Exp. Cell Res.* 50: 151–158
- Hajdukiewicz P, Svab Z & Maliga P (1994) The small versatile pZP family of *Agrobacterium* binary vectors for plant transformation. *Plant Mol. Biol.* 25: 989–994
- Hinchey MAW, Ward DVC, Newell CA, McDonnell RE, Sato SJ, Gasser CS, Fischhoff DA, Re DB, Fraley RT & Horsch RB (1988) Production of transgenic soybean plants using *Agrobacterium*-mediated DNA transfer. *Bio/Technol.* 6: 915–922
- Hood EE, Helmer GL, Fraley RT & Chilton MD (1986) The hypervirulence of *Agrobacterium tumefaciens* A281 is encoded in a region of pTiBo542 outside of T-DNA. *J. Bacteriol.* 168: 1291–1301
- Hood EE, Gelvin SB, Melchers LS & Hoekema A (1993) New *Agrobacterium* helper plasmids for gene transfer to plants. *Transgenic Research.* 2: 208–218
- Janssen BJ & Gardner RC (1993) The use of transient GUS expression to develop an *Agrobacterium*-mediated gene transfer system for kiwifruit. *Plant Cell Rep.* 13: 28–31
- Jefferson RA, Kavanagh TA & Bevan MW (1987) GUS fusions:  $\beta$ -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO Journal* 6: 3901–3907
- Mitsuhara I, Ugaki M, Hirochika H, Ohshima M, Murakami T, Gotoh Y, Katayose Y, Nakamura S, Honkura R, Nishimiya S, Ueno K, Mochizuki A, Tanimoto H, Tsugawa H, Otsuki Y & Ohashi Y (1996) Efficient promoter cassettes for enhanced expression of foreign genes in dicotyledonous, and monocotyledonous plants. *Plant Cell Physiol.* 37: 49–59
- Mullins MG, Archie-Tang FC & Facciotti D (1990) *Agrobacterium*-mediated genetic transformation of grapevines: transgenic plants of *Vitis rupestris* scheele and buds of *Vitis vinifera* L. *Bio/Technol.* 8: 1041–1045
- Murashige T & Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15: 473–479
- Padgett SR, Kolacz KH, Delannay X, Re DB, LaVallee BJ, Tinius CN, Rhodes WK, Otero YI, Barry GF, Eicholtz DA, Peschke VM, Nida DL, Taylor NB & Kishore GM (1995) Development, identification, and characterization of a glyphosate-tolerant soybean line. *Crop Science* 35: 1451–1461
- Rhee Y & Staswick PE (1992) Nucleotide sequence of a soybean vegetative storage protein *vspB* gene. *Plant Physiol.* 98: 795–796
- Shelp BJ, Swanton CJ & Hall JC (1992) Glufosinate (Phosphinothricin) mobility in young soybean shoots. *J. Plant Physiol.* 139: 626–628
- Southern EM (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98: 503–517
- Stewart CN, Adang MJ, All JN, Boerma HR, Cardineau G, Tucker D & Parrott WA (1996) Genetic transformation, recovery and characterization of fertile soybean transgenic for synthetic *Bacillus thuringiensis* cryIIAc gene. *Plant Physiol.* 112: 121–129
- Thompson CJ, Movva NR, Tichard R, Cramer R, Davies JE & Lauwereys M (1987) Characterization of the herbicide-resistance gene *bar* from *Streptomyces hygroscopicus*. *EMBO J.* 6: 2519–2523
- Trick, HN, Dinkins ER, Santarem ER, Di R, Samoylov C, Meurer D, Walker D, Parrott WA, Finer JJ & Collins GB 1997 Recent advances in soybean transformation. *Plant Tiss. Cult. Biotechnol.* 3: 9–26
- Zhang Z, Coyne DP & Mita A (1997) Factors affecting *Agrobacterium*-mediated transformation of common bean. *J. Amer. Soc. Hort. Sci.* 122: 300–305