Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2008/058183

International filing date: 26 March 2008 (26.03.2008)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/909,315
Filing date: 30 March 2007 (30.03.2007)

Date of receipt at the International Bureau: 13 April 2008 (13.04.2008)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)

World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

63351

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

April 13, 2008

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: *60/909,315*
FILING DATE: *March 30, 2007*
RELATED PCT APPLICATION NUMBER: *PCT/US08/58183*

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS *US60/909,315*

Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

PTO/SB/16 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET – Page 1 of 2

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. _____

INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
MICHAEL	SOFIA, J.	DOYLESTOWN, PA
JINFA	DU	NEW HOPE, PA
PEIYUAN	WANG	GLEN ROCK, NJ

Additional inventors are being named on the _____ *separately numbered sheets attached hereto*

TITLE OF THE INVENTION (500 characters max):

NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS

Direct all correspondence to: **CORRESPONDENCE ADDRESS**

☑ The address corresponding to Customer Number: 23552

OR

☐ Firm or Individual Name

Address

City	State	Zip
Country	Telephone	Email

ENCLOSED APPLICATION PARTS *(check all that apply)*

☑ Application Data Sheet. See 37 CFR 1.76

☐ CD(s), Number of CDs _____

☐ Drawing(s) *Number of Sheets* _____

☐ Other (specify) _____

☑ Specification (e.g. description of the invention) *Number of Pages* 645

Fees Due: Filing Fee of $200 ($100 for small entity). If the specification and drawings exceed 100 sheets of paper, an application size fee is also due, which is $250 ($125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).

METHOD OF PAYMENT OF THE FILING FEE AND APPLICATION SIZE FEE FOR THIS PROVISIONAL APPLICATION FOR PATENT

☑ Applicant claims small entity status. See 37 CFR 1.27.

☐ A check or money order is enclosed to cover the filing fee and application size fee (if applicable).

☐ Payment by credit card. Form PTO-2038 is attached

☑ The Director is hereby authorized to charge the filing fee and application size fee (if applicable) or credit any overpayment to Deposit Account Number: 13-2725. A duplicative copy of this form is enclosed for fee processing.

1,462.50

TOTAL FEE AMOUNT ($)

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PTO/SB/16 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☑ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: _____

_____

WARNING:

Petitioner/applicant is cautioned to avoid submitting personal information in documents filed in a patent application that may contribute to identity theft. Personal information such as social security numbers, bank account numbers, or credit card numbers (other than a check or credit card authorization form PTO-2038 submitted for payment purposes) is never required by the USPTO to support a petition or an application. If this type of personal information is included in documents submitted to the USPTO, petitioners/applicants should consider redacting such personal information from the documents before submitting them to the USPTO. Petitioner/applicant is advised that the record of a patent application is available to the public after publication of the application (unless a non-publication request in compliance with 37 CFR 1.213(a) is made in the application) or issuance of a patent. Furthermore, the record from an abandoned application may also be available to the public if the application is referenced in a published application or an issued patent (see 37 CFR 1.14). Checks and credit card authorization forms PTO-2038 submitted for payment purposes are not retained in the application file and therefore are not publicly available.

SIGNATURE _____ Date 2007/03/30

TYPED or PRINTED NAME DANIEL R. EVANS REGISTRATION NO. 55,868 *(if appropriate)*

TELEPHONE (404) 954-5061 Docket Number: 60137.0034USP1

PTO/SB/14 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76	Attorney Docket Number	60137.0034USP1
	Application Number	
Title of Invention	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS	

The application data sheet is part of the provisional or nonprovisional application for which it is being submitted. The following form contains the bibliographic data arranged in a format specified by the United States Patent and Trademark Office as outlined in 37 CFR 1.76.
This document may be completed electronically and submitted to the Office in electronic format using the Electronic Filing System (EFS) or the document may be printed and included in a paper filed application.

Secrecy Order 37 CFR 5.2

☐ Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2 (Paper filers only. Applications that fall under Secrecy Order may not be filed electronically.)

Applicant Information:

Applicant 1 Remove

Applicant Authority ◉ Inventor ○ Legal Representative under 35 U.S.C. 117 ○ Party of Interest under 35 U.S.C. 118

Prefix	Given Name	Middle Name	Family Name	Suffix
	MICHAEL	J.	SOFIA	

Residence Information (Select One) ◉ US Residency ○ Non US Residency ○ Active US Military Service

City	DOYLESTOWN	State/Province	PA	Country of Residence[i]	US

Citizenship under 37 CFR 1.41(b)[i] US

Mailing Address of Applicant:

Address 1	3066 ANTLER DRIVE		
Address 2			
City	DOYLESTOWN	State/Province	PA
Postal Code	18902	Country[i]	US

Applicant 2 Remove

Applicant Authority ◉ Inventor ○ Legal Representative under 35 U.S.C. 117 ○ Party of Interest under 35 U.S.C. 118

Prefix	Given Name	Middle Name	Family Name	Suffix
	JINFA		DU	

Residence Information (Select One) ◉ US Residency ○ Non US Residency ○ Active US Military Service

City	NEW HOPE	State/Province	PA	Country of Residence[i]	US

Citizenship under 37 CFR 1.41(b)[i] US

Mailing Address of Applicant:

Address 1	1206 REINS CIRCLE		
Address 2			
City	NEW HOPE	State/Province	PA
Postal Code	18938	Country[i]	US

Applicant 3 Remove

Applicant Authority ◉ Inventor ○ Legal Representative under 35 U.S.C. 117 ○ Party of Interest under 35 U.S.C. 118

Prefix	Given Name	Middle Name	Family Name	Suffix
	PEIYUAN		WANG	

Residence Information (Select One) ◉ US Residency ○ Non US Residency ○ Active US Military Service

City	GLEN ROCK	State/Province	NJ	Country of Residence[i]	US

PTO/SB/14 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76	Attorney Docket Number	60137.0034USP1
	Application Number	
Title of Invention	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS	

Citizenship under 37 CFR 1.41(b) [i]	CN		
Mailing Address of Applicant:			
Address 1	20 RADBURN ROAD		
Address 2			
City	GLEN ROCK	**State/Province**	NJ
Postal Code	07452	**Country**[i]	US

All Inventors Must Be Listed - Additional Inventor Information blocks may be generated within this form by selecting the **Add** button. Add

Correspondence Information:

Enter either Customer Number or complete the Correspondence Information section below.
For further information see 37 CFR 1.33(a).

☐ **An Address is being provided for the correspondence Information of this application.**

Customer Number	23552		
Email Address	devans@merchantgould.com	Add Email	Remove Email

Application Information:

Title of the Invention	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS		
Attorney Docket Number	60137.0034USP1	**Small Entity Status Claimed**	☒
Application Type	Provisional		
Subject Matter	Utility		
Suggested Class (if any)		**Sub Class (if any)**	
Suggested Technology Center (if any)			
Total Number of Drawing Sheets (if any)		**Suggested Figure for Publication (if any)**	

Publication Information:

☐ Request Early Publication (Fee required at time of Request 37 CFR 1.219)

☐ Request Not to Publish. I hereby request that the attached application not be published under 35 U.S.C. 122(b) and certify that the invention disclosed in the attached application has not been and will not be the subject of an application filed in another country, or under a multilateral agreement, that requires publication at eighteen months after filing.

Representative Information:

Representative information should be provided for all practitioners having a power of attorney in the application. Providing this information in the Application Data Sheet does not constitute a power of attorney in the application (see 37 CFR 1.32).
Enter either Customer Number or complete the Representative Name section below. If both sections are completed the Customer Number will be used for the Representative Information during processing.

Please Select One:	◉ Customer Number	○ US Patent Practitioner	○ US Representative (37 CFR 11.9)

PTO/SB/14 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76	Attorney Docket Number	60137.0034USP1
	Application Number	
Title of Invention	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS	

Customer Number	23552

Domestic Priority Information:

This section allows for the applicant to claim benefit under 35 U.S.C. 119(e), 120, 121, or 365(c). Providing this information in the application data sheet constitutes the specific reference required by 35 U.S.C. 119(e) or 120, and 37 CFR 1.78(a)(2) or CFR 1.78(a)(4), and need not otherwise be made part of the specification.

Prior Application Status			Remove
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)

Additional Domestic Priority Data may be generated within this form by selecting the **Add** button. Add

Foreign Priority Information:

This section allows for the applicant to claim benefit of foreign priority and to identify any prior foreign application for which priority is not claimed. Providing this information in the application data sheet constitutes the claim for priority as required by 35 U.S.C. 119(b) and 37 CFR 1.55(a).

Remove

Application Number	Country [i]	Parent Filing Date (YYYY-MM-DD)	Priority Claimed
			◉ Yes ○ No

Additional Foreign Priority Data may be generated within this form by selecting the **Add** button. Add

Assignee Information:

Providing this information in the application data sheet does not substitute for compliance with any requirement of part 3 of Title 37 of the CFR to have an assignment recorded in the Office.

Assignee 1 Remove

If the Assignee is an Organization check here. ☒

Organization Name	PHARMASSET, INC.		
Mailing Address Information:			
Address 1	303-A COLLEGE ROAD EAST		
Address 2			
City	PRINCETON	**State/Province**	NJ
Country [i] US		Postal Code	08540
Phone Number	(609) 4134100	Fax Number	(609) 6134150
Email Address			

Additional Assignee Data may be generated within this form by selecting the **Add** button. Add

Signature:

A signature of the applicant or representative is required in accordance with 37 CFR 1.33 and 10.18. Please see 37 CFR 1.4(d) for the form of the signature.

PTO/SB/14 (02-07)
Approved for use through 02/28/2007. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76	Attorney Docket Number	60137.0034USP1
	Application Number	
Title of Invention	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS	

Signature	/Daniel R. Evans/			Date (YYYY-MM-DD)	2007-03-30
First Name	Daniel	Last Name	Evans	Registration Number	55868

This collection of information is required by 37 CFR 1.76. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 23 minutes to complete, including gathering, preparing, and submitting the completed application data sheet form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. Accordingly, pursuant to the requirements of the Act, please be advised that: (1) the general authority for the collection of this information is 35 U.S.C. 2(b)(2); (2) furnishing of the information solicited is voluntary; and (3) the principal purpose for which the information is used by the U.S. Patent and Trademark Office is to process and/or examine your submission related to a patent application or patent. If you do not furnish the requested information, the U.S. Patent and Trademark Office may not be able to process and/or examine your submission, which may result in termination of proceedings or abandonment of the application or expiration of the patent.

The information provided by you in this form will be subject to the following routine uses:

1. The information on this form will be treated confidentially to the extent allowed under the Freedom of Information Act (5 U.S.C. 552) and the Privacy Act (5 U.S.C. 552a). Records from this system of records may be disclosed to the Department of Justice to determine whether the Freedom of Information Act requires disclosure of these records.

2. A record from this system of records may be disclosed, as a routine use, in the course of presenting evidence to a court, magistrate, or administrative tribunal, including disclosures to opposing counsel in the course of settlement negotiations.

3. A record in this system of records may be disclosed, as a routine use, to a Member of Congress submitting a request involving an individual, to whom the record pertains, when the individual has requested assistance from the Member with respect to the subject matter of the record.

4. A record in this system of records may be disclosed, as a routine use, to a contractor of the Agency having need for the information in order to perform a contract. Recipients of information shall be required to comply with the requirements of the Privacy Act of 1974, as amended, pursuant to 5 U.S.C. 552a(m).

5. A record related to an International Application filed under the Patent Cooperation Treaty in this system of records may be disclosed, as a routine use, to the International Bureau of the World Intellectual Property Organization, pursuant to the Patent Cooperation Treaty.

6. A record in this system of records may be disclosed, as a routine use, to another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)).

7. A record from this system of records may be disclosed, as a routine use, to the Administrator, General Services, or his/her designee, during an inspection of records conducted by GSA as part of that agency's responsibility to recommend improvements in records management practices and programs, under authority of 44 U.S.C. 2904 and 2906. Such disclosure shall be made in accordance with the GSA regulations governing inspection of records for this purpose, and any other relevant (i.e., GSA or Commerce) directive. Such disclosure shall not be used to make determinations about individuals.

8. A record from this system of records may be disclosed, as a routine use, to the public after either publication of the application pursuant to 35 U.S.C. 122(b) or issuance of a patent pursuant to 35 U.S.C. 151. Further, a record may be disclosed, subject to the limitations of 37 CFR 1.14, as a routine use, to the public if the record was filed in an application which became abandoned or in which the proceedings were terminated and which application is referenced by either a published application, an application open to public inspections or an issued patent.

9. A record from this system of records may be disclosed, as a routine use, to a Federal, State, or local law enforcement agency, if the USPTO becomes aware of a violation or potential violation of law or regulation.

Docket No. 60137.0034USP1

NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS

Field of Invention

The present invention pertains to nucleoside phosphoramidates and their use as agents for treating viral diseases. These compounds are inhibitors of RNA-dependent RNA viral replication and are useful as inhibitors of HCV NS5B polymerase, as inhibitors of HCV replication and for treatment of hepatitis C infection in mammals. The invention provides novel chemical compounds, and the use of these compounds alone or in combination with other antiviral agents for treating HCV infection.

Background

Hepatitis C virus (HCV) infection is a major health problem that leads to chronic liver disease, such as cirrhosis and hepatocellular carcinoma, in a substantial number of infected individuals, estimated to be 2-15% of the world's population. There are an estimated 4.5 million infected people in the United States alone, according to the U.S. Center for Disease Control. According to the World Health Organization, there are more than 200 million infected individuals worldwide, with at least 3 to 4 million people being infected each year. Once infected, about 20% of people clear the virus, but the rest can harbor HCV the rest of their lives. Ten to twenty percent of chronically infected individuals eventually develop liver-destroying cirrhosis or cancer. The viral disease is transmitted parenterally by contaminated blood and blood products, contaminated needles, or sexually and vertically from infected mothers or carrier mothers to their offspring. Current treatments for HCV infection, which are restricted to immunotherapy

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 10 of 657

with recombinant interferon-α alone or in combination with the nucleoside analog ribavirin, are of limited clinical benefit as resistance develops rapidly. Moreover, there is no established vaccine for HCV. Consequently, there is an urgent need for improved therapeutic agents that effectively combat chronic HCV infection.

The HCV virion is an enveloped positive-strand RNA virus with a single oligoribonucleotide genomic sequence of about 9600 bases which encodes a polyprotein of about 3,010 amino acids. The protein products of the HCV gene consist of the structural proteins C, E1, and E2, and the non-structural proteins NS2, NS3, NS4A and NS4B, and NS5A and NS5B. The nonstructural (NS) proteins are believed to provide the catalytic machinery for viral replication. The NS3 protease releases NS5B, the RNA-dependent RNA polymerase from the polyprotein chain. HCV NS5B polymerase is required for the synthesis of a double-stranded RNA from a single-stranded viral RNA that serves as a template in the replication cycle of HCV. Therefore, NS5B polymerase is considered to be an essential component in the HCV replication complex (K. Ishi, et al, *Heptology*, 1999, 29: 1227-1235; V. Lohmann, et al., *Virology*, 1998, 249: 108-118). Inhibition of HCV NS5B polymerase prevents formation of the double-stranded HCV RNA and therefore constitutes an attractive approach to the development of HCV-specific antiviral therapies.

HCV belongs to a much larger family of viruses that share many common features.

Flaviviridae Viruses

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 11 of 657

The Flaviviridae family of viruses comprises at least three distinct genera: *pestiviruses*, which cause disease in cattle and pigs; *flavivruses*, which are the primary cause of diseases such as dengue fever and yellow fever; and *hepaciviruses*, whose sole member is HCV. The flavivirus genus includes more than 68 members separated into groups on the basis of serological relatedness (Calisher et al., *J. Gen. Virol*, 1993,70,37-43). Clinical symptoms vary and include fever, encephalitis and hemorrhagic fever (*Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 1996, Chapter 31, 931-959). Flaviviruses of global concern that are associated with human disease include the Dengue Hemorrhagic Fever viruses (DHF), yellow fever virus, shock syndrome and Japanese encephalitis virus (Halstead, S. B., *Rev. Infect. Dis.*, 1984, 6, 251-264; Halstead, S. B., *Science*, 239:476-481, 1988; Monath, T. P., *New Eng. J. Med*, 1988, 319, 64 1-643).

The pestivirus genus includes bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, also called hog cholera virus) and border disease virus (BDV) of sheep (Moennig, V. et al. *Adv. Vir. Res.* 1992, 41, 53-98). Pestivirus infections of domesticated livestock (cattle, pigs and sheep) cause significant economic losses worldwide. BVDV causes mucosal disease in cattle and is of significant economic importance to the livestock industry (Meyers, G. and Thiel, H.J., Advances in Virus Research, 1996, 47, 53-118; Moennig V., et al, Adv. Vir. Res. 1992, 41, 53-98). Human pestiviruses have not been as extensively characterized as the animal pestiviruses. However, serological surveys indicate considerable pestivirus exposure in humans.

Pestiviruses and hepaciviruses are closely related virus groups within the Flaviviridae family. Other closely related viruses in this family include the GB virus A,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 12 of 657

GB virus A-like agents, GB virus-B and GB virus-C (also called hepatitis G virus, HGV). The hepacivirus group (hepatitis C virus; HCV) consists of a number of closely related but genotypically distinguishable viruses that infect humans. There are at least 6 HCV genotypes and more than 50 subtypes. Due to the similarities between pestiviruses and hepaciviruses, combined with the poor ability of hepaciviruses to grow efficiently in cell culture, bovine viral diarrhea virus (BVDV) is often used as a surrogate to study the HCV virus.

The genetic organization of pestiviruses and hepaciviruses is very similar. These positive stranded RNA viruses possess a single large open reading frame (ORF) encoding all the viral proteins necessary for virus replication. These proteins are expressed as a polyprotein that is co- and post-translationally processed by both cellular and virus-encoded proteinases to yield the mature viral proteins. The viral proteins responsible for the replication of the viral genome RNA are located within approximately the carboxy-terminal. Two-thirds of the ORF are termed nonstructural (NS) proteins. The genetic organization and polyprotein processing of the nonstructural protein portion of the ORF for pestiviruses and hepaciviruses is very similar. For both the pestiviruses and hepaciviruses, the mature nonstructural (NS) proteins, in sequential order from the amino-terminus of the nonstructural protein coding region to the carboxy-terminus of the ORF, consist of p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

The NS proteins of pestiviruses and hepaciviruses share sequence domains that are characteristic of specific protein functions. For example, the NS3 proteins of viruses in both groups possess amino acid sequence motifs characteristic of serine proteinases and of helicases (Gorbalenya et al., *Nature,* 1988, 333, 22; Bazan and Fletterick *Virology*

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 13 of 657

, 1989,171,637-639; Gorbalenya et al., *Nucleic Acid Res.*, 1989, 17, 3889-3897). Similarly, the NS5B proteins of pestiviruses and hepaciviruses have the motifs characteristic of RNA-directed RNA polymerases (Koonin, E.V. and Dolja, V.V., *Crir. Rev. Biochem. Molec. Biol.* 1993, 28, 375-430).

The actual roles and functions of the NS proteins of pestiviruses and hepaciviruses in the lifecycle of the viruses are directly analogous. In both cases, the NS3 serine proteinase is responsible for all proteolytic processing of polyprotein precursors downstream of its position in the ORF (Wiskerchen and Collett, *Virology*, 1991, 184, 341-350; Bartenschlager et al., *J. Virol.* 1993, 67, 3835-3844; Eckart et al. *Biochem. Biophys. Res. Comm.* 1993,192, 399-406; Grakoui et al., *J. Virol.* 1993, 67, 2832-2843; Grakoui et al., *Proc. Natl. Acad Sci. USA* 1993, 90, 10583-10587; Hijikata et al., *J. Virol.* 1993, 67, 4665-4675; Tome et al., *J. Virol.*, 1993, 67, 4017-4026). The NS4A protein, in both cases, acts as a cofactor with the NS3 serine protease (Bartenschlager et al., *J. Virol.* 1994, 68, 5045-5055; Failla et al., *J. Virol.* 1994, 68, 3753-3760; Xu et al., *J. Virol.*, 1997, 71:53 12-5322). The NS3 protein of both viruses also functions as a helicase (Kim et al., *Biochem. Biophys. Res. Comm.*, 1995, 215, 160-166; Jin and Peterson, *Arch. Biochem. Biophys.*, 1995, 323, 47-53; Warrener and Collett, *J. Virol.* 1995, 69,1720-1726). Finally, the NS5B proteins of pestiviruses and hepaciviruses have the predicted RNA-directed RNA polymerases activity (Behrens et al., EMBO, 1996, 15, 12-22; Lechmann et al., *J. Virol.*, 1997, 71, 8416-8428; Yuan et al., *Biochem. Biophys. Res. Comm.* 1997, 232, 231-235; Hagedorn, PCT WO 97/12033; Zhong et al, *J. Virol.*, 1998, 72, 9365-9369).

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 14 of 657

Currently, there are limited treatment options for individuals infected with hepatitis C virus. The current approved therapeutic option is the use of immunotherapy with recombinant interferon-α alone or in combination with the nucleoside analog ribavirin. This therapy is limited in its clinical effectiveness and only 50% of treated patients respond to therapy. Therefore, there is significant need for more effective and novel therapies to address the unmet medical need posed by HCV infection.

A number of potential molecular targets for drug development of direct acting antivirals as anti -HCV therapeutics have now been identified including, but not limited to, the NS2-NS3 autoprotease, the N3 protease, the N3 helicase and the NS5B polymerase. The RNA-dependent RNA polymerase is absolutely essential for replication of the single-stranded, positive sense, RNA genome and this enzyme has elicited significant interest among medicinal chemists.

Inhibitors of HCV NS5B as potential therapies for HCV infection have been reviewed: Tan, S.-L., et al., *Nature Rev. Drug Discov.*, 2002, 1, 867-881; Walker, M.P. et al., *Exp. Opin. Investigational Drugs*, 2003, 12, 1269-1280; Ni, Z-J., et al., *Current Opinion in Drug Discovery and Development*, 2004, 7, 446-459; Beaulieu, P. L., et al., *Current Opinion in Investigational Drugs*, 2004, 5, 838-850; Wu, J., et al., *Current Drug Targets-Infectious Disorders*, 2003, 3, 207-219; Griffith, R.C., et al, *Annual Reports in Medicinal Chemistry*, 2004, 39, 223-237; Carrol, S., et al., *Infectious Disorders-Drug Targets*, 2006, 6, 17-29. The potential for the emergence of resistant HCV strains and the need to identify agents with broad genotype coverage supports the need for continuing efforts to identify novel and more effective nucleosides as HCV NS5B inhibitors.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 15 of 657

Nucleoside inhibitors of NS5B polymerase can act either as a non-natural substrate that results in chain termination or as a competitive inhibitor which competes with nucleotide binding to the polymerase. To function as a chain terminator the nucleoside analog must be taken up by the cell and converted *in vivo* to a triphosphate to compete for the polymerase nucleotide binding site. This conversion to the triphosphate is commonly mediated by cellular kinases which imparts additional structural requirements on a potential nucleoside polymerase inhibitor. Unfortunately, this limits the direct evaluation of nucleosides as inhibitors of HCV replication to cell-based assays capable of *in situ* phosphorylation.


In some cases, the biological activity of a nucleoside is hampered by its poor substrate characteristics for one or more of the kinases needed to convert it to the active triphosphate form. Formation of the monophosphate by a nucleoside kinase is generally viewed as the rate limiting step of the three phosphorylation events. To circumvent the need for the initial phosphorylation step in the metabolism of a nucleoside to the active triphosphate analog, the preparation of stable phosphate prodrugs has been reported. Nucleoside phosphoramidate prodrugs have been shown to be precursors of the active nucleoside triphosphate and to inhibit viral replication when administered to viral infected whole cells (McGuigan, C., et al., *J. Med. Chem.*, 1996, 39, 1748-1753; Valette, G., et al., *J. Med. Chem.*, 1996, 39, 1981-1990; Balzarini, J., et al., *Proc. National Acad Sci USA*, 1996, 93, 7295-7299; Siddiqui, A. Q., et al., *J. Med. Chem.*, 1999, 42, 4122-4128; Eisenberg, E. J., et al., *Nucleosides, Nucleotides and Nucleic Acids*, 2001, 20, 1091-1098; Lee, W.A., et al., *Antimicrobial Agents and Chemotheryapy*, 2005, 49, 1898)

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 16 of 657

Also limiting the utility of nucleosides as viable therapeutic agents is their sometimes poor physicochemical and pharmacokinetic properties. These poor properties can limit the intestinal absorption of an agent and limit uptake into the target tissue or cell. To improve on their properties prodrugs of nucleosides have been employed. It has been demonstrated that preparation of nucleoside phosphoramidates improves the systemic absorption of a nucleoside and furthermore, the phosphoramidate moiety of these "pronucleotides" is masked with neutral lipophilic groups to obtain a suitable partition coefficient to optimize uptake and transport into the cell dramatically enhancing the intracellular concentration of the nucleoside monophosphate analog relative to administering the parent nucleoside alone. Enzyme-mediated hydrolysis of the phosphate ester moiety produces a nucleoside monophosphate wherein the rate limiting initial phosphorylation is unnecessary.

SUMMARY OF THE INVENTION

The present invention is directed toward novel phosphoramidate prodrugs of nucleoside derivatives for the treatment of viral infections in mammals, which is a compound, its stereoisomers, salts (acid or basic addition salts), hydrates, solvates, or crystalline forms thereof, represented by the following structure:



I

wherein

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 17 of 657

R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $–N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazolimidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazolimidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 18 of 657

R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

R^5 is H, an optionally substituted alkyl (including lower alkyl), cyano (CN), CH_3, vinyl, O-alkyl, O-(lower alkyl), including OCH_3, OCH_2CH_3, hydroxyl alkyl, i.e., -$(CH_2)_oOH$, wherein o is 1 – 10, hydroxyl lower alkyl, i.e., -$(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), fluoromethyl (CH_2F), azido (N_3), CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H.

R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


X is H, OH, F, OMe, halogen, NH_2, or N_3

Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, or N(C_{1-18} acyl)$_2$,

wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$.

Base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 19 of 657



a **b** **c** **d**

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl, which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl);

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, wherein R' is defined above.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 20 of 657

DEFINITIONS

The phrase "a" or "an" entity as used herein refers to one or more of that entity; for example, a compound refers to one or more compounds or at least one compound. As such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.

The phrase "as defined herein above" refers to the first definition provided in the Summary of the Invention.

The terms "optional" or "optionally" as used herein means that a subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional bond" means that the bond may or may not be present, and that the description includes single, double, or triple bonds.

The term "independently" is used herein to indicate that a variable is applied in any one instance without regard to the presence or absence of a variable having that same or a different definition within the same compound. Thus, in a compound in which R appears twice and is defined as "independently carbon or nitrogen", both R's can be carbon, both R's can be nitrogen, or one R' can be carbon and the other nitrogen.

The term "alkenyl" refers to an unsubstituted hydrocarbon chain radical having from 2 to 10 carbon atoms having one or two olefinic double bonds, preferably one olefinic double bond. The term "$C_{2\text{-}N}$ alkenyl" refers to an alkenyl comprising 2 to N carbon atoms, where N is an integer having the following values: 3, 4, 5, 6, 7, 8, 9, or 10. The term "$C_{2\text{-}10}$ alkenyl" refers to an alkenyl comprising 2 to 10 carbon atoms. The term "$C_{2\text{-}4}$ alkenyl" refers to an alkenyl comprising 2 to 4 carbon atoms. Examples include, but are not limited to, vinyl, 1-propenyl, 2-propenyl (allyl) or 2-butenyl (crotyl).

The term "halogenated alkenyl" refers to an alkenyl comprising at least one of F, Cl, Br, and I.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 21 of 657

The term "alkyl" refers to an unbranched or branched chain, saturated, monovalent hydrocarbon residue containing 1 to 30 carbon atoms. The term "C_{1-M} alkyl" refers to an alkyl comprising 2 to M carbon atoms, where M is an integer having the following values: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30. The term "C_{1-4} alkyl" refers to an alkyl containing 1 to 4 carbon atoms. The term "lower alkyl" denotes a straight or branched chain hydrocarbon residue comprising 1 to 6 carbon atoms. "C_{1-20} alkyl" as used herein refers to an alkyl comprising 1 to 20 carbon atoms. "C_{1-10} alkyl" as used herein refers to an alkyl comprising 1 to 10 carbons. Examples of alkyl groups include, but are not limited to, lower alkyl groups include methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl. The term (ar)alkyl or (heteroaryl)alkyl indicate the alkyl group is optionally substituted by an aryl or a heteroaryl group respectively.

The term "halogenated alkyl" (or "haloalkyl") refers to an unbranched or branched chain alkyl comprising at least one of F, Cl, Br, and I. The term "C_{1-3} haloalkyl" refers to a haloalkyl comprising 1 to 3 carbons and at least one of F, Cl, Br, and I. The term "halogenated lower alkyl" refers to a haloalkyl comprising 1 to 6 carbon atoms and at least one of F, Cl, Br, and I. Examples include, but are not limited to, fluoromethyl, chloromethyl, bromomethyl, iodomethyl, difluoromethyl, dichloromethyl, dibromomethyl, diiodomethyl, trifluoromethyl, trichloromethyl, tribromomethyl, triiodomethyl, 1-fluoroethyl, 1-chloroethyl, 1-bromoethyl, 1-iodoethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-difluoroethyl, 2,2-dichloroethyl, 2,2-dibromomethyl, 2-2-diiodomethyl, 3-fluoropropyl, 3-chloropropyl, 3-bromopropyl, 2,2,2-trifluoroethyl or 1,1,2,2,2-pentafluoroethyl.

The term "alkynyl" refers to an unbranched or branched hydrocarbon chain radical having from 2 to 10 carbon atoms, preferably 2 to 5 carbon atoms, and having one triple bond. The term "C_{2-N} alkynyl" refers to an alkynyl comprising 2 to N carbon atoms, where N is an integer having the following values: 3, 4, 5, 6, 7, 8, 9, or 10. The term "C C_{2-4} alkynyl" refers to an alkynyl comprising 2 to 4 carbon atoms. The term "C_{2-10}

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 22 of 657

alkynyl" refers to an alkynyl comprising 2 to 10 carbons. Examples include, but are limited to, ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl or 3-butynyl.

The term "halogenated alkynyl" refers to an unbranched or branched hydrocarbon chain radical having from 2 to 10 carbon atoms, preferably 2 to 5 carbon atoms, and having one triple bond and at least one of F, Cl, Br, and I.

The term "cycloalkyl" refers to a saturated carbocyclic ring comprising 3 to 8 carbon atoms, i.e. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl. The term "C_{3-7} cycloalkyl" as used herein refers to a cycloalkyl comprising 3 to 7 carbons in the carbocyclic ring.

The term "alkoxy" refers to an -O-alkyl group, wherein alkyl is as defined above. Examples include, but are not limited to, methoxy, ethoxy, *n*-propyloxy, *i*-propyloxy, *n*-butyloxy, *i*-butyloxy, *t*-butyloxy. "Lower alkoxy" as used herein denotes an alkoxy group with a "lower alkyl" group as previously defined. "C_{1-10} alkoxy" refers to an-O-alkyl wherein alkyl is C_{1-10}.

The term "halogenated alkoxy" refers to an –O-alkyl group in which the alkyl group comprises at least one of F, Cl, Br, and I.

The term "halogenated lower alkoxy" refers to an –O-(lower alkyl) group in which the lower alkyl group comprises at least one of F, Cl, Br, and I.

The term "amino acid" includes naturally occurring and synthetic α, β γ or δ amino acids, and includes but is not limited to, amino acids found in proteins, i.e. glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucinyl, isoleucinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glycinyl, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl, glutaroyl, lysinyl, argininyl, histidinyl, β-alanyl, β-valinyl, β-leucinyl, β-isoleucinyl, β-prolinyl, β-phenylalaninyl, β-tryptophanyl, β-methioninyl, β-glycinyl, β-serinyl, β-threoninyl, β-

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 23 of 657

cysteinyl, β-tyrosinyl, β-asparaginyl, β-glutaminyl, β-aspartoyl, β-glutaroyl, β-lysinyl, β-argininyl or β-histidinyl. When the term amino acid is used, it is considered to be a specific and independent disclosure of each of the esters of α, β γ or δ glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine in the D and L-configurations.

The terms "alkylamino" or "arylamino" refer to an amino group that has one or two alkyl or aryl substituents, respectively.

The term "protected," as used herein and unless otherwise defined, refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis. Non-limiting examples include: C(O)-alkyl, C(O)Ph, C(O)aryl, CH_3, CH_2-alkyl, CH_2-alkenyl, CH_2Ph, CH_2-aryl, CH_2O-alkyl, CH_2O-aryl, SO_2-alkyl, SO_2-aryl, *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl, and 1,3-(1,1,3,3-tetraisopropyldisiloxanylidene).

The term "aryl," as used herein, and unless otherwise specified, refers to substituted or unsubstituted phenyl (Ph), biphenyl, or naphthyl, preferably the term aryl refers to substituted or unsubstituted phenyl. The aryl group can be substituted with one or more moieties selected from among hydroxyl, F, Cl, Br, I, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in T.W. Greene and P.G. M. Wuts, "Protective Groups in Organic Synthesis," 3rd ed., John Wiley & Sons, 1999.

The terms "alkaryl" or "alkylaryl" refer to an alkyl group with an aryl substituent. The terms "aralkyl" or "arylalkyl" refer to an aryl group with an alkyl substituent.

The term "halo," as used herein, includes chloro, bromo, iodo and fluoro.

The term "acyl" refers to a substituent containing a carbonyl moiety and a non-carbonyl moiety. The carbonyl moiety contains a double-bond between the carbonyl

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 24 of 657

carbon and a heteroatom, where the heteroatom is selected from among O, N and S. When the heteroatom is N, the N is substituted by a lower alkyl. The non-carbonyl moiety is selected from straight, branched, or cyclic alkyl, which includes, but is not limited to, a straight, branched, or cyclic C_{1-20} alkyl, C_{1-10} alkyl, or lower alkyl; alkoxyalkyl, including methoxymethyl; aralkyl, including benzyl; aryloxyalkyl, such as phenoxymethyl; or aryl, including phenyl optionally substituted with halogen (F, Cl, Br, I), hydroxyl, C_1 to C_4 alkyl, or C_1 to C_4 alkoxy, sulfonate esters, such as alkyl or aralkyl sulphonyl, including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, trialkylsilyl (e.g. dimethyl-t-butylsilyl) or diphenylmethylsilyl. When at least one aryl group is present in the non-carbonyl moiety, it is preferred that the aryl group comprises a phenyl group.

The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

The term "purine" or "pyrimidine" base includes, but is not limited to, adenine, N^6-alkylpurines, N^6-acylpurines (wherein acyl is C(O)(alkyl, aryl, alkylaryl, or arylalkyl), N^6-benzylpurine, N^6-halopurine, N^6-vinylpurine, N^6-acetylenic purine, N^6-acyl purine, N^6-hydroxyalkyl purine, N^6-allcylaminopurine, N^6-thioallcyl purine, N^2-alkylpurines, N^2-alkyl-6-thiopurines, thymine, cytosine, 5-fluorocytosine, 5-methylcytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrmidine, uracil, 5-halouracil, including 5-fluorouracil, C^5-alkylpyrimidines, C^5-benzylpyrimidines, C^5-halopyrimidines, C^5-vinylpyrimidine, C^5-acetylenic pyrimidine, C^5-acyl pyrimidine, C^5-hydroxyalkyl purine, C^5-amidopyrimidine, C^5-cyanopyrimidine, ,C^5-iodopyrimidine, C^6-lodo-pyrimidine, C^5-Br-vinyl pyrimidine, C^6-Br-vinyl pyrimidine, C^5-nitropyrimidine, C^5-amino-pyrimidine, N^2-alkylpurines, N^2-alkyl-6-thiopurines, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and pyrazolopyrimidinyl. Purine bases include, but are not limited to, guanine, adenine, hypoxanthine, 2,6-diaminopurine, and 6-chloropurine. Functional oxygen and nitrogen groups on the base can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 25 of 657


t-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl, alkyl groups, and acyl groups such as acetyl and propionyl, methanesulfonyl, and p-toluenesulfonyl.

The term "tautomerism" and "tautomers" have their accepted plain meanings.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 26 of 657

DETAILED DESCRIPTION OF THE INVENTION

An aspect of the invention is directed to a compound, its salts, hydrates, solvates, crystalline forms, and the like represented by formula I:



wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_e(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 27 of 657

adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl),

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 28 of 657

$OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl$)_2$, $N(C_{1-18}$ acyl$)_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl$)_2$, $N(C_{1-4}$ acyl$)_2$.

The base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl ,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 29 of 657


which includes, but is not limited to, an optionally substituted $C_{1\text{-}20}$ alkyl, an optionally substituted $C_{1\text{-}10}$ alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)($C_{1\text{-}20}$ alkyl), C(O)($C_{1\text{-}10}$ alkyl), or C(O)(lower alkyl).

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

As can be appreciated from the structure represented by formula I above, there are myriad ways to express the several embodiments and aspects of each embodiment of the present invention. As seen below, the inventors have disclosed certain embodiments directed to the compound of formula **I**, each having several aspects, based on the identity of the modified purine or pyrimidine base. This is not intended to be an explicit or implicit admission that the three embodiments are independent or distinct nor should it be interpreted as such. Rather, it is intended to convey information so that the full breadth of the present invention can be understood. Furthermore, the following embodiments, and aspects thereof, are not meant to be limiting on the full breadth of the invention as recited by the structure of formula **I**.

A first embodiment of the invention is directed to a compound represented by formula **I-1**:

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 30 of 657



I-1

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b}

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 31 of 657

together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, an alkyl and an optionally substituted alkyl (including lower alkyl), cyano (CN), CH_3, vinyl, O-alkyl, O-(lower alkyl), including OCH_3, OCH_2CH_3, hydroxyl alkyl, i.e., $-(CH_2)_oOH$, wherein o is 1 – 10, hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), fluoromethyl (CH_2F), azido (N_3), CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 32 of 657

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, vinyl, N_3, CN, Cl, Br, F, I, $O(C_{1-6}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$.

(i) R^7, R^8,R^9 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

A first aspect of the first embodiment is directed to a compound represented by formula **I-1**

wherein

(a) R^1 is hydrogen, n-alkyl or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl;

(b) R^2 is hydrogen, C_{1-6} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi)

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 33 of 657

R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, lower alkyl, aryl, or substituted aryl by lower alkyl, C(O)(lower alkyl), wherein said aryl is phenyl;

(e) R^5 is H, CN, CH_3, vinyl, OCH_3, OCH_2CH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, vinyl, N_3, CN, Cl, Br, F, I, O(C_{1-6} acyl), O(C_{1-4} alkyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$;

(i) R^7, R^8,R^9 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 34 of 657

CO_2R', $CONH_2$, $CONHR'$, $CONR'_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

A second aspect of the first embodiment is directed to a compound represented by formula **I-1**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a C_{1-3} alkyl, a C_{1-3} alkoxy, F, Cl, Br, I, nitro, cyano, and a C_{1-3} haloalkyl;

(b) R^2 is hydrogen, CH_3, R^{3a} or R^{3b} and R^2 together are $(CH_2)_3$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 35 of 657

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A third aspect of the first embodiment is directed to a compound represented by formula **I-1**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 36 of 657

$CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, -$CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;


(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a $C_{1\text{-}20}$ alkyl; a $C_{1\text{-}20}$ cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl,

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A fourth aspect of the first embodiment is directed to a compound represented by formula **I-2**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 37 of 657



I-2

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 38 of 657

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl;

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A fifth aspect of the first embodiment is directed to a compound represented by formula **I-2**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 39 of 657

$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH;

(d) R^4 is hydrogen, CH_3, Et, ^iPr, ^nPr, ^nBu, 2-butyl, ^tBu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl,

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A sixth aspect of the first embodiment is directed to a compound represented by formula **I-2**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 40 of 657

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, OMe, CN, CH_2F, F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, or N_3;

(h) Y is an H, OH, CH_3, F, Cl, Br, I, or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A seventh aspect of the first embodiment is directed to a compound represented by formula **I-2**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 41 of 657

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, or N_3;

(h) Y is an OH, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) R^9 is selected from among OH, OCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from among 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

An eighth aspect of the first embodiment is directed to a compound represented by formula **I-2**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, or N_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 42 of 657


(h) Y is an OH, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) R^9 is selected from among OH, OCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from among 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A second embodiment of the invention is directed to a compound represented by formula **I** in which the base is a structure represented by formula **b** above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, Y, R^7, and $R^8$are defined in the Summary of the Invention section above.

A first aspect of the second embodiment is directed to a compound represented by formula **I-3**



I-3

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 43 of 657

optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $–N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 44 of 657

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, an alkyl and an optionally substituted alkyl (including lower alkyl), cyano (CN), CH_3, vinyl, O-alkyl, O-(lower alkyl), including OCH_3, OCH_2CH_3, hydroxyl alkyl, i.e., $-(CH_2)_oOH$, wherein o is 1 – 10, hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), fluoromethyl (CH_2F), azido (N_3), CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, vinyl, N_3, CN, Cl, Br, F, I, $O(C_{1-6}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The second aspect of the second embodiment is directed to a compound represented by formula **I-3**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 45 of 657

wherein

(a) R^1 is hydrogen, n-alkyl or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl;

(b) R^2 is hydrogen, C_{1-6} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, lower alkyl, aryl, or substituted aryl by lower alkyl, C(O)(lower alkyl), wherein said aryl is phenyl;

(e) R^5 is H, CN, CH_3, vinyl, OCH_3, OCH_2CH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 46 of 657

(h) Y is an OH, H, C_{1-4} alkyl, vinyl, N_3, CN, Cl, Br, F, I, $O(C_{1-6}$ acyl), $O(C_{1-4}$ alkyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The third aspect of the second embodiment is directed to a compound represented by formula **I-3**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a C_{1-3} alkyl, a C_{1-3} alkoxy, F, Cl, Br, I, nitro, cyano, and a C_{1-3} haloalkyl;

(b) R^2 is hydrogen, CH_3, R^{3a} or R^{3b} and R^2 together are $(CH_2)_3$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi)

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 47 of 657

R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

The fourth aspect of the second embodiment is directed to a compound represented by formula **I-3**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 48 of 657

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a $C_{1\text{-}20}$ alkyl; a $C_{1\text{-}20}$ cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The fifth aspect of the second embodiment is directed to a compound represented by formula **I-4**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 49 of 657



I-4

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 50 of 657

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl,

The sixth aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 51 of 657

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The seventh aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, OMe, CN, CH_2F, F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is an H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 52 of 657

The eighth aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

The ninth aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 53 of 657

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;


(g) X is H, OH, OCH_3, F, or N_3;

(h) Y is an OH, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

A third embodiment of the invention is directed to a compound represented by formula **I** in which the base is a structure represented by formula **c** above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, Y, Z, R^{10}, R^{11}, and R^{12} are defined in the Summary of the Invention section above.

The first aspect of the third embodiment is directed to a compound represented by formula **I-5**



I-5

wherein

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 54 of 657

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^9, R^{10} and R^{11} are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl,

(j) Z is N or CR^9.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 55 of 657

The second aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^9, R^{10}, and R^{11} are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl;

(j) Z is N or CR^9

The third aspect of the third embodiment is directed to a compound represented

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 56 of 657

by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is an H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^9, R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

The fourth aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 57 of 657

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

The fifth aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 58 of 657

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;


(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^9, R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

The sixth aspect of the third embodiment is directed to a compound represented by formula **I-6**



I-6

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 59 of 657

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^9, R^{10} and R^{11} are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl;

(j) Z is N or CR^9.

The seventh aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 60 of 657

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, or CH_2SH;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^9, R^{10} and R^{11} are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a $C_{1\text{-}20}$ alkyl; a $C_{1\text{-}20}$ cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl;

(j) Z is N or CR^9.

The eighth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 61 of 657

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is an H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^9, R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

The ninth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 62 of 657

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^9, R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

The tenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 63 of 657

(i) R^9, R^{10} and R^{11} are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

(j) Z is N or CR^9.

A fourth embodiment of the invention is directed to a compound represented by formula **I** in which the base is a structure represented by formula **d** above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, and Y are defined in the Summary of the Invention section above.

The first aspect of the fourth embodiment is directed to a compound represented by formula **I-7**



I-7

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 64 of 657

$CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, -$CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

The second aspect of the fourth embodiment is directed to a compound represented by formula **I-7**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, -$CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 65 of 657

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

The third aspect of the fourth embodiment is directed to a compound represented by formula **I-7**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is an H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

The fourth aspect of the fourth embodiment is directed to a compound represented by formula **I-7**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 66 of 657

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

The fifth aspect of the fourth embodiment is directed to a compound represented by formula **I-7**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 67 of 657


(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

The sixth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**



I-8

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 68 of 657

Ph), or CH_2SH or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

The seventh aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 69 of 657

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is an OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^9, R^{10} and R^{11} are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl;

(j) Z is N or CR^9.

The eighth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 70 of 657

(h) Y is an H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

The ninth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

The tenth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 71 of 657

(c) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, or benzyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;


(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is an OH, NH_2, OCH_3, or $OC(O)CH_3$;

The following tables contain numeric identifiers associated with various substituent designators that should be viewed in light of the accompanying structure. These structures are contemplated species of the various aspects of the disclosed embodiments and are not intended to be limiting on full breadth of the contemplated compound represented by the structure of formula **I**. In each of the presented tables, the phosphoramidate substituent containing the substituents R^{3a} and R^{3b} are depicted without reference to stereochemical structure (*cf.* structures **I-1**, **I-3**, and **I-5** above). It is contemplated that the compounds recited below embody compounds in which R^{3a} projects toward the viewer while R^{3b} projects away from the viewer (cf. structures **I-2**, **I-4**, and **I-6**). Moreover, it is contemplated that the compounds recited below also embody compounds in which R^{3a} projects away from the viewer while R^{3b} projects towards the viewer. Not meant to be limiting, however, the inventors of the present invention contemplate that preferred compounds are those in which R^{3a} projects towards the viewer and R^{3b} projects away from the viewer such that the natural L-amino acid (S)-configuration is presented. Additionally, the inventors recognize that the phosphorus atom of the phosphoramidate moiety is another source of chirality. Although the structures below do not specifically depict chirality at phosphorus, the inventors recognize that stereochemical configurations are possible such that in a staggered (or zig-zag) line structure the oxo-substitutent projects towards the viewer while the OR^1

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 72 of 657

substitutent projects away from the viewer, and vice versa. Therefore, the structures below include all possible stereochemical configurations possible for phosphorus.



II

Table II-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-1-1	CH_3	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-8	CH_3	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-2-1	Et	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-2	Et	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 73 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-8	Et	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-5-1	Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 74 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-8	Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 75 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 76 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-11-1	CH_3	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-2	CH_3	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-8	CH_3	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-12-1	Et	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-2	Et	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-8	Et	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-13-1	^{i}Pr	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-8	^{i}Pr	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 77 of 657

Table II-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-14-1	^{t}Bu	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-8	^{t}Bu	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-15-1	Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-2	Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-8	Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 78 of 657

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-17-1	p-F-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-8	p-F-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 79 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-20-1	p-I-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-8	p-I-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-22-1	Et	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 80 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-8	Et	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 81 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 82 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-31.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 83 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-32-1	Et	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-8	Et	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 84 of 657

Table II-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 85 of 657

Table II-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 86 of 657

Table II-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-41-1	CH_3	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-8	CH_3	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-42-1	Et	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-2	Et	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-8	Et	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 87 of 657

Table II-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-45-1	Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-2	Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-8	Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 88 of 657

Table II-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 89 of 657

Table II-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 90 of 657



III

Table III-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 91 of 657

Table III-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 92 of 657

Table III-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 93 of 657

Table III-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 94 of 657

Table III-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-12-1	Et	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-8	Et	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-14-1	^{t}Bu	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 95 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-14-8	^{t}Bu	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-15-1	Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-8	Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 96 of 657

Table III-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 97 of 657

Table III-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 98 of 657

Table III-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 99 of 657

Table III-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 100 of 657

Table III-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 101 of 657

Table III-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 102 of 657

Table III-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 103 of 657

Table III-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 104 of 657

Table III-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-42-1	Et	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-8	Et	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 105 of 657

Table III-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 106 of 657

Table III-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 107 of 657

Table III-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



IV

Table IV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-1-1	CH_3	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 108 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-1-8	CH_3	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-2-1	Et	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-8	Et	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 109 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-5-1	Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-8	Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 110 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 111 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-11-1	CH_3	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-11-8	CH_3	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-12-1	Et	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-12-2	Et	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-12-8	Et	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 112 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-15-1	Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 113 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-15-2	Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-15-8	Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-18.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 114 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 115 of 657

Table IV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 116 of 657

Table IV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 117 of 657

Table IV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 118 of 657

Table IV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 119 of 657

Table IV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 120 of 657

Table IV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 121 of 657

Table IV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-41-1	CH_3	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-41-8	CH_3	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 122 of 657

Table IV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-42-1	Et	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-42-2	Et	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-42-8	Et	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 123 of 657

Table IV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-45-1	Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-45-8	Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 124 of 657

Table IV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 125 of 657



V

Table V-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-1-1	CH_3	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-1-8	CH_3	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-2-1	Et	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-2-8	Et	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 126 of 657

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-5-1	Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 127 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-5-8	Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 128 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-11-1	CH_3	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 129 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-11-8	CH_3	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-12-1	Et	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-12-2	Et	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-12-8	Et	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 130 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-15-1	Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-15-2	Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-15-8	Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 131 of 657

Table V-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 132 of 657

Table V-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 133 of 657

Table V-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 134 of 657

Table V-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 135 of 657

Table V-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 136 of 657

Table V-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 137 of 657

Table V-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 138 of 657

Table V-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 139 of 657

Table V-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-41-1	CH_3	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-41-8	CH_3	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-42-1	Et	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-42-2	Et	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-42-8	Et	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 140 of 657

Table V-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-45-1	Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-45-8	Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 141 of 657

Table V-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 142 of 657

Table V-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



VI

Table VI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-1-1	CH_3	H	H	H	CH_3	H	F	H	OH	H	H
VI-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 143 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-1-8	CH_3	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-2-1	Et	H	H	H	CH_3	H	F	H	OH	H	H
VI-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-2-8	Et	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH	H	H
VI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH	H	H
VI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 144 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-5-1	Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-5-8	Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 145 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 146 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-11-1	CH_3	H	H	H	Et	H	F	H	OH	H	H
VI-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH	H	H
VI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-11-8	CH_3	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-12-1	Et	H	H	H	Et	H	F	H	OH	H	H
VI-12-2	Et	H	H	CH_3	Et	H	F	H	OH	H	H
VI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-12-8	Et	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 147 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH	H	H
VI-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH	H	H
VI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH	H	H
VI-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH	H	H
VI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-15-1	Ph	H	H	H	Et	H	F	H	OH	H	H
VI-15-2	Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-15-8	Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 148 of 657

Table VI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 149 of 657

Table VI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 150 of 657

Table VI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 151 of 657

Table VI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 152 of 657

Table VI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 153 of 657

Table VI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 154 of 657

Table VI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 155 of 657

Table VI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 156 of 657

Table VI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-41-1	CH_3	H	H	H	Bz	H	F	H	OH	H	H
VI-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-41-8	CH_3	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-42-1	Et	H	H	H	Bz	H	F	H	OH	H	H
VI-42-2	Et	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-42-8	Et	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 157 of 657

Table VI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH	H	H
VI-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH	H	H
VI-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-45-1	Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-45-8	Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 158 of 657

Table VI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 159 of 657

Table VI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 160 of 657



VII

Table VII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-1-1	CH_3	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-8	CH_3	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-2-1	Et	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-8	Et	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 161 of 657

Table VII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-5-1	Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-8	Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 162 of 657

Table VII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 163 of 657

Table VII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-11-1	CH_3	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-11-8	CH_3	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 164 of 657

Table VII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-12-1	Et	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-12-2	Et	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-12-8	Et	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 165 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-15-1	Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-15-2	Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-15-8	Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 166 of 657

Table VII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 167 of 657

Table VII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 168 of 657

Table VII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 169 of 657

Table VII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 170 of 657

Table VII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 171 of 657

Table VII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 172 of 657

Table VII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 173 of 657

Table VII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 174 of 657

Table VII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-41-1	CH_3	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-41-8	CH_3	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-42-1	Et	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-42-2	Et	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-42-8	Et	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 175 of 657

Table VII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-45-1	Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-45-8	Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 176 of 657

Table VII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 177 of 657

Table VII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH	H	H	NH_2
VII-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H	NH_2
VII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H	NH_2
VII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H	NH_2
VII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H	NH_2
VII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H	NH_2
VII-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



VIII

Table VIII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-1-1	CH_3	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 178 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-1-8	CH_3	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-2-1	Et	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-2-2	Et	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-2-8	Et	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 179 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-5-1	Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-5-8	Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 180 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H
VIII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 181 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H
VIII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H
VIII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H
VIII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H
VIII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H
VIII-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-11-1	CH_3	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-11-2	CH_3	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-11-8	CH_3	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-12-1	Et	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-12-2	Et	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-12-8	Et	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 182 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-13-1	^{i}Pr	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-13-8	^{i}Pr	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-14-1	^{t}Bu	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-14-8	^{t}Bu	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-15-1	Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-15-2	Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-15-8	Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 183 of 657

Table VIII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-17-1	p-F-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-17-8	p-F-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 184 of 657

Table VIII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-20-1	p-I-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H
VIII-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H
VIII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H
VIII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H
VIII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H
VIII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H
VIII-20-8	p-I-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 185 of 657

Table VIII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-22-1	Et	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-22-8	Et	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 186 of 657

Table VIII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 187 of 657

Table VIII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H
VIII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 188 of 657

Table VIII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-32-1	Et	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-32-8	Et	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 189 of 657

Table VIII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 190 of 657

Table VIII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 191 of 657

Table VIII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H
VIII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-41-1	CH_3	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-41-8	CH_3	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-42-1	Et	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-42-2	Et	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-42-8	Et	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 192 of 657

Table VIII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-45-1	Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-45-2	Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-45-8	Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 193 of 657

Table VIII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 194 of 657

Table VIII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VIII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VIII-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H
VIII-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H
VIII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H
VIII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H
VIII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H
VIII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H
VIII-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 195 of 657



IX

Table IX-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 196 of 657

Table IX-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 197 of 657

Table IX-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 198 of 657

Table IX-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H
IX-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H
IX-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H
IX-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H
IX-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H
IX-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H
IX-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH	H	H
IX-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 199 of 657

Table IX-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-12-1	Et	H	H	H	Et	H	CH_3	F	OH	H	H
IX-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-12-8	Et	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH	H	H
IX-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-14-1	^{t}Bu	H	H	H	Et	H	CH_3	F	OH	H	H
IX-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-14-8	^{t}Bu	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 200 of 657

Table IX-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-15-1	Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-15-8	Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 201 of 657

Table IX-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH	H	H
IX-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H
IX-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H
IX-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H
IX-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H
IX-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H
IX-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 202 of 657

Table IX-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 203 of 657

Table IX-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 204 of 657

Table IX-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 205 of 657

Table IX-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H
IX-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 206 of 657

Table IX-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 207 of 657

Table IX-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 208 of 657

Table IX-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H
IX-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 209 of 657

Table IX-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-42-1	Et	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-42-8	Et	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 210 of 657

Table IX-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 211 of 657

Table IX-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IX-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
IX-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H
IX-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H
IX-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H
IX-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H
IX-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H
IX-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H
IX-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 212 of 657



X

Table X-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-1-1	CH_3	H	H	H	CH_3	H	F	F	OH	H	H
X-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-1-8	CH_3	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-2-1	Et	H	H	H	CH_3	H	F	F	OH	H	H
X-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 213 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-2-8	Et	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH	H	H
X-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-4-1	^{t}Bu	H	H	H	CH_3	H	F	F	OH	H	H
X-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-4-8	^{t}Bu	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-5-1	Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 214 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-5-8	Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 215 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH	H	H
X-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H
X-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H
X-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H
X-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H
X-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H
X-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-11-1	CH_3	H	H	H	Et	H	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 216 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH	H	H
X-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-11-8	CH_3	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-12-1	Et	H	H	H	Et	H	F	F	OH	H	H
X-12-2	Et	H	H	CH_3	Et	H	F	F	OH	H	H
X-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-12-8	Et	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH	H	H
X-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH	H	H
X-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-14.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 217 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH	H	H
X-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH	H	H
X-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-15-1	Ph	H	H	H	Et	H	F	F	OH	H	H
X-15-2	Ph	H	H	CH_3	Et	H	F	F	OH	H	H
X-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-15-8	Ph	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH	H	H
X-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH	H	H
X-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 218 of 657

Table X-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH	H	H
X-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH	H	H
X-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH	H	H
X-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH	H	H
X-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH	H	H
X-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH	H	H
X-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H
X-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H
X-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H
X-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H
X-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 219 of 657

Table X-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7
X-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH	H
X-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH	H
X-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H
X-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H
X-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H
X-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H
X-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H
X-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 220 of 657

Table X-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 221 of 657

Table X-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 222 of 657

Table X-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H
X-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H
X-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H
X-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H
X-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H
X-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H
X-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 223 of 657

Table X-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 224 of 657

Table X-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 225 of 657

Table X-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H
X-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H
X-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H
X-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H
X-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H
X-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H
X-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 226 of 657

Table X-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-41-1	CH_3	H	H	H	Bz	H	F	F	OH	H	H
X-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH	H	H
X-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-41-8	CH_3	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-42-1	Et	H	H	H	Bz	H	F	F	OH	H	H
X-42-2	Et	H	H	CH_3	Bz	H	F	F	OH	H	H
X-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-42-8	Et	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH	H	H
X-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH	H	H
X-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 227 of 657

Table X-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH	H	H
X-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH	H	H
X-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-45-1	Ph	H	H	H	Bz	H	F	F	OH	H	H
X-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-45-8	Ph	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH	H	H
X-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 228 of 657

Table X-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH	H	H
X-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH	H	H
X-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table X-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH	H	H
X-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 229 of 657

Table X-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
X-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH	H	H
X-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H
X-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H
X-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H
X-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H
X-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H
X-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



XI

Table XI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-1-1	CH_3	H	H	H	CH_3	H	H	F	OH	H	H
XI-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 230 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-1-8	CH_3	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-2-1	Et	H	H	H	CH_3	H	H	F	OH	H	H
XI-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-2-8	Et	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH	H	H
XI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH	H	H
XI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 231 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-5-1	Ph	H	H	H	CH_3	H	H	F	OH	H	H
XI-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-5-8	Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH	H	H
XI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH	H	H
XI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 232 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH	H	H
XI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH	H	H
XI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 233 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H
XI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H
XI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H
XI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H
XI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H
XI-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-11-1	CH_3	H	H	H	Et	H	H	F	OH	H	H
XI-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH	H	H
XI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-11-8	CH_3	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-12-1	Et	H	H	H	Et	H	H	F	OH	H	H
XI-12-2	Et	H	H	CH_3	Et	H	H	F	OH	H	H
XI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-12-8	Et	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 234 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH	H	H
XI-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH	H	H
XI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH	H	H
XI-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH	H	H
XI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-15-1	Ph	H	H	H	Et	H	H	F	OH	H	H
XI-15-2	Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-15-8	Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 235 of 657

Table XI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH	H	H
XI-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH	H	H
XI-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH	H	H
XI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 236 of 657

Table XI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH	H	H
XI-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH	H	H
XI-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH	H	H
XI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H
XI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H
XI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H
XI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H
XI-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 237 of 657

Table XI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 238 of 657

Table XI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 239 of 657

Table XI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H
XI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H
XI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H
XI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H
XI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H
XI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H
XI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 240 of 657

Table XI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 241 of 657

Table XI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 242 of 657

Table XI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 243 of 657

Table XI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H
XI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H
XI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H
XI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H
XI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H
XI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H
XI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-41-1	CH_3	H	H	H	Bz	H	H	F	OH	H	H
XI-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-41-8	CH_3	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-42-1	Et	H	H	H	Bz	H	H	F	OH	H	H
XI-42-2	Et	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-42-8	Et	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 244 of 657

Table XI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH	H	H
XI-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH	H	H
XI-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-45-1	Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-45-8	Ph	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 245 of 657

Table XI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 246 of 657

Table XI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XI-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH	H	H
XI-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H
XI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H
XI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H
XI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H
XI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H
XI-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 247 of 657



XII

Table XII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-1-1	CH_3	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-1-2	CH_3	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-1-5	CH_3	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-1-8	CH_3	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-2-1	Et	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-2-2	Et	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-2-5	Et	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-2-8	Et	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 248 of 657

Table XII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-3-1	^{i}Pr	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-3-2	^{i}Pr	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-3-8	^{i}Pr	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-4-1	^{t}Bu	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-4-2	^{t}Bu	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-4-8	^{t}Bu	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-5-1	Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-5-2	Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-5-5	Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-5-8	Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 249 of 657

Table XII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-6-1	p-Me-Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-6-2	p-Me-Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-6-8	p-Me-Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-7-1	p-F-Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-7-2	p-F-Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-7-20	p-F-Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-8-1	p-Cl-Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-8-8	p-Cl-Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 250 of 657

Table XII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-9-1	p-Br-Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-9-2	p-Br-Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-9-20	p-Br-Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-10-1	p-I-Ph	H	H	H	CH_3	N_3	F	F	OH	H	H
XII-10-2	p-I-Ph	H	H	CH_3	CH_3	N_3	F	F	OH	H	H
XII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	F	OH	H	H
XII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	N_3	F	F	OH	H	H
XII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	F	OH	H	H
XII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	F	OH	H	H
XII-10-8	p-I-Ph	*	H	*	CH_3	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-11-1	CH_3	H	H	H	Et	N_3	F	F	OH	H	H
XII-11-2	CH_3	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-11-5	CH_3	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-11-8	CH_3	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 251 of 657

Table XII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-12-1	Et	H	H	H	Et	N_3	F	F	OH	H	H
XII-12-2	Et	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-12-3	Et	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-12-5	Et	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-12-6	Et	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-12-8	Et	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-13-1	^{i}Pr	H	H	H	Et	N_3	F	F	OH	H	H
XII-13-2	^{i}Pr	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-13-5	^{i}Pr	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-13-8	^{i}Pr	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-14-1	^{t}Bu	H	H	H	Et	N_3	F	F	OH	H	H
XII-14-2	^{t}Bu	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 252 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-14-5	^{t}Bu	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-14-8	^{t}Bu	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-15-1	Ph	H	H	H	Et	N_3	F	F	OH	H	H
XII-15-2	Ph	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-15-5	Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-15-8	Ph	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-16-1	p-Me-Ph	H	H	H	Et	N_3	F	F	OH	H	H
XII-16-2	p-Me-Ph	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-16-8	p-Me-Ph	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 253 of 657

Table XII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-17-1	p-F-Ph	H	H	H	Et	N_3	F	F	OH	H	H
XII-17-2	p-F-Ph	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-17-5	p-F-Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-17-8	p-F-Ph	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-18-1	p-Cl-Ph	H	H	H	Et	N_3	F	F	OH	H	H
XII-18-2	p-Cl-Ph	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-18-8	p-Cl-Ph	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-19-1	p-Br-Ph	H	H	H	Et	N_3	F	F	OH	H	H
XII-19-2	p-Br-Ph	H	H	CH_3	Et	N_3	F	F	OH	H	H
XII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H	H
XII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H	H
XII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H	H
XII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H	H
XII-19-8	p-Br-Ph	*	H	*	Et	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 254 of 657

Table XII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7
XII-20-1	p-I-Ph	H	H	H	Et	N_3	F	F	OH	H
XII-20-2	p-I-Ph	H	H	CH_3	Et	N_3	F	F	OH	H
XII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	F	OH	H
XII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	F	OH	H
XII-20-5	p-I-Ph	H	H	CH_2Ph	Et	N_3	F	F	OH	H
XII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	F	OH	H
XII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	F	OH	H
XII-20-8	p-I-Ph	*	H	*	Et	N_3	F	F	OH	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-21-1	CH_3	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-21-2	CH_3	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-21-8	CH_3	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-22-1	Et	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-22-2	Et	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-22-5	Et	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-22-8	Et	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 255 of 657

Table XII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-23-1	^{i}Pr	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-23-8	^{i}Pr	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-24-1	^{t}Bu	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-24-8	^{t}Bu	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-25-1	Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-25-2	Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-25-8	Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 256 of 657

Table XII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-26-1	p-Me-Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-26-8	p-Me-Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-27-1	p-F-Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-27-8	p-F-Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 257 of 657

Table XII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-29-1	p-Br-Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-29-8	p-Br-Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-30-1	p-I-Ph	H	H	H	^{i}Pr	N_3	F	F	OH	H	H
XII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	N_3	F	F	OH	H	H
XII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	F	OH	H	H
XII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	F	OH	H	H
XII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	F	OH	H	H
XII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	F	OH	H	H
XII-30-8	p-I-Ph	*	H	*	^{i}Pr	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-31-1	CH_3	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-31-2	CH_3	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-31-8	CH_3	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 258 of 657

Table XII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-32-1	Et	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-32-2	Et	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-32-5	Et	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-32-8	Et	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-33-1	^{i}Pr	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-33-8	^{i}Pr	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-34-1	^{t}Bu	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-34-8	^{t}Bu	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 259 of 657

Table XII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-35-1	Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-35-2	Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-35-8	Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-36-1	p-Me-Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-36-8	p-Me-Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-37-1	p-F-Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-37-8	p-F-Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 260 of 657

Table XII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-39-1	p-Br-Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-39-8	p-Br-Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-40-1	p-I-Ph	H	H	H	^{n}Bu	N_3	F	F	OH	H	H
XII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	N_3	F	F	OH	H	H
XII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	F	OH	H	H
XII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	F	OH	H	H
XII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	F	OH	H	H
XII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	F	OH	H	H
XII-40-8	p-I-Ph	*	H	*	^{n}Bu	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 261 of 657

Table XII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-41-1	CH_3	H	H	H	Bz	N_3	F	F	OH	H	H
XII-41-2	CH_3	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-41-5	CH_3	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-41-8	CH_3	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-42-1	Et	H	H	H	Bz	N_3	F	F	OH	H	H
XII-42-2	Et	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-42-5	Et	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-42-8	Et	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-43-1	^{i}Pr	H	H	H	Bz	N_3	F	F	OH	H	H
XII-43-2	^{i}Pr	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-43-8	^{i}Pr	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 262 of 657

Table XII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-44-1	^{t}Bu	H	H	H	Bz	N_3	F	F	OH	H	H
XII-44-2	^{t}Bu	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-44-8	^{t}Bu	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-45-1	Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-45-2	Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-45-5	Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-45-8	Ph	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-46-1	p-Me-Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-46-2	p-Me-Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-46-8	p-Me-Ph	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 263 of 657

Table XII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-47-1	p-F-Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-47-2	p-F-Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-47-8	p-F-Ph	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-48-1	p-Cl-Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-48-2	p-Cl-Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-48-8	p-Cl-Ph	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-49-1	p-Br-Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-49-2	p-Br-Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-49-8	p-Br-Ph	*	H	*	Bz	N_3	F	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 264 of 657

Table XII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XII-50-1	p-I-Ph	H	H	H	Bz	N_3	F	F	OH	H	H
XII-50-2	p-I-Ph	H	H	CH_3	Bz	N_3	F	F	OH	H	H
XII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	F	OH	H	H
XII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	N_3	F	F	OH	H	H
XII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	F	OH	H	H
XII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	F	OH	H	H
XII-50-8	p-I-Ph	*	H	*	Bz	N_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



XIII

Table XIII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-1-1	CH_3	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-1-2	CH_3	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-1-5	CH_3	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 265 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-1-8	CH_3	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-2-1	Et	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-2-2	Et	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-2-5	Et	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-2-8	Et	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-3-1	^{i}Pr	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-3-2	^{i}Pr	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-3-8	^{i}Pr	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-4-1	^{t}Bu	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-4-2	^{t}Bu	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 266 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-4-8	^{t}Bu	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-5-1	Ph	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-5-2	Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-5-5	Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-5-8	Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-6-1	p-Me-Ph	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-6-2	p-Me-Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-6-8	p-Me-Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-7-1	p-F-Ph	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-7-2	p-F-Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 267 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-7-20	p-F-Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-8-1	p-Cl-Ph	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-8-8	p-Cl-Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-9-1	p-Br-Ph	H	H	H	CH_3	N_3	H	F	OH	H	H
XIII-9-2	p-Br-Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-9-20	p-Br-Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-10-1	p-I-Ph	H	H	H	CH_3	N_3	H	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 268 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-10-2	p-I-Ph	H	H	CH_3	CH_3	N_3	H	F	OH	H	H
XIII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	H	F	OH	H	H
XIII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	N_3	H	F	OH	H	H
XIII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	H	F	OH	H	H
XIII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	H	F	OH	H	H
XIII-10-8	p-I-Ph	*	H	*	CH_3	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-11-1	CH_3	H	H	H	Et	N_3	H	F	OH	H	H
XIII-11-2	CH_3	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-11-5	CH_3	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-11-8	CH_3	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-12-1	Et	H	H	H	Et	N_3	H	F	OH	H	H
XIII-12-2	Et	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-12-3	Et	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-12-5	Et	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-12-6	Et	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-12-8	Et	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 269 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-13-1	^{i}Pr	H	H	H	Et	N_3	H	F	OH	H	H
XIII-13-2	^{i}Pr	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-13-5	^{i}Pr	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-13-8	^{i}Pr	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-14-1	^{t}Bu	H	H	H	Et	N_3	H	F	OH	H	H
XIII-14-2	^{t}Bu	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-14-5	^{t}Bu	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-14-8	^{t}Bu	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-15-1	Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-15-2	Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-15-5	Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-15-8	Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 270 of 657

Table XIII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-16-1	p-Me-Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-16-2	p-Me-Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-16-8	p-Me-Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-17-1	p-F-Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-17-2	p-F-Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-17-5	p-F-Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-17-8	p-F-Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-18-1	p-Cl-Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-18-2	p-Cl-Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-18-8	p-Cl-Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 271 of 657

Table XIII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-19-1	p-Br-Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-19-2	p-Br-Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-19-8	p-Br-Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-20-1	p-I-Ph	H	H	H	Et	N_3	H	F	OH	H	H
XIII-20-2	p-I-Ph	H	H	CH_3	Et	N_3	H	F	OH	H	H
XIII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	H	F	OH	H	H
XIII-20-5	p-I-Ph	H	H	CH_2Ph	Et	N_3	H	F	OH	H	H
XIII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	N_3	H	F	OH	H	H
XIII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	H	F	OH	H	H
XIII-20-8	p-I-Ph	*	H	*	Et	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-21-1	CH_3	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-2	CH_3	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-21-8	CH_3	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 272 of 657

Table XIII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-22-1	Et	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-2	Et	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-5	Et	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-22-8	Et	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-23-1	^{i}Pr	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-23-8	^{i}Pr	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-24-1	^{t}Bu	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-24-8	^{t}Bu	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 273 of 657

Table XIII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-25-1	Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-2	Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-25-8	Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-26-1	p-Me-Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-26-8	p-Me-Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-27-1	p-F-Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-27-8	p-F-Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 274 of 657

Table XIII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-29-1	p-Br-Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-29-8	p-Br-Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-30-1	p-I-Ph	H	H	H	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	H	F	OH	H	H
XIII-30-8	p-I-Ph	*	H	*	^{i}Pr	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 275 of 657

Table XIII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-31-1	CH_3	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-2	CH_3	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-31-8	CH_3	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-32-1	Et	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-2	Et	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-5	Et	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-32-8	Et	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-33-1	^{i}Pr	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-33-8	^{i}Pr	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 276 of 657

Table XIII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-34-1	^{t}Bu	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-34-8	^{t}Bu	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-35-1	Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-2	Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-35-8	Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-36-1	p-Me-Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-36-8	p-Me-Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 277 of 657

Table XIII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-37-1	p-F-Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-37-8	p-F-Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-39-1	p-Br-Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-39-8	p-Br-Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 278 of 657

Table XIII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-40-1	p-I-Ph	H	H	H	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	H	F	OH	H	H
XIII-40-8	p-I-Ph	*	H	*	^{n}Bu	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-41-1	CH_3	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-41-2	CH_3	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-41-5	CH_3	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-41-8	CH_3	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-42-1	Et	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-42-2	Et	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-42-5	Et	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-42-8	Et	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 279 of 657

Table XIII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-43-1	^{i}Pr	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-43-2	^{i}Pr	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-43-8	^{i}Pr	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-44-1	^{t}Bu	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-44-2	^{t}Bu	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-44-8	^{t}Bu	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-45-1	Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-45-2	Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-45-5	Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-45-8	Ph	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 280 of 657

Table XIII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-46-1	p-Me-Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-46-2	p-Me-Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-46-8	p-Me-Ph	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-47-1	p-F-Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-47-2	p-F-Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-47-8	p-F-Ph	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-48-1	p-Cl-Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-48-2	p-Cl-Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-48-8	p-Cl-Ph	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 281 of 657

Table XIII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-49-1	p-Br-Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-49-2	p-Br-Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-49-8	p-Br-Ph	*	H	*	Bz	N_3	H	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIII-50-1	p-I-Ph	H	H	H	Bz	N_3	H	F	OH	H	H
XIII-50-2	p-I-Ph	H	H	CH_3	Bz	N_3	H	F	OH	H	H
XIII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	H	F	OH	H	H
XIII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	N_3	H	F	OH	H	H
XIII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	H	F	OH	H	H
XIII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	H	F	OH	H	H
XIII-50-8	p-I-Ph	*	H	*	Bz	N_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 282 of 657



XIV

Table XIV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-1-1	CH_3	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-1-2	CH_3	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-1-5	CH_3	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-1-8	CH_3	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-2-1	Et	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-2-2	Et	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-2-5	Et	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-2-8	Et	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 283 of 657

Table XIV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-3-1	^{i}Pr	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-3-2	^{i}Pr	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-3-8	^{i}Pr	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-4-1	^{t}Bu	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-4-2	^{t}Bu	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-4-8	^{t}Bu	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-5-1	Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-5-2	Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-5-5	Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-5-8	Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 284 of 657

Table XIV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-6-1	p-Me-Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-6-2	p-Me-Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-6-8	p-Me-Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-7-1	p-F-Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-7-2	p-F-Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-7-20	p-F-Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-8-1	p-Cl-Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-8-8	p-Cl-Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 285 of 657

Table XIV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-9-1	p-Br-Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-9-2	p-Br-Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-9-20	p-Br-Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-10-1	p-I-Ph	H	H	H	CH_3	N_3	F	H	OH	H	H
XIV-10-2	p-I-Ph	H	H	CH_3	CH_3	N_3	F	H	OH	H	H
XIV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	N_3	F	H	OH	H	H
XIV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	N_3	F	H	OH	H	H
XIV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	N_3	F	H	OH	H	H
XIV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	N_3	F	H	OH	H	H
XIV-10-8	p-I-Ph	*	H	*	CH_3	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-11-1	CH_3	H	H	H	Et	N_3	F	H	OH	H	H
XIV-11-2	CH_3	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-11-5	CH_3	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-11-8	CH_3	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 286 of 657

Table XIV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-12-1	Et	H	H	H	Et	N_3	F	H	OH	H	H
XIV-12-2	Et	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-12-3	Et	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-12-5	Et	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-12-6	Et	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-12-8	Et	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-13-1	^{i}Pr	H	H	H	Et	N_3	F	H	OH	H	H
XIV-13-2	^{i}Pr	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-13-5	^{i}Pr	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-13-8	^{i}Pr	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-14-1	^{t}Bu	H	H	H	Et	N_3	F	H	OH	H	H
XIV-14-2	^{t}Bu	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-14-5	^{t}Bu	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-14-8	^{t}Bu	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 287 of 657

Table XIV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-15-1	Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-15-2	Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-15-5	Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-15-8	Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-16-1	p-Me-Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-16-2	p-Me-Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-16-8	p-Me-Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-17-1	p-F-Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-17-2	p-F-Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-17-5	p-F-Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-17-8	p-F-Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 288 of 657

Table XIV-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-18-1	p-Cl-Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-18-2	p-Cl-Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-18-8	p-Cl-Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-19-1	p-Br-Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-19-2	p-Br-Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-19-8	p-Br-Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-20-1	p-I-Ph	H	H	H	Et	N_3	F	H	OH	H	H
XIV-20-2	p-I-Ph	H	H	CH_3	Et	N_3	F	H	OH	H	H
XIV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	N_3	F	H	OH	H	H
XIV-20-5	p-I-Ph	H	H	CH_2Ph	Et	N_3	F	H	OH	H	H
XIV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	N_3	F	H	OH	H	H
XIV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	N_3	F	H	OH	H	H
XIV-20-8	p-I-Ph	*	H	*	Et	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 289 of 657

Table XIV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-21-1	CH_3	H	H	H	^iPr	N_3	F	H	OH	H	H
XIV-21-2	CH_3	H	H	CH_3	^iPr	N_3	F	H	OH	H	H
XIV-21-3	CH_3	H	H	$CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-21-5	CH_3	H	H	CH_2Ph	^iPr	N_3	F	H	OH	H	H
XIV-21-6	CH_3	H	H	CH_2-indol-3-yl	^iPr	N_3	F	H	OH	H	H
XIV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^iPr	N_3	F	H	OH	H	H
XIV-21-8	CH_3	*	H	*	^iPr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-22-1	Et	H	H	H	^iPr	N_3	F	H	OH	H	H
XIV-22-2	Et	H	H	CH_3	^iPr	N_3	F	H	OH	H	H
XIV-22-3	Et	H	H	$CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-22-5	Et	H	H	CH_2Ph	^iPr	N_3	F	H	OH	H	H
XIV-22-6	Et	H	H	CH_2-indol-3-yl	^iPr	N_3	F	H	OH	H	H
XIV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^iPr	N_3	F	H	OH	H	H
XIV-22-8	Et	*	H	*	^iPr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-23-1	^iPr	H	H	H	^iPr	N_3	F	H	OH	H	H
XIV-23-2	^iPr	H	H	CH_3	^iPr	N_3	F	H	OH	H	H
XIV-23-3	^iPr	H	H	$CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-23-4	^iPr	H	H	$CH_2CH(CH_3)_2$	^iPr	N_3	F	H	OH	H	H
XIV-23-5	^iPr	H	H	CH_2Ph	^iPr	N_3	F	H	OH	H	H
XIV-23-6	^iPr	H	H	CH_2-indol-3-yl	^iPr	N_3	F	H	OH	H	H
XIV-23-7	^iPr	H	H	$CH_2CH_2SCH_3$	^iPr	N_3	F	H	OH	H	H
XIV-23-8	^iPr	*	H	*	^iPr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 290 of 657

Table XIV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-24-1	^{t}Bu	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-24-8	^{t}Bu	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-25-1	Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-2	Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-25-8	Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-26-1	p-Me-Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-26-8	p-Me-Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 291 of 657

Table XIV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-27-1	p-F-Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-27-8	p-F-Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-29-1	p-Br-Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-29-8	p-Br-Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 292 of 657

Table XIV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-30-1	p-I-Ph	H	H	H	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	N_3	F	H	OH	H	H
XIV-30-8	p-I-Ph	*	H	*	^{i}Pr	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-31-1	CH_3	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-2	CH_3	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-31-8	CH_3	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-32-1	Et	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-2	Et	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-5	Et	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-32-8	Et	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 293 of 657

Table XIV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-33-1	^{i}Pr	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-33-8	^{i}Pr	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-34-1	^{t}Bu	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-34-8	^{t}Bu	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-35-1	Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-2	Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-35-8	Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 294 of 657

Table XIV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-36-1	p-Me-Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-36-8	p-Me-Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-37-1	p-F-Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-37-8	p-F-Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 295 of 657

Table XIV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-39-1	p-Br-Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-39-8	p-Br-Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-40-1	p-I-Ph	H	H	H	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	N_3	F	H	OH	H	H
XIV-40-8	p-I-Ph	*	H	*	^{n}Bu	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-41-1	CH_3	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-41-2	CH_3	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-41-5	CH_3	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-41-8	CH_3	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 296 of 657

Table XIV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-42-1	Et	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-42-2	Et	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-42-5	Et	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-42-8	Et	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-43-1	^{i}Pr	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-43-2	^{i}Pr	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-43-8	^{i}Pr	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-44-1	^{t}Bu	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-44-2	^{t}Bu	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-44-8	^{t}Bu	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 297 of 657

Table XIV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-45-1	Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-45-2	Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-45-5	Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-45-8	Ph	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-46-1	p-Me-Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-46-2	p-Me-Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-46-8	p-Me-Ph	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-47-1	p-F-Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-47-2	p-F-Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-47-8	p-F-Ph	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 298 of 657

Table XIV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-48-1	p-Cl-Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-48-2	p-Cl-Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-48-8	p-Cl-Ph	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-49-1	p-Br-Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-49-2	p-Br-Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-49-8	p-Br-Ph	*	H	*	Bz	N_3	F	H	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIV-50-1	p-I-Ph	H	H	H	Bz	N_3	F	H	OH	H	H
XIV-50-2	p-I-Ph	H	H	CH_3	Bz	N_3	F	H	OH	H	H
XIV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	N_3	F	H	OH	H	H
XIV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	N_3	F	H	OH	H	H
XIV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	N_3	F	H	OH	H	H
XIV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	N_3	F	H	OH	H	H
XIV-50-8	p-I-Ph	*	H	*	Bz	N_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 299 of 657



XV

Table XV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-1-1	CH_3	H	H	H	CH_3	F	F	H	OH	H	H
XV-1-2	CH_3	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-1-5	CH_3	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-1-8	CH_3	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-2-1	Et	H	H	H	CH_3	F	F	H	OH	H	H
XV-2-2	Et	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-2-5	Et	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 300 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-2-8	Et	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-3-1	^{i}Pr	H	H	H	CH_3	F	F	H	OH	H	H
XV-3-2	^{i}Pr	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-3-8	^{i}Pr	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-4-1	^{t}Bu	H	H	H	CH_3	F	F	H	OH	H	H
XV-4-2	^{t}Bu	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-4-8	^{t}Bu	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-5-1	Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-5-2	Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-5-5	Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 301 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-5-8	Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-6-1	p-Me-Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-6-2	p-Me-Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-6-8	p-Me-Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-7-1	p-F-Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-7-2	p-F-Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-7-20	p-F-Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-8-1	p-Cl-Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 302 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-8-8	p-Cl-Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-9-1	p-Br-Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-9-2	p-Br-Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-9-20	p-Br-Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-10-1	p-I-Ph	H	H	H	CH_3	F	F	H	OH	H	H
XV-10-2	p-I-Ph	H	H	CH_3	CH_3	F	F	H	OH	H	H
XV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	H	OH	H	H
XV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	F	F	H	OH	H	H
XV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	H	OH	H	H
XV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	H	OH	H	H
XV-10-8	p-I-Ph	*	H	*	CH_3	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-11-1	CH_3	H	H	H	Et	F	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 303 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-11-2	CH_3	H	H	CH_3	Et	F	F	H	OH	H	H
XV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-11-5	CH_3	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-11-8	CH_3	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-12-1	Et	H	H	H	Et	F	F	H	OH	H	H
XV-12-2	Et	H	H	CH_3	Et	F	F	H	OH	H	H
XV-12-3	Et	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-12-5	Et	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-12-6	Et	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-12-8	Et	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-13-1	^{i}Pr	H	H	H	Et	F	F	H	OH	H	H
XV-13-2	^{i}Pr	H	H	CH_3	Et	F	F	H	OH	H	H
XV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-13-5	^{i}Pr	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-13-8	^{i}Pr	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-14.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 304 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-14-1	^{t}Bu	H	H	H	Et	F	F	H	OH	H	H
XV-14-2	^{t}Bu	H	H	CH_3	Et	F	F	H	OH	H	H
XV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-14-5	^{t}Bu	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-14-8	^{t}Bu	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-15-1	Ph	H	H	H	Et	F	F	H	OH	H	H
XV-15-2	Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-15-5	Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-15-8	Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-16-1	p-Me-Ph	H	H	H	Et	F	F	H	OH	H	H
XV-16-2	p-Me-Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-16-8	p-Me-Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 305 of 657

Table XV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-17-1	p-F-Ph	H	H	H	Et	F	F	H	OH	H	H
XV-17-2	p-F-Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-17-5	p-F-Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-17-8	p-F-Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-18-1	p-Cl-Ph	H	H	H	Et	F	F	H	OH	H	H
XV-18-2	p-Cl-Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-18-8	p-Cl-Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-19-1	p-Br-Ph	H	H	H	Et	F	F	H	OH	H	H
XV-19-2	p-Br-Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-19-8	p-Br-Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 306 of 657

Table XV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-20-1	p-I-Ph	H	H	H	Et	F	F	H	OH	H	H
XV-20-2	p-I-Ph	H	H	CH_3	Et	F	F	H	OH	H	H
XV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	H	OH	H	H
XV-20-5	p-I-Ph	H	H	CH_2Ph	Et	F	F	H	OH	H	H
XV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	F	F	H	OH	H	H
XV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	H	OH	H	H
XV-20-8	p-I-Ph	*	H	*	Et	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-21-1	CH_3	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-21-2	CH_3	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-21-8	CH_3	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-22-1	Et	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-22-2	Et	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-22-5	Et	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-22-8	Et	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 307 of 657

Table XV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-23-1	^{i}Pr	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-23-8	^{i}Pr	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-24-1	^{t}Bu	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-24-8	^{t}Bu	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-25-1	Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-25-2	Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-25-8	Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 308 of 657

Table XV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-26-1	p-Me-Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-26-8	p-Me-Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-27-1	p-F-Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-27-8	p-F-Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 309 of 657

Table XV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-29-1	p-Br-Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-29-8	p-Br-Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-30-1	p-I-Ph	H	H	H	^{i}Pr	F	F	H	OH	H	H
XV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	F	F	H	OH	H	H
XV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	H	OH	H	H
XV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	F	F	H	OH	H	H
XV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	H	OH	H	H
XV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	H	OH	H	H
XV-30-8	p-I-Ph	*	H	*	^{i}Pr	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-31-1	CH_3	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-31-2	CH_3	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-31-8	CH_3	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 310 of 657

Table XV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-32-1	Et	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-32-2	Et	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-32-5	Et	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-32-8	Et	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-33-1	^{i}Pr	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-33-8	^{i}Pr	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-34-1	^{t}Bu	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-34-8	^{t}Bu	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 311 of 657

Table XV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-35-1	Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-35-2	Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-35-8	Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-36-1	p-Me-Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-36-8	p-Me-Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-37-1	p-F-Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-37-8	p-F-Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 312 of 657

Table XV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-39-1	p-Br-Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-39-8	p-Br-Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-40-1	p-I-Ph	H	H	H	^{n}Bu	F	F	H	OH	H	H
XV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	F	F	H	OH	H	H
XV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	H	OH	H	H
XV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	F	F	H	OH	H	H
XV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	H	OH	H	H
XV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	H	OH	H	H
XV-40-8	p-I-Ph	*	H	*	^{n}Bu	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 313 of 657

Table XV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-41-1	CH_3	H	H	H	Bz	F	F	H	OH	H	H
XV-41-2	CH_3	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-41-5	CH_3	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-41-8	CH_3	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-42-1	Et	H	H	H	Bz	F	F	H	OH	H	H
XV-42-2	Et	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-42-5	Et	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-42-8	Et	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-43-1	^{i}Pr	H	H	H	Bz	F	F	H	OH	H	H
XV-43-2	^{i}Pr	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-43-8	^{i}Pr	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 314 of 657

Table XV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-44-1	^{t}Bu	H	H	H	Bz	F	F	H	OH	H	H
XV-44-2	^{t}Bu	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-44-8	^{t}Bu	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-45-1	Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-45-2	Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-45-5	Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-45-8	Ph	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-46-1	p-Me-Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-46-2	p-Me-Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-46-8	p-Me-Ph	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 315 of 657

Table XV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-47-1	p-F-Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-47-2	p-F-Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-47-8	p-F-Ph	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-48-1	p-Cl-Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-48-2	p-Cl-Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-48-8	p-Cl-Ph	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-49-1	p-Br-Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-49-2	p-Br-Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-49-8	p-Br-Ph	*	H	*	Bz	F	F	H	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 316 of 657

Table XV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XV-50-1	p-I-Ph	H	H	H	Bz	F	F	H	OH	H	H
XV-50-2	p-I-Ph	H	H	CH_3	Bz	F	F	H	OH	H	H
XV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	H	OH	H	H
XV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	F	F	H	OH	H	H
XV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	H	OH	H	H
XV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	H	OH	H	H
XV-50-8	p-I-Ph	*	H	*	Bz	F	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



XVI

Table XVI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-1-1	CH_3	H	H	H	CH_3	F	F	F	OH	H	H
XVI-1-2	CH_3	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-1-5	CH_3	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 317 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-1-8	CH_3	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-2-1	Et	H	H	H	CH_3	F	F	F	OH	H	H
XVI-2-2	Et	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-2-5	Et	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-2-8	Et	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-3-1	^{i}Pr	H	H	H	CH_3	F	F	F	OH	H	H
XVI-3-2	^{i}Pr	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-3-8	^{i}Pr	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-4-1	^{t}Bu	H	H	H	CH_3	F	F	F	OH	H	H
XVI-4-2	^{t}Bu	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 318 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-4-8	^{t}Bu	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-5-1	Ph	H	H	H	CH_3	F	F	F	OH	H	H
XVI-5-2	Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-5-5	Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-5-8	Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-6-1	p-Me-Ph	H	H	H	CH_3	F	F	F	OH	H	H
XVI-6-2	p-Me-Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-6-8	p-Me-Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-7-1	p-F-Ph	H	H	H	CH_3	F	F	F	OH	H	H
XVI-7-2	p-F-Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 319 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-7-20	p-F-Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-8-1	p-Cl-Ph	H	H	H	CH_3	F	F	F	OH	H	H
XVI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-8-8	p-Cl-Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-9-1	p-Br-Ph	H	H	H	CH_3	F	F	F	OH	H	H
XVI-9-2	p-Br-Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-9-20	p-Br-Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-10-1	p-I-Ph	H	H	H	CH_3	F	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 320 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-10-2	p-I-Ph	H	H	CH_3	CH_3	F	F	F	OH	H	H
XVI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	F	F	OH	H	H
XVI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	F	F	F	OH	H	H
XVI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	F	F	F	OH	H	H
XVI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	F	F	OH	H	H
XVI-10-8	p-I-Ph	*	H	*	CH_3	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-11-1	CH_3	H	H	H	Et	F	F	F	OH	H	H
XVI-11-2	CH_3	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-11-5	CH_3	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-11-8	CH_3	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-12-1	Et	H	H	H	Et	F	F	F	OH	H	H
XVI-12-2	Et	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-12-3	Et	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-12-5	Et	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-12-6	Et	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-12-8	Et	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 321 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-13-1	^{i}Pr	H	H	H	Et	F	F	F	OH	H	H
XVI-13-2	^{i}Pr	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-13-5	^{i}Pr	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-13-8	^{i}Pr	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-14-1	^{t}Bu	H	H	H	Et	F	F	F	OH	H	H
XVI-14-2	^{t}Bu	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-14-5	^{t}Bu	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-14-8	^{t}Bu	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-15-1	Ph	H	H	H	Et	F	F	F	OH	H	H
XVI-15-2	Ph	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-15-5	Ph	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-15-8	Ph	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 322 of 657

Table XVI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-16-1	p-Me-Ph	H	H	H	Et	F	F	F	OH	H	H
XVI-16-2	p-Me-Ph	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-16-8	p-Me-Ph	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-17-1	p-F-Ph	H	H	H	Et	F	F	F	OH	H	H
XVI-17-2	p-F-Ph	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-17-5	p-F-Ph	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-17-8	p-F-Ph	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-18-1	p-Cl-Ph	H	H	H	Et	F	F	F	OH	H	H
XVI-18-2	p-Cl-Ph	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-18-8	p-Cl-Ph	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 323 of 657

Table XVI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-19-1	p-Br-Ph	H	H	H	Et	F	F	F	OH	H	H
XVI-19-2	p-Br-Ph	H	H	CH_3	Et	F	F	F	OH	H	H
XVI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H	H
XVI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	F	F	F	OH	H	H
XVI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H	H
XVI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H	H
XVI-19-8	p-Br-Ph	*	H	*	Et	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7
XVI-20-1	p-I-Ph	H	H	H	Et	F	F	F	OH	H
XVI-20-2	p-I-Ph	H	H	CH_3	Et	F	F	F	OH	H
XVI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	F	F	F	OH	H
XVI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	F	F	OH	H
XVI-20-5	p-I-Ph	H	H	CH_2Ph	Et	F	F	F	OH	H
XVI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	F	F	F	OH	H
XVI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	F	F	OH	H
XVI-20-8	p-I-Ph	*	H	*	Et	F	F	F	OH	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-21-1	CH_3	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-21-2	CH_3	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-21-8	CH_3	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 324 of 657

Table XVI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-22-1	Et	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-22-2	Et	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-22-5	Et	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-22-8	Et	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-23-1	^{i}Pr	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-23-8	^{i}Pr	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-24-1	^{t}Bu	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-24-8	^{t}Bu	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 325 of 657

Table XVI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-25-1	Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-25-2	Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-25-8	Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-26-1	p-Me-Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-26-8	p-Me-Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-27-1	p-F-Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-27-8	p-F-Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 326 of 657

Table XVI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-29-1	p-Br-Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-29-8	p-Br-Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-30-1	p-I-Ph	H	H	H	^{i}Pr	F	F	F	OH	H	H
XVI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	F	F	F	OH	H	H
XVI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	F	F	OH	H	H
XVI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	F	F	F	OH	H	H
XVI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	F	F	OH	H	H
XVI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	F	F	OH	H	H
XVI-30-8	p-I-Ph	*	H	*	^{i}Pr	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 327 of 657

Table XVI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-31-1	CH_3	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-31-2	CH_3	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-31-8	CH_3	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-32-1	Et	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-32-2	Et	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-32-5	Et	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-32-8	Et	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-33-1	^{i}Pr	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-33-8	^{i}Pr	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 328 of 657

Table XVI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-34-1	^{t}Bu	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-34-8	^{t}Bu	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-35-1	Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-35-2	Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-35-8	Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-36-1	p-Me-Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-36-8	p-Me-Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 329 of 657

Table XVI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-37-1	p-F-Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-37-8	p-F-Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-39-1	p-Br-Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-39-8	p-Br-Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 330 of 657

Table XVI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-40-1	p-I-Ph	H	H	H	^{n}Bu	F	F	F	OH	H	H
XVI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	F	F	F	OH	H	H
XVI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	F	F	OH	H	H
XVI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	F	F	F	OH	H	H
XVI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	F	F	OH	H	H
XVI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	F	F	OH	H	H
XVI-40-8	p-I-Ph	*	H	*	^{n}Bu	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-41-1	CH_3	H	H	H	Bz	F	F	F	OH	H	H
XVI-41-2	CH_3	H	H	CH_3	Bz	F	F	F	OH	H	H
XVI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-41-5	CH_3	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVI-41-8	CH_3	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-42-1	Et	H	H	H	Bz	F	F	F	OH	H	H
XVI-42-2	Et	H	H	CH_3	Bz	F	F	F	OH	H	H
XVI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-42-5	Et	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVI-42-8	Et	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 331 of 657

Table XVI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-43-1	^{i}Pr	H	H	H	Bz	F	F	F	OH	H	H
XVI-43-2	^{i}Pr	H	H	CH_3	Bz	F	F	F	OH	H	H
XVI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVI-43-8	^{i}Pr	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVI-44-1	^{t}Bu	H	H	H	Bz	F	F	F	OH	H	H
XVI-44-2	^{t}Bu	H	H	CH_3	Bz	F	F	F	OH	H	H
XVI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVI-44-8	^{t}Bu	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-45-1	Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-45-2	Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-45-5	Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-45-8	Ph	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 332 of 657

Table XVI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-46-1	p-Me-Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-46-2	p-Me-Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-46-8	p-Me-Ph	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-47-1	p-F-Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-47-2	p-F-Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-47-8	p-F-Ph	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-48-1	p-Cl-Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-48-2	p-Cl-Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-48-8	p-Cl-Ph	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 333 of 657

Table XVI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-49-1	p-Br-Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-49-2	p-Br-Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-49-8	p-Br-Ph	*	H	*	Bz	F	F	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-50-1	p-I-Ph	H	H	H	Bz	F	F	F	OH	H	H
XVII-50-2	p-I-Ph	H	H	CH_3	Bz	F	F	F	OH	H	H
XVII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	F	F	OH	H	H
XVII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	F	F	F	OH	H	H
XVII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	F	F	F	OH	H	H
XVII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	F	F	OH	H	H
XVII-50-8	p-I-Ph	*	H	*	Bz	F	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 334 of 657



XVII

Table XVII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-1-1	CH_3	H	H	H	CH_3	F	H	F	OH	H	H
XVII-1-2	CH_3	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-1-5	CH_3	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-1-8	CH_3	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-2-1	Et	H	H	H	CH_3	F	H	F	OH	H	H
XVII-2-2	Et	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-2-5	Et	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-2-8	Et	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 335 of 657

Table XVII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-3-1	^{i}Pr	H	H	H	CH_3	F	H	F	OH	H	H
XVII-3-2	^{i}Pr	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-3-8	^{i}Pr	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-4-1	^{t}Bu	H	H	H	CH_3	F	H	F	OH	H	H
XVII-4-2	^{t}Bu	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-4-8	^{t}Bu	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-5-1	Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-5-2	Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-5-5	Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-5-8	Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 336 of 657

Table XVII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-6-1	p-Me-Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-6-2	p-Me-Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-6-8	p-Me-Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-7-1	p-F-Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-7-2	p-F-Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-7-20	p-F-Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-8-1	p-Cl-Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-8-8	p-Cl-Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 337 of 657

Table XVII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-9-1	p-Br-Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-9-2	p-Br-Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-9-20	p-Br-Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-10-1	p-I-Ph	H	H	H	CH_3	F	H	F	OH	H	H
XVII-10-2	p-I-Ph	H	H	CH_3	CH_3	F	H	F	OH	H	H
XVII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	F	H	F	OH	H	H
XVII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	F	H	F	OH	H	H
XVII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	F	H	F	OH	H	H
XVII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	F	H	F	OH	H	H
XVII-10-8	p-I-Ph	*	H	*	CH_3	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-11-1	CH_3	H	H	H	Et	F	H	F	OH	H	H
XVII-11-2	CH_3	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-11-5	CH_3	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-11-8	CH_3	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 338 of 657

Table XVII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-12-1	Et	H	H	H	Et	F	H	F	OH	H	H
XVII-12-2	Et	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-12-3	Et	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-12-5	Et	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-12-6	Et	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-12-8	Et	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-13-1	^{i}Pr	H	H	H	Et	F	H	F	OH	H	H
XVII-13-2	^{i}Pr	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-13-5	^{i}Pr	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-13-8	^{i}Pr	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-14-1	^{t}Bu	H	H	H	Et	F	H	F	OH	H	H
XVII-14-2	^{t}Bu	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-14-5	^{t}Bu	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-14-8	^{t}Bu	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 339 of 657

Table XVII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-15-1	Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-15-2	Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-15-5	Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-15-8	Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-16-1	p-Me-Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-16-2	p-Me-Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-16-8	p-Me-Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-17-1	p-F-Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-17-2	p-F-Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-17-5	p-F-Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-17-8	p-F-Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 340 of 657

Table XVII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-18-1	p-Cl-Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-18-2	p-Cl-Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-18-8	p-Cl-Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-19-1	p-Br-Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-19-2	p-Br-Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-19-8	p-Br-Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-20-1	p-I-Ph	H	H	H	Et	F	H	F	OH	H	H
XVII-20-2	p-I-Ph	H	H	CH_3	Et	F	H	F	OH	H	H
XVII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	F	H	F	OH	H	H
XVII-20-5	p-I-Ph	H	H	CH_2Ph	Et	F	H	F	OH	H	H
XVII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	F	H	F	OH	H	H
XVII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	F	H	F	OH	H	H
XVII-20-8	p-I-Ph	*	H	*	Et	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 341 of 657

Table XVII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-21-1	CH_3	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-21-2	CH_3	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-21-8	CH_3	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-22-1	Et	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-22-2	Et	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-22-5	Et	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-22-8	Et	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-23-1	^{i}Pr	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-23-8	^{i}Pr	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 342 of 657

Table XVII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-24-1	^{t}Bu	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-24-8	^{t}Bu	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-25-1	Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-25-2	Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-25-8	Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-26-1	p-Me-Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-26-8	p-Me-Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 343 of 657

Table XVII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-27-1	p-F-Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-27-8	p-F-Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-29-1	p-Br-Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-29-8	p-Br-Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 344 of 657

Table XVII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-30-1	p-I-Ph	H	H	H	^{i}Pr	F	H	F	OH	H	H
XVII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	F	H	F	OH	H	H
XVII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	F	H	F	OH	H	H
XVII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	F	H	F	OH	H	H
XVII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	F	H	F	OH	H	H
XVII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	F	H	F	OH	H	H
XVII-30-8	p-I-Ph	*	H	*	^{i}Pr	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-31-1	CH_3	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-31-2	CH_3	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-31-8	CH_3	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-32-1	Et	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-32-2	Et	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-32-5	Et	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-32-8	Et	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 345 of 657

Table XVII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-33-1	^{i}Pr	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-33-8	^{i}Pr	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-34-1	^{t}Bu	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-34-8	^{t}Bu	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-35-1	Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-35-2	Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-35-8	Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 346 of 657

Table XVII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-36-1	p-Me-Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-36-8	p-Me-Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-37-1	p-F-Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-37-8	p-F-Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 347 of 657

Table XVII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-39-1	p-Br-Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-39-8	p-Br-Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-40-1	p-I-Ph	H	H	H	^{n}Bu	F	H	F	OH	H	H
XVII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	F	H	F	OH	H	H
XVII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	F	H	F	OH	H	H
XVII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	F	H	F	OH	H	H
XVII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	F	H	F	OH	H	H
XVII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	F	H	F	OH	H	H
XVII-40-8	p-I-Ph	*	H	*	^{n}Bu	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-41-1	CH_3	H	H	H	Bz	F	H	F	OH	H	H
XVII-41-2	CH_3	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-41-5	CH_3	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-41-8	CH_3	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 348 of 657

Table XVII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-42-1	Et	H	H	H	Bz	F	H	F	OH	H	H
XVII-42-2	Et	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-42-5	Et	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-42-8	Et	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-43-1	^{i}Pr	H	H	H	Bz	F	H	F	OH	H	H
XVII-43-2	^{i}Pr	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-43-8	^{i}Pr	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-44-1	^{t}Bu	H	H	H	Bz	F	H	F	OH	H	H
XVII-44-2	^{t}Bu	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-44-8	^{t}Bu	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 349 of 657

Table XVII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-45-1	Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-45-2	Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-45-5	Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-45-8	Ph	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-46-1	p-Me-Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-46-2	p-Me-Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-46-8	p-Me-Ph	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-47-1	p-F-Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-47-2	p-F-Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-47-8	p-F-Ph	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 350 of 657

Table XVII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-48-1	p-Cl-Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-48-2	p-Cl-Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-48-8	p-Cl-Ph	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-49-1	p-Br-Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-49-2	p-Br-Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-49-8	p-Br-Ph	*	H	*	Bz	F	H	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVII-50-1	p-I-Ph	H	H	H	Bz	F	H	F	OH	H	H
XVII-50-2	p-I-Ph	H	H	CH_3	Bz	F	H	F	OH	H	H
XVII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	F	H	F	OH	H	H
XVII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	F	H	F	OH	H	H
XVII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	F	H	F	OH	H	H
XVII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	F	H	F	OH	H	H
XVII-50-8	p-I-Ph	*	H	*	Bz	F	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 351 of 657



XVIII

Table XVIII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-1-1	CH_3	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-2	CH_3	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-5	CH_3	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-1-8	CH_3	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-2-1	Et	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-2	Et	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-5	Et	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 352 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-2-8	Et	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-3-1	^{i}Pr	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-2	^{i}Pr	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-3-8	^{i}Pr	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-4-1	^{t}Bu	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-2	^{t}Bu	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-4-8	^{t}Bu	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-5-1	Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-2	Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-5	Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 353 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-5-8	Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-6-1	p-Me-Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-2	p-Me-Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-6-8	p-Me-Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-7-1	p-F-Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-2	p-F-Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-7-20	p-F-Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-8-1	p-Cl-Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 354 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-8-8	p-Cl-Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-9-1	p-Br-Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-2	p-Br-Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-9-20	p-Br-Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-10-1	p-I-Ph	H	H	H	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-2	p-I-Ph	H	H	CH_3	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	H	OH	H	H
XVIII-10-8	p-I-Ph	*	H	*	CH_3	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-11-1	CH_3	H	H	H	Et	OCH_3	F	H	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 355 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-11-2	CH_3	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-11-5	CH_3	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-11-8	CH_3	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-12-1	Et	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-12-2	Et	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-12-3	Et	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-12-5	Et	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-12-6	Et	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-12-8	Et	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-13-1	^{i}Pr	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-13-2	^{i}Pr	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-13-5	^{i}Pr	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-13-8	^{i}Pr	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-14.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 356 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-14-1	^{t}Bu	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-14-2	^{t}Bu	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-14-5	^{t}Bu	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-14-8	^{t}Bu	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-15-1	Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-15-2	Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-15-5	Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-15-8	Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-16-1	p-Me-Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-16-2	p-Me-Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-16-8	p-Me-Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 357 of 657

Table XVIII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-17-1	p-F-Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-17-2	p-F-Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-17-5	p-F-Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-17-8	p-F-Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-18-1	p-Cl-Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-18-2	p-Cl-Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-18-8	p-Cl-Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-19-1	p-Br-Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-19-2	p-Br-Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-19-8	p-Br-Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 358 of 657

Table XVIII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-20-1	p-I-Ph	H	H	H	Et	OCH_3	F	H	OH	H	H
XVIII-20-2	p-I-Ph	H	H	CH_3	Et	OCH_3	F	H	OH	H	H
XVIII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	H	OH	H	H
XVIII-20-5	p-I-Ph	H	H	CH_2Ph	Et	OCH_3	F	H	OH	H	H
XVIII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	H	OH	H	H
XVIII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	H	OH	H	H
XVIII-20-8	p-I-Ph	*	H	*	Et	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-21-1	CH_3	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-2	CH_3	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-21-8	CH_3	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-22-1	Et	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-2	Et	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-5	Et	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-22-8	Et	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 359 of 657

Table XVIII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-23-1	^{i}Pr	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-23-8	^{i}Pr	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-24-1	^{t}Bu	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-24-8	^{t}Bu	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-25-1	Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-2	Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-25-8	Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 360 of 657

Table XVIII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-26-1	p-Me-Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-26-8	p-Me-Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-27-1	p-F-Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-27-8	p-F-Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 361 of 657

Table XVIII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-29-1	p-Br-Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-29-8	p-Br-Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-30-1	p-I-Ph	H	H	H	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	H	OH	H	H
XVIII-30-8	p-I-Ph	*	H	*	^{i}Pr	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-31-1	CH_3	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-2	CH_3	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-31-8	CH_3	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 362 of 657

Table XVIII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-32-1	Et	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-2	Et	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-5	Et	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-32-8	Et	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-33-1	^{i}Pr	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-33-8	^{i}Pr	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-34-1	^{t}Bu	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-34-8	^{t}Bu	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 363 of 657

Table XVIII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-35-1	Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-2	Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-35-8	Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-36-1	p-Me-Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-36-8	p-Me-Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-37-1	p-F-Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-37-8	p-F-Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 364 of 657

Table XVIII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-39-1	p-Br-Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-39-8	p-Br-Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-40-1	p-I-Ph	H	H	H	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	H	OH	H	H
XVIII-40-8	p-I-Ph	*	H	*	^{n}Bu	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 365 of 657

Table XVIII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-41-1	CH_3	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-41-2	CH_3	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-41-5	CH_3	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-41-8	CH_3	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-42-1	Et	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-42-2	Et	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-42-5	Et	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-42-8	Et	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-43-1	^{i}Pr	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-43-2	^{i}Pr	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-43-8	^{i}Pr	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 366 of 657

Table XVIII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-44-1	^{t}Bu	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-44-2	^{t}Bu	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-44-8	^{t}Bu	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-45-1	Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-45-2	Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-45-5	Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-45-8	Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-46-1	p-Me-Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-46-2	p-Me-Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-46-8	p-Me-Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 367 of 657

Table XVIII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-47-1	p-F-Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-47-2	p-F-Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-47-8	p-F-Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-48-1	p-Cl-Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-48-2	p-Cl-Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-48-8	p-Cl-Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XVIII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-49-1	p-Br-Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-49-2	p-Br-Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-49-8	p-Br-Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 368 of 657

Table XVIII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XVIII-50-1	p-I-Ph	H	H	H	Bz	OCH_3	F	H	OH	H	H
XVIII-50-2	p-I-Ph	H	H	CH_3	Bz	OCH_3	F	H	OH	H	H
XVIII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	H	OH	H	H
XVIII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	OCH_3	F	H	OH	H	H
XVIII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	H	OH	H	H
XVIII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	H	OH	H	H
XVIII-50-8	p-I-Ph	*	H	*	Bz	OCH_3	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



XIX

Table XIX-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-1-1	CH_3	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-1-2	CH_3	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-1-5	CH_3	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 369 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-1-8	CH_3	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-2-1	Et	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-2-2	Et	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-2-5	Et	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-2-8	Et	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-3-1	^{i}Pr	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-3-2	^{i}Pr	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-3-8	^{i}Pr	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-4-1	^{t}Bu	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-4-2	^{t}Bu	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 370 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-4-8	^{t}Bu	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-5-1	Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-5-2	Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-5-5	Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-5-8	Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-6-1	p-Me-Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-6-2	p-Me-Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-6-8	p-Me-Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-7-1	p-F-Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-7-2	p-F-Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 371 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-7-20	p-F-Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-8-1	p-Cl-Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-8-2	p-Cl-Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-8-8	p-Cl-Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-9-1	p-Br-Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H
XIX-9-2	p-Br-Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-9-20	p-Br-Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-10-1	p-I-Ph	H	H	H	CH_3	OCH_3	F	F	OH	H	H

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 372 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-10-2	p-I-Ph	H	H	CH_3	CH_3	OCH_3	F	F	OH	H	H
XIX-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	F	F	OH	H	H
XIX-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	OCH_3	F	F	OH	H	H
XIX-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	F	F	OH	H	H
XIX-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	F	F	OH	H	H
XIX-10-8	p-I-Ph	*	H	*	CH_3	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-11-1	CH_3	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-11-2	CH_3	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-11-5	CH_3	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-11-8	CH_3	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-12-1	Et	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-12-2	Et	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-12-3	Et	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-12-5	Et	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-12-6	Et	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-12-8	Et	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-13.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 373 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-13-1	^{i}Pr	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-13-2	^{i}Pr	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-13-5	^{i}Pr	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-13-8	^{i}Pr	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-14-1	^{t}Bu	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-14-2	^{t}Bu	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-14-5	^{t}Bu	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-14-8	^{t}Bu	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-15-1	Ph	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-15-2	Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-15-3	Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-15-5	Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-15-6	Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-15-8	Ph	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 374 of 657

Table XIX-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-16-1	p-Me-Ph	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-16-2	p-Me-Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-16-5	p-Me-Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-16-8	p-Me-Ph	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-17-1	p-F-Ph	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-17-2	p-F-Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-17-5	p-F-Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-17-8	p-F-Ph	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-18-1	p-Cl-Ph	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-18-2	p-Cl-Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-18-8	p-Cl-Ph	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 375 of 657

Table XIX-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-19-1	p-Br-Ph	H	H	H	Et	OCH_3	F	F	OH	H	H
XIX-19-2	p-Br-Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H	H
XIX-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H	H
XIX-19-5	p-Br-Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H	H
XIX-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H	H
XIX-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H	H
XIX-19-8	p-Br-Ph	*	H	*	Et	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7
XIX-20-1	p-I-Ph	H	H	H	Et	OCH_3	F	F	OH	H
XIX-20-2	p-I-Ph	H	H	CH_3	Et	OCH_3	F	F	OH	H
XIX-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	F	F	OH	H
XIX-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	F	F	OH	H
XIX-20-5	p-I-Ph	H	H	CH_2Ph	Et	OCH_3	F	F	OH	H
XIX-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	F	F	OH	H
XIX-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	F	F	OH	H
XIX-20-8	p-I-Ph	*	H	*	Et	OCH_3	F	F	OH	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-21-1	CH_3	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-2	CH_3	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-21-8	CH_3	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 376 of 657

Table XIX-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-22-1	Et	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-2	Et	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-5	Et	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-22-8	Et	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-23-1	^{i}Pr	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-23-8	^{i}Pr	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-24-1	^{t}Bu	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-24-8	^{t}Bu	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 377 of 657

Table XIX-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-25-1	Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-2	Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-5	Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-25-8	Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-26-1	p-Me-Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-26-8	p-Me-Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-27-1	p-F-Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-27-8	p-F-Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 378 of 657

Table XIX-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-28-1	p-Cl-Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-28-8	p-Cl-Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-29-1	p-Br-Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-29-8	p-Br-Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-30-1	p-I-Ph	H	H	H	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	F	F	OH	H	H
XIX-30-8	p-I-Ph	*	H	*	^{i}Pr	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 379 of 657

Table XIX-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-31-1	CH_3	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-2	CH_3	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-31-8	CH_3	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-32-1	Et	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-2	Et	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-5	Et	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-32-8	Et	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-33-1	^{i}Pr	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-33-8	^{i}Pr	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 380 of 657

Table XIX-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-34-1	^{t}Bu	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-34-8	^{t}Bu	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-35-1	Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-2	Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-5	Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-35-8	Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-36-1	p-Me-Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-36-8	p-Me-Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 381 of 657

Table XIX-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-37-1	p-F-Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-37-8	p-F-Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-38-1	p-Cl-Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-38-8	p-Cl-Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-39-1	p-Br-Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-39-8	p-Br-Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 382 of 657

Table XIX-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-40-1	p-I-Ph	H	H	H	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	F	F	OH	H	H
XIX-40-8	p-I-Ph	*	H	*	^{n}Bu	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-41-1	CH_3	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-41-2	CH_3	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-41-5	CH_3	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-41-8	CH_3	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-42-1	Et	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-42-2	Et	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-42-3	Et	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-42-5	Et	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-42-6	Et	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-42-8	Et	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 383 of 657

Table XIX-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-43-1	^{i}Pr	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-43-2	^{i}Pr	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-43-5	^{i}Pr	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-43-8	^{i}Pr	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-44-1	^{t}Bu	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-44-2	^{t}Bu	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-44-5	^{t}Bu	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-44-8	^{t}Bu	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-45-1	Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-45-2	Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-45-5	Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-45-8	Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 384 of 657

Table XIX-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-46-1	p-Me-Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-46-2	p-Me-Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-46-8	p-Me-Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-47-1	p-F-Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-47-2	p-F-Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-47-5	p-F-Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-47-8	p-F-Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-48-1	p-Cl-Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-48-2	p-Cl-Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-48-8	p-Cl-Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 385 of 657

Table XIX-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-49-1	p-Br-Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-49-2	p-Br-Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-49-8	p-Br-Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XIX-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XIX-50-1	p-I-Ph	H	H	H	Bz	OCH_3	F	F	OH	H	H
XIX-50-2	p-I-Ph	H	H	CH_3	Bz	OCH_3	F	F	OH	H	H
XIX-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	F	F	OH	H	H
XIX-50-5	p-I-Ph	H	H	CH_2Ph	Bz	OCH_3	F	F	OH	H	H
XIX-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	F	F	OH	H	H
XIX-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	F	F	OH	H	H
XIX-50-8	p-I-Ph	*	H	*	Bz	OCH_3	F	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 386 of 657



XX

Table XX-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-1-1	CH_3	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-1-2	CH_3	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-1-5	CH_3	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-1-8	CH_3	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-2-1	Et	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-2-2	Et	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-2-5	Et	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-2-8	Et	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 387 of 657

Table XX-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-3-1	^{i}Pr	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-3-2	^{i}Pr	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-3-8	^{i}Pr	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-4-1	^{t}Bu	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-4-2	^{t}Bu	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-4-8	^{t}Bu	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-5-1	Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-5-2	Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-5-5	Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-5-8	Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 388 of 657

Table XX-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-6-1	p-Me-Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-6-2	p-Me-Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-6-8	p-Me-Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-7-1	p-F-Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-7-2	p-F-Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-7-20	p-F-Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-8-1	p-Cl-Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-8-2	p-Cl-Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-8-8	p-Cl-Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 389 of 657

Table XX-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-9-1	p-Br-Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-9-2	p-Br-Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-9-20	p-Br-Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-10-1	p-I-Ph	H	H	H	CH_3	OCH_3	H	F	OH	H	H
XX-10-2	p-I-Ph	H	H	CH_3	CH_3	OCH_3	H	F	OH	H	H
XX-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	OCH_3	H	F	OH	H	H
XX-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	OCH_3	H	F	OH	H	H
XX-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	OCH_3	H	F	OH	H	H
XX-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	OCH_3	H	F	OH	H	H
XX-10-8	p-I-Ph	*	H	*	CH_3	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-11-1	CH_3	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-11-2	CH_3	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-11-5	CH_3	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-11-8	CH_3	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 390 of 657

Table XX-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-12-1	Et	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-12-2	Et	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-12-3	Et	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-12-5	Et	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-12-6	Et	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-12-8	Et	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-13-1	^{i}Pr	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-13-2	^{i}Pr	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-13-5	^{i}Pr	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-13-8	^{i}Pr	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-14-1	^{t}Bu	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-14-2	^{t}Bu	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-14-5	^{t}Bu	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-14-8	^{t}Bu	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 391 of 657

Table XX-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-15-1	Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-15-2	Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-15-3	Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-15-5	Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-15-6	Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-15-8	Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-16-1	p-Me-Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-16-2	p-Me-Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-16-5	p-Me-Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-16-8	p-Me-Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-17-1	p-F-Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-17-2	p-F-Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-17-5	p-F-Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-17-8	p-F-Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 392 of 657

Table XX-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-18-1	p-Cl-Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-18-2	p-Cl-Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-18-8	p-Cl-Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-19-1	p-Br-Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-19-2	p-Br-Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-19-5	p-Br-Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-19-8	p-Br-Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-20-1	p-I-Ph	H	H	H	Et	OCH_3	H	F	OH	H	H
XX-20-2	p-I-Ph	H	H	CH_3	Et	OCH_3	H	F	OH	H	H
XX-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	OCH_3	H	F	OH	H	H
XX-20-5	p-I-Ph	H	H	CH_2Ph	Et	OCH_3	H	F	OH	H	H
XX-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	OCH_3	H	F	OH	H	H
XX-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	OCH_3	H	F	OH	H	H
XX-20-8	p-I-Ph	*	H	*	Et	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 393 of 657

Table XX-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-21-1	CH_3	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-2	CH_3	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-21-8	CH_3	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-22-1	Et	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-2	Et	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-5	Et	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-22-8	Et	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-23-1	^{i}Pr	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-23-8	^{i}Pr	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 394 of 657

Table XX-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-24-1	^{t}Bu	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-24-8	^{t}Bu	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-25-1	Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-2	Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-5	Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-25-8	Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-26-1	p-Me-Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-26-8	p-Me-Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 395 of 657

Table XX-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-27-1	p-F-Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-27-8	p-F-Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-28-1	p-Cl-Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-28-8	p-Cl-Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-29-1	p-Br-Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-29-8	p-Br-Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 396 of 657

Table XX-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-30-1	p-I-Ph	H	H	H	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	OCH_3	H	F	OH	H	H
XX-30-8	p-I-Ph	*	H	*	^{i}Pr	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-31-1	CH_3	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-2	CH_3	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-31-8	CH_3	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-32-1	Et	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-2	Et	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-5	Et	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-32-8	Et	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 397 of 657

Table XX-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-33-1	^{i}Pr	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-33-8	^{i}Pr	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-34-1	^{t}Bu	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-34-8	^{t}Bu	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-35-1	Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-2	Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-5	Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-35-8	Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 398 of 657

Table XX-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-36-1	p-Me-Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-36-8	p-Me-Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-37-1	p-F-Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-37-8	p-F-Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-38-1	p-Cl-Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-38-8	p-Cl-Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 399 of 657

Table XX-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-39-1	p-Br-Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-39-8	p-Br-Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-40-1	p-I-Ph	H	H	H	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	OCH_3	H	F	OH	H	H
XX-40-8	p-I-Ph	*	H	*	^{n}Bu	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-41-1	CH_3	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-41-2	CH_3	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-41-5	CH_3	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-41-8	CH_3	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 400 of 657

Table XX-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-42-1	Et	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-42-2	Et	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-42-3	Et	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-42-5	Et	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-42-6	Et	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-42-8	Et	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-43-1	^{i}Pr	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-43-2	^{i}Pr	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-43-5	^{i}Pr	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-43-8	^{i}Pr	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-44-1	^{t}Bu	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-44-2	^{t}Bu	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-44-5	^{t}Bu	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-44-8	^{t}Bu	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 401 of 657

Table XX-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-45-1	Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-45-2	Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-45-5	Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-45-8	Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-46-1	p-Me-Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-46-2	p-Me-Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-46-8	p-Me-Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-47-1	p-F-Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-47-2	p-F-Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-47-5	p-F-Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-47-8	p-F-Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 402 of 657

Table XX-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-48-1	p-Cl-Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-48-2	p-Cl-Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-48-8	p-Cl-Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-49-1	p-Br-Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-49-2	p-Br-Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-49-8	p-Br-Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XX-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
XX-50-1	p-I-Ph	H	H	H	Bz	OCH_3	H	F	OH	H	H
XX-50-2	p-I-Ph	H	H	CH_3	Bz	OCH_3	H	F	OH	H	H
XX-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	OCH_3	H	F	OH	H	H
XX-50-5	p-I-Ph	H	H	CH_2Ph	Bz	OCH_3	H	F	OH	H	H
XX-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	OCH_3	H	F	OH	H	H
XX-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	OCH_3	H	F	OH	H	H
XX-50-8	p-I-Ph	*	H	*	Bz	OCH_3	H	F	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 403 of 657



XXI

Table XXI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH
XXI-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH
XXI-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 404 of 657

Table XXI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH
XXI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH
XXI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 405 of 657

Table XXI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 406 of 657

Table XXI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXI-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH
XXI-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH
XXI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 407 of 657

Table XXI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-12-1	Et	H	H	H	Et	H	CH_3	F	OH
XXI-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH
XXI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-12-8	Et	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH
XXI-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH
XXI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-14-1	^{t}Bu	H	H	H	Et	H	CH_3	F	OH
XXI-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH
XXI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-14-8	^{t}Bu	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 408 of 657

Table XXI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-15-1	Ph	H	H	H	Et	H	CH_3	F	OH
XXI-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-15-8	Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH
XXI-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH
XXI-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 409 of 657

Table XXI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH
XXI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH
XXI-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH
XXI-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXI-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 410 of 657

Table XXI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 411 of 657

Table XXI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 412 of 657

Table XXI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 413 of 657

Table XXI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 414 of 657

Table XXI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 415 of 657

Table XXI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 416 of 657

Table XXI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH
XXI-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 417 of 657

Table XXI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-42-1	Et	H	H	H	Bz	H	CH_3	F	OH
XXI-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-42-8	Et	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH
XXI-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH
XXI-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 418 of 657

Table XXI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 419 of 657

Table XXI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXI-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH
XXI-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXI-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 420 of 657



XXII

Table XXII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-1-1	CH_3	H	H	H	CH_3	H	F	H	OH
XXII-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH
XXII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-1-8	CH_3	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-2-1	Et	H	H	H	CH_3	H	F	H	OH
XXII-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH
XXII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-2-8	Et	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 421 of 657

Table XXII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH
XXII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH
XXII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH
XXII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH
XXII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-5-1	Ph	H	H	H	CH_3	H	F	H	OH
XXII-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-5-8	Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 422 of 657

Table XXII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH
XXII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH
XXII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH
XXII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 423 of 657

Table XXII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH
XXII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH
XXII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXII-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-11-1	CH_3	H	H	H	Et	H	F	H	OH
XXII-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH
XXII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH
XXII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-11-8	CH_3	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 424 of 657

Table XXII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-12-1	Et	H	H	H	Et	H	F	H	OH
XXII-12-2	Et	H	H	CH_3	Et	H	F	H	OH
XXII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH
XXII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-12-8	Et	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH
XXII-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH
XXII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH
XXII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH
XXII-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH
XXII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH
XXII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 425 of 657

Table XXII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-15-1	Ph	H	H	H	Et	H	F	H	OH
XXII-15-2	Ph	H	H	CH_3	Et	H	F	H	OH
XXII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-15-8	Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH
XXII-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH
XXII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH
XXII-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH
XXII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 426 of 657

Table XXII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH
XXII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH
XXII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH
XXII-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH
XXII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH
XXII-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH
XXII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXII-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 427 of 657

Table XXII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH
XXII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH
XXII-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH
XXII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 428 of 657

Table XXII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH
XXII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 429 of 657

Table XXII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 430 of 657

Table XXII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH
XXII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH
XXII-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 431 of 657

Table XXII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH
XXII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH
XXII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 432 of 657

Table XXII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 433 of 657

Table XXII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-41-1	CH_3	H	H	H	Bz	H	F	H	OH
XXII-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH
XXII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-41-8	CH_3	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 434 of 657

Table XXII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-42-1	Et	H	H	H	Bz	H	F	H	OH
XXII-42-2	Et	H	H	CH_3	Bz	H	F	H	OH
XXII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-42-8	Et	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH
XXII-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH
XXII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH
XXII-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH
XXII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 435 of 657

Table XXII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-45-1	Ph	H	H	H	Bz	H	F	H	OH
XXII-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-45-8	Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH
XXII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH
XXII-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 436 of 657

Table XXII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH
XXII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH
XXII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXII-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH
XXII-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH
XXII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXII-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 437 of 657



XXIII

Table XXIII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-1-1	CH_3	H	H	H	CH_3	H	F	F	OH
XXIII-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH
XXIII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-1-8	CH_3	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-2-1	Et	H	H	H	CH_3	H	F	F	OH
XXIII-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH
XXIII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-2-8	Et	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 438 of 657

Table XXIII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH
XXIII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH
XXIII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-4-1	^{t}Bu	H	H	H	CH_3	H	F	F	OH
XXIII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	F	OH
XXIII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-4-8	^{t}Bu	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-5-1	Ph	H	H	H	CH_3	H	F	F	OH
XXIII-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-5-8	Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 439 of 657

Table XXIII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH
XXIII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH
XXIII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH
XXIII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 440 of 657

Table XXIII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH
XXIII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH
XXIII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXIII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXIII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXIII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXIII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXIII-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-11-1	CH_3	H	H	H	Et	H	F	F	OH
XXIII-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH
XXIII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-11-8	CH_3	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 441 of 657

Table XXIII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-12-1	Et	H	H	H	Et	H	F	F	OH
XXIII-12-2	Et	H	H	CH_3	Et	H	F	F	OH
XXIII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-12-8	Et	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH
XXIII-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH
XXIII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH
XXIII-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH
XXIII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 442 of 657

Table XXIII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-15-1	Ph	H	H	H	Et	H	F	F	OH
XXIII-15-2	Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-15-8	Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH
XXIII-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH
XXIII-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 443 of 657

Table XXIII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH
XXIII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH
XXIII-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH
XXIII-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH
XXIII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXIII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXIII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXIII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXIII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXIII-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 444 of 657

Table XXIII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH
XXIII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH
XXIII-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH
XXIII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 445 of 657

Table XXIII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH
XXIII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 446 of 657

Table XXIII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 447 of 657

Table XXIII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXIII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXIII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXIII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXIII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXIII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXIII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH
XXIII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH
XXIII-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 448 of 657

Table XXIII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH
XXIII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH
XXIII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 449 of 657

Table XXIII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 450 of 657

Table XXIII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXIII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXIII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXIII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXIII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXIII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXIII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-41-1	CH_3	H	H	H	Bz	H	F	F	OH
XXIII-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH
XXIII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-41-8	CH_3	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 451 of 657

Table XXIII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-42-1	Et	H	H	H	Bz	H	F	F	OH
XXIII-42-2	Et	H	H	CH_3	Bz	H	F	F	OH
XXIII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-42-8	Et	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH
XXIII-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH
XXIII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH
XXIII-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH
XXIII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 452 of 657

Table XXIII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-45-1	Ph	H	H	H	Bz	H	F	F	OH
XXIII-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-45-8	Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH
XXIII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH
XXIII-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 453 of 657

Table XXIII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH
XXIII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH
XXIII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIII-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH
XXIII-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH
XXIII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXIII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXIII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXIII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXIII-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 454 of 657



XXIV

Table XXIV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-1-1	CH_3	H	H	H	CH_3	H	H	F	OH
XXIV-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH
XXIV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-1-8	CH_3	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-2-1	Et	H	H	H	CH_3	H	H	F	OH
XXIV-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH
XXIV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-2-8	Et	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 455 of 657

Table XXIV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH
XXIV-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH
XXIV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH
XXIV-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH
XXIV-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-5-1	Ph	H	H	H	CH_3	H	H	F	OH
XXIV-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-5-8	Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 456 of 657

Table XXIV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH
XXIV-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH
XXIV-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH
XXIV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 457 of 657

Table XXIV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH
XXIV-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH
XXIV-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXIV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXIV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXIV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXIV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXIV-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-11-1	CH_3	H	H	H	Et	H	H	F	OH
XXIV-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH
XXIV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-11-8	CH_3	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 458 of 657

Table XXIV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-12-1	Et	H	H	H	Et	H	H	F	OH
XXIV-12-2	Et	H	H	CH_3	Et	H	H	F	OH
XXIV-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-12-8	Et	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH
XXIV-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH
XXIV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH
XXIV-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH
XXIV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 459 of 657

Table XXIV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-15-1	Ph	H	H	H	Et	H	H	F	OH
XXIV-15-2	Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-15-8	Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH
XXIV-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH
XXIV-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 460 of 657

Table XXIV-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH
XXIV-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH
XXIV-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH
XXIV-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH
XXIV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXIV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXIV-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXIV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXIV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXIV-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 461 of 657

Table XXIV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH
XXIV-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH
XXIV-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH
XXIV-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 462 of 657

Table XXIV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH
XXIV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 463 of 657

Table XXIV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 464 of 657

Table XXIV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXIV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXIV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXIV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXIV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXIV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXIV-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH
XXIV-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH
XXIV-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 465 of 657

Table XXIV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH
XXIV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH
XXIV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 466 of 657

Table XXIV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 467 of 657

Table XXIV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXIV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXIV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXIV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXIV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXIV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXIV-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-41-1	CH_3	H	H	H	Bz	H	H	F	OH
XXIV-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH
XXIV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-41-8	CH_3	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 468 of 657

Table XXIV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-42-1	Et	H	H	H	Bz	H	H	F	OH
XXIV-42-2	Et	H	H	CH_3	Bz	H	H	F	OH
XXIV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-42-8	Et	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH
XXIV-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH
XXIV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH
XXIV-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH
XXIV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 469 of 657

Table XXIV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-45-1	Ph	H	H	H	Bz	H	H	F	OH
XXIV-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-45-8	Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH
XXIV-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH
XXIV-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 470 of 657

Table XXIV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH
XXIV-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH
XXIV-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIV-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH
XXIV-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH
XXIV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXIV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXIV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXIV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXIV-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 471 of 657



XXV

Table XXV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH
XXV-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH
XXV-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 472 of 657

Table XXV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH
XXV-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH
XXV-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 473 of 657

Table XXV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 474 of 657

Table XXV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXV-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXV-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH
XXV-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH
XXV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 475 of 657

Table XXV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-12-1	Et	H	H	H	Et	H	CH_3	F	OH
XXV-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH
XXV-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-12-8	Et	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH
XXV-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH
XXV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-14-1	^{t}Bu	H	H	H	Et	H	CH_3	F	OH
XXV-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH
XXV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-14-8	^{t}Bu	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 476 of 657

Table XXV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-15-1	Ph	H	H	H	Et	H	CH_3	F	OH
XXV-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-15-8	Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH
XXV-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH
XXV-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 477 of 657

Table XXV-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH
XXV-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH
XXV-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH
XXV-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXV-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXV-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 478 of 657

Table XXV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 479 of 657

Table XXV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 480 of 657

Table XXV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 481 of 657

Table XXV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXV-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 482 of 657

Table XXV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 483 of 657

Table XXV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 484 of 657

Table XXV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXV-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH
XXV-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 485 of 657

Table XXV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-42-1	Et	H	H	H	Bz	H	CH_3	F	OH
XXV-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-42-8	Et	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH
XXV-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH
XXV-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 486 of 657

Table XXV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 487 of 657

Table XXV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXV-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH
XXV-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXV-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 488 of 657



XXVI

Table XXVI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-1-1	CH_3	H	H	H	CH_3	H	F	H	OH
XXVI-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH
XXVI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-1-8	CH_3	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-2-1	Et	H	H	H	CH_3	H	F	H	OH
XXVI-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH
XXVI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-2-8	Et	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 489 of 657

Table XXVI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH
XXVI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH
XXVI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH
XXVI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH
XXVI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-5-1	Ph	H	H	H	CH_3	H	F	H	OH
XXVI-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-5-8	Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 490 of 657

Table XXVI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH
XXVI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH
XXVI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH
XXVI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 491 of 657

Table XXVI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH
XXVI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH
XXVI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXVI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXVI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXVI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXVI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXVI-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-11-1	CH_3	H	H	H	Et	H	F	H	OH
XXVI-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH
XXVI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-11-8	CH_3	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 492 of 657

Table XXVI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-12-1	Et	H	H	H	Et	H	F	H	OH
XXVI-12-2	Et	H	H	CH_3	Et	H	F	H	OH
XXVI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-12-8	Et	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH
XXVI-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH
XXVI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH
XXVI-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH
XXVI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 493 of 657

Table XXVI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-15-1	Ph	H	H	H	Et	H	F	H	OH
XXVI-15-2	Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-15-8	Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH
XXVI-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH
XXVI-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 494 of 657

Table XXVI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH
XXVI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH
XXVI-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH
XXVI-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH
XXVI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXVI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXVI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXVI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXVI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXVI-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 495 of 657

Table XXVI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH
XXVI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH
XXVI-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH
XXVI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 496 of 657

Table XXVI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH
XXVI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 497 of 657

Table XXVI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 498 of 657

Table XXVI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXVI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXVI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXVI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXVI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXVI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXVI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH
XXVI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH
XXVI-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 499 of 657

Table XXVI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH
XXVI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH
XXVI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 500 of 657

Table XXVI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 501 of 657

Table XXVI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXVI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXVI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXVI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXVI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXVI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXVI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-41-1	CH_3	H	H	H	Bz	H	F	H	OH
XXVI-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH
XXVI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-41-8	CH_3	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 502 of 657

Table XXVI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-42-1	Et	H	H	H	Bz	H	F	H	OH
XXVI-42-2	Et	H	H	CH_3	Bz	H	F	H	OH
XXVI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-42-8	Et	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH
XXVI-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH
XXVI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH
XXVI-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH
XXVI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 503 of 657

Table XXVI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-45-1	Ph	H	H	H	Bz	H	F	H	OH
XXVI-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-45-8	Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH
XXVI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH
XXVI-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 504 of 657

Table XXVI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH
XXVI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH
XXVI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVI-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH
XXVI-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH
XXVI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXVI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXVI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXVI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXVI-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 505 of 657



XXVII

Table XXVII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-1-1	CH_3	H	H	H	CH_3	H	F	F	OH
XXVII-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH
XXVII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-1-8	CH_3	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-2-1	Et	H	H	H	CH_3	H	F	F	OH
XXVII-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH
XXVII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-2-8	Et	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 506 of 657

Table XXVII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH
XXVII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH
XXVII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-4-1	^{t}Bu	H	H	H	CH_3	H	F	F	OH
XXVII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	F	OH
XXVII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-4-8	^{t}Bu	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-5-1	Ph	H	H	H	CH_3	H	F	F	OH
XXVII-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-5-8	Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 507 of 657

Table XXVII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH
XXVII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH
XXVII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH
XXVII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 508 of 657

Table XXVII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH
XXVII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH
XXVII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXVII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXVII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXVII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXVII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXVII-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-11-1	CH_3	H	H	H	Et	H	F	F	OH
XXVII-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH
XXVII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-11-8	CH_3	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 509 of 657

Table XXVII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-12-1	Et	H	H	H	Et	H	F	F	OH
XXVII-12-2	Et	H	H	CH_3	Et	H	F	F	OH
XXVII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-12-8	Et	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH
XXVII-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH
XXVII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH
XXVII-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH
XXVII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 510 of 657

Table XXVII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-15-1	Ph	H	H	H	Et	H	F	F	OH
XXVII-15-2	Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-15-8	Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH
XXVII-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH
XXVII-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 511 of 657

Table XXVII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH
XXVII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH
XXVII-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH
XXVII-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH
XXVII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXVII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXVII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXVII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXVII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXVII-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 512 of 657

Table XXVII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH
XXVII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH
XXVII-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH
XXVII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 513 of 657

Table XXVII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH
XXVII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 514 of 657

Table XXVII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 515 of 657

Table XXVII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXVII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXVII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXVII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXVII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXVII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXVII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH
XXVII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH
XXVII-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 516 of 657

Table XXVII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH
XXVII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH
XXVII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 517 of 657

Table XXVII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 518 of 657

Table XXVII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXVII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXVII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXVII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXVII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXVII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXVII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-41-1	CH_3	H	H	H	Bz	H	F	F	OH
XXVII-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH
XXVII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-41-8	CH_3	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 519 of 657

Table XXVII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-42-1	Et	H	H	H	Bz	H	F	F	OH
XXVII-42-2	Et	H	H	CH_3	Bz	H	F	F	OH
XXVII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-42-8	Et	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH
XXVII-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH
XXVII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH
XXVII-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH
XXVII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 520 of 657

Table XXVII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-45-1	Ph	H	H	H	Bz	H	F	F	OH
XXVII-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-45-8	Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH
XXVII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH
XXVII-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 521 of 657

Table XXVII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH
XXVII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH
XXVII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVII-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH
XXVII-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH
XXVII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXVII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXVII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXVII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXVII-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 522 of 657



XXVIII

Table XXVIII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-1-1	CH_3	H	H	H	CH_3	H	H	F	OH
XXVIII-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-1-8	CH_3	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-2-1	Et	H	H	H	CH_3	H	H	F	OH
XXVIII-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-2-8	Et	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 523 of 657

Table XXVIII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH
XXVIII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH
XXVIII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-5-1	Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-5-8	Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 524 of 657

Table XXVIII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 525 of 657

Table XXVIII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH
XXVIII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXVIII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXVIII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXVIII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXVIII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXVIII-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-11-1	CH_3	H	H	H	Et	H	H	F	OH
XXVIII-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH
XXVIII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-11-8	CH_3	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 526 of 657

Table XXVIII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-12-1	Et	H	H	H	Et	H	H	F	OH
XXVIII-12-2	Et	H	H	CH_3	Et	H	H	F	OH
XXVIII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-12-8	Et	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH
XXVIII-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH
XXVIII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH
XXVIII-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH
XXVIII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 527 of 657

Table XXVIII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-15-1	Ph	H	H	H	Et	H	H	F	OH
XXVIII-15-2	Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-15-8	Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH
XXVIII-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH
XXVIII-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 528 of 657

Table XXVIII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH
XXVIII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH
XXVIII-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH
XXVIII-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH
XXVIII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXVIII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXVIII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXVIII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXVIII-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 529 of 657

Table XXVIII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 530 of 657

Table XXVIII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 531 of 657

Table XXVIII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 532 of 657

Table XXVIII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXVIII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXVIII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXVIII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXVIII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXVIII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXVIII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 533 of 657

Table XXVIII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 534 of 657

Table XXVIII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 535 of 657

Table XXVIII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXVIII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXVIII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXVIII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXVIII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXVIII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXVIII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-41-1	CH_3	H	H	H	Bz	H	H	F	OH
XXVIII-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH
XXVIII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-41-8	CH_3	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 536 of 657

Table XXVIII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-42-1	Et	H	H	H	Bz	H	H	F	OH
XXVIII-42-2	Et	H	H	CH_3	Bz	H	H	F	OH
XXVIII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-42-8	Et	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH
XXVIII-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH
XXVIII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH
XXVIII-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH
XXVIII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 537 of 657

Table XXVIII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-45-1	Ph	H	H	H	Bz	H	H	F	OH
XXVIII-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-45-8	Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH
XXVIII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH
XXVIII-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 538 of 657

Table XXVIII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH
XXVIII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH
XXVIII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXVIII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXVIII-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH
XXVIII-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH
XXVIII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXVIII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXVIII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXVIII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXVIII-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 539 of 657



XXIX

Table XXIX-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH
XXIX-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH
XXIX-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 540 of 657

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH
XXIX-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH
XXIX-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 541 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 542 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH
XXIX-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH
XXIX-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH
XXIX-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH
XXIX-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH
XXIX-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH
XXIX-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH
XXIX-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 543 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-12-1	Et	H	H	H	Et	H	CH_3	F	OH
XXIX-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-12-8	Et	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH
XXIX-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-14-	^{t}Bu	H	H	H	Et	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 544 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
1									
XXIX-14-	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH
2									
XXIX-14-	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
3									
XXIX-14-	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
4									
XXIX-14-	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH
5									
XXIX-14-	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
6									
XXIX-14-	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
7									
XXIX-14-	^{t}Bu	*	H	*	Et	H	CH_3	F	OH
8									

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-15-1	Ph	H	H	H	Et	H	CH_3	F	OH
XXIX-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-15-8	Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH
XXIX-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 545 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH
XXIX-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH
XXIX-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 546 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH
XXIX-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH
XXIX-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH
XXIX-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH
XXIX-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH
XXIX-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH
XXIX-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-22.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 547 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 548 of 657

Table XXIX-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 549 of 657

Table XXIX-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH
XXIX-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH
XXIX-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH
XXIX-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH
XXIX-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH
XXIX-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH
XXIX-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 550 of 657

Table XXIX-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 551 of 657

Table XXIX-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 552 of 657

Table XXIX-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 553 of 657

Table XXIX-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH
XXIX-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH
XXIX-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH
XXIX-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH
XXIX-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH
XXIX-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH
XXIX-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH
XXIX-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-42-1	Et	H	H	H	Bz	H	CH_3	F	OH
XXIX-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-42-8	Et	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 554 of 657

Table XXIX-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH
XXIX-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH
XXIX-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 555 of 657

Table XXIX-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 556 of 657

Table XXIX-49.


№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXIX-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXIX-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH
XXIX-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH
XXIX-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH
XXIX-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH
XXIX-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH
XXIX-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH
XXIX-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



XXX

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 557 of 657

Table XXX-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-1-1	CH_3	H	H	H	CH_3	H	F	H	OH
XXX-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH
XXX-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-1-8	CH_3	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-2-1	Et	H	H	H	CH_3	H	F	H	OH
XXX-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH
XXX-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-2-8	Et	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH
XXX-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH
XXX-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 558 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH
XXX-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH
XXX-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-5-1	Ph	H	H	H	CH_3	H	F	H	OH
XXX-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-5-8	Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH
XXX-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 559 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH
XXX-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH
XXX-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH
XXX-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 560 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH
XXX-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH
XXX-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH
XXX-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH
XXX-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH
XXX-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH
XXX-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-11-1	CH_3	H	H	H	Et	H	F	H	OH
XXX-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH
XXX-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH
XXX-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-11-8	CH_3	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-12-1	Et	H	H	H	Et	H	F	H	OH

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 561 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-12-2	Et	H	H	CH_3	Et	H	F	H	OH
XXX-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH
XXX-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-12-8	Et	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH
XXX-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH
XXX-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH
XXX-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH
XXX-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH
XXX-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH
XXX-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-15.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 562 of 657

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-15-1	Ph	H	H	H	Et	H	F	H	OH
XXX-15-2	Ph	H	H	CH_3	Et	H	F	H	OH
XXX-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-15-8	Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH
XXX-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH
XXX-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH
XXX-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH
XXX-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 563 of 657

Table XXX-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH
XXX-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH
XXX-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH
XXX-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH
XXX-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH
XXX-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH
XXX-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH
XXX-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH
XXX-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH
XXX-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH
XXX-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH
XXX-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 564 of 657

Table XXX-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH
XXX-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH
XXX-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH
XXX-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 565 of 657

Table XXX-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH
XXX-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 566 of 657

Table XXX-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 567 of 657

Table XXX-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH
XXX-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH
XXX-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH
XXX-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH
XXX-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH
XXX-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH
XXX-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH
XXX-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH
XXX-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 568 of 657

Table XXX-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH
XXX-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH
XXX-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 569 of 657

Table XXX-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 570 of 657

Table XXX-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH
XXX-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH
XXX-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH
XXX-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH
XXX-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH
XXX-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH
XXX-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-41-1	CH_3	H	H	H	Bz	H	F	H	OH
XXX-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH
XXX-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-41-8	CH_3	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 571 of 657

Table XXX-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-42-1	Et	H	H	H	Bz	H	F	H	OH
XXX-42-2	Et	H	H	CH_3	Bz	H	F	H	OH
XXX-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-42-8	Et	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH
XXX-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH
XXX-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH
XXX-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH
XXX-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 572 of 657

Table XXX-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-45-1	Ph	H	H	H	Bz	H	F	H	OH
XXX-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-45-8	Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH
XXX-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH
XXX-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 573 of 657

Table XXX-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH
XXX-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH
XXX-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXX-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXX-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH
XXX-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH
XXX-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH
XXX-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH
XXX-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH
XXX-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH
XXX-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH
XXX-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 574 of 657



XXXI

Table XXXI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-1-1	CH_3	H	H	H	CH_3	H	F	F	OH
XXXI-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH
XXXI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-1-8	CH_3	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-2-1	Et	H	H	H	CH_3	H	F	F	OH
XXXI-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH
XXXI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-2-8	Et	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 575 of 657

Table XXXI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH
XXXI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH
XXXI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-4-1	^{t}Bu	H	H	H	CH_3	H	F	F	OH
XXXI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	F	OH
XXXI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-4-8	^{t}Bu	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-5-1	Ph	H	H	H	CH_3	H	F	F	OH
XXXI-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-5-8	Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 576 of 657

Table XXXI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH
XXXI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH
XXXI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH
XXXI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 577 of 657

Table XXXI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH
XXXI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH
XXXI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH
XXXI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH
XXXI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH
XXXI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH
XXXI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH
XXXI-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-11-1	CH_3	H	H	H	Et	H	F	F	OH
XXXI-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH
XXXI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-11-8	CH_3	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 578 of 657

Table XXXI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-12-1	Et	H	H	H	Et	H	F	F	OH
XXXI-12-2	Et	H	H	CH_3	Et	H	F	F	OH
XXXI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-12-8	Et	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH
XXXI-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH
XXXI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH
XXXI-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH
XXXI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 579 of 657

Table XXXI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-15-1	Ph	H	H	H	Et	H	F	F	OH
XXXI-15-2	Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-15-8	Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH
XXXI-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH
XXXI-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 580 of 657

Table XXXI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH
XXXI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH
XXXI-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH
XXXI-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH
XXXI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH
XXXI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH
XXXI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH
XXXI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH
XXXI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH
XXXI-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 581 of 657

Table XXXI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH
XXXI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH
XXXI-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH
XXXI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 582 of 657

Table XXXI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH
XXXI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 583 of 657

Table XXXI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 584 of 657

Table XXXI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH
XXXI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH
XXXI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH
XXXI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH
XXXI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH
XXXI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH
XXXI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH
XXXI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH
XXXI-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 585 of 657

Table XXXI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH
XXXI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH
XXXI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 586 of 657

Table XXXI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 587 of 657

Table XXXI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH
XXXI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH
XXXI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH
XXXI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH
XXXI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH
XXXI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH
XXXI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-41-1	CH_3	H	H	H	Bz	H	F	F	OH
XXXI-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH
XXXI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-41-8	CH_3	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 588 of 657

Table XXXI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-42-1	Et	H	H	H	Bz	H	F	F	OH
XXXI-42-2	Et	H	H	CH_3	Bz	H	F	F	OH
XXXI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-42-8	Et	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH
XXXI-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH
XXXI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH
XXXI-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH
XXXI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 589 of 657

Table XXXI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-45-1	Ph	H	H	H	Bz	H	F	F	OH
XXXI-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-45-8	Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH
XXXI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH
XXXI-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 590 of 657

Table XXXI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH
XXXI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH
XXXI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXI-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH
XXXI-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH
XXXI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH
XXXI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH
XXXI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH
XXXI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH
XXXI-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 591 of 657



XXXII

Table XXXII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-1-1	CH_3	H	H	H	CH_3	H	H	F	OH
XXXII-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH
XXXII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-1-8	CH_3	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-2-1	Et	H	H	H	CH_3	H	H	F	OH
XXXII-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH
XXXII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-2-8	Et	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 592 of 657

Table XXXII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH
XXXII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH
XXXII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH
XXXII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH
XXXII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-5-1	Ph	H	H	H	CH_3	H	H	F	OH
XXXII-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-5-8	Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 593 of 657

Table XXXII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH
XXXII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH
XXXII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH
XXXII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 594 of 657

Table XXXII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH
XXXII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH
XXXII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH
XXXII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH
XXXII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH
XXXII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH
XXXII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH
XXXII-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-11-1	CH_3	H	H	H	Et	H	H	F	OH
XXXII-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH
XXXII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-11-8	CH_3	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 595 of 657

Table XXXII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-12-1	Et	H	H	H	Et	H	H	F	OH
XXXII-12-2	Et	H	H	CH_3	Et	H	H	F	OH
XXXII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-12-8	Et	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH
XXXII-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH
XXXII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH
XXXII-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH
XXXII-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 596 of 657

Table XXXII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-15-1	Ph	H	H	H	Et	H	H	F	OH
XXXII-15-2	Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-15-8	Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH
XXXII-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH
XXXII-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 597 of 657

Table XXXII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH
XXXII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH
XXXII-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH
XXXII-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH
XXXII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH
XXXII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH
XXXII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH
XXXII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH
XXXII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH
XXXII-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 598 of 657

Table XXXII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH
XXXII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH
XXXII-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH
XXXII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 599 of 657

Table XXXII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH
XXXII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 600 of 657

Table XXXII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 601 of 657

Table XXXII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH
XXXII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH
XXXII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH
XXXII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH
XXXII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH
XXXII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH
XXXII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH
XXXII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH
XXXII-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 602 of 657

Table XXXII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH
XXXII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH
XXXII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 603 of 657

Table XXXII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 604 of 657

Table XXXII-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH
XXXII-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH
XXXII-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH
XXXII-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH
XXXII-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH
XXXII-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH
XXXII-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-41-1	CH_3	H	H	H	Bz	H	H	F	OH
XXXII-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH
XXXII-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-41-8	CH_3	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 605 of 657

Table XXXII-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-42-1	Et	H	H	H	Bz	H	H	F	OH
XXXII-42-2	Et	H	H	CH_3	Bz	H	H	F	OH
XXXII-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-42-8	Et	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH
XXXII-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH
XXXII-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH
XXXII-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH
XXXII-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 606 of 657

Table XXXII-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-45-1	Ph	H	H	H	Bz	H	H	F	OH
XXXII-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-45-8	Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH
XXXII-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH
XXXII-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 607 of 657

Table XXXII-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH
XXXII-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH
XXXII-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table XXXII-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y
XXXII-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH
XXXII-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH
XXXII-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH
XXXII-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH
XXXII-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH
XXXII-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH
XXXII-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 608 of 657

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 609 of 657

DOSAGE, ADMINISTRATION, AND USE

A fifth embodiment of the present invention is directed to a composition for the treatment of any of the viral agents disclosed herein said composition comprising a pharmaceutically acceptable medium selected from among an excipient, carrier, diluent, or equivalent medium and a compound, that is intended to include its salts (acid or basic addition salts), hydrates, solvates, and crystalline forms can be obtained, represented by formula **I**.

It is contemplated that the formulation of the fifth embodiment can contain any of the compounds contemplated in any of the aspects of the first, second, third, and fourth embodiments or those specifically recited in the tables above, either alone or in combination with another compound of the present invention.

The compounds of the present invention may be formulated in a wide variety of oral administration dosage forms and carriers. Oral administration can be in the form of tablets, coated tablets, hard and soft gelatin capsules, solutions, emulsions, syrups, or suspensions. Compounds of the present invention are efficacious when administered by suppository administration, among other routes of administration. The most convenient manner of administration is generally oral using a convenient daily dosing regimen which can be adjusted according to the severity of the disease and the patient's response to the antiviral medication.

A compound or compounds of the present invention, as well as their pharmaceutically acceptable salts, together with one or more conventional excipients, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds and the unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as suspensions, emulsions, or filled capsules for oral use; or in the form of suppositories for

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 610 of 657

rectal or vaginal administration. A typical preparation will contain from about 5% to about 95% active compound or compounds (w/w). The term "preparation" or "dosage form" is intended to include both solid and liquid formulations of the active compound and one skilled in the art will appreciate that an active ingredient can exist in different preparations depending on the desired dose and pharmacokinetic parameters.

The term "excipient" as used herein refers to a compound that is used to prepare a pharmaceutical composition, and is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipients that are acceptable for veterinary use as well as human pharmaceutical use. The compounds of this invention can be administered alone but will generally be administered in admixture with one or more suitable pharmaceutical excipients, diluents or carriers selected with regard to the intended route of administration and standard pharmaceutical practice.

A "pharmaceutically acceptable salt" form of an active ingredient may also initially confer a desirable pharmacokinetic property on the active ingredient which were absent in the non-salt form, and may even positively affect the pharmacodynamics of the active ingredient with respect to its therapeutic activity in the body. The phrase "pharmaceutically acceptable salt" of a compound as used herein means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as glycolic acid, pyruvic acid, lactic acid, malonic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, salicylic acid, muconic acid, and the like or (2) basic addition salts formed with the conjugate bases of any of the inorganic acids listed above, wherein the conjugate bases comprise a cationic component selected from among Na^+, K^+, Mg^{2+}, Ca^{2+}, $NH_gR'''_{4-g}{}^+$, in which R''' is a C_{1-3} alkyl and g is a number selected from among 0, 1, 2, 3, or 4. It should

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 611 of 657

be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same acid addition salt.

Solid form preparations include powders, tablets, pills, capsules, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. Solid form preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

Liquid formulations also are suitable for oral administration include liquid formulation including emulsions, syrups, elixirs and aqueous suspensions. These include solid form preparations which are intended to be converted to liquid form preparations shortly before use. Emulsions may be prepared in solutions, for example, in aqueous propylene glycol solutions or may contain emulsifying agents such as lecithin, sorbitan monooleate, or acacia. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known suspending agents.

The compounds of the present invention may be formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, for example,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 612 of 657

by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Suitable formulations along with pharmaceutical carriers, diluents and expcipients are described in *Remington: The Science and Practice of Pharmacy* **1995,** edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania, which is hereby incorporated by reference. A skilled formulation scientist may modify the formulations within the teachings of the specification to provide numerous formulations for a particular route of administration without rendering the compositions of the present invention unstable or compromising their therapeutic activity.

The modification of the present compounds to render them more soluble in water or other vehicle, for example, may be easily accomplished by minor modifications (e.g., salt formulation), which are well within the ordinary skill in the art. It is also well within the ordinary skill of the art to modify the route of administration and dosage regimen of a particular compound in order to manage the pharmacokinetics of the present compounds for maximum beneficial effect in patients.

A sixth embodiment of the present invention is directed to a use of the compound represented by formula **I** in the manufacture of a medicament for the treatment of any condition the result of an infection by any one of the following viral agents: hepatitis C virus, West Nile virus, yellow fever virus, dengue virus, rhinovirus, polio virus, hepatitis A virus, bovine viral diarrhea virus and Japanese encephalitis virus.

The term "medicament" means a substance used in a method of treatment and/or prophylaxis of a subject in need thereof, wherein the substance includes, but is not limited to, a composition, a formulation, a dosage form, and the like, comprising the compound of formula **I**. It is contemplated that the compound of the use of the compound represented by formula **I** in the manufacture of a medicament for the treatment

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 613 of 657

of any of the antiviral conditions disclosed herein of the sixth embodiment can be any of the compounds contemplated in any of the aspects of the first, second, third, and fourth embodiments or those specifically recited in the tables above, either alone or in combination with another compound of the present invention. A medicament includes, but is not limited to, any one of the compositions contemplated by the fifth embodiment of the present invention.

A seventh embodiment of the present invention is directed to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering a therapeutically effective amount of the compound represented by formula I to the subject.

A first aspect of the seventh embodiment is directed to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering a therapeutically effective of at least two or more different compounds falling within the scope of the compound represented by formula I to the subject.

A second aspect of the seventh embodiment is directed to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises alternatively or concurrently administering a therapeutically effective of at least two compounds falling within the scope of the compound represented by formula I to the subject.

It is intended that a subject in need thereof is one that has any condition the result of an infection by any of the viral agents disclosed herein, which includes, but is not limited to, hepatitis C virus, West Nile virus, yellow fever virus, dengue virus, rhinovirus, polio virus, hepatitis A virus, bovine viral diarrhea virus or Japanese encephalitis virus, flaviviridae viruses or pestiviruses or hepaciviruses or a viral agent causing symptoms equivalent or comparable to any of the above-listed viruses.

The term "subject" means a mammal, which includes, but is not limited to, cattle, pigs, sheep, chicken, turkey, buffalo, llama, ostrich, dogs, cats, and humans, preferably the subject is a human. It is contemplated that in the method of treating a subject thereof of the sixth embodiment can be any of the compounds contemplated in any of the aspects

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 614 of 657

of the first, second, and third embodiments or those specifically recited in the tables above, either alone or in combination with another compound of the present invention.

The term "therapeutically effective amount" as used herein means an amount required to reduce symptoms of the disease in an individual. The dose will be adjusted to the individual requirements in each particular case. That dosage can vary within wide limits depending upon numerous factors such as the severity of the disease to be treated, the age and general health condition of the patient, other medicaments with which the patient is being treated, the route and form of administration and the preferences and experience of the medical practitioner involved. For oral administration, a daily dosage of between about 0.1 and about 10 g, including all values in between, such as 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, and 9.5, per day should be appropriate in monotherapy and/or in combination therapy. A preferred daily dosage is between about 0.5 and about 7.5 g per day, more preferred 1.5 and about 6.0 g per day. Generally, treatment is initiated with a large initial "loading dose" to rapidly reduce or eliminate the virus following by a decreasing the dose to a level sufficient to prevent resurgence of the infection. One of ordinary skill in treating diseases described herein will be able, without undue experimentation and in reliance on personal knowledge, experience and the disclosures of this application, to ascertain a therapeutically effective amount of the compounds of the present invention for a given disease and patient.

Therapeutic efficacy can be ascertained from tests of liver function including, but not limited to protein levels such as serum proteins (e.g., albumin, clotting factors, alkaline phosphatase, aminotransferases (e.g., alanine transaminase, aspartate transaminase), 5'-nucleosidase, γ-glutaminyltranspeptidase, etc.), synthesis of bilirubin, synthesis of cholesterol, and synthesis of bile acids; a liver metabolic function, including, but not limited to, carbohydrate metabolism, amino acid and ammonia metabolism. Alternatively the therapeutic effectiveness may be monitored by measuring HCV-RNA. The results of these tests will allow the dose to be optimized.

A third aspect of the seventh embodiment, to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering to the

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 615 of 657

subject a therapeutically effective of a compound represented by formula **I** and a therapeutically effective amount of another antiviral agent; wherein the administration is concurrent or alternative. It is understood that the time between alternative administration can range between 1-24 hours, which includes any sub-range in between including, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, and 23 hours.

A fourth aspect of the seventh embodiment, to a method of treatment in a subject in need thereof said method comprises alternatively or concurrently administering a therapeutically effective of a compound represented by formula **I** and another antiviral agent to the subject. It is understood that the time between alternative administration can range between 1-24 hours, which includes any sub-range in between including, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, and 23 hours.


It is contemplated that the another antiviral agent includes, but is not limited to interferon-α, interferon-β, pegylated interferon-α, ribavirin, levovirin, viramidine, another nucleoside HCV polymerase inhibitor, a HCV non-nucleoside polymerase inhibitor, a HCV protease inhibitor, a HCV helicase inhibitor or a HCV fusion inhibitor. When the active compound or its derivative or salt are administered in combination with another antiviral agent the activity may be increased over the parent compound. When the treatment is combination therapy, such administration may be concurrent or sequential with respect to that of the nucleoside derivatives. "Concurrent administration" as used herein thus includes administration of the agents at the same time or at different times. Administration of two or more agents at the same time can be achieved by a single formulation containing two or more active ingredients or by substantially simultaneous administration of two or more dosage forms with a single active agent.

It will be understood that references herein to treatment extend to prophylaxis as well as to the treatment of existing conditions. Furthermore, the term "treatment" of a HCV infection, as used herein, also includes treatment or prophylaxis of a disease or a condition associated with or mediated by HCV infection, or the clinical symptoms thereof.


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 616 of 657

PROCESS FOR PREPARATION

An eighth embodiment of the present invention is directed to a process for preparing the compound of formula I, which comprises reacting a suitably substituted phosphochloridate compound **4** with a nucleoside analog **5**



4




5

wherein the substituents R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, X, Y, R^6, and base have their meanings as disclosed in the Detailed Description of the Invention.


This reaction is performed in an anhydrous aprotic solvent such tetrahydrofuran, dioxane, or both tetrahydrofuran and dioxane, or any functional equivalent thereof, with tetrahydrofuran being the preferred solvent. The reaction is typically initiated at a temperature range from -78°C to 40°C with the preferred reaction temperature being between 0°C and room temperature. The nucleoside is first stirred with a base (5 to 12 equivalents) such as N-methylimidazole, collidine, pyridine, 2,6-lutidine, 2, 6-^{t}Bu-pyridine, etc. a tertiary amine base, such as triethylamine, diisopropylethylamine, etc., or an alkyl Grignard reagent, such as tBuMgCl, tBuMgBr, MeMgCl, MeMgBr, etc. The phosphorochloridate (3-10 equivalents) is dissolved in the reaction solvent and added to the mixture of the nucleoside and base. The reaction is then allowed to stir over a period of time at a temperature between room temperature and 40°C for a period of 30 min to 24 hr. with the preferred reaction temperature being room temperature and time being 24 hr. The solvent is removed from the reaction mixture and the product is purified by chromatography on silica gel.

A ninth embodiment of the present invention is directed to a product obtained by a process which comprises reacting a suitably substituted phosphochloridate compound **4** with a nucleoside analog **5**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 617 of 657



4



5

wherein the substituents R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, X, Y, R^6, and base have their meanings as disclosed in the Detailed Description of the Invention.

This reaction can be performed in an anhydrous aprotic solvent or other suitable solvent, such as tetrahydrofuran, dioxane, or a mixture of tetrahydrofuran and dioxane, with tetrahydrofuran being the preferred solvent. The reaction is typically initiated at a temperature range from -78°C to 40°C with the preferred reaction temperature being between 0°C and room temperature. The nucleoside is first stirred with a base (5 to 12 equivalents) such as N-methylimidazole, a tertiary amine base or tButyl Magnesium Chloride. The phosphorochloridate (3-10 equivalents) is dissolved in the reaction solvent and added to the mixture of the nucleoside and base. The reaction is then allowed to stir over a period of time at a temperature between room temperature and 40°C for a period of 30 min to 24 hr. with the preferred reaction temperature being room temperature and time being 24 hr. The solvent is removed from the reaction mixture and the product is purified by chromatography on silica gel.


Compounds and Preparation

Phosphoramidate compounds of the present invention can be prepared by condensation of a nucleoside analog **5** with a suitably substituted phosphochloridate compound **4** (Scheme 1). Thus, a suitably substituted phenol can be reacted with phosphorus oxychloride (**1**) to afford an aryloxy phosphorodichloridate **2** (see Example 1) which is subsequently treated with a acid addition salt of an α-amino acid ester in the presence of TEA to afford an aryloxy phosphorochloridate **4**. This arylalkoxy-amino acid phosphoramidate is reacted with the nucleoside analog to provide the product **I** (for

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 618 of 657

procedure see, e.g., C. McGuigan et al. Antiviral Res. 1992 17:311-321; D. Curley et al. Antiviral Res. 1990 14:345-356; McGuigan et al. Antiviral Chem. Chemother 1990 1(2):107-113).

Scheme 1



The preparation of nucleoside phosphoramidates requires reacting an appropriately substituted phosphochloridate with a nucleoside containing a free 5'-hydroxyl moiety. In cases where only one hydroxyl group is present, preparation of the phosphoramidate usually proceeds smoothly when the phosphochloridate is reacted with the desired nucleoside. In cases where the nucleoside contains more than one free hydroxyl group, preparation of the appropriately protected nucleoside might be required. Silyl, acetonide or other alcohol protecting groups known in the art might be warranted for protection of the sugar moiety. For protection of the nucleoside base, protecting a free amino group may require amidine protection strategy.

Condensation of the phosphochloridate can be carried out on the unprotected


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 619 of 657

nucleoside. Since the 5'-OH group of a nucleoside is much less hindered than the 3'-OH group, selective phosphoramidation is possible under carefully controlled conditions. After condensation to form a protected phosphoramidate nucleoside, deprotection to obtain the free phosphoramidate nucleoside can be carried out using standard protocols for nucleic acid chemistry. In many cases, the desired product is readily separated from the starting material using column chromatography on silica gel. The synthetic scheme is summarized in Scheme 1.

A further understanding of the disclosed embodiments will be appreciated by consideration of the following examples, which are only meant to be illustrative, and not limit the disclosed invention.

EXAMPLE 1

General Procedure for Preparation of phosphorodichloridates




A solution of the appropriate phenol **R^1-OH** (1eq) and triethylamine (1 eq.) in anhydrous ether was added dropwise to a stirred solution of phosphoryl trichloride **1** (1eq) at 0 °C over a period of 3 hours under nitrogen. Then the temperature was warmed to room temperature, and the reaction was stirred overnight. The triethylamine salt was quickly removed with suction filtration and the filtrate concentrated *in vacuo* to dryness to afford **2** as an oil which was used without further purification.

EXAMPLE 2

General Procedure for Preparation of phosphorochloridates

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 620 of 657




2 3 4

A solution of triethylamine (2eq) in anhydrous dichloromethane was added dropwise to a solution of aryloxy-phosphodichloridate **2** (1 eq) and the appropriate amino ester **3** (1 eq) in anhydrous dichloromethane with vigorous stirring at -78 °C over a period of 30 to 120 minutes. Then the reaction temperature was allowed to warm to room temperature and stirred over night. Solvent was removed. The residue was washed with ethyl ether and filtered, the filtrate was dried over reduced pressure to give **4**.

EXAMPLE 3

General Procedures for nucleoside phosphoramidate derivatives



4 5 I


A solution of the appropriate phosphorochloridate **4** (6.5 equivalents) in anhydrous tetrahydrofuran (THF) was added to a mixture of nucleoside **5** (1 equivalent) and N-methylimidazole (8 equivalents) in anhydrous THF with vigorous stirring at room temperature and the reaction mixture was stirred overnight. The solvent was removed *in*

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 621 of 657

vacuo and the crude was purified by column chromatography and/or preparative thin layer chromatography to give **I**.

EXAMPLE 4

Preparation of 2'-deoxy-2'-fluoro-2'-*C*-methyluridine



2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine (1.0g, 1 eq) (Clark, J., et al., J. Med. Chem., 2005, 48, 5504-5508) was dissolved in 10 ml of anhydrous pyridine and concentrated to dryness in vacuo. The resulting syrup was dissolved in 20 ml of anhydrous pyridine under nitrogen and cooled to 0°C with stirring. The brown solution was treated with benzoyl chloride (1.63g, 3eq) dropwise over 10 min. The ice bath was removed and stirring continued for 1.5h whereby thin-layer chromatography (TLC) showed no remaining starting material. The mixture was quenched by addition of water (0.5 ml) and concentrated to dryness. The residue was dissolved in 50 mL of dichloromethane (DCM) and washed with saturated $NaHCO_3$ aqueous solution and H_2O. The organic phase was dried over $NaSO_4$ and filtered, concentrated to dryness to give N^4,3',5'-tribenzoyl-2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine (2.0 g, Yield: 91%).


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 622 of 657

N^4,3',5'-tribenzoyl-2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine (2.0g, 1 eq) was refluxed in 80% aqueous AcOH overnight. After cooling and standing at room temperature (15 °C), most of the product precipitated and then was filtered through a sintered funnel. White precipitate was washed with water and co-evaporated with toluene to give a white solid. The filtrate was concentrated and co-evaporated with toluene to give additional product which was washed with water to give a white solid. Combining the two batches of white solid gave 1.50g of 3',5'-dibenzoyl-2'-Deoxy-2'-fluoro-2'-*C*-methyluridine (Yield: 91%).

To a solution of 3',5'-dibenzoyl-2'-Deoxy-2'-fluoro-2'-*C*-methyluridine (1.5 g, 1eq) in MeOH (10 mL) was added a solution of saturated ammonia in MeOH (20mL). The reaction mixture was stirred at 0 °C for 30 min, and then warmed to room temperature slowly. After the reaction mixture was stirred for another 18 hours, the reaction mixture was evaporated under reduced pressure to give the residue, which was purified by column chromatography to afford pure compound 2'-deoxy-2'-fluoro-2'-*C*-methyluridine (500 mg, Yield: 60 %).

EXAMPLE 5

Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine 5'-(phenyl methoxy-alanyl phosphate)




Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 623 of 657

Phenyl methoxyalaninyl phosphorochloridate (1 g, 6.5 eq) dissolved in 3 mL of THF was added to a mixture of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine (0.15 g,1 eq) and N-methylimidazole (0.3 g, 8 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product (50.1 mg, 15.6%). ^{1}H NMR (DMSO-d_6) δ 1.20-1.27 (m, 6H), 3.58 (d, J = 16.0 Hz, 3H), 3.75-3.92 (m, 2H), 4.015-4.379 (m, 2H), 5.54 (t, J = 10.2 Hz, 1H), 5.83-5.91 (m, 1H), 6.00-6.16 (m, 1H), 7.18 (d, J = 8.0 Hz, 2H), 7.22 (s, 1H), 7.35 (t, J = 4.4 Hz, 2H), 7.55 (s, 1H), 11.52 (s, 1H); MS, *m/e* 502 $(M+1)^+$.

EXAMPLE 6

Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine 5'-(phenyl methoxy-valyl phosphate)




Phenyl methoxy-valyl phosphorochloridate (0.6 g, 3.6 eq) dissolved in 3 mL of THF was added to a mixture of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine (0.15 g,1 eq) and N-methylimidazole (0.44 g, 9 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product (60 mg, 20%). ^{1}H NMR (DMSO-d_6) δ 0.74-0.847 (m, 6H), 1.20-1.28 (m,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 624 of 657

3H), 1.89-1.92 (m, 1H), 3.50-3.54 (m, 1H), 3.58 (d, J = 10.4Hz, 3H), 3.72-3.95 (m, 1H), 4.03-4.05 (m, 1H), 4.23-4.43 (m, 2H), 5.56 (t, J = 16.0 Hz, 1H), 5.85-5.92 (m, 1H), 6.01-6.07 (m, 1H), 7.16-7.21 (m, 3H), 7.37 (t, J = 8 Hz, 2H), 7.55-7.60 (m, 1H), 11.52 (s, 1H); MS, *m/e* 530 $(M+1)^+$.

EXAMPLE 7

Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine 5'-(4-bromophenyl methoxy-valyl phosphate)




4-Bromophenyl methoxy-valyl phosphorochloridate (1 g, 3.4 eq) dissolved in 3 mL of THF was added to a mixture of 2'-deoxy-2'-fluoro-2'-*C*-methyluridine (0.2 g,1 eq) and N-methylimidazole (0.35 g, 6 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed reduced pressure to give the desired product (120 mg, 26%). ^{1}H NMR (DMSO-d_6) δ 0.72-0.82 (m, 6H), 1.19-1.26 (m, 3H), 1.86-1.92 (m, 1H), 3.48-3.50 (m, 1H), 3.56 (d, J = 12.0 Hz, 3H), 3.72-3.89 (m, 1H), 3.96-4.03 (m, 1H), 4.22-4.37 (m, 2H), 5.54-5.60 (m, 1H), 5.85-5.91 (m, 1H), 5.98-6.13 (m, 1H), 7.15 (d, J = 8.0 Hz, 2H), 7.49-7.56 (m, 3H), 11.53 (s, 1H); MS, *m/e* 608 $(M+1)^-$.

EXAMPLE 8

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 625 of 657


Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methyluridine 5'-(4-bromophenyl methoxy-alanyl phosphate)



4-Bromophenyl methoxy-alanyl phosphorochloridate (0.6 g, 5 eq) dissolved in 3 mL of THF was added to a mixture of 2'-deoxy-2'-fluoro-2'-*C*-methyluridine (0.15 g,1 eq)and N-methylimidazole (0.3 g, 7.8 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product (40 mg, 12 %); ^{1}H NMR (DMSO-d_6) δ 1.20-1.26 (m, 6H), 3.57 (d, J = 2.8 Hz, 3H), 3.84 (s, 1H), 3.97-4.03 (m, 1H), 4.21-4.25 (m, 1H), 4.33-4.37 (m, 2H), 5.54-5.60 (m, 1H), 5.83-5.89 (m, 1H), 5.98-6.19 (m, 1H), 7.16 (t, J = 10.2 Hz, 2H), 7.52-7.57 (m, 3H), 11.52 (s, 1 H); MS, *m/e* 580(M+1)$^+$.

EXAMPLE 9

Preparation of N^4-(*N*,*N*-dimethylformamidinyl)-2'-deoxy-2'-fluoro-2'-*C*-methylcytidine




2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine (500 mg, 1.9 mmol) was stirred with dimethylformamide dimethyl acetal in DMF (10 mL). The resulting mixture was stirred at room temperature overnight. After solvent removal the crude product was used

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 626 of 657

for next step without further purification.

EXAMPLE 10

Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine 5'-(phenyl methoxy-alanyl phosphate)




Phenyl methoxyalaninyl phosphorochloridate (0.6 g, 6 eq) dissolved in 3 mL of THF was added to a mixture of N^4-(*N,N*-dimethylformamidinyl)-2'-deoxy-2'-fluoro-2'-*C*-methylcytidine (0.15 g,1 eq) and N-methylimidazole (0.3 g, 7.8 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product (62 mg, 20.6%). ^{1}H NMR (DMSO-d_6) δ 1.16 (d, J = 23.2 Hz, 3H), 1.22 (d, J = 7.2 Hz, 3H), 3.56 (S, 3H), 3.69-3.75 (d, J = 25.6 Hz, 1H), 3.82-3.86 (m, 1H), 3.96-3.98 (m, 1H), 4.21-4.34 (m, 2H), 5.68 (d, J = 7.2 Hz, 1H), 5.75-5.77 (m, 1H), 6.07-6.16 (m, 1H), 7.15-7.19 (m, 3H), 7.2 (d, J = 9.2 Hz, 2H), 7.39 (t, J = 7.8 Hz, 2H), 7.48 (d, J = 9.2 Hz, 1H); MS, *m/e* 501(M+1)$^+$.

EXAMPLE 11

Preparation of 2'-Deoxy-2'-fluoro-2'-*C*-methylcytidine 5'-(4-bromophenyl methoxy-valyl phosphate)


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 627 of 657



4-Bromophenyl methoxy-valyl phosphorochloridate (1.0 g, 3.4 eq.) dissolved in 3 mL of THF was added to a mixture of N^4-(*N,N*-dimethylformamidinyl)-2'-deoxy-2'-fluoro-2'-*C*-methylcytidine (0.2 g,1 eq.) and N-methylimidazole (0.35 g, 6 eq.) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure. The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product as a white solid (59 mg, 13%); ^{1}H NMR (DMSO-d_6) δ 0.74-0.83 (m, 6H), 1.12-1.20 (m, 3H), 1.89-1.92 (m, 1H), 3.49-3.51 (m, 1H), 3.55 (s, 3H), 3.59-3.68 (m, 1H), 3.72-.383 (m, 1H), 4.21-4.39 (m, 2H), 5.70-5.72 (m, 1H), 5.76-5.83 (m, 1H), 6.04-6.16 (m, 1H), 7.15 (d, J = 13.0 Hz, 2H), 7.26 (s, 1H), 7.33 (s, 1H), 7.46-7.55 (m, 1H), 7.56 (d, J = 4.4 Hz, 2H) ; MS, *m/e* 607 $(M+1)^+$.

EXAMPLE 12

Preparation of 2'-deoxy-2'-fluoro-2'-*C*-methylcytidine 5'-(phenyl methoxy-valyl phosphate)



Phenyl methoxy-valyl phosphorochloridate (0.6 g, 6 eq) dissolved in 3 mL of THF was added to a mixture of N^4-(*N,N*-dimethylformamidinyl)-2'-deoxy-2'-fluoro-2'-*C*-

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 628 of 657






methylcytidine (0.15 g,1 eq) and N-methylimidazole (0.3 g, 7.8 eq) in 3 mL THF with vigorous stirring at room temperature, then the reaction was stirred overnight. Solvent was removed by reduced pressure The resulting crude product was dissolved in methanol purified by prep-HPLC on a YMC 25x30X2 mm column using a water / acetonitrile gradient elution mobile phase. The acetonitrile and water were removed under reduced pressure to give the desired product as a white solid (86 mg, 42.9 %). ^{1}H NMR (DMSO-d_6) δ 0.72-0.80 (m, 6H), 1.09-1.18 (m, 3H), 1.87-1.92 (m, 1H), 3.47-3.51 (m, 1H), 3.58 (s, 3H), 3.71-3.75 (m, 1H), 3.97 (t, J = 11.2 Hz,1H), 4.22-4.37 (m, 2H), 5.70 (d, J = 8.0 Hz,1H), 5.76-5.84 (m, 1H), 6.01-6.15 (m, 1H), 7.13-7.18 (m, 3H), 7.27 (s, 2H), 7.34 (d, J = 4.0 Hz, 2H), 7.46-7.50 (m, 1H); MS, *m/e* 529 $(M+1)^+$.

EXAMPLE 13



HCV replicon assay. HCV replicon RNA-containing Huh7 cells (clone A cells; Apath, LLC, St. Louis, Mo.) were kept at exponential growth in Dulbecco's modified Eagle's medium (high glucose) containing 10% fetal bovine serum, 4 mM L-glutamine and 1 mM sodium pyruvate, 1× nonessential amino acids, and G418 (1,000 μg/ml). Antiviral assays were performed in the same medium without G418. Cells were seeded in a 96-well plate at 1,500 cells per well, and test compounds were added immediately after seeding. Incubation time 4 days. At the end of the incubation step, total cellular RNA was isolated (RNeasy 96 kit; Qiagen). Replicon RNA and an internal control (TaqMan rRNA control reagents; Applied Biosystems) were amplified in a single-step multiplex RT-PCR protocol as recommended by the manufacturer. The HCV primers and probe were designed with Primer Express software (Applied Biosystems) and covered highly conserved 5′-untranslated region (UTR) sequences (sense, 5′-AGCCATGGCGTTAGTA(T)GAGTGT-3′, and antisense, 5′-TTCCGCAGACCACTATGG-3′; probe, 5′-FAM-CCTCCAGGACCCCCCCTCCC-TAMRA-3′).

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 629 of 657

To express the antiviral effectiveness of a compound, the threshold RT-PCR cycle of the test compound was subtracted from the average threshold RT-PCR cycle of the no-drug control (ΔCt_{HCV}). A ΔCt of 3.3 equals a 1-log 10 reduction (equal to the 90% effective concentration [EC_{90}]) in replicon RNA levels. The cytotoxicity of the test compound could also be expressed by calculating the ΔCt_{rRNA} values. The $\Delta\Delta Ct$ specificity parameter could then be introduced ($\Delta Ct_{HCV} - \Delta Ct_{rRNA}$), in which the levels of HCV RNA are normalized for the rRNA levels and calibrated against the no-drug control.

Compound	Log10 Reduction at 50µM	EC90 (µM)
	-1.21	3.0
	-0.45	ND
	0.31	ND
	0.31	ND
	-0.55	ND

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 630 of 657

Compound	Log10 Reduction at 50μM	EC90 (μM)
	-1.25	19.15
	-1.48	2.11

[1](4-BrPh): 4-bromo-phenyl.

All the references contained herein are hereby incorporated by reference in so far as needed to supplement the present disclosure. In the event that the incorporated reference contains a term that conflicts with a term disclosed herein, the meaning of the term contained herein controls provided that the overall meaning of the incorporated subject matter is not lost.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 631 of 657

Electronic Acknowledgement Receipt

EFS ID:	1642729
Application Number:	60909315
International Application Number:	
Confirmation Number:	1346
Title of Invention:	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS
First Named Inventor/Applicant Name:	MICHAEL J SOFIA
Customer Number:	23552
Filer:	Daniel R. Evans
Filer Authorized By:	
Attorney Docket Number:	60137.0034USP1
Receipt Date:	30-MAR-2007
Filing Date:	
Time Stamp:	18:41:06
Application Type:	Provisional

Payment information:

Submitted with Payment	yes
Payment was successfully received in RAM	$1100
RAM confirmation Number	1393
Deposit Account	132725
The Director of the USPTO is hereby authorized to charge indicated fees and credit any overpayment as follows: Charge any Additional Fees required under 37 C.F.R. Section 1.16 and 1.17	

File Listing:

Document Number	Document Description	File Name	File Size(Bytes)	Multi Part /.zip	Pages (if appl.)
1	Transmittal of New Application	601370034usp1_sb0016.pdf	224044	no	2
Warnings:					
Information:					
2	Application Data Sheet	601370034usp1_US_ADS_Form_SB_14.pdf	1127300	no	5
Warnings:					
Information:					
3	Specification	601370034USP1_prov_appln_FINAL_033007_Part_01.pdf	15887205	no	200
Warnings:					
Information:					
4	Specification	601370034USP1_prov_appln_FINAL_033007_Part_02.pdf	15653430	no	200
Warnings:					
Information:					
5	Specification	601370034USP1_prov_appln_FINAL_033007_Part_03.pdf	19175763	no	245
Warnings:					
Information:					
6	Fee Worksheet (PTO-06)	fee-info.pdf	8264	no	2
Warnings:					
Information:					
		Total Files Size (in bytes):	52076006		

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for a filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application.

National Stage of an International Application under 35 U.S.C. 371


If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

We claim:

1. A compound, its stereoisomer, salt, hydrate, solvate, or crystalline form thereof, represented by formula I:



I

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b}

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 635 of 657

together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 636 of 657

alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein

Z is N or CR^{12};


R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 637 of 657

C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl);

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

2. A composition for the treatment and/or prophylaxis of any of the viral agents disclosed herein said composition comprising a pharmaceutically acceptable medium selected from among an excipient, carrier, diluent, or equivalent medium and a compound, its stereoisomer, salt, hydrate, solvate, or crystalline form thereof, represented by formula I:



I

wherein

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 638 of 657

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $–N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 639 of 657

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;


(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 640 of 657



a **b** **c** **d**

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted $C_{1\text{-}20}$ alkyl, an optionally substituted $C_{1\text{-}10}$ alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)($C_{1\text{-}20}$ alkyl), C(O)($C_{1\text{-}10}$ alkyl), or C(O)(lower alkyl);


R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

3. A use of the compound represented by formula I in the manufacture of a medicament for the treatment and/or prophylaxis of any condition the result of an

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 641 of 657

infection by hepatitis C virus, West Nile virus, yellow fever virus, dengue virus, rhinovirus, polio virus, hepatitis A virus, bovine viral diarrhea virus or Japanese encephalitis virus.

wherein the compound, its stereoisomer, salt, hydrate, solvate, or crystalline form thereof, are represented by formula I:



I

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 642 of 657

C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;


(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein


Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl);

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

4. A method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering a therapeutically effective amount of the compound represented by formula I to the subject;

wherein the compound or its stereoisomer, salt, hydrate, solvate, or crystalline form thereof represented by formula I:



I

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 645 of 657

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $–N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 646 of 657

hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3"}$ is –OR' or -$N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., -$(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;


(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$;

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 647 of 657

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein

Z is N or CR^{12};


R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'_2, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl);

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 648 of 657

Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, $CONHR'$, $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

5. A process for preparing a compound, its stereoisomer, salt, hydrate, solvate, or crystalline form thereof, represented by formula I:



I

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or -$N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, -$(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b}

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 649 of 657

together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 650 of 657

alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-4} acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 651 of 657

C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl);


R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$;

said process comprising:

reacting a substituted phosphochloridate compound **4** with a nucleoside analog **5**



4




5


6. A product its stereoisomer, salt, hydrate, solvate or crystalline form thereof, prepared a process comprising:

reacting a substituted phosphochloridate compound **4** with a nucleoside analog **5**

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 652 of 657



4



5

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl;. ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl or aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro or cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $–N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is independently selected from H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$,

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 653 of 657

CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, -$CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or -$N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, aryl or substituted aryl wherein said aryl is phenyl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., -$(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, ethynyl alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is an OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-10} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, C(O)O(C_{1-4} alkyl), C(O)O(C_{1-4} alkyl), C(O)O(C_{2-4} alkynyl), C(O)O(C_{2-4} alkenyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl),

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 654 of 657

$S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a **b** **c** **d**

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$ wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, $C(O)(C_{1-20}$ alkyl), $C(O)(C_{1-10}$ alkyl), or C(O)(lower alkyl);


Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 655 of 657

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'_2, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 656 of 657

ABSTRACT

Disclosed herein are phosphoramidate prodrugs of nucleoside derivatives for the treatment of viral infections in mammals, which is a compound, its stereoisomer, salt (acid or basic addition salt), hydrate, solvate, or crystalline form thereof, represented by the following structure:



I

Also disclosed are methods of treatment, uses, and processes for preparing each of which utilize the compound represented by formula **I**.

Atea Pharmaceuticals, Inc.
Exhibit 1003 Page 657 of 657