PTO/AIA/15 (03-13)
Approved for use through 01/31/2014. OMB 0651-0032
U.S. Patent and Trademark Office. U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(ONLY FOR NEW NONPROVISIONAL APPLICATIONS UNDER

Attorney Docket No.	PH3080.PC6-TRK1
First Named Inventor	Jinfa Du
Title	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

1. [ ] **Fee Transmittal Form** (PTO/SB/17 or equivalent)
2. [ ] **Applicant asserts small entity status.** See 37 CFR 1.27
3. [ ] **Applicant certifies micro entity status.** See 37 CFR 1.29. Applicant must attach form PTO/SB/15A or B or equivalent.
4. [X] **Specification** [*Total Pages* 736]
   Both the claims and abstract must start on a new page.
   (See MPEP § 608.01(a) for information on the preferred arrangement)
5. [ ] **Drawing(s)** *(35 U.S.C. 113)* [*Total Sheets*]
6. **Inventor's Oath or Declaration** [*Total Pages* 4]
   (including substitute statements under 37 CFR 1.64 and assignments serving as an oath or declaration under 37 CFR 1.63(e))
   a. [ ] Newly executed (original or copy)
   b. [X] A copy from a prior application (37 CFR 1.63(d))
7. [X] **Application Data Sheet** **See note below.*
   See 37 CFR 1.76 (PTO/AIA/14 or equivalent)
8. [ ] **CD-ROM or CD-R**
   In duplicate, large table, or Computer Program *(Appendix)*
   [ ] Landscape Table on CD
9. **Nucleotide and/or Amino Acid Sequence Submission**
   (if applicable, items a. – c. are required)
   a. [ ] Computer Readable Form (CRF)
   b. Specification Sequence Listing on:
      i. [ ] CD-ROM or CD-R (2 copies); or ii. [ ] Paper
   c. [ ] Statements verifying identity of above copies

ADDRESS TO:

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

ACCOMPANYING APPLICATION PARTS

10. [ ] **Assignment Papers**
    (cover sheet & document(s))
    Name of Assignee
11. [ ] **37 CFR 3.73(c) Statement** *(when there is an assignee)* [X] **Power of Attorney**
12. [ ] **English Translation Document** *(if applicable)*
13. [ ] **Information Disclosure Statement** (PTO/SB/08 or PTO-1449)
    [ ] Copies of citations attached
14. [X] **Preliminary Amendment**
15. [ ] **Return Receipt Postcard** *(MPEP § 503) (Should be specifically itemized)*
16. [ ] **Certified Copy of Priority Document(s)** *(if foreign priority is claimed)*
17. [ ] **Nonpublication Request** Under 35 U.S.C.122 (b)(2)(B)(i). Applicant must attach form PTO/SB/35 or equivalent.
18. [ ] Other:

***Note:** (1) Benefit claims under 37 CFR 1.78 and foreign priority claims under 1.55 **must** be included in an Application Data Sheet (ADS).
(2) For applications filed under 35 U.S.C. 111, the application must contain an ADS specifying the applicant if the applicant is an assignee, person to whom the inventor is under an obligation to assign, or person who otherwise shows sufficient proprietary interest in the matter. See 37 CFR 1.46(b).

18. CORRESPONDENCE ADDRESS

[X] The address associated with Customer Number: 25000 ***OR*** [ ] Correspondence address below

Name					
Address					
City		State		Zip Code	
Country		Telephone		Email	

Signature	Lawrence F. Rozsnyai	Date	April 16, 2014
Name (Print/Type)	Lawrence F. Rozsnyai	Registration No. (Attorney/Agent)	58,427

Corrected Application Data Sheet

Inventor Information

Inventor Number::	1
Name Prefix::	Mr.
Given Name::	Jinfa
Family Name::	Du
City of Residence::	Redwood City
State or Province of Residence::	CALIFORNIA
Country of Residence::	US
Street of mailing address::	c/o Gilead Sciences, Inc. 333 Lakeside Drive
City of mailing address::	Foster City
State or Province of mailing address::	CALIFORNIA
Postal or Zip Code of mailing address::	94404

Inventor Number::	2
Given Name::	Dhanapalan
Family Name::	Nagarathnam
City of Residence::	Bethany
State or Province of Residence::	CONNECTICUT
Country of Residence::	US
Street of mailing address::	52 Virginia Rail Drive
City of mailing address::	Bethany

Gilead Exhibit 2026
PGR2023-00046 - Page 002

State or Province of mailing address::	CONNECTICUT
Postal or Zip Code of mailing address::	06524
Inventor Number::	3
Name Prefix::	Mr.
Given Name::	Michael
Middle Name::	Joseph
Family Name::	Sofia
City of Residence::	Doylestown
State or Province of Residence::	PENNSYLVANIA
Country of Residence::	US
Street of mailing address::	3066 Antler Drive
City of mailing address::	Doylestown
State or Province of mailing address::	PENNSYLVANIA
Postal or Zip Code of mailing address::	18902
Inventor Number::	4
Name Prefix::	Mr.
Given Name::	Peiyuan
Family Name::	Wang
City of Residence::	Foster City
State or Province of Residence::	CALIFORNIA
Country of Residence::	US

	c/o Gilead Sciences, Inc.
Street of mailing address::	333 Lakeside Drive
City of mailing address::	Foster City
State or Province of mailing address::	CALIFORNIA
Postal or Zip Code of mailing address::	94404

Correspondence Information

Correspondence Customer Number::	25000

Application Information

Application Number::	Not Yet Assigned
Application Type::	Regular
Subject Matter::	Utility
CD-ROM or CD-R?::	None
Sequence submission?::	None
Computer Readable Form (CRF)?::	No
Title::	NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS
Attorney Docket Number::	PH3080.PC6-TRK1
Request for Early Publication?::	No
Request for Non-Publication?::	No
Small Entity?::	No
Petition included?::	No

Gilead Exhibit 2026
PGR2023-00046 - Page 004

Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2::	No
Authorization to Permit Access to the Instant Application by the Participating Offices::	No
This application (1) claims priority to or the benefit of an application filed before March 16, 2013 and (2) also contains, or contained at any time, a claim to a claimed invention that has an effective filing date on or after March 16, 2013::	No

Representative Information

Representative Customer Number:: 25000

Domestic Priority Information

This application is a continuation application of U.S. Patent Application No. 14/057,675, filed October 18, 2013, which is a continuation application of U.S. Patent Application No. 13/609,614, filed September 11, 2012, which issued as U.S. Patent No. 8,580,765 on November 12, 2013, which is a continuation of U.S. Patent Application No. 13/099,671, filed May 3, 2011, which issued as U.S. Patent No. 8,334,270 on December 18, 2012, which is a continuation of U.S. Patent Application No. 12/053,015, filed March 21, 2008, which issued as U.S. Patent No. 7,964,580 on June 21, 2011, which claims the benefit of U.S. Provisional Application No. 60/982,309, filed October 24, 2007, and U.S. Provisional Application No. 60/909,315, filed March 30, 2007.

Foreign Priority Information

N/A


Gilead Exhibit 2026
PGR2023-00046 - Page 005

Applicant Information

Applicant Number::	1
Applicant Type::	Assignee
Organization Name::	Gilead Pharmasset LLC
Street of mailing address::	333 Lakeside Drive
City of mailing address::	Foster City
State or Province of mailing address::	CA
Postal or Zip Code of mailing address::	94404
Phone Number::	(650) 574-3000
Fax Number::	(650) 522-5575

Non-Applicant Assignee Information

Signature:

A signature of the applicant or representative is required in accordance with 37 CFR 1.33 and 10.18. Please see 37 CFR 1.4(d) for the form of the signature.			
Signature		Date	April 16, 2014
Name (Print/Type)	Lawrence F. Rozsnyai	Registration No. (Attorney/Agent)	58,427

Gilead Exhibit 2026
PGR2023-00046 - Page 006

Docket PH3080.PC6-TRK1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application No.: Not yet assigned	)	
	:	Examiner: Not yet assigned
First Named Inventor:	)	
	:	Group Art Unit: Not yet assigned
JINFA DU	)	
	:	Confirmation No.: Not yet assigned
Filed: April 16, 2014	)	
	:	
For: NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS	)	
	:	
	)	

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

PRELIMINARY AMENDMENT

Sir:

Prior to examination on the merits of the above-identified application, please amend the application as follows and consider the following remarks.

331919.1

Gilead Exhibit 2026
PGR2023-00046 - Page 007

AMENDMENTS TO THE SPECIFICATION

At page 1, line 1, immediately after the title of invention and before the section titled "Field of Invention," please add the following section heading and paragraph:

CROSS-REFERENCE TO RELATED APPLICATIONS


This application is a continuation application of U.S. Patent Application No. 14/057,675, filed October 18, 2013, which is a continuation application of U.S. Patent Application No. 13/609,614, filed September 11, 2012, which issued as U.S. Patent No. 8,580,765 on November 12, 2013, which is a continuation of U.S. Patent Application No. 13/099,671, filed May 3, 2011, which issued as U.S. Patent No. 8,334,270 on December 18, 2012, which is a continuation of U.S. Patent Application No. 12/053,015, filed March 21, 2008, which issued as U.S. Patent No. 7,964,580 on June 21, 2011, which claims the benefit of U.S. Provisional Application No. 60/982,309, filed October 24, 2007, and U.S. Provisional Application No. 60/909,315, filed March 30, 2007. The entire contents of the aforementioned applications are incorporated herein by reference in their entireties.

AMENDMENTS TO THE CLAIMS


The following listing of the claims replaces all prior versions and listings.

1.-3. (Cancelled).

4. (New) A method of treating a human infected by hepatitis C virus, comprising administering to the subject an effective amount of an NS3 inhibitor and an effective amount of a compound represented by the following formula:



5. (New) The method of claim 4 wherein the NS3 inhibitor is a compound represented by the following formula:



REMARKS

Claims 1-3 have been cancelled without prejudice or disclaimer of subject matter. Claims 4 and 5 have been added, with claim 4 being in independent form. Support for the new claims may be found in the subject specification as filed. Support for claim 4 is found, for example, at page 647, lines 20-25 of the specification as filed. Support for claim 5 is found, for example, at page 76 of WO 2007/014926 (shown as compound 47), the disclosure of which has been incorporated by reference into the subject specification.


The specification has been amended to provide a cross-reference to related applications. Applicant submits that no new matter has been added.

Early and favorable action on the merits is respectfully requested.

Should the Examiner have any questions, he is invited to contact the undersigned attorney at (650) 522-1871.

Dated: April 16, 2014

Respectfully submitted,



Lawrence F. Rozsnyai
Registration No. 58,427
Attorney for the Applicant
Gilead Sciences, Inc.
333 Lakeside Drive
Foster City, California 94494
(650) 522-1871
(650) 522-5575 – Fax

Attorney Docket No. 60137.0034USU1

MERCHANT & GOULD P.C.

United States Patent Application

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: NUCLEOSIDE PHOSPHARAMIDATE PRODRUGS

The specification of which
a. ☐ is attached hereto
b. ☒ was filed on MARCH 21, 2008 as application serial no. 12/053,015 and was amended on
(if applicable) (in the case of a PCT-filed application) described and claimed in international no. filed
and as amended on (if any), which I have reviewed and for which I solicit a United States patent.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

a. ☒ no such applications have been filed.
b. ☐ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

U.S. PROVISIONAL APPLICATION NUMBER	DATE OF FILING (Day, Month, Year)
60/909,315	30 MARCH 2007
60/982,309	24 OCTOBER 2007

I acknowledge the duty to disclose information that is material to patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (reprinted below):

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

(1) prior art cited in search reports of a foreign patent office in a counterpart application, and

(2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

(1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;

or

(2) It refutes, or is inconsistent with, a position the applicant takes in:

(i) Opposing an argument of unpatentability relied on by the Office, or

(ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

(1) Each inventor named in the application:

(2) Each attorney or agent who prepares or prosecutes the application; and

(3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby appoint the attorney(s) and/or patent agent(s) associated with the following customer number to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith.

23552
PATENT TRADEMARK OFFICE

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant & Gould P.C. to the contrary.

I understand that the execution of this document, and the grant of a power of attorney, does not in itself establish an attorney-client relationship between the undersigned and the law firm Merchant & Gould P.C., or any of its attorneys.

Please direct all correspondence in this case to customer number 23552.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor	Family Name SOFIA	First Given Name MICHAEL	Second Given Name JOSEPH
°0	Residence & Citizenship	City DOYLESTOWN	State or Foreign Country PA	Country of Citizenship US
1	Mailing Address	Address 3066 ANTLER DRIVE	City PA	State & Zip Code/Country 18902
Signature of Inventor 201: Michael J. Sofia				Date: 3/26/08

2	Full Name Of Inventor	Family Name DU	First Given Name JINFA	Second Given Name
0	Residence & Citizenship	City NEW HOPE	State or Foreign Country PA	Country of Citizenship US
2	Mailing Address	Address 1206 REINS CIRCLE	City PA	State & Zip Code/Country 18938
Signature of Inventor 202:				Date: 3/26/08

2	Full Name Of Inventor	Family Name WANG	First Given Name PEIYUAN	Second Given Name
0	Residence & Citizenship	City GLEN ROCK	State or Foreign Country NJ	Country of Citizenship ~~CN~~ US
3	Mailing Address	Address 20 RADBURN ROAD	City GLEN ROCK	State & Zip Code/Country 07452
Signature of Inventor 203: Peiyuan Wang				Date: 3/26/08

2	Full Name Of Inventor	Family Name NAGARATHNAM	First Given Name DHANAPALAN	Second Given Name
0	Residence & Citizenship	City BETHANY	State or Foreign Country CT	Country of Citizenship US
4	Mailing Address	Address 52 VIRGINIA RAIL DRIVE	City BETHANY	State & Zip Code/Country 06524
Signature of Inventor 204:				Date: 3/31/08

NUCLEOSIDE PHOSPHORAMIDATE PRODRUGS

Field of Invention

The present invention pertains to nucleoside phosphoramidates and their use as agents for treating viral diseases. These compounds are inhibitors of RNA-dependent RNA viral replication and are useful as inhibitors of HCV NS5B polymerase, as inhibitors of HCV replication and for treatment of hepatitis C infection in mammals. The invention provides novel chemical compounds, and the use of these compounds alone or in combination with other antiviral agents for treating HCV infection.

Background

Hepatitis C virus (HCV) infection is a major health problem that leads to chronic liver disease, such as cirrhosis and hepatocellular carcinoma, in a substantial number of infected individuals, estimated to be 2-15% of the world's population. There are an estimated 4.5 million infected people in the United States alone, according to the U.S. Center for Disease Control. According to the World Health Organization, there are more than 200 million infected individuals worldwide, with at least 3 to 4 million people being infected each year. Once infected, about 20% of people clear the virus, but the rest can harbor HCV the rest of their lives. Ten to twenty percent of chronically infected individuals eventually develop liver-destroying cirrhosis or cancer. The viral disease is transmitted parenterally by contaminated blood and blood products, contaminated needles, or sexually and vertically from infected mothers or carrier mothers to their offspring. Current treatments for HCV infection, which are restricted to immunotherapy

with recombinant interferon-α alone or in combination with the nucleoside analog ribavirin, are of limited clinical benefit. Moreover, there is no established vaccine for HCV. Consequently, there is an urgent need for improved therapeutic agents that effectively combat chronic HCV infection.

The HCV virion is an enveloped positive-strand RNA virus with a single oligoribonucleotide genomic sequence of about 9600 bases which encodes a polyprotein of about 3,010 amino acids. The protein products of the HCV gene consist of the structural proteins C, E1, and E2, and the non-structural proteins NS2, NS3, NS4A and NS4B, and NS5A and NS5B. The nonstructural (NS) proteins are believed to provide the catalytic machinery for viral replication. The NS3 protease releases NS5B, the RNA-dependent RNA polymerase from the polyprotein chain. HCV NS5B polymerase is required for the synthesis of a double-stranded RNA from a single-stranded viral RNA that serves as a template in the replication cycle of HCV. Therefore, NS5B polymerase is considered to be an essential component in the HCV replication complex (K. Ishi, et al, *Heptology*, 1999, 29: 1227-1235; V. Lohmann, et al., *Virology*, 1998, 249: 108-118). Inhibition of HCV NS5B polymerase prevents formation of the double-stranded HCV RNA and therefore constitutes an attractive approach to the development of HCV-specific antiviral therapies.

HCV belongs to a much larger family of viruses that share many common features.

Flaviviridae Viruses

Gilead Exhibit 2026
PGR2023-00046 - Page 0016

The Flaviviridae family of viruses comprises at least three distinct genera: *pestiviruses*, which cause disease in cattle and pigs; *flavivruses*, which are the primary cause of diseases such as dengue fever and yellow fever; and *hepaciviruses*, whose sole member is HCV. The flavivirus genus includes more than 68 members separated into groups on the basis of serological relatedness (Calisher et al., *J. Gen. Virol*, 1993,70,37-43). Clinical symptoms vary and include fever, encephalitis and hemorrhagic fever (*Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 1996, Chapter 31, 931-959). Flaviviruses of global concern that are associated with human disease include the Dengue Hemorrhagic Fever viruses (DHF), yellow fever virus, shock syndrome and Japanese encephalitis virus (Halstead, S. B., *Rev. Infect. Dis.*, 1984, 6, 251-264; Halstead, S. B., *Science*, 239:476-481, 1988; Monath, T. P., *New Eng. J. Med*, 1988, 319, 64 1-643).

The pestivirus genus includes bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, also called hog cholera virus) and border disease virus (BDV) of sheep (Moennig, V. et al. *Adv. Vir. Res.* 1992, 41, 53-98). Pestivirus infections of domesticated livestock (cattle, pigs and sheep) cause significant economic losses worldwide. BVDV causes mucosal disease in cattle and is of significant economic importance to the livestock industry (Meyers, G. and Thiel, H.J., Advances in Virus Research, 1996, 47, 53-118; Moennig V., et al, Adv. Vir. Res. 1992, 41, 53-98). Human pestiviruses have not been as extensively characterized as the animal pestiviruses. However, serological surveys indicate considerable pestivirus exposure in humans.

Pestiviruses and hepaciviruses are closely related virus groups within the Flaviviridae family. Other closely related viruses in this family include the GB virus A,

Gilead Exhibit 2026
PGR2023-00046 - Page 0017

GB virus A-like agents, GB virus-B and GB virus-C (also called hepatitis G virus, HGV). The hepacivirus group (hepatitis C virus; HCV) consists of a number of closely related but genotypically distinguishable viruses that infect humans. There are at least 6 HCV genotypes and more than 50 subtypes. Due to the similarities between pestiviruses and hepaciviruses, combined with the poor ability of hepaciviruses to grow efficiently in cell culture, bovine viral diarrhea virus (BVDV) is often used as a surrogate to study the HCV virus.

The genetic organization of pestiviruses and hepaciviruses is very similar. These positive stranded RNA viruses possess a single large open reading frame (ORF) encoding all the viral proteins necessary for virus replication. These proteins are expressed as a polyprotein that is co- and post-translationally processed by both cellular and virus-encoded proteinases to yield the mature viral proteins. The viral proteins responsible for the replication of the viral genome RNA are located within approximately the carboxy-terminal. Two-thirds of the ORF are termed nonstructural (NS) proteins. The genetic organization and polyprotein processing of the nonstructural protein portion of the ORF for pestiviruses and hepaciviruses is very similar. For both the pestiviruses and hepaciviruses, the mature nonstructural (NS) proteins, in sequential order from the amino-terminus of the nonstructural protein coding region to the carboxy-terminus of the ORF, consist of p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

The NS proteins of pestiviruses and hepaciviruses share sequence domains that are characteristic of specific protein functions. For example, the NS3 proteins of viruses in both groups possess amino acid sequence motifs characteristic of serine proteinases and of helicases (Gorbalenya et al., *Nature*, 1988, 333, 22; Bazan and Fletterick *Virology*

Gilead Exhibit 2026
PGR2023-00046 - Page 0018

, 1989,171,637-639; Gorbalenya et al., *Nucleic Acid Res.*, 1989, 17, 3889-3897). Similarly, the NS5B proteins of pestiviruses and hepaciviruses have the motifs characteristic of RNA-directed RNA polymerases (Koonin, E.V. and Dolja, V.V., *Crir. Rev. Biochem. Molec. Biol.* 1993, 28, 375-430).

The actual roles and functions of the NS proteins of pestiviruses and hepaciviruses in the lifecycle of the viruses are directly analogous. In both cases, the NS3 serine proteinase is responsible for all proteolytic processing of polyprotein precursors downstream of its position in the ORF (Wiskerchen and Collett, *Virology*, 1991, 184, 341-350; Bartenschlager et al., *J. Virol.* 1993, 67, 3835-3844; Eckart et al. *Biochem. Biophys. Res. Comm.* 1993,192, 399-406; Grakoui et al., *J. Virol.* 1993, 67, 2832-2843; Grakoui et al., *Proc. Natl. Acad Sci. USA* 1993, 90, 10583-10587; Hijikata et al., *J. Virol.* 1993, 67, 4665-4675; Tome et al., *J. Virol.*, 1993, 67, 4017-4026). The NS4A protein, in both cases, acts as a cofactor with the NS3 serine protease (Bartenschlager et al., *J. Virol.* 1994, 68, 5045-5055; Failla et al., *J. Virol.* 1994, 68, 3753-3760; Xu et al., *J. Virol.*, 1997, 71:53 12-5322). The NS3 protein of both viruses also functions as a helicase (Kim et al., *Biochem. Biophys. Res. Comm.*, 1995, 215, 160-166; Jin and Peterson, *Arch. Biochem. Biophys.*, 1995, 323, 47-53; Warrener and Collett, *J. Virol.* 1995, 69,1720-1726). Finally, the NS5B proteins of pestiviruses and hepaciviruses have the predicted RNA-directed RNA polymerases activity (Behrens et al., EMBO, 1996, 15, 12-22; Lechmann et al., *J. Virol.*, 1997, 71, 8416-8428; Yuan et al., *Biochem. Biophys. Res. Comm.* 1997, 232, 231-235; Hagedorn, PCT WO 97/12033; Zhong et al, *J. Virol.*, 1998, 72, 9365-9369).

Gilead Exhibit 2026
PGR2023-00046 - Page 0019

Currently, there are limited treatment options for individuals infected with hepatitis C virus. The current approved therapeutic option is the use of immunotherapy with recombinant interferon-α alone or in combination with the nucleoside analog ribavirin. This therapy is limited in its clinical effectiveness and only 50% of treated patients respond to therapy. Therefore, there is significant need for more effective and novel therapies to address the unmet medical need posed by HCV infection.

A number of potential molecular targets for drug development of direct acting antivirals as anti -HCV therapeutics have now been identified including, but not limited to, the NS2-NS3 autoprotease, the N3 protease, the N3 helicase and the NS5B polymerase. The RNA-dependent RNA polymerase is absolutely essential for replication of the single-stranded, positive sense, RNA genome and this enzyme has elicited significant interest among medicinal chemists.

Inhibitors of HCV NS5B as potential therapies for HCV infection have been reviewed: Tan, S.-L., et al., *Nature Rev. Drug Discov.*, 2002, 1, 867-881; Walker, M.P. et al., *Exp. Opin. Investigational Drugs*, 2003, 12, 1269-1280; Ni, Z-J., et al., *Current Opinion in Drug Discovery and Development*, 2004, 7, 446-459; Beaulieu, P. L., et al., *Current Opinion in Investigational Drugs*, 2004, 5, 838-850; Wu, J., et al., *Current Drug Targets-Infectious Disorders*, 2003, 3, 207-219; Griffith, R.C., et al, *Annual Reports in Medicinal Chemistry*, 2004, 39, 223-237; Carrol, S., et al., *Infectious Disorders-Drug Targets*, 2006, 6, 17-29. The potential for the emergence of resistant HCV strains and the need to identify agents with broad genotype coverage supports the need for continuing efforts to identify novel and more effective nucleosides as HCV NS5B inhibitors.

Gilead Exhibit 2026
PGR2023-00046 - Page 0020

Nucleoside inhibitors of NS5B polymerase can act either as a non-natural substrate that results in chain termination or as a competitive inhibitor which competes with nucleotide binding to the polymerase. To function as a chain terminator the nucleoside analog must be taken up by the cell and converted *in vivo* to a triphosphate to compete for the polymerase nucleotide binding site. This conversion to the triphosphate is commonly mediated by cellular kinases which imparts additional structural requirements on a potential nucleoside polymerase inhibitor. Unfortunately, this limits the direct evaluation of nucleosides as inhibitors of HCV replication to cell-based assays capable of *in situ* phosphorylation.


In some cases, the biological activity of a nucleoside is hampered by its poor substrate characteristics for one or more of the kinases needed to convert it to the active triphosphate form. Formation of the monophosphate by a nucleoside kinase is generally viewed as the rate limiting step of the three phosphorylation events. To circumvent the need for the initial phosphorylation step in the metabolism of a nucleoside to the active triphosphate analog, the preparation of stable phosphate prodrugs has been reported. Nucleoside phosphoramidate prodrugs have been shown to be precursors of the active nucleoside triphosphate and to inhibit viral replication when administered to viral infected whole cells (McGuigan, C., et al., *J. Med. Chem.*, 1996, 39, 1748-1753; Valette, G., et al., *J. Med. Chem.*, 1996, 39, 1981-1990; Balzarini, J., et al., *Proc. National Acad Sci USA*, 1996, 93, 7295-7299; Siddiqui, A. Q., et al., *J. Med. Chem.*, 1999, 42, 4122-4128; Eisenberg, E. J., et al., *Nucleosides, Nucleotides and Nucleic Acids*, 2001, 20, 1091-1098; Lee, W.A., et al., *Antimicrobial Agents and Chemotherapy*, 2005, 49, 1898); US 2006/0241064; and WO 2007/095269.

Gilead Exhibit 2026
PGR2023-00046 - Page 0021

Also limiting the utility of nucleosides as viable therapeutic agents is their sometimes poor physicochemical and pharmacokinetic properties. These poor properties can limit the intestinal absorption of an agent and limit uptake into the target tissue or cell. To improve on their properties prodrugs of nucleosides have been employed. It has been demonstrated that preparation of nucleoside phosphoramidates improves the systemic absorption of a nucleoside and furthermore, the phosphoramidate moiety of these "pronucleotides" is masked with neutral lipophilic groups to obtain a suitable partition coefficient to optimize uptake and transport into the cell dramatically enhancing the intracellular concentration of the nucleoside monophosphate analog relative to administering the parent nucleoside alone. Enzyme-mediated hydrolysis of the phosphate ester moiety produces a nucleoside monophosphate wherein the rate limiting initial phosphorylation is unnecessary.

SUMMARY OF THE INVENTION

The present invention is directed toward novel phosphoramidate prodrugs of nucleoside derivatives for the treatment of viral infections in mammals, which is a compound, its stereoisomers, salts (acid or basic addition salts), hydrates, solvates, or crystalline forms thereof, represented by the following structure:



I

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0022

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is $-OR^{1'}$ or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b};

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or

Gilead Exhibit 2026
PGR2023-00046 - Page 0023

alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, R^{3*} is –OR' or -$N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy, di(lower alkyl)-amino, or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., -$(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;


(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

Gilead Exhibit 2026
PGR2023-00046 - Page 0024

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:



a b c d

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

Gilead Exhibit 2026
PGR2023-00046 - Page 0025

R^{12} is H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that when base is represented by the structure c with R^{11} being hydrogen, R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

DEFINITIONS

The phrase "a" or "an" entity as used herein refers to one or more of that entity; for example, a compound refers to one or more compounds or at least one compound. As such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.

The phrase "as defined herein above" refers to the first definition provided in the Summary of the Invention.

The terms "optional" or "optionally" as used herein means that a subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional bond" means that the bond may or may not be present, and that the description includes single, double, or triple bonds.

The term "independently" is used herein to indicate that a variable is applied in any one instance without regard to the presence or absence of a variable having that same or a different definition within the same compound. Thus, in a compound in which R appears twice and is defined as "independently carbon or nitrogen", both R's can be carbon, both R's can be nitrogen, or one R' can be carbon and the other nitrogen.

The term "alkenyl" refers to an unsubstituted hydrocarbon chain radical having from 2 to 10 carbon atoms having one or two olefinic double bonds, preferably one olefinic double bond. The term "$C_{2\text{-}N}$ alkenyl" refers to an alkenyl comprising 2 to N

Gilead Exhibit 2026
PGR2023-00046 - Page 0026

carbon atoms, where N is an integer having the following values: 3, 4, 5, 6, 7, 8, 9, or 10. The term "C_{2-10} alkenyl" refers to an alkenyl comprising 2 to 10 carbon atoms. The term "C_{2-4} alkenyl" refers to an alkenyl comprising 2 to 4 carbon atoms. Examples include, but are not limited to, vinyl, 1-propenyl, 2-propenyl (allyl) or 2-butenyl (crotyl).

The term "halogenated alkenyl" refers to an alkenyl comprising at least one of F, Cl, Br, and I.

The term "alkyl" refers to an unbranched or branched chain, saturated, monovalent hydrocarbon residue containing 1 to 30 carbon atoms. The term "C_{1-M} alkyl" refers to an alkyl comprising 1 to M carbon atoms, where M is an integer having the following values: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30. The term "C_{1-4} alkyl" refers to an alkyl containing 1 to 4 carbon atoms. The term "lower alkyl" denotes a straight or branched chain hydrocarbon residue comprising 1 to 6 carbon atoms. "C_{1-20} alkyl" as used herein refers to an alkyl comprising 1 to 20 carbon atoms. "C_{1-10} alkyl" as used herein refers to an alkyl comprising 1 to 10 carbons. Examples of alkyl groups include, but are not limited to, lower alkyl groups include methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl. The term (ar)alkyl or (heteroaryl)alkyl indicate the alkyl group is optionally substituted by an aryl or a heteroaryl group respectively.


The term "cycloalkyl" refers to an unsubstituted or substituted carbocycle, in which the carbocycle contains 3 to 10 carbon atoms; preferably 3 to 8 carbon atoms; more preferably 3 to 6 carbon atoms (i.e., lower cycloalkyls). Examples of cycloalkyl groups include, but are not limited to, cyclopropyl, 2-methyl-cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

The term "cycloalkyl alkyl" refers to an additionally unsubstituted or substituted alkyl substituted by a lower cycloalkyl. Examples of cycloalkyl alkyls include, but are not limited to, any one of methyl, ethyl, propyl, i-propyl, n-butyl, i-butyl, t-butyl or pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl that is substituted with cyclopropyl, 2-methyl-cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

Gilead Exhibit 2026
PGR2023-00046 - Page 0027

The term "cycloheteroalkyl" refers to an unsubstituted or substituted heterocycle, in which the heterocycle contains 2 to 9 carbon atoms; preferably 2 to 7 carbon atoms; more preferably 2 to 5 carbon atoms. Examples of cycloheteroalkyls include, but are not limited to, aziridin-2-yl, *N*-C_{1-3}-alkyl-aziridin-2-yl, azetidinyl, *N*-C_{1-3}-alkyl-azetidin-m'-yl, pyrrolidin-m'-yl, *N*-C_{1-3}-alkyl-pyrrolidin-m'-yl, piperidin-m'-yl, and *N*-C_{1-3}-alkyl-piperidin-m'-yl, where m' is 2, 3, or 4 depending on the cycloheteroalkyl. Specific examples of N-C_{1-3}-alkyl-cycloheteroalkyls include, but are not limited to, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-2-yl, N-methyl-piperidin-3-yl, and N-methyl-piperidin-4-yl. In the instance of R^4, the point of attachment between the cycloheteroalkyl ring carbon and the oxygen occurs at any one of m'

The term "heterocycle" refers to an unsubstituted or substituted heterocycle containing carbon, hydrogen, and at least one of N, O, and S, where the C and N can be trivalent or tetravalent, i.e., sp^2- or sp^3-hybridized. Examples of heterocycles include, but are not limited to, aziridine, azetidine, pyrrolidine, piperidine, imidazole, oxazole, piperazine, etc. In the instance of piperazine, as related to R^{10} for NR'_2, the corresponding opposite nitrogen atom of the piperazinyl is substituted by a lower alkyl represented by the following structure:



Preferably, the opposite nitrogen of the piperazinyl is substituted by a methyl group.

The term "halogenated alkyl" (or "haloalkyl") refers to an unbranched or branched chain alkyl comprising at least one of F, Cl, Br, and I. The term "C_{1-M} haloalkyl" refers to an alkyl comprising 1 to M carbon atoms that comprises at least one of F, Cl, Br, and I, where M is an integer having the following values: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30. "C_{1-3} haloalkyl" refers to a haloalkyl comprising 1 to 3 carbons and at least one of F, Cl, Br, and I. The term "halogenated lower alkyl" (or "lower haloalkyl") refers to a haloalkyl comprising 1 to 6

Gilead Exhibit 2026
PGR2023-00046 - Page 0028

carbon atoms and at least one of F, Cl, Br, and I. Examples include, but are not limited to, fluoromethyl, chloromethyl, bromomethyl, iodomethyl, difluoromethyl, dichloromethyl, dibromomethyl, diiodomethyl, trifluoromethyl, trichloromethyl, tribromomethyl, triiodomethyl, 1-fluoroethyl, 1-chloroethyl, 1-bromoethyl, 1-iodoethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-difluoroethyl, 2,2-dichloroethyl, 2,2-dibromomethyl, 2-2-diiodomethyl, 3-fluoropropyl, 3-chloropropyl, 3-bromopropyl, 2,2,2-trifluoroethyl or 1,1,2,2,2-pentafluoroethyl.

The term "alkynyl" refers to an unbranched or branched hydrocarbon chain radical having from 2 to 10 carbon atoms, preferably 2 to 5 carbon atoms, and having one triple bond. The term "C_{2-N} alkynyl" refers to an alkynyl comprising 2 to N carbon atoms, where N is an integer having the following values: 3, 4, 5, 6, 7, 8, 9, or 10. The term "C C_{2-4} alkynyl" refers to an alkynyl comprising 2 to 4 carbon atoms. The term "C_{2-10} alkynyl" refers to an alkynyl comprising 2 to 10 carbons. Examples include, but are limited to, ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl or 3-butynyl.

The term "halogenated alkynyl" refers to an unbranched or branched hydrocarbon chain radical having from 2 to 10 carbon atoms, preferably 2 to 5 carbon atoms, and having one triple bond and at least one of F, Cl, Br, and I.

The term "cycloalkyl" refers to a saturated carbocyclic ring comprising 3 to 8 carbon atoms, i.e. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl. The term "C_{3-7} cycloalkyl" as used herein refers to a cycloalkyl comprising 3 to 7 carbons in the carbocyclic ring.

The term "alkoxy" refers to an -O-alkyl group or an –O-cycloalkyl group, wherein alkyl and cycloalkyl are as defined above. Examples of –O-alkyl groups include, but are not limited to, methoxy, ethoxy, *n*-propyloxy, *i*-propyloxy, *n*-butyloxy, *i*-butyloxy, *t*-butyloxy. "Lower alkoxy" as used herein denotes an alkoxy group with a "lower alkyl" group as previously defined. "C_{1-10} alkoxy" refers to an-O-alkyl wherein alkyl is C_{1-10}. Examples of –O-cycloalkyl groups include, but are not limited to, -O-*c*-propyl, -O-*c*-butyl, -O-*c*-pentyl, and –O-*c*-hexyl.

Gilead Exhibit 2026
PGR2023-00046 - Page 0029

The term "halogenated alkoxy" refers to an –O-alkyl group in which the alkyl group comprises at least one of F, Cl, Br, and I.

The term "halogenated lower alkoxy" refers to an –O-(lower alkyl) group in which the lower alkyl group comprises at least one of F, Cl, Br, and I.

The term "amino acid" includes naturally occurring and synthetic α, β γ or δ amino acids, and includes but is not limited to, amino acids found in proteins, i.e. glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucinyl, isoleucinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glycinyl, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl, glutaroyl, lysinyl, argininyl, histidinyl, β-alanyl, β-valinyl, β-leucinyl, β-isoleucinyl, β-prolinyl, β-phenylalaninyl, β-tryptophanyl, β-methioninyl, β-glycinyl, β-serinyl, β-threoninyl, β-cysteinyl, β-tyrosinyl, β-asparaginyl, β-glutaminyl, β-aspartoyl, β-glutaroyl, β-lysinyl, β-argininyl or β-histidinyl. When the term amino acid is used, it is considered to be a specific and independent disclosure of each of the esters of α, β γ or δ glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine in the D and L-configurations.

The term "aminoacyl" includes N,N-unsubstituted, N,N-monosubstituted, and N,N-disubstituted derivatives of naturally occurring and synthetic α, β γ or δ amino acyls, where the amino acyls are derived from amino acids. The amino-nitrogen can be substituted or unsubstituted. When the amino-nitrogen is substituted, the nitrogen is either mono- or di-substituted, where the substituent bound to the amino-nitrogen is a lower alkyl or an alkaryl. In the instance of its use for Y, the expression "O(aminoacyl)" is used. It is understood that the C3' carbon of the ribose is bound to the oxygen "O", which is then bound to the carbonyl carbon of the aminoacyl.

Gilead Exhibit 2026
PGR2023-00046 - Page 0030

The terms "alkylamino" or "arylamino" refer to an amino group that has one or two alkyl or aryl substituents, respectively.

The term "protected," as used herein and unless otherwise defined, refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis. Non-limiting examples include: C(O)-alkyl, C(O)Ph, C(O)aryl, CH_3, CH_2-alkyl, CH_2-alkenyl, CH_2Ph, CH_2-aryl, CH_2O-alkyl, CH_2O-aryl, SO_2-alkyl, SO_2-aryl, *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl, and 1,3-(1,1,3,3-tetraisopropyldisiloxanylidene).

The term "aryl," as used herein, and unless otherwise specified, refers to substituted or unsubstituted phenyl (Ph), biphenyl, or naphthyl, preferably the term aryl refers to substituted or unsubstituted phenyl. The aryl group can be substituted with one or more moieties selected from among hydroxyl, F, Cl, Br, I, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, and phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in T.W. Greene and P.G. M. Wuts, "Protective Groups in Organic Synthesis," 3rd ed., John Wiley & Sons, 1999.

The terms "alkaryl" or "alkylaryl" refer to an alkyl group with an aryl substituent, such as benzyl. The terms "aralkyl" or "arylalkyl" refer to an aryl group with an alkyl substituent.

The term "di(lower alkyl)amino-lower alkyl" refers to a lower alkyl substituted by an amino group that is itself substituted by two lower alkyl groups. Examples include, but are not limited to, $(CH_3)_2NCH_2$, $(CH_3)_2NCH_2CH_2$, $(CH_3)_2NCH_2CH_2CH_2$, etc. The examples above show lower alkyls substituted at the terminus carbon atom with an N,N-dimethyl-amino substituent. These are intended as examples only and are not intended to limit the meaning of the term "di(lower alkyl)amino-lower alkyl" so as to require the same. It is contemplated that the lower alkyl chain can be substituted with an N,N-di(lower alkyl)-amino at any point along the chain, e.g., CH_3CH(N-(lower alkyl)$_2$)CH_2CH_2.

Gilead Exhibit 2026
PGR2023-00046 - Page 0031

The term "halo," as used herein, includes chloro, bromo, iodo and fluoro.

The term "acyl" refers to a substituent containing a carbonyl moiety and a non-carbonyl moiety. The carbonyl moiety contains a double-bond between the carbonyl carbon and a heteroatom, where the heteroatom is selected from among O, N and S. When the heteroatom is N, the N is substituted by a lower alkyl. The non-carbonyl moiety is selected from straight, branched, and cyclic alkyl, which includes, but is not limited to, a straight, branched, or cyclic C_{1-20} alkyl, C_{1-10} alkyl, or lower alkyl; alkoxyalkyl, including methoxymethyl; aralkyl, including benzyl; aryloxyalkyl, such as phenoxymethyl; or aryl, including phenyl optionally substituted with halogen (F, Cl, Br, I), hydroxyl, C_1 to C_4 alkyl, or C_1 to C_4 alkoxy, sulfonate esters, such as alkyl or aralkyl sulphonyl, including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, trialkylsilyl (e.g. dimethyl-t-butylsilyl) or diphenylmethylsilyl. When at least one aryl group is present in the non-carbonyl moiety, it is preferred that the aryl group comprises a phenyl group.

The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

The term "purine" or "pyrimidine" base includes, but is not limited to, adenine, N^6-alkylpurines, N^6-acylpurines (wherein acyl is C(O)(alkyl, aryl, alkylaryl, or arylalkyl), N^6-benzylpurine, N^6-halopurine, N^6-vinylpurine, N^6-acetylenic purine, N^6-acyl purine, N^6-hydroxyalkyl purine, N^6-allcylaminopurine, N^6-thioallcyl purine, N^2-alkylpurines, N^2-alkyl-6-thiopurines, thymine, cytosine, 5-fluorocytosine, 5-methylcytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrmidine, uracil, 5-halouracil, including 5-fluorouracil, C^5-alkylpyrimidines, C^5-benzylpyrimidines, C^5-halopyrimidines, C^5-vinylpyrimidine, C^5-acetylenic pyrimidine, C^5-acyl pyrimidine, C^5-hydroxyalkyl purine, C^5-amidopyrimidine, C^5-cyanopyrimidine, ,C^5-iodopyrimidine, C^6-lodo-pyrimidine, C^5-Br-vinyl pyrimidine, C^6-Br-vinyl pyrimidine, C^5-nitropyrimidine, C^5-amino-pyrimidine, N^2-alkylpurines, N^2-alkyl-6-thiopurines, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, and pyrazolopyrimidinyl. Purine bases include, but are not limited to, guanine, adenine,

Gilead Exhibit 2026
PGR2023-00046 - Page 0032


hypoxanthine, 2,6-diaminopurine, and 6-chloropurine. Functional oxygen and nitrogen groups on the base can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl, alkyl groups, and acyl groups such as acetyl and propionyl, methanesulfonyl, and p-toluenesulfonyl.

The term "tautomerism" and "tautomers" have their accepted plain meanings.

The term "P*" means that the phosphorous atom is chiral and that it has a corresponding Cahn-Ingold-Prelog designation of "R" or "S" which have their accepted plain meanings. It is contemplated that compounds of the formula I are racemic because the chirality at phosphorous. Applicants contemplate use of the racemate and/or the resolved enantiomers. In some instances, an asterisk does not appear next to the phosphoroamidate phosphorous atom. In these instances, it is understood that the phosphorous atom is chiral and that one of ordinary skill understands this to be so unless the substituents bound to the phosphorous exclude the possibility of chirality at phosphorous, such as in $P(O)Cl_3$.

DETAILED DESCRIPTION OF THE INVENTION

An aspect of the invention is directed to a compound, its salts, hydrates, solvates, crystalline forms, and the like represented by formula I:



wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are

Gilead Exhibit 2026
PGR2023-00046 - Page 0033

optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b};

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

Gilead Exhibit 2026
PGR2023-00046 - Page 0034

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy, di(lower alkyl)-amino, or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;


(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

the base is a naturally occurring or modified purine or pyrimidine base represented by the following structures:

Gilead Exhibit 2026
PGR2023-00046 - Page 0035



a b c d

wherein

Z is N or CR^{12};

R^7, R^8,R^9, R^{10}, and $R^{11}$are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=$CHCO_2H$, or CH=$CHCO_2R'$,

wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, C(O)(C_{1-20} alkyl), C(O)(C_{1-10} alkyl), or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and


R^{12} is H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6,

Gilead Exhibit 2026
PGR2023-00046 - Page 0036

halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, $CONHR'$, $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$; with the proviso that when base is represented by the structure c with R^{11} being hydrogen, R^{12} is not a: (i) $-C{\equiv}C-H$, (ii) $-C{=}CH_2$, or (iii) $-NO_2$.

As can be appreciated from the structure represented by formula I above, there are myriad ways to express the several embodiments and aspects of each embodiment of the present invention. As seen below, the inventors have disclosed certain embodiments directed to the compound of formula I, each having several aspects, based on the identity of the modified purine or pyrimidine base. This is not intended to be an explicit or implicit admission that the three embodiments are independent or distinct nor should it be interpreted as such. Rather, it is intended to convey information so that the full breadth of the present invention can be understood. Furthermore, the following embodiments, and aspects thereof, are not meant to be limiting on the full breadth of the invention as recited by the structure of formula I.

A first embodiment of the invention is directed to a compound represented by formula I-1:



I-1

Gilead Exhibit 2026
PGR2023-00046 - Page 0037

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen, C_{1-10} alkyl, R^{3a} or R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b};

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_e(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-

Gilead Exhibit 2026
PGR2023-00046 - Page 0038

Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy, di(lower alkyl)-amino, or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)-amino, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$

Gilead Exhibit 2026
PGR2023-00046 - Page 0039

alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^7, R^8,R^9 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

A first aspect of the first embodiment is directed to a compound represented by formula **I-1**

wherein

(a) R^1 is hydrogen, n-alkyl or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, -$CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

Gilead Exhibit 2026
PGR2023-00046 - Page 0040

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy, di(lower alkyl)-amino, or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, CN, CH_3, vinyl, OCH_3, OCH_2CH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, vinyl, N_3, CN, Cl, Br, F, I, O(C_{1-6} acyl), O(C_{1-4} alkyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, N(C_{1-18} acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, OC(O)O(C_{1-4} alkyl), OC(O)O(C_{1-4} alkyl), OC(O)O(C_{2-4} alkynyl), OC(O)O(C_{2-4} alkenyl), OC_{1-4} haloalkyl, O(aminoacyl), O(C_{1-4} acyl), O(C_{1-4} alkyl), O(C_{2-4} alkenyl), S(C_{1-4} acyl), S(C_{1-4} alkyl), S(C_{2-4} alkynyl), S(C_{2-4} alkenyl), SO(C_{1-4} acyl), SO(C_{1-4} alkyl), SO(C_{2-4} alkynyl), SO(C_{2-4} alkenyl), SO_2(C_{1-4} acyl), SO_2(C_{1-4} alkyl), SO_2(C_{2-4} alkynyl), SO_2(C_{2-4} alkenyl), $OS(O)_2$(C_{1-4} acyl), $OS(O)_2$(C_{1-4} alkyl), $OS(O)_2$(C_{2-4} alkenyl), NH_2, NH(C_{1-4} alkyl), NH(C_{2-4} alkenyl), NH(C_{2-4} alkynyl), NH(C_{1-4} acyl), N(C_{1-4} alkyl)$_2$, or N(C_{1-4} acyl)$_2$;

(i) R^7, R^8,R^9 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

A second aspect of the first embodiment is directed to a compound represented by formula I-1

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0041

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a C_{1-3} alkyl, a C_{1-3} alkoxy, F, Cl, Br, I, nitro, cyano, and a C_{1-3} haloalkyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, di(lower alkyl)amino-lower alkyl, or aminoacyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, NH_2, or N_3;

Gilead Exhibit 2026
PGR2023-00046 - Page 0042

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$; and

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A third aspect of the first embodiment is directed to a compound represented by formula **I-1**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl and R^{3b} is H;

Gilead Exhibit 2026
PGR2023-00046 - Page 0043

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, lower haloalkyl, di(lower alkyl)amino-lower alkyl, or aminoacyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, NH_2, or N_3;


(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl'; and

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A fourth aspect of the first embodiment is directed to a compound represented by formula I-2

Gilead Exhibit 2026
PGR2023-00046 - Page 0044



I-2

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H;

Gilead Exhibit 2026
PGR2023-00046 - Page 0045

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH , base is cytosine and R^6 is H, R^5 cannot be N_3;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl; and

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, OC(O)(C1-20 alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, NHC(O)(C1-20 alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and and $N(C(O)(CH_2)_sCH_3$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A fifth aspect of the first embodiment is directed to a compound represented by formula **I-2**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0046

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, lower haloalkyl, di(lower alkyl)amino-lower alkyl, or aminoacyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a $C_{1\text{-}20}$ alkyl; a $C_{1\text{-}20}$ cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl; and

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1\text{-}20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1\text{-}20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A sixth aspect of the first embodiment is directed to a compound represented by formula **I-2**

Gilead Exhibit 2026
PGR2023-00046 - Page 0047

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, OMe, CN, CH_2F, F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, or N_3, OCH_3, $OC(O)CH_3$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$; and

(j) R^9 is selected from among OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A seventh aspect of the first embodiment is directed to a compound represented by formula I-2

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0048

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, or N_3;

(h) Y is OH, OCH_3, $OC(O)CH_3$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$; and

(j) R^9 is selected from among OH, OCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from among 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

An eighth aspect of the first embodiment is directed to a compound represented by formula I-2

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0049

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, or N_3;

(h) Y is OH, OCH_3, $OC(O)CH_3$, or O(aminoacyl);

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;


(j) R^9 is selected from among OH, OCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, $OC(O)(C_{1-20}$ alkyl), which include but are not limited to $OC(O)(CH_2)_sCH_3$, $NHC(O)(C_{1-20}$ alkyl), which include but are not limited to $NHC(O)(CH_2)_sCH_3$, and $N(C(O)(CH_2)_sCH_3)_2$, which include but is not limited to $N(C(O)(CH_2)_sCH_3)_2$, where s is an integer selected from among 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

A second embodiment of the invention is directed to a compound represented by formula **I** in which the base is a structure represented by formula **b** above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, Y, R^7, and R^8 are defined in the Summary of the Invention section above.

A first aspect of the second embodiment is directed to a compound represented by

Gilead Exhibit 2026
PGR2023-00046 - Page 0050

formula I-3



I-3

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are

Gilead Exhibit 2026
PGR2023-00046 - Page 0051

$(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl),

Gilead Exhibit 2026
PGR2023-00046 - Page 0052

$S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, or $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, or $N(C_{1-4}$ acyl)$_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The second aspect of the second embodiment is directed to a compound represented by formula **I-3**

wherein

(a) R^1 is hydrogen, n-alkyl or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi)

Gilead Exhibit 2026
PGR2023-00046 - Page 0053

R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, or CH_2SH and R^{3b} is H;

(d) R^4 is hydrogen, $C_{1\text{-}10}$ alkyl, $C_{1\text{-}10}$ alkyl optionally substituted with a lower alkyl, alkoxy or halogen, $C_{1\text{-}10}$ haloalkyl, $C_{3\text{-}10}$ cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, CN, CH_3, vinyl, OCH_3, OCH_2CH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, $C_{1\text{-}4}$ alkyl, vinyl, N_3, CN, Cl, Br, F, I, $O(C_{1\text{-}6}$ acyl), $O(C_{1\text{-}4}$ alkyl), NH_2, $NH(C_{1\text{-}4}$ alkyl), $NH(C_{2\text{-}4}$ alkenyl), $NH(C_{2\text{-}4}$ alkynyl), $NH(C_{1\text{-}4}$ acyl), $N(C_{1\text{-}4}$ alkyl)$_2$, $N(C_{1\text{-}18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $OC(O)O(C_{1\text{-}4}$ alkyl), $OC(O)O(C_{1\text{-}4}$ alkyl), $OC(O)O(C_{2\text{-}4}$ alkynyl), $OC(O)O(C_{2\text{-}4}$ alkenyl), $OC_{1\text{-}10}$ haloalkyl, O(aminoacyl), $O(C_{1\text{-}4}$ acyl), $O(C_{1\text{-}4}$ alkyl), $O(C_{2\text{-}4}$ alkenyl), $S(C_{1\text{-}4}$ acyl), $S(C_{1\text{-}4}$ alkyl), $S(C_{2\text{-}4}$ alkynyl), $S(C_{2\text{-}4}$ alkenyl), $SO(C_{1\text{-}4}$ acyl), $SO(C_{1\text{-}4}$ alkyl), $SO(C_{2\text{-}4}$ alkynyl), $SO(C_{2\text{-}4}$ alkenyl), $SO_2(C_{1\text{-}4}$ acyl), $SO_2(C_{1\text{-}4}$ alkyl), $SO_2(C_{2\text{-}4}$ alkynyl), $SO_2(C_{2\text{-}4}$ alkenyl), $OS(O)_2(C_{1\text{-}4}$ acyl), $OS(O)_2(C_{1\text{-}4}$ alkyl), $OS(O)_2(C_{2\text{-}4}$ alkenyl), NH_2, $NH(C_{1\text{-}4}$ alkyl), $NH(C_{2\text{-}4}$ alkenyl), $NH(C_{2\text{-}4}$ alkynyl), $NH(C_{1\text{-}4}$ acyl), $N(C_{1\text{-}4}$ alkyl)$_2$, $N(C_{1\text{-}4}$ acyl)$_2$;

Gilead Exhibit 2026
PGR2023-00046 - Page 0054

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl, halogenated (F, Cl, Br, I) lower alkyl, lower alkenyl of C_2-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The third aspect of the second embodiment is directed to a compound represented by formula **I-3**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a C_{1-3} alkyl, a C_{1-3} alkoxy, F, Cl, Br, I, nitro, cyano, and a C_{1-3} haloalkyl;

(b) R^2 is hydrogen, CH_3, R^{3a} or R^{3b} and R^2 together are $(CH_2)_3$ so as to form a cyclic ring that includes the adjoining N and C atoms, $C(O)CR^{3a}R^{3b}NHR^1$, where n is 2 to 4 and R^1, R^{3a}, and R^{3b} are as defined herein;

(c) R^{3a} and R^{3b} are independently (i) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 3 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-

Gilead Exhibit 2026
PGR2023-00046 - Page 0055

methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

The fourth aspect of the second embodiment is directed to a compound represented by formula **I-3**

wherein

(a) R^1 is hydrogen, n-alkyl or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, -

Gilead Exhibit 2026
PGR2023-00046 - Page 0056

$CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2(halo), such as CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;


(g) X is H, OH, F, OCH_3, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The fifth aspect of the second embodiment is directed to a compound represented by formula **I-4**

Gilead Exhibit 2026
PGR2023-00046 - Page 0057



I-4

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0058

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl,

The sixth aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0059

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^7 and R^8 are independently H, F, Cl, Br, I, OH, OCH_3, SH, SCH_3, NH_2, $NHCH_3$, $N(CH_3)_2$, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2H, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$, wherein R' is a C_{1-20} alkyl; a C_{1-20} cycloalkyl; a C_2-C_6 alkenyl, a C_2-C_6 alkynyl.

The seventh aspect of the second embodiment is directed to a compound represented by formula I-4

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, OMe, CN, CH_2F, F, Cl, Br, or I;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

Gilead Exhibit 2026
PGR2023-00046 - Page 0060

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$;

The eighth aspect of the second embodiment is directed to a compound represented by formula **I-4**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

The ninth aspect of the second embodiment is directed to a compound represented by formula **I-4**

Gilead Exhibit 2026
PGR2023-00046 - Page 0061

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, or N_3;


(h) Y is OH, OCH_3, or $OC(O)CH_3$;

(i) R^7 and R^8 are independently H, F, Br, SCH_3, CH_3, $CH_{3-q}X_q$, where X is F, Cl, Br, or I and q is 1 to 3, vinyl, CO_2CH_3, $CONH_2$, $CONHCH_3$, or $CON(CH_3)_2$.

A third embodiment of the invention is directed to a compound represented by formula I in which the base is a structure represented by formula c above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, Y, Z, R^{10}, R^{11}, and R^{12} are defined in the Summary of the Invention section above; with the proviso that R^{11} is not H.

A first aspect of the third embodiment is directed to a compound represented by formula I-5

Gilead Exhibit 2026
PGR2023-00046 - Page 0062



I-5

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen

Gilead Exhibit 2026
PGR2023-00046 - Page 0063

or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is from H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl),

Gilead Exhibit 2026
PGR2023-00046 - Page 0064

$SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^{10} and R^{11} are independently H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO$_2$H, or CH=CHCO$_2$R', with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, $C(O)(C_{1-20}$ alkyl), $C(O)(C_{1-10}$ alkyl), or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F,

Gilead Exhibit 2026
PGR2023-00046 - Page 0065

Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, $CONHR'$, $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A second aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, $CONHR'$, $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

Gilead Exhibit 2026
PGR2023-00046 - Page 0066

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

A third aspect of the third embodiment is directed to a compound represented by formula I-5

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

Gilead Exhibit 2026
PGR2023-00046 - Page 0067

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A fourth aspect of the third embodiment is directed to a compound represented by formula I-5

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

Gilead Exhibit 2026
PGR2023-00046 - Page 0068

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A fifth aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0069

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A sixth aspect of the third embodiment is directed to a compound represented by formula I-5

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0070

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

A seventh aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

Gilead Exhibit 2026
PGR2023-00046 - Page 0071

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

An eighth aspect of the third embodiment is directed to a compound represented by formula I-5

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0072

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R';

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R'.

A ninth aspect of the third embodiment is directed to a compound represented by formula **I-5**

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0073

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, or N-methyl-pyrrolidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$;


wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A tenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

Gilead Exhibit 2026
PGR2023-00046 - Page 0074



I-6

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen

Gilead Exhibit 2026
PGR2023-00046 - Page 0075

or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl),

Gilead Exhibit 2026
PGR2023-00046 - Page 0076

$SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R', with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R'.

An eleventh aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH,

Gilead Exhibit 2026
PGR2023-00046 - Page 0077

$CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A twelvth aspect of the third embodiment is directed to a compound represented by formula I-6

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0078

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is a H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

A thirteenth aspect of the third embodiment is directed to a compound represented

Gilead Exhibit 2026
PGR2023-00046 - Page 0079

by formula I-6

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

Gilead Exhibit 2026
PGR2023-00046 - Page 0080

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A fourteenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$, with the proviso that when R^{10} is OH and R^{11} is not NH_2;

Gilead Exhibit 2026
PGR2023-00046 - Page 0081

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A fifteenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_eCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen

Gilead Exhibit 2026
PGR2023-00046 - Page 0082

or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl),

Gilead Exhibit 2026
PGR2023-00046 - Page 0083

$SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R';

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R'.

An sixteenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH,

Gilead Exhibit 2026
PGR2023-00046 - Page 0084

$CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'-OH)-Ph)$, CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^iPr, ^nPr, ^nBu, 2-butyl, ^tBu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$.

A seventeenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0085

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

An eighteenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 0086

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R';

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R'.

A nineteenth aspect of the third embodiment is directed to a compound represented by formula **I-6**

Gilead Exhibit 2026
PGR2023-00046 - Page 0087

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is NH_2 and R^{11} are independently H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, CH=$CHCO_2H$, or CH=$CHCO_2R'$;

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and


(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, CH=$CHCO_2H$, or CH=$CHCO_2R'$.

Gilead Exhibit 2026
PGR2023-00046 - Page 0088

A fourth embodiment of the invention is directed to a compound represented by formula I in which the base is a structure represented by formula c above, where R^{11} is H and R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, and Y are defined in the Summary of the Invention section above.

A first aspect of the fourth embodiment is directed to a compound represented by formula I-7



I-7

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, -

Gilead Exhibit 2026
PGR2023-00046 - Page 0089

$(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

Gilead Exhibit 2026
PGR2023-00046 - Page 0090

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^{10} is H, F, Cl, Br, I, OH, OR', SH, SR', NH_2, NHR', NR'$_2$, lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6, lower alkynyl of C_2-C_6 such as C≡CH, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is an optionally substituted alkyl , which includes, but is not limited to, an optionally substituted C_{1-20} alkyl, an optionally substituted C_{1-10} alkyl, an optionally substituted lower alkyl; an optionally substituted cycloalkyl; an optionally substituted alkynyl of C_2-C_6, an optionally substituted lower alkenyl of C_2-C_6, or optionally substituted acyl, which includes but is not limited to C(O) alkyl, $C(O)(C_{1-20}$ alkyl), $C(O)(C_{1-10}$ alkyl), or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

Gilead Exhibit 2026
PGR2023-00046 - Page 0091

(j) Z is N or CR^{12}; and

R^{12} is H, halogen (including F, Cl, Br, I), OH, OR', SH, SR', NH_2, NHR', NR'$_2$, NO_2 lower alkyl of C_1-C_6, halogenated (F, Cl, Br, I) lower alkyl of C_1-C_6, lower alkenyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkenyl of C_2-C_6 , lower alkynyl of C_2-C_6, halogenated (F, Cl, Br, I) lower alkynyl of C_2-C_6, lower alkoxy of C_1-C_6, halogenated (F, Cl, Br, I) lower alkoxy of C_1-C_6, CO_2H, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$; with the proviso that R^{12} is not a: (i) –C≡C–H, (ii) –$C{=}CH_2$, or (iii) $-NO_2$.

A second aspect of the fourth embodiment is directed to a compound represented by formula I-7

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

Gilead Exhibit 2026
PGR2023-00046 - Page 0092

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

A third aspect of the third embodiment is directed to a compound represented by formula **I-7**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0093

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

A fourth aspect of the fourth embodiment is directed to a compound represented by formula I-7

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-

Gilead Exhibit 2026
PGR2023-00046 - Page 0094

methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} and R^{11} H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

A fifth aspect of the fourth embodiment is directed to a compound represented by formula **I-7**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 0095

(d) R^4 is hydrogen, CH_3, Et, ^iPr, $^n Pr$, $^n Bu$, 2-butyl, $^t Bu$, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$,


wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$.

A sixth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

Gilead Exhibit 2026
PGR2023-00046 - Page 0096



I-8

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$ alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is $-OR^1$ or $-N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen

Gilead Exhibit 2026
PGR2023-00046 - Page 0097

or C_{1-6} alkyl and $R^{3'}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., $-(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl),

Gilead Exhibit 2026
PGR2023-00046 - Page 0098

$OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R',

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, CH=CHCO_2H, or CH=CHCO_2R'; with the proviso that when base is represented by the structure c with R^{11} being hydrogen, R^{12} is not a: (i) –C≡C–H, (ii) –C=CH_2, or (iii) –NO_2.

A seventh aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

Gilead Exhibit 2026
PGR2023-00046 - Page 0099

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R',

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2$R'; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

An eighth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 00100

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl , N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$,

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'$_2$, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH{=}CHCO_2H$, or $CH{=}CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C{\equiv}C-H$, (ii) $-C{=}CH_2$, or (iii) $-NO_2$.

A ninth aspect of the fourth embodiment is directed to a compound represented by

Gilead Exhibit 2026
PGR2023-00046 - Page 00101

formula I-8

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'_2, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

Gilead Exhibit 2026
PGR2023-00046 - Page 00102

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'$_2$, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

A tenth aspect of the fourth embodiment is directed to a compound represented by formula **I-8**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

(i) R^{10} is H, F, Br, I, OH, OR', NH_2, NHR', NR'$_2$, CO_2R', $CONH_2$, CONHR', CONR'$_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$,

Gilead Exhibit 2026
PGR2023-00046 - Page 00103


wherein R' is a lower alkyl, a lower cycloalkyl, or C(O)(lower alkyl) or alternatively, in the instance of NR'_2, each R' comprise at least one C atom that are joined to form a heterocycle comprising at least two carbon atoms; and

(j) Z is N or CR^{12}; and

R^{12} is an H, halogen (including F, Cl, Br, I), OR', NH_2, NHR', NR'_2, NO_2, lower alkyl of C_1-C_6, CO_2R', $CONH_2$, CONHR', $CONR'_2$, $CH=CHCO_2H$, or $CH=CHCO_2R'$; with the proviso that R^{12} is not a: (i) $-C\equiv C-H$, (ii) $-C=CH_2$, or (iii) $-NO_2$.

A fifth embodiment of the invention is directed to a compound represented by formula I in which the base is a structure represented by formula d above, wherein R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, and Y are defined in the Summary of the Invention section above.

The first aspect of the fifth embodiment is directed to a compound represented by formula I-9



I-9

wherein

(a) R^1 is hydrogen, n-alkyl; branched alkyl; cycloalkyl; or aryl, which includes, but is not limited to, phenyl or naphthyl, where phenyl or naphthyl are optionally substituted with at least one of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, F, Cl, Br, I, nitro, cyano, C_{1-6} haloalkyl, $-N(R^{1'})_2$, C_{1-6} acylamino, $-NHSO_2C_{1-6}$

Gilead Exhibit 2026
PGR2023-00046 - Page 00104

alkyl, $-SO_2N(R^{1'})_2$, $COR^{1''}$, and $-SO_2C_{1-6}$ alkyl; ($R^{1'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{1''}$ is –OR' or -$N(R^{1'})_2$);

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} and R^{3b} are (i) independently selected from hydrogen, C_{1-10} alkyl, cycloalkyl, $-(CH_2)_c(NR^{3'})_2$, C_{1-6} hydroxyalkyl, $-CH_2SH$, $-(CH_2)_2S(O)_dMe$, $-(CH_2)_3NHC(=NH)NH_2$, (1H-indol-3-yl)methyl, (1H-imidazol-4-yl)methyl, $-(CH_2)_cCOR^{3''}$, aryl and aryl C_{1-3} alkyl, said aryl groups optionally substituted with a group selected from hydroxyl, C_{1-10} alkyl, C_{1-6} alkoxy, halogen, nitro and cyano; (ii) R^{3a} and R^{3b} both are C_{1-6} alkyl; (iii) R^{3a} and R^{3b} together are$(CH_2)_f$ so as to form a spiro ring; (iv) R^{3a} is hydrogen and R^{3b} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms (v) R^{3b} is hydrogen and R^{3a} and R^2 together are $(CH_2)_n$ so as to form a cyclic ring that includes the adjoining N and C atoms, where c is 1 to 6, d is 0 to 2, e is 0 to 3, f is 2 to 5, n is 2 to 4, and where $R^{3'}$ is independently hydrogen or C_{1-6} alkyl and $R^{3''}$ is -OR' or $-N(R^{3'})_2$); (vi) R^{3a} is H and R^{3b} is H, CH_3, CH_2CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl; or (viii) R^{3a} is CH_3, $-CH_2CH_3$, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl and R^{3b} is H, where $R^{3'}$ is independently hydrogen or alkyl, which includes, but is not limited to, C_{1-20} alkyl, C_{1-10} alkyl, or C_{1-6} alkyl, $R^{3''}$ is –OR' or $-N(R^{3'})_2$);

(d) R^4 is hydrogen, C_{1-10} alkyl, C_{1-10} alkyl optionally substituted with a lower alkyl, alkoxy or halogen, C_{1-10} haloalkyl, C_{3-10} cycloalkyl, cycloalkyl alkyl, cycloheteroalkyl, aminoacyl, di(lower alkyl)amino-lower alkyl, aryl, such as phenyl, heteroaryl, such as, pyridinyl, substituted aryl, or substituted heteroaryl;

Gilead Exhibit 2026
PGR2023-00046 - Page 00105

(e) R^5 is H, a lower alkyl, CN, vinyl, O-(lower alkyl), hydroxyl lower alkyl, i.e., -$(CH_2)_pOH$, where p is 1 -6, including hydroxyl methyl (CH_2OH), CH_2F, N_3, CH_2CN, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, alkyne (optionally substituted), or halogen, including F, Cl, Br, or I, with the provisos that when X is OH, base is cytosine and R^6 is H, R^5 cannot be N_3 and when X is OH, R^6 is CH_3 or CH_2F and B is a purine base, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OMe, halogen, NH_2, or N_3;

(h) Y is OH, H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, vinyl, N_3, CN, Cl, Br, F, I, NO_2, $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{1-4}$ alkyl), $OC(O)O(C_{2-4}$ alkynyl), $OC(O)O(C_{2-4}$ alkenyl), OC_{1-10} haloalkyl, O(aminoacyl), $O(C_{1-10}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-18}$ acyl)$_2$, wherein alkyl, alkynyl, alkenyl and vinyl are optionally substituted by N_3, CN, one to three halogen (Cl, Br, F, I), NO_2, $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{1-4}$ alkyl), $C(O)O(C_{2-4}$ alkynyl), $C(O)O(C_{2-4}$ alkenyl), $O(C_{1-4}$ acyl), $O(C_{1-4}$ alkyl), $O(C_{2-4}$ alkenyl), $S(C_{1-4}$ acyl), $S(C_{1-4}$ alkyl), $S(C_{2-4}$ alkynyl), $S(C_{2-4}$ alkenyl), $SO(C_{1-4}$ acyl), $SO(C_{1-4}$ alkyl), $SO(C_{2-4}$ alkynyl), $SO(C_{2-4}$ alkenyl), $SO_2(C_{1-4}$ acyl), $SO_2(C_{1-4}$ alkyl), $SO_2(C_{2-4}$ alkynyl), $SO_2(C_{2-4}$ alkenyl), $OS(O)_2(C_{1-4}$ acyl), $OS(O)_2(C_{1-4}$ alkyl), $OS(O)_2(C_{2-4}$ alkenyl), NH_2, $NH(C_{1-4}$ alkyl), $NH(C_{2-4}$ alkenyl), $NH(C_{2-4}$ alkynyl), $NH(C_{1-4}$ acyl), $N(C_{1-4}$ alkyl)$_2$, $N(C_{1-4}$ acyl)$_2$.

A second aspect of the fifth embodiment is directed to a compound represented by formula **I-9**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 00106

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H.

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, halogen, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3;

A third aspect of the fifth embodiment is directed to a compound represented by formula **I-9**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-

Gilead Exhibit 2026
PGR2023-00046 - Page 00107

methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I; with the proviso that X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 N_3, OCH_3, or $OC(O)CH_3$;

A fourth aspect of the fifth embodiment is directed to a compound represented by formula I-9

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

Gilead Exhibit 2026
PGR2023-00046 - Page 00108

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

A fifth aspect of the fifth embodiment is directed to a compound represented by formula **I-9**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;


(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

A sixth aspect of the fifth embodiment is directed to a compound represented by formula **I-10**

Gilead Exhibit 2026
PGR2023-00046 - Page 00109



I-10

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, or a substituted or unsubstituted phenyl, where the substitutent of the substituted phenyl is at least one of a CH_3, OCH_3, F, Cl, Br, I, nitro, cyano, and a $CH_{3-q}X_q$, where X is F, Cl, Br, or I, and q is 1-3;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl or R^{3a} is CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, $CH_2((4'\text{-}OH)\text{-}Ph)$, CH_2SH, or lower cycloalkyl and R^{3b} is H;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

Gilead Exhibit 2026
PGR2023-00046 - Page 00110

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, N_3, CH_2CN, CH_2N_3, CH_2NH_2, CH_2NHCH_3, $CH_2N(CH_3)_2$, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3;

(h) Y is OH, H, CH_3, vinyl, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3, NH_2, $NHCH_3$, NH(vinyl), NH(acetyl), $NH(C(O)CH_3)$, $N(CH_3)_2$, $N(C(O)CH_3)_2$;

A seventh aspect of the fifth embodiment is directed to a compound represented by formula **I-10**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, CH_2-indol-3-yl, $-CH_2CH_2SCH_3$, CH_2CO_2H, $CH_2C(O)NH_2$, CH_2CH_2COOH, $CH_2CH_2C(O)NH_2$, $CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2NHC(NH)NH_2$, CH_2-imidazol-4-yl, CH_2OH, $CH(OH)CH_3$, CH_2((4'-OH)-Ph), CH_2SH, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_3, OCH_3, CH_2OH, CH_2F, halogen, including F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, F, or CN;

Gilead Exhibit 2026
PGR2023-00046 - Page 00111

(g) X is H, OH, F, OCH_3, Cl, Br, I, NH_2, or N_3; and

(h) Y is OH, H, CH_3, vinyl, NH_2, N_3, CN, Cl, Br, F, I, $OC(O)CH_3$, OCH_3.

An eighth aspect of the fifth embodiment is directed to a compound represented by formula **I-10**

wherein

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen or CH_3;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, CN, CH_2F, F, Cl, Br, or I, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, F, OCH_3, F, Cl, Br, I, NH_2 or N_3;

(h) Y is H, OH, CH_3, F, Cl, Br, I, NH_2 or N_3, OCH_3, or $OC(O)CH_3$;

A ninth aspect of the fifth embodiment is directed to a compound represented by formula **I-10**

wherein

Gilead Exhibit 2026
PGR2023-00046 - Page 00112

(a) R^1 is hydrogen, methyl, ethyl, *n*-propyl, *i*-propyl, phenyl, p-tolyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;

(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$;

A tenth aspect of the fifth embodiment is directed to a compound represented by formula **I-10**

wherein

(a) R^1 is hydrogen, methyl, phenyl, p-bromo-phenyl, p-chloro-phenyl, p-fluorophenyl;

(b) R^2 is hydrogen;

(c) R^{3a} is H and R^{3b} is H, CH_3, $CH(CH_3)_2$, $CH_2CH(CH_3)_2$, $CH(CH_3)CH_2CH_3$, CH_2Ph, or lower cycloalkyl;

(d) R^4 is hydrogen, CH_3, Et, ^{i}Pr, ^{n}Pr, ^{n}Bu, 2-butyl, ^{t}Bu, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, N-methyl-aziridin-2-yl, N-methyl-azetidin-3-yl, N-

Gilead Exhibit 2026
PGR2023-00046 - Page 00113

methyl-pyrrolidin-3-yl, N-methyl-pyrrolidin-4-yl, N-methyl-piperidin-4-yl, lower haloalkyl, or di(lower alkyl)amino-lower alkyl;

(e) R^5 is H, with the provisos that when X is OH, R^6 is CH_3 or CH_2F, R^5 cannot be H;

(f) R^6 is H, CH_3, CH_2F, CHF_2, CF_3, or F;


(g) X is H, OH, OCH_3, F, NH_2 or N_3;

(h) Y is OH, NH_2, OCH_3, or $OC(O)CH_3$.

The following tables contain numeric identifiers associated with various substituent designators that should be viewed in light of the accompanying structure. These structures are contemplated species of the various aspects of the disclosed embodiments and are not intended to be limiting on full breadth of the contemplated compound represented by the structure of formula **I**. However, it is contemplated that any one of the exemplified nucleoside bases can be used in combination with any one of contemplated species that specify a particular combination of R^1, R^2, R^{3a}, R^{3b}, R^4, R^5, R^6, X, and Y. In each of the presented tables, the phosphoramidate substituent containing the substituents R^{3a} and R^{3b} are depicted without reference to stereochemical structure (*cf.* structures **I-1**, **I-3**, **I-5**, **I-7**, and **I-9** above). It is contemplated that the compounds recited below embody compounds in which R^{3a} projects toward the viewer while R^{3b} projects away from the viewer (cf. structures **I-2**, **I-4**, **I-6**, **I-8**, and **I-10**). Moreover, it is contemplated that the compounds recited below also embody compounds in which R^{3a} projects away from the viewer while R^{3b} projects towards the viewer. Not meant to be limiting, however, it is contemplated that preferred compounds are those in which R^{3a} projects towards the viewer and R^{3b} projects away from the viewer such that the natural L-amino acid (S)-configuration is presented. Additionally, the inventors recognize that the phosphorus atom of the phosphoramidate moiety is another source of chirality. Although the structures below do not specifically depict chirality at phosphorus, the inventors recognize that stereochemical configurations are possible such that in a staggered (or zig-zag) line structure the oxo-substitutent projects towards the viewer

Gilead Exhibit 2026
PGR2023-00046 - Page 00114

while the OR^1 substitutent projects away from the viewer, and vice versa, i.e., where the Cahn-Ingold-Prelog stereochemical designation of phosphorous is either R or S. Therefore, the structures below include all possible stereochemical configurations possible for phosphorus.



II

Table II-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-1-1	CH_3	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-7	CH_3	H	H	$CH_3CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-1-8	CH_3	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-2-1	Et	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-2	Et	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00115

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-2-8	Et	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-5-1	Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00116

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-5-8	Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Gilead Exhibit 2026
PGR2023-00046 - Page 00117

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	OH	OH	H	H	NH_2
II-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00118

Table II-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-11-1	CH_3	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-2	CH_3	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-11-8	CH_3	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-12-1	Et	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-2	Et	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-12-8	Et	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-13-1	^iPr	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-2	^iPr	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-3	^iPr	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-4	^iPr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-5	^iPr	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-6	^iPr	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-7	^iPr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-13-8	^iPr	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00119

Table II-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-14-1	^{t}Bu	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-14-8	^{t}Bu	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-15-1	Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-2	Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-15-8	Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00120

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-17-1	p-F-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-17-8	p-F-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00121

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-20-1	p-I-Ph	H	H	H	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	OH	OH	H	H	NH_2
II-20-8	p-I-Ph	*	H	*	Et	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-22-1	Et	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00122

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-22-8	Et	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00123

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2
II-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00124

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-28-2	p-Cl-Ph	H	H	CH_3	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-5	p-Cl-Ph	H	H	CH_2Ph	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-28-8	p-Cl-Ph	*	H	*	^iPr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-29-1	p-Br-Ph	H	H	H	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-2	p-Br-Ph	H	H	CH_3	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-5	p-Br-Ph	H	H	CH_2Ph	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-29-8	p-Br-Ph	*	H	*	^iPr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-30-1	p-I-Ph	H	H	H	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-2	p-I-Ph	H	H	CH_3	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-5	p-I-Ph	H	H	CH_2Ph	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^iPr	H	CH_3	OH	OH	H	H	NH_2
II-30-8	p-I-Ph	*	H	*	^iPr	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-31.

Gilead Exhibit 2026
PGR2023-00046 - Page 00125

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-32-1	Et	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-32-8	Et	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00126

Table II-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00127

Table II-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00128

Table II-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2
II-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-41-1	CH_3	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-41-8	CH_3	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-42-1	Et	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-2	Et	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-42-8	Et	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00129

Table II-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-45-1	Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-2	Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-45-8	Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00130

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00131

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table II-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
II-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	OH	OH	H	H	NH_2
II-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	OH	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00132



III

Table III-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-1-1	CH_3	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-2	CH_3	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-5	CH_3	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-1-8	CH_3	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-2-1	Et	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-2	Et	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-5	Et	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-2-8	Et	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00133

Table III-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-3-1	^{i}Pr	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-2	^{i}Pr	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-3-8	^{i}Pr	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-4-1	^{t}Bu	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-2	^{t}Bu	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-4-8	^{t}Bu	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-5-1	Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-2	Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-5	Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-5-8	Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00134

Table III-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-6-1	p-Me-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-6-8	p-Me-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-7-1	p-F-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-2	p-F-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-7-20	p-F-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-8-1	p-Cl-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-8-8	p-Cl-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00135

Table III-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-9-1	p-Br-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-9-20	p-Br-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-10-1	p-I-Ph	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-2	p-I-Ph	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
III-10-8	p-I-Ph	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-11-1	CH_3	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-11-2	CH_3	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-5	CH_3	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-11-8	CH_3	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00136

Table III-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-12-1	Et	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-12-2	Et	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-5	Et	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-12-8	Et	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-13-1	^{i}Pr	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-13-2	^{i}Pr	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-13-8	^{i}Pr	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-14-1	^{t}Bu	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-14-2	^{t}Bu	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00137

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-14-8	^{t}Bu	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-15-1	Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-15-2	Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-5	Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-15-8	Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-16-1	p-Me-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-16-2	p-Me-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-16-8	p-Me-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00138

Table III-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-17-1	p-F-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-17-2	p-F-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-17-8	p-F-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-18-1	p-Cl-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-18-2	p-Cl-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-18-8	p-Cl-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-19-1	p-Br-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-19-2	p-Br-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-19-8	p-Br-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00139

Table III-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-20-1	p-I-Ph	H	H	H	Et	H	CH_3	F	OH	H	H	NH_2
III-20-2	p-I-Ph	H	H	CH_3	Et	H	CH_3	F	OH	H	H	NH_2
III-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	CH_3	F	OH	H	H	NH_2
III-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	CH_3	F	OH	H	H	NH_2
III-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	CH_3	F	OH	H	H	NH_2
III-20-8	p-I-Ph	*	H	*	Et	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-21-1	CH_3	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-2	CH_3	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-21-8	CH_3	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-22-1	Et	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-2	Et	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-22-8	Et	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00140

Table III-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-23-1	^{i}Pr	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-23-8	^{i}Pr	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-24-1	^{t}Bu	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-24-8	^{t}Bu	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-25-1	Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-2	Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-25-8	Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00141

Table III-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-27-1	p-F-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-27-8	p-F-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00142

Table III-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-30-1	p-I-Ph	H	H	H	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	CH_3	F	OH	H	H	NH_2
III-30-8	p-I-Ph	*	H	*	^{i}Pr	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-31-1	CH_3	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-2	CH_3	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-31-8	CH_3	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00143

Table III-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-32-1	Et	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-2	Et	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-32-8	Et	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-33-1	^{i}Pr	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-33-8	^{i}Pr	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-34-1	^{t}Bu	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-34-8	^{t}Bu	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00144

Table III-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-35-1	Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-2	Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-35-8	Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-37-1	p-F-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-37-8	p-F-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00145

Table III-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-40-1	p-I-Ph	H	H	H	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	CH_3	F	OH	H	H	NH_2
III-40-8	p-I-Ph	*	H	*	^{n}Bu	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00146

Table III-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-41-1	CH_3	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-2	CH_3	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-5	CH_3	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-41-8	CH_3	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-42-1	Et	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-2	Et	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-5	Et	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-42-8	Et	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-43-1	^{i}Pr	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-2	^{i}Pr	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-43-8	^{i}Pr	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00147

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-44-1	^{t}Bu	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-2	^{t}Bu	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-44-8	^{t}Bu	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-45-1	Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-2	Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-5	Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-45-8	Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-46-1	p-Me-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-2	p-Me-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00148

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-46-8	p-Me-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-47-1	p-F-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-2	p-F-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-47-8	p-F-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-48-1	p-Cl-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-48-8	p-Cl-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-49-1	p-Br-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-2	p-Br-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00149

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-49-8	p-Br-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table III-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
III-50-1	p-I-Ph	H	H	H	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-2	p-I-Ph	H	H	CH_3	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	CH_3	F	OH	H	H	NH_2
III-50-8	p-I-Ph	*	H	*	Bz	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

5



IV

Table IV-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-1-1	CH_3	H	H	H	CH_3	H	F	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00150

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-1-2	CH_3	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-1-8	CH_3	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-2-1	Et	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-2-2	Et	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-5	Et	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-2-8	Et	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-3-1	^{i}Pr	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-3-8	^{i}Pr	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-4.

Gilead Exhibit 2026
PGR2023-00046 - Page 00151

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-4-1	'Bu	H	H	H	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-2	'Bu	H	H	CH_3	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-3	'Bu	H	H	$CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-4	'Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-5	'Bu	H	H	CH_2Ph	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-6	'Bu	H	H	CH_2-indol-3-yl	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-7	'Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	CH_3	F	OH	H	H	NH_2
IV-4-8	'Bu	*	H	*	CH_3	H	CH_3	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-5-1	Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-5-2	Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-5-8	Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-6-1	p-Me-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-6-8	p-Me-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00152

Table IV-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-7-1	p-F-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-7-20	p-F-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-9-1	p-Br-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-9-20	p-Br-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00153

Table IV-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-10-1	p-I-Ph	H	H	H	CH_3	H	F	F	OH	H	H	NH_2
IV-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	F	OH	H	H	NH_2
IV-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	F	OH	H	H	NH_2
IV-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	F	OH	H	H	NH_2
IV-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	F	OH	H	H	NH_2
IV-10-8	p-I-Ph	*	H	*	CH_3	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-11-1	CH_3	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-11-2	CH_3	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-11-5	CH_3	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-11-8	CH_3	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-12-1	Et	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-12-2	Et	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-12-5	Et	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-12-8	Et	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00154

Table IV-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-13-1	^{i}Pr	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-13-2	^{i}Pr	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-13-8	^{i}Pr	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-14-1	^{t}Bu	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-14-2	^{t}Bu	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-14-8	^{t}Bu	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Gilead Exhibit 2026
PGR2023-00046 - Page 00155

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-15-1	Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-15-2	Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-15-5	Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-15-8	Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-16-1	p-Me-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-16-8	p-Me-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-17-1	p-F-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-17-2	p-F-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-17-8	p-F-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00156

Table IV-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-18-1	p-Cl-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-18-8	p-Cl-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-19-1	p-Br-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-19-8	p-Br-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-20-1	p-I-Ph	H	H	H	Et	H	F	F	OH	H	H	NH_2
IV-20-2	p-I-Ph	H	H	CH_3	Et	H	F	F	OH	H	H	NH_2
IV-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	F	OH	H	H	NH_2
IV-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	F	OH	H	H	NH_2
IV-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	F	OH	H	H	NH_2
IV-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	F	OH	H	H	NH_2
IV-20-8	p-I-Ph	*	H	*	Et	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00157

Table IV-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-21-1	CH_3	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-21-8	CH_3	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-22-1	Et	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-2	Et	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-22-8	Et	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00158

Table IV-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-25-1	Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-25-8	Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00159

Table IV-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00160

Table IV-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	F	OH	H	H	NH_2
IV-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-31-1	CH_3	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-31-8	CH_3	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-32-1	Et	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-2	Et	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-32-8	Et	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00161

Table IV-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-35-1	Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-35-8	Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00162

Table IV-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00163

Table IV-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	F	OH	H	H	NH_2
IV-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-41-1	CH_3	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-41-2	CH_3	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-41-8	CH_3	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00164

Table IV-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-42-1	Et	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-42-2	Et	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-42-5	Et	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-42-8	Et	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-43-1	^{i}Pr	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-43-8	^{i}Pr	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-44-1	^{t}Bu	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-44-8	^{t}Bu	*	H	*	Bz	H	F	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00165

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-45-1	Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-45-2	Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-45-5	Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-45-8	Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-46-1	p-Me-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-46-8	p-Me-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-47-1	p-F-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00166

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-47-8	p-F-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-48-1	p-Cl-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-48-8	p-Cl-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table IV-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-49-1	p-Br-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2
IV-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-49-8	p-Br-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.


Table IV-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-50-1	p-I-Ph	H	H	H	Bz	H	F	F	OH	H	H	NH_2
IV-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	F	OH	H	H	NH_2
IV-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	F	OH	H	H	NH_2
IV-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00167

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
IV-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	F	OH	H	H	NH_2
IV-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	F	OH	H	H	NH_2
IV-50-8	p-I-Ph	*	H	*	Bz	H	F	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



V

Table V-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-1-1	CH_3	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-1-2	CH_3	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-1-5	CH_3	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-1-8	CH_3	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-2-1	Et	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-2-2	Et	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00168

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-2-5	Et	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-2-8	Et	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-3-1	^{i}Pr	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-3-2	^{i}Pr	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-3-8	^{i}Pr	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-4-1	^{t}Bu	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-4-2	^{t}Bu	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-4-8	^{t}Bu	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Gilead Exhibit 2026
PGR2023-00046 - Page 00169

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-5-1	Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-5-2	Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-5-5	Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-5-8	Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-6-1	p-Me-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-6-8	p-Me-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-7-1	p-F-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-7-2	p-F-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-7-20	p-F-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00170

Table V-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-8-1	p-Cl-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-8-8	p-Cl-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-9-1	p-Br-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-9-20	p-Br-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-10-1	p-I-Ph	H	H	H	CH_3	H	H	F	OH	H	H	NH_2
V-10-2	p-I-Ph	H	H	CH_3	CH_3	H	H	F	OH	H	H	NH_2
V-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	H	F	OH	H	H	NH_2
V-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	H	F	OH	H	H	NH_2
V-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	H	F	OH	H	H	NH_2
V-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	H	F	OH	H	H	NH_2
V-10-8	p-I-Ph	*	H	*	CH_3	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00171

Table V-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-11-1	CH_3	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-11-2	CH_3	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-11-5	CH_3	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-11-8	CH_3	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-12-1	Et	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-12-2	Et	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-12-5	Et	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-12-8	Et	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-13-1	^{i}Pr	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-13-2	^{i}Pr	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-13-8	^{i}Pr	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00172

Table V-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-14-1	^{t}Bu	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-14-2	^{t}Bu	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-14-8	^{t}Bu	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-15-1	Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-15-2	Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-15-5	Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-15-8	Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-16-1	p-Me-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-16-2	p-Me-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-16-8	p-Me-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00173

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-17-1	p-F-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-17-2	p-F-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-17-8	p-F-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-18-1	p-Cl-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-18-2	p-Cl-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-18-8	p-Cl-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-19-1	p-Br-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-19-2	p-Br-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00174

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-19-8	p-Br-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-20-1	p-I-Ph	H	H	H	Et	H	H	F	OH	H	H	NH_2
V-20-2	p-I-Ph	H	H	CH_3	Et	H	H	F	OH	H	H	NH_2
V-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	H	F	OH	H	H	NH_2
V-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	H	F	OH	H	H	NH_2
V-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	H	F	OH	H	H	NH_2
V-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	H	F	OH	H	H	NH_2
V-20-8	p-I-Ph	*	H	*	Et	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-21-1	CH_3	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-2	CH_3	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-21-8	CH_3	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-22-1	Et	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-2	Et	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00175

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-22-8	Et	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-23-1	^{i}Pr	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-23-8	^{i}Pr	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-24-1	^{t}Bu	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-24-8	^{t}Bu	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-25-1	Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-2	Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00176

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-25-8	Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-27-1	p-F-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-27-8	p-F-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00177

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-30-1	p-I-Ph	H	H	H	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	H	F	OH	H	H	NH_2
V-30-8	p-I-Ph	*	H	*	^{i}Pr	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-31.

Gilead Exhibit 2026
PGR2023-00046 - Page 00178

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-31-1	CH_3	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-2	CH_3	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-31-8	CH_3	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-32-1	Et	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-2	Et	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-32-8	Et	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-33-1	^{i}Pr	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-33-8	^{i}Pr	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00179

Table V-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-34-1	^{t}Bu	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-34-8	^{t}Bu	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-35-1	Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-2	Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-35-8	Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00180

Table V-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-37-1	p-F-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-37-8	p-F-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00181

Table V-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-40-1	p-I-Ph	H	H	H	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	H	F	OH	H	H	NH_2
V-40-8	p-I-Ph	*	H	*	^{n}Bu	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-41-1	CH_3	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-41-2	CH_3	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-41-5	CH_3	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-41-8	CH_3	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-42-1	Et	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-42-2	Et	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-42-5	Et	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-42-8	Et	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00182

Table V-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-43-1	^{i}Pr	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-43-2	^{i}Pr	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-43-8	^{i}Pr	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-44-1	^{t}Bu	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-44-2	^{t}Bu	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-44-8	^{t}Bu	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-45-1	Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-45-2	Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-45-5	Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-45-8	Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00183

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-46-1	p-Me-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-46-2	p-Me-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-46-8	p-Me-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-47-1	p-F-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-47-2	p-F-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-47-8	p-F-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-48-1	p-Cl-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00184

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-48-8	p-Cl-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-49-1	p-Br-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-49-2	p-Br-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-49-8	p-Br-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2


*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table V-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
V-50-1	p-I-Ph	H	H	H	Bz	H	H	F	OH	H	H	NH_2
V-50-2	p-I-Ph	H	H	CH_3	Bz	H	H	F	OH	H	H	NH_2
V-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	H	F	OH	H	H	NH_2
V-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	H	F	OH	H	H	NH_2
V-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	H	F	OH	H	H	NH_2
V-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	H	F	OH	H	H	NH_2
V-50-8	p-I-Ph	*	H	*	Bz	H	H	F	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00185



VI

Table VI-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-1-1	CH_3	H	H	H	CH_3	H	F	H	OH	H	H
VI-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-1-8	CH_3	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-2-1	Et	H	H	H	CH_3	H	F	H	OH	H	H
VI-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-2-8	Et	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00186

Table VI-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH	H	H
VI-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH	H	H
VI-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-5.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-5-1	Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-5-8	Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00187

Table VI-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00188

Table VI-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH	H	H
VI-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H
VI-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H
VI-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H
VI-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H
VI-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H
VI-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-11-1	CH_3	H	H	H	Et	H	F	H	OH	H	H
VI-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH	H	H
VI-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-11-8	CH_3	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00189

Table VI-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-12-1	Et	H	H	H	Et	H	F	H	OH	H	H
VI-12-2	Et	H	H	CH_3	Et	H	F	H	OH	H	H
VI-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-12-8	Et	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH	H	H
VI-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH	H	H
VI-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-14-1	^{t}Bu	H	H	H	Et	H	F	H	OH	H	H
VI-14-2	^{t}Bu	H	H	CH_3	Et	H	F	H	OH	H	H
VI-14-3	^{t}Bu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-14-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-14-5	^{t}Bu	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-14-6	^{t}Bu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-14-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-14-8	^{t}Bu	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00190

Table VI-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-15-1	Ph	H	H	H	Et	H	F	H	OH	H	H
VI-15-2	Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-15-8	Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00191

Table VI-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH	H	H
VI-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH	H	H
VI-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H
VI-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H
VI-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H
VI-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H
VI-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00192

Table VI-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00193

Table VI-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00194

Table VI-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00195

Table VI-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H
VI-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H
VI-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H
VI-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H
VI-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H
VI-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H
VI-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00196

Table VI-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00197

Table VI-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-37.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-37-1	p-F-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-37-2	p-F-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-37-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-37-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-37-5	p-F-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-37-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-37-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-37-8	p-F-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-38.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-38-1	p-Cl-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-38-2	p-Cl-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-38-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-38-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-38-5	p-Cl-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-38-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-38-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-38-8	p-Cl-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00198

Table VI-39.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-39-1	p-Br-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-39-2	p-Br-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-39-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-39-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-39-5	p-Br-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-39-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-39-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-39-8	p-Br-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-40.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-40-1	p-I-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H
VI-40-2	p-I-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H
VI-40-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-40-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H
VI-40-5	p-I-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H
VI-40-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H
VI-40-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H
VI-40-8	p-I-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-41.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-41-1	CH_3	H	H	H	Bz	H	F	H	OH	H	H
VI-41-2	CH_3	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-41-3	CH_3	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-41-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-41-5	CH_3	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-41-6	CH_3	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-41-7	CH_3	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-41-8	CH_3	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00199

Table VI-42.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-42-1	Et	H	H	H	Bz	H	F	H	OH	H	H
VI-42-2	Et	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-42-3	Et	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-42-4	Et	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-42-5	Et	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-42-6	Et	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-42-7	Et	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-42-8	Et	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-43.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-43-1	^{i}Pr	H	H	H	Bz	H	F	H	OH	H	H
VI-43-2	^{i}Pr	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-43-3	^{i}Pr	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-43-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-43-5	^{i}Pr	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-43-6	^{i}Pr	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-43-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-43-8	^{i}Pr	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-44.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-44-1	^{t}Bu	H	H	H	Bz	H	F	H	OH	H	H
VI-44-2	^{t}Bu	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-44-3	^{t}Bu	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-44-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-44-5	^{t}Bu	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-44-6	^{t}Bu	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-44-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-44-8	^{t}Bu	*	H	*	Bz	H	F	H	OH	H	H

Gilead Exhibit 2026
PGR2023-00046 - Page 00200

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-45.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-45-1	Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-45-2	Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-45-3	Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-45-4	Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-45-5	Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-45-6	Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-45-7	Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-45-8	Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-46.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-46-1	p-Me-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-46-2	p-Me-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-46-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-46-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-46-5	p-Me-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-46-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-46-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-46-8	p-Me-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-47.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-47-1	p-F-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-47-2	p-F-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-47-3	p-F-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-47-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-47-5	p-F-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-47-6	p-F-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-47-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H

Gilead Exhibit 2026
PGR2023-00046 - Page 00201

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-47-8	p-F-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-48.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-48-1	p-Cl-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-48-2	p-Cl-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-48-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-48-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-48-5	p-Cl-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-48-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-48-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-48-8	p-Cl-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VI-49.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-49-1	p-Br-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-49-2	p-Br-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-49-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-49-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-49-5	p-Br-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H
VI-49-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-49-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-49-8	p-Br-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.


Table VI-50.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-50-1	p-I-Ph	H	H	H	Bz	H	F	H	OH	H	H
VI-50-2	p-I-Ph	H	H	CH_3	Bz	H	F	H	OH	H	H
VI-50-3	p-I-Ph	H	H	$CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-50-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Bz	H	F	H	OH	H	H
VI-50-5	p-I-Ph	H	H	CH_2Ph	Bz	H	F	H	OH	H	H

Gilead Exhibit 2026
PGR2023-00046 - Page 00202

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8
VI-50-6	p-I-Ph	H	H	CH_2-indol-3-yl	Bz	H	F	H	OH	H	H
VI-50-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Bz	H	F	H	OH	H	H
VI-50-8	p-I-Ph	*	H	*	Bz	H	F	H	OH	H	H

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.



VII

Table VII-1.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-1-1	CH_3	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-1-2	CH_3	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-1-3	CH_3	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-4	CH_3	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-5	CH_3	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-1-6	CH_3	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-1-7	CH_3	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-1-8	CH_3	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-2.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-2-1	Et	H	H	H	CH_3	H	F	H	OH	H	H	NH_2

Gilead Exhibit 2026
PGR2023-00046 - Page 00203

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-2-2	Et	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-2-3	Et	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-4	Et	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-5	Et	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-2-6	Et	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-2-7	Et	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-2-8	Et	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-3.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-3-1	^{i}Pr	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-3-2	^{i}Pr	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-3-3	^{i}Pr	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-5	^{i}Pr	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-3-6	^{i}Pr	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-3-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-3-8	^{i}Pr	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-4.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-4-1	^{t}Bu	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-4-2	^{t}Bu	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-4-3	^{t}Bu	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-5	^{t}Bu	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-4-6	^{t}Bu	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-4-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-4-8	^{t}Bu	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-5.

Gilead Exhibit 2026
PGR2023-00046 - Page 00204

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-5-1	Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-5-2	Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-5-3	Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-4	Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-5	Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-5-6	Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-5-7	Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-5-8	Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-6.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-6-1	p-Me-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-6-2	p-Me-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-6-3	p-Me-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-5	p-Me-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-6-6	p-Me-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-6-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-6-8	p-Me-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-7.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-7-1	p-F-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-7-2	p-F-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-7-3	p-F-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-6	p-F-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-7-7	p-F-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-7-8	p-F-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-7-20	p-F-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00205

Table VII-8.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-8-1	p-Cl-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-8-2	p-Cl-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-8-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-5	p-Cl-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-8-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-8-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-8-8	p-Cl-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-9.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-9-1	p-Br-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-9-2	p-Br-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-9-3	p-Br-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-6	p-Br-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-9-7	p-Br-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-9-8	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-9-20	p-Br-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-10.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-10-1	p-I-Ph	H	H	H	CH_3	H	F	H	OH	H	H	NH_2
VII-10-2	p-I-Ph	H	H	CH_3	CH_3	H	F	H	OH	H	H	NH_2
VII-10-3	p-I-Ph	H	H	$CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-5	p-I-Ph	H	H	CH_2Ph	CH_3	H	F	H	OH	H	H	NH_2
VII-10-6	p-I-Ph	H	H	CH_2-indol-3-yl	CH_3	H	F	H	OH	H	H	NH_2
VII-10-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	CH_3	H	F	H	OH	H	H	NH_2
VII-10-8	p-I-Ph	*	H	*	CH_3	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00206

Table VII-11.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-11-1	CH_3	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-11-2	CH_3	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-11-3	CH_3	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-11-4	CH_3	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-11-5	CH_3	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-11-6	CH_3	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-11-7	CH_3	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-11-8	CH_3	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-12.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-12-1	Et	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-12-2	Et	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-12-3	Et	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-12-4	Et	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-12-5	Et	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-12-6	Et	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-12-7	Et	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-12-8	Et	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-13.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-13-1	^{i}Pr	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-13-2	^{i}Pr	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-13-3	^{i}Pr	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-13-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-13-5	^{i}Pr	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-13-6	^{i}Pr	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-13-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-13-8	^{i}Pr	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00207

Table VII-14.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-14-1	tBu	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-14-2	tBu	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-14-3	tBu	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-14-4	tBu	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-14-5	tBu	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-14-6	tBu	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-14-7	tBu	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-14-8	tBu	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-15.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-15-1	Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-15-2	Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-15-3	Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-15-4	Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-15-5	Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-15-6	Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-15-7	Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-15-8	Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-16.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9

Gilead Exhibit 2026
PGR2023-00046 - Page 00208

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-16-1	p-Me-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-16-2	p-Me-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-16-3	p-Me-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-16-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-16-5	p-Me-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-16-6	p-Me-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-16-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-16-8	p-Me-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-17.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-17-1	p-F-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-17-2	p-F-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-17-3	p-F-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-17-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-17-5	p-F-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-17-6	p-F-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-17-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-17-8	p-F-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-18.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-18-1	p-Cl-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-18-2	p-Cl-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-18-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-18-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-18-5	p-Cl-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-18-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-18-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-18-8	p-Cl-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00209

Table VII-19.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-19-1	p-Br-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-19-2	p-Br-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-19-3	p-Br-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-19-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-19-5	p-Br-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-19-6	p-Br-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-19-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-19-8	p-Br-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-20.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-20-1	p-I-Ph	H	H	H	Et	H	F	H	OH	H	H	NH_2
VII-20-2	p-I-Ph	H	H	CH_3	Et	H	F	H	OH	H	H	NH_2
VII-20-3	p-I-Ph	H	H	$CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-20-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	Et	H	F	H	OH	H	H	NH_2
VII-20-5	p-I-Ph	H	H	CH_2Ph	Et	H	F	H	OH	H	H	NH_2
VII-20-6	p-I-Ph	H	H	CH_2-indol-3-yl	Et	H	F	H	OH	H	H	NH_2
VII-20-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	Et	H	F	H	OH	H	H	NH_2
VII-20-8	p-I-Ph	*	H	*	Et	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-21.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-21-1	CH_3	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-2	CH_3	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-3	CH_3	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-5	CH_3	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-6	CH_3	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-21-8	CH_3	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00210

Table VII-22.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-22-1	Et	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-2	Et	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-3	Et	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-4	Et	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-5	Et	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-6	Et	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-7	Et	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-22-8	Et	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-23.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-23-1	^{i}Pr	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-2	^{i}Pr	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-5	^{i}Pr	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-23-8	^{i}Pr	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-24.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-24-1	^{t}Bu	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-2	^{t}Bu	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-5	^{t}Bu	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-24-8	^{t}Bu	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00211

Table VII-25.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-25-1	Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-2	Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-3	Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-5	Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-6	Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-7	Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-25-8	Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-26.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-26-1	p-Me-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-2	p-Me-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-5	p-Me-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-26-8	p-Me-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-27.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-27-1	p-F-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-2	p-F-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-3	p-F-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-4	p-F-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-5	p-F-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-6	p-F-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-7	p-F-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-27-8	p-F-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00212

Table VII-28.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-28-1	p-Cl-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-2	p-Cl-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-3	p-Cl-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-4	p-Cl-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-5	p-Cl-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-6	p-Cl-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-7	p-Cl-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-28-8	p-Cl-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-29.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-29-1	p-Br-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-2	p-Br-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-3	p-Br-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-4	p-Br-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-5	p-Br-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-6	p-Br-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-7	p-Br-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-29-8	p-Br-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-30.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-30-1	p-I-Ph	H	H	H	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-2	p-I-Ph	H	H	CH_3	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-3	p-I-Ph	H	H	$CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-4	p-I-Ph	H	H	$CH_2CH(CH_3)_2$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-5	p-I-Ph	H	H	CH_2Ph	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-6	p-I-Ph	H	H	CH_2-indol-3-yl	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-7	p-I-Ph	H	H	$CH_2CH_2SCH_3$	^{i}Pr	H	F	H	OH	H	H	NH_2
VII-30-8	p-I-Ph	*	H	*	^{i}Pr	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00213

Table VII-31.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-31-1	CH_3	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-2	CH_3	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-3	CH_3	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-4	CH_3	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-5	CH_3	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-6	CH_3	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-7	CH_3	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-31-8	CH_3	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-32.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-32-1	Et	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-2	Et	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-3	Et	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-4	Et	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-5	Et	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-6	Et	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-7	Et	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-32-8	Et	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-33.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-33-1	^{i}Pr	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-2	^{i}Pr	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-3	^{i}Pr	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-4	^{i}Pr	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-5	^{i}Pr	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-6	^{i}Pr	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-7	^{i}Pr	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-33-8	^{i}Pr	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00214

Table VII-34.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-34-1	^{t}Bu	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-2	^{t}Bu	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-3	^{t}Bu	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-4	^{t}Bu	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-5	^{t}Bu	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-6	^{t}Bu	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-7	^{t}Bu	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-34-8	^{t}Bu	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-35.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-35-1	Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-2	Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-3	Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-4	Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-5	Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-6	Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-7	Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-35-8	Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Table VII-36.

№	R^1	R^2	R^{3a}	R^{3b}	R^4	R^5	R^6	X	Y	R^7	R^8	R^9
VII-36-1	p-Me-Ph	H	H	H	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-2	p-Me-Ph	H	H	CH_3	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-3	p-Me-Ph	H	H	$CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-4	p-Me-Ph	H	H	$CH_2CH(CH_3)_2$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-5	p-Me-Ph	H	H	CH_2Ph	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-6	p-Me-Ph	H	H	CH_2-indol-3-yl	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-7	p-Me-Ph	H	H	$CH_2CH_2SCH_3$	^{n}Bu	H	F	H	OH	H	H	NH_2
VII-36-8	p-Me-Ph	*	H	*	^{n}Bu	H	F	H	OH	H	H	NH_2

*R^2 and R^{3b} joined together by $(CH_2)_3$ to form five-membered ring.

Gilead Exhibit 2026
PGR2023-00046 - Page 00215