

L-CYSTEINE HYDROCHL

+

▼

←

→

↻

🏠

🔒

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cysteine

🔖

☆

🔍

🔗

⋮

INTERNET ARCHIVE

Wayback Machine

2 captures

24 Aug 2016 - 3 Apr 2017

Go

AUG

APR 03

MAY

2016

2017

2018

About this capture

www.drugsDB.eu

With permission of U.S. National Library of Medicine

DRUGS DataBase

You are here: Home > Prescription(RX) Drugs > L > L-cysteine Hydrochloride (Sandoz Inc)

Loading

Name:	L-cysteine Hydrochloride
Manufacturer:	Sandoz Inc
Category:	Prescription Marketed Drugs

Prescription Marketed Drugs Alphabetically

A| B| C| D| E| F| G| H| I| J| K| L| M| N| O| P| Q| R| S| T| U| V| W| X| Y| Z| 0-9

Categories:

Prescription(RX) Drugs

Over-the-counter (OTC) Drugs

Homeopathic Drugs

Animal Drugs

Feedback

L-CYSTEINE HYDROCHLORIDE - cysteine hydrochloride injection, solution

Sandoz Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

L-Cysteine Hydrochloride Injection, USP

0.5 g/10 mL (50 mg/mL)

PHARMACY BULK PACKAGE

NOT FOR DIRECT INFUSION

DESCRIPTION

L-Cysteine Hydrochloride Injection, USP, 50 mg/mL, is a sterile, nonpyrogenic solution. Each mL contains: 50 mg of L-Cysteine Hydrochloride Monohydrate USP; Water for Injection, USP q.s.; Air replaced with Nitrogen. pH 1.0-2.5

L-Cysteine is a sulfur-containing amino acid. In premixed solutions of crystalline amino acids, cysteine is relatively unstable over time, eventually converting to insoluble cystine. To avoid such precipitation, L-Cysteine Hydrochloride Injection USP is intended to be used as an additive with crystalline Amino Acid Injections immediately prior to administration to the patient.

The structural formula of Cysteine Hydrochloride Monohydrate USP is:

$$\begin{array}{c}
 \text{H} \\
 | \\
 \text{HSCH}_2 - \text{C} - \text{COOH} \cdot \text{HCl} \cdot \text{H}_2\text{O} \\
 | \\
 \text{NH}_2
 \end{array}$$

Molecular Weight

175.63

Molecular Formula

C₃H₇NO₂S·HCl·H₂O

Nexus Ex. 1005
Page 2 of 11

Nexus Ex. 1005
Page 3 of 11

Internet Archive Wayback Machine

2 captures

24 Aug 2016 - 3 Apr 2017

entries increase the potential of microbial and particulate contamination. The withdrawal of container contents should be accomplished without delay using aseptic technique. However, should this not be possible, a maximum time of 4 hours from initial closure entry is permitted to complete fluid transfer operations.

RECOMMENDED STORAGE CONDITIONS AFTER OPENING
Keep under laminar flow hood at room temperature. Any unused portion of the vial must be discarded within 4 hours after initial entry.

HOW SUPPLIED
L-Cysteine Hydrochloride Injection, USP (50 mg/mL) is supplied as follows:

PHARMACY BULK PACKAGE

NDC Number	Volume
66758-005-01	50 mL
66758-005-02	5 x 50 mL

Also available as:

SINGLE DOSE VIAL

NDC Number	Volume
66758-004-01	10 mL
66758-004-02	10 x 10 mL

Store at controlled room temperature 15°-30°C (59°-86°F) Do not freeze.
For Sandoz Inc. Customer Service, call [1-800-525-8747](tel:1-800-525-8747)

Rx only
Manufactured for:
SANDOZ
Princeton, NJ 08540
Revised: November 2009
L-028-00

Package Label - Principal Display Panel - 50 mL Vial
SANDOZ
Rx only **NDC 66758-005-01**
L-Cysteine
Hydrochloride Injection, USP
PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION
50 mg/mL
For IV Use Only After Dilution
Do Not Dispense As A Unit
50 mL
L-018-00
PHARMACY BULK PACKAGE

Internet Archive Wayback Machine

2 captures
24 Aug 2016 - 3 Apr 2017

Go

2016 APR 03 2017

https://web.archive.org/web/20170403170533/http://drugsdb.eu/drug.php?d=L-cystein

http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=Sandoz%20Inc&id=083386d8-0437-4ee0-90d4-440a5b5d03b5.x

SANDOZ
Rx only NDC 66758-005-01
L-Cysteine
Hydrochloride Injection, USP
PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION
50 mg/mL
For IV Use Only After Dilution
Do Not Dispense As A Unit
50 mL
L-018-03
SANDOZ VIAL 50ML 0.5000000000000000

Each mL contains: 50 mg L-Cysteine Hydrochloride Monohydrate, USP; Water for Injection, USP, q.s.; Air replaced with Nitrogen, pH 1.0-2.5
Directions for Use: See package insert.
Warning: For IV use only. Must be diluted before use. Contents should be dispensed promptly after initial closure puncture. Discard unused contents after 4 hours.
Store at controlled room temperature 15°-30°C (59°-86°F). Do not freeze.
Contains no more than 5,000 mcg/L of aluminum.
Manufactured for: Sandoz Inc., Princeton, NJ 08540
Date/Time Entered: NDC(2016) 00503386D8 0437-4EE0-90D4-440A5B5D03B5

Package Label - Principal Display Panel - 50 mL Vial Carton
SANDOZ
5 x 50 mL Vials NDC 66758-005-02
L-Cysteine Hydrochloride
Injection, USP
50 mg/mL
Pharmacy Bulk Package
Not For Direct Infusion
For IV Use Only After Dilution
Rx only

5 x 50 mL Vials
SANDOZ NDC 66758-005-02
L-Cysteine Hydrochloride
Injection, USP
50 mg/mL
Pharmacy Bulk Package
Not For Direct Infusion
For IV Use Only After Dilution
Rx only

L-CYSTEINE HYDROCHLORIDE
L-cysteine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66758-005
Route of Administration	INTRAVENOUS	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CYSTEINE HYDROCHLORIDE (CYSTEINE)	CYSTEINE HYDROCHLORIDE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER	

Product Characteristics

Color		Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#Item Code	Package Description	Multilevel Packaging
1 NDC:66758-005-02	5 VIAL, PHARMACY BULK PACKAGE (VIAL) in 1 CARTON	contains a VIAL, PHARMACY BULK PACKAGE (66758-005-01)
1 NDC:66758-005-01	50 mL in 1 VIAL, PHARMACY BULK PACKAGE	This package is contained within the CARTON (66758-005-02)

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		12/01/2008	

Labeler - Sandoz Inc (005387188)

Revised: 06/2010 Sandoz Inc

Source: <http://dailymed.nlm.nih.gov>
Reproduced with permission of U.S. National Library of Medicine

Copyright © 2017 Drugsdb.eu by Dionisios Fentas || [Terms of Use](#)

https://web.archive.org/web/20200301000000/http://drugsdb.eu/drug.php?d=L-cysteine%20

Internet Archive Wayback Machine

2 captures
24 Aug 2016 - 3 Apr 2017

2015 2016 2017

24

2016

2017

About this capture

www.drugsDB.eu

DRUGS DataBase

With permission of
U.S. National Library
of Medicine

You are here: Home > Prescription(RX) Drugs > L > L-cysteine Hydrochloride (Sandoz Inc)

Loading

Name: L-cysteine Hydrochloride

Manufacturer: Sandoz Inc

Category: Prescription Marketed Drugs

Prescription Marketed Drugs Alphabetically
A| B| C| D| E| F| G| H| I| J| K| L| M| N| O| P| Q| R|
S| T| U| V| W| X| Y| Z| 0-9

Categories:

Prescription(RX) Drugs
Over-the-counter (OTC) Drugs
Homeopathic Drugs
Animal Drugs
Feedback

L-CYSTEINE HYDROCHLORIDE - cysteine hydrochloride injection, solution
Sandoz Inc
Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

L-Cysteine Hydrochloride Injection, USP
0.5 g/10 mL (50 mg/mL)

PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION

DESCRIPTION
L-Cysteine Hydrochloride Injection, USP, 50 mg/mL, is a sterile, nonpyrogenic solution. Each mL contains: 50 mg of L-Cysteine Hydrochloride Monohydrate USP; Water for Injection, USP q.s.; Air replaced with Nitrogen. pH 1.0-2.5
L-Cysteine is a sulfur-containing amino acid. In premixed solutions of crystalline amino acids, cysteine is relatively unstable over time, eventually converting to insoluble cystine. To avoid such precipitation, L-Cysteine Hydrochloride Injection USP is intended to be used as an additive with crystalline Amino Acid Injections immediately prior to administration to the patient.
The structural formula of Cysteine Hydrochloride Monohydrate USP is:

$$\begin{array}{c} \text{H} \\ | \\ \text{HSCH}_2 - \text{C} - \text{COOH} \cdot \text{HCl} \cdot \text{H}_2\text{O} \end{array}$$

L-CYSTEINE HYDROCHL

https://web.archive.org/web/20160824090050/http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=5

Wayback Machine 2 captures 24 Aug 2016 - 3 Apr 2017

$$\text{HSCH}_2 - \text{C} - \text{COOH} \cdot \text{HCl} \cdot \text{H}_2\text{O}$$

$$\quad \quad \quad |$$

$$\quad \quad \quad \text{NH}_2$$

Molecular Weight 175.63 **Molecular Formula** $\text{C}_3\text{H}_7\text{NO}_2\text{S} \cdot \text{HCl} \cdot \text{H}_2\text{O}$

CLINICAL PHARMACOLOGY
 L-Cysteine is synthesized from methionine via the trans-sulfuration pathway in the adult, but newborn infants lack the enzyme necessary to effect this conversion. Therefore, L-Cysteine is generally considered to be an essential amino acid in infants.

INDICATIONS AND USAGE
 L-Cysteine Hydrochloride Injection, USP is intended for use only after dilution as an additive to Crystalline Amino Acid Injections to meet the intravenous amino acid nutritional requirements of infants receiving total parenteral nutrition.

CONTRAINDICATIONS
 This preparation should not be used in patients with hepatic coma or metabolic disorders involving impaired nitrogen utilization.

WARNINGS
 Peripheral intravenous infusion of amino acids may induce a rise in blood urea nitrogen (BUN) especially in patients with impaired hepatic or renal function. Appropriate laboratory tests should be performed periodically and infusion discontinued if BUN levels exceed normal postprandial limits and continue to rise. It should be noted that a modest rise in BUN normally occurs as a result of increased protein intake.
 Administration of amino acid solutions to a patient with hepatic insufficiency may result in serum amino acid imbalances, metabolic alkalosis, prerenal azotemia, hyperammonemia, stupor and coma.
 Administration of amino acid solutions in the presence of impaired renal function may augment an increasing BUN, as does any protein dietary component.
 Solutions containing sodium ion should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency, and in clinical states in which there exists edema with sodium retention.
 Solutions which contain potassium ion should be used with great care, if at all, in patients with hyperkalemia, severe renal failure and in conditions in which potassium retention is present.
 Solutions containing acetate ion should be used with great care in patients with metabolic or respiratory alkalosis. Acetate should be administered with great care in those conditions in which there is an increased level or an impaired utilization of this ion such as severe hepatic insufficiency.
 Hyperammonemia is of special significance in infants, as it can result in mental retardation. Therefore it is essential that blood ammonia levels be measured frequently in infants.
 Instances of asymptomatic hyperammonemia have been reported in patients without overt liver dysfunction. The mechanisms of this reaction are not clearly defined but may involve genetic defects and immature or subclinically impaired liver function.

Frequent Clinical Evaluation and Laboratory Determinations are Necessary for Proper Monitoring During Administration. Blood studies should include glucose, urea nitrogen, serum

L-CYSTEINE HYDROCHL x + v

https://web.archive.org/web/20160824090050/http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=S

Internet Archive Wayback Machine

2 captures
24 Aug 2016 - 3 Apr 2017

2015 2016 2017

24

electrolytes, ammonia, cholesterol, acid-base balance, serum proteins, kidney and liver function tests, osmolality and hemogram. White blood count and blood cultures are to be determined if indicated. Urinary osmolality and glucose should be determined frequently. Safe use during pregnancy has not been established, therefore, infusion of amino acids should be undertaken during pregnancy only when this is deemed essential to the patients' welfare, as judged by the physician.

WARNING: *This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum. Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration*

PRECAUTIONS

Special care must be taken when administering hypertonic glucose to provide calories in diabetic or prediabetic patients. Because of its antianabolic activity, concurrent administration of tetracycline may reduce the nitrogen sparing effects of infused amino acids. Do not withdraw venous blood for blood chemistries through the peripheral infusion site, as interference with estimations of nitrogen containing substances may occur. Intravenous feeding regimens which include amino acids should be used with caution in patients with a history of renal disease, pulmonary disease, or with cardiac insufficiency so as to avoid excessive fluid accumulation. The effect of infusion of amino acids, without dextrose, upon carbohydrate metabolism of children is not known at this time. Nitrogen intake should be carefully monitored in patients with impaired renal function. For long-term total nutrition, or if a patient has inadequate fat stores, it is essential to provide adequate exogenous calories concurrently with the amino acids. Concentrated dextrose solutions are an effective source of such calories. Such strongly hypertonic nutrient solutions should be administered through an indwelling intravenous catheter with the tip located in the superior vena cava.

ADVERSE REACTIONS

Local reactions consisting of a warm sensation, erythema, phlebitis and thrombosis at the infusion site have occurred with peripheral intravenous infusion of amino acids, particularly if the other substances, such as antibiotics, are also administered through the same site. In such cases the infusion site should be changed promptly to another vein. Use of large peripheral veins, inline filters, and slowing the rate of infusion may reduce the incidence of local venous irritation. Electrolyte additives should be spread throughout the day. Irritating additive medications may need to be injected at another venous site. Generalized flushing, fever and nausea also have been reported during peripheral infusions of amino acid solutions.

Drug Abuse and Dependence

None known.

DOSAGE AND ADMINISTRATION

L-Cysteine Hydrochloride Injection USP is intended for use only after dilution in Crystalline Amino Acid

L-CYSTEINE HYDROCHL
+

[https://web.archive.org/web/20160824090050/http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=S](https://web.archive.org/web/20160824090050/http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=Sandoz%20Inc&id=083366d6-0437-4ee0-90d4-440a5b5d03b5.x)
Go
JUL
AUG
APR
24
2015
2016
2017
About this capture

<http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=Sandoz%20Inc&id=083366d6-0437-4ee0-90d4-440a5b5d03b5.x>

2 captures
24 Aug 2016 - 3 Apr 2017

Injection. Each 0.5 gram of L-Cysteine Hydrochloride Monohydrate should be combined aseptically with 12.5 grams of Crystalline Amino Acid Injection, such as that present in 250 mL of 5% Crystalline Amino Acid Injection. The admixture is then diluted with 250 mL of dextrose 50% or such lesser volume as indicated. Equal volumes of 5% Crystalline Amino Acid Injection and dextrose 50% produce a final solution which contains Crystalline Amino Acid Injection 2.5% in dextrose 25%, which is suitable for administration by central venous infusion. Administration of the final admixture should begin within one hour of mixing. Otherwise, the admixture should be refrigerated immediately and used within 24 hours of the time of mixing. For the recommended rate of administration, see the Crystalline Amino Acid Injection package insert.
Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

DIRECTIONS FOR PROPER USE OF PHARMACY BULK PACKAGE
The pharmacy bulk package is for use in a Pharmacy Admixture Service only.
Use of this product is restricted to a suitable work area, such as a laminar flow hood. Prior to entering the vial, remove the flip-off seal and cleanse the rubber closure with a suitable antiseptic agent.
The container closure may be penetrated only one time, utilizing a suitable sterile transfer device or dispensing set which allows measured distribution of the contents. The date and time the vial was initially opened should be recorded in the space provided on the label. Transfer individual doses(s) to appropriate intravenous infusion solutions. Use of a syringe with needle is not recommended. Multiple entries increase the potential of microbial and particulate contamination.
The withdrawal of container contents should be accomplished without delay using aseptic technique. However, should this not be possible, a maximum time of 4 hours from initial closure entry is permitted to complete fluid transfer operations.

RECOMMENDED STORAGE CONDITIONS AFTER OPENING
Keep under laminar flow hood at room temperature. Any unused portion of the vial must be discarded within 4 hours after initial entry.

HOW SUPPLIED
L-Cysteine Hydrochloride Injection, USP (50 mg/mL) is supplied as follows:

PHARMACY BULK PACKAGE

NDC Number	Volume
66758-005-01	50 mL
66758-005-02	5 × 50 mL

Also available as:

SINGLE DOSE VIAL

NDC Number	Volume
66758-004-01	10 mL
66758-004-02	10 × 10 mL


Store at controlled room temperature 15°-30°C (59°-86°F) Do not freeze.
For Sandoz Inc. Customer Service, call [1-800-525-8747](tel:1-800-525-8747)

Rx only
Manufactured for:
SANDOZ
Princeton, NJ 08540

Internet Archive
Wayback Machine
2 captures
24 Aug 2016 - 3 Apr 2017

Revised: November 2009
L-028-00

Package Label - Principal Display Panel - 50 mL Vial
SANDOZ
Rx only NDC 66758-005-01
L-Cysteine
Hydrochloride Injection, USP
PHARMACY BULK PACKAGE
NOT FOR DIRECT INFUSION
50 mg/mL
For IV Use Only After Dilution
Do Not Dispense As A Unit
50 mL
L-018-00
PHARMACY BULK PACKAGE



Each mL contains: 50 mg L-Cysteine Hydrochloride Monohydrate, USP; Water for Injection, USP, q.s.; Air replaced with Nitrogen, pH 1.0-2.5
Directions for Use: See package insert.
Warning: For IV use only. Must be diluted before use. Contents should be dispensed promptly after initial closure puncture. Discard unused contents after 4 hours.
Store at controlled room temperature 15°-30°C (59°-86°F). Do not freeze.
Contains no more than 5,000 mcg/L of aluminum.
Manufactured for: Sandoz Inc., Princeton, NJ 08540
Date/Time Entered: _____

Package Label - Principal Display Panel - 50 mL Vial Carton
SANDOZ
5 × 50 mL Vials NDC 66758-005-02
L-Cysteine Hydrochloride
Injection, USP
50 mg/mL
Pharmacy Bulk Package
Not For Direct Infusion
For IV Use Only After Dilution
Rx only

Internet Archive
Wayback Machine


2 captures
24 Aug 2016 - 5 Apr 2017

https://web.archive.org/web/20160824090050/http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=S

http://drugsdb.eu/drug.php?d=L-cysteine%20Hydrochloride&m=Sandoz%20Inc&id=083366d6-0437-4ee0-90d4-440a5b5d03b5.x

JUL 2015 AUG 2016 APR 2017

For IV Use Only After Dilution
Rx only



L-CYSTEINE HYDROCHLORIDE
l-cysteine hydrochloride injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66758-005
Route of Administration	INTRAVENOUS	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CYSTEINE HYDROCHLORIDE (CYSTEINE)	CYSTEINE HYDROCHLORIDE	50 mg in 1 mL	
Inactive Ingredients			
Ingredient Name	Strength		
WATER			
Product Characteristics			
Color	Score		
Shape	Size		
Flavor	Imprint Code		
Contains			
Packaging			
#Item Code	Package Description	Multilevel Packaging	
1 NDC:66758-005-02	5 VIAL, PHARMACY BULK PACKAGE (VIAL) in 1 CARTON	contains a VIAL, PHARMACY BULK PACKAGE (66758-005-01)	
1 NDC:66758-005-01	50 mL in 1 VIAL, PHARMACY BULK PACKAGE	This package is contained within the CARTON (66758-005-02)	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		12/01/2008	
Labeler - Sandoz Inc (005387188)			
Revised: 06/2010		Sandoz Inc	
Source: http://dailymed.nlm.nih.gov Reproduced with permission of U.S. National Library of Medicine			
Copyright © 2016 Drugsdb.eu by Dionisios Fentas Terms of Use			