



US005689034A

United States Patent [19]
Fullerton

[11] **Patent Number:** **5,689,034**
[45] **Date of Patent:** **Nov. 18, 1997**

[54] **INBRED MAIZE LINE PH24E**

[75] **Inventor:** Samuel Gregg Fullerton, Huron, S. Dak.

[73] **Assignee:** Pioneer Hi-Bred International, Inc., Des Moines, Iowa

[21] **Appl. No.:** 523,660

[22] **Filed:** Sep. 5, 1995

[51] **Int. Cl.⁶** A01H 5/00; A01H 4/00; A01H 1/00; C12N 5/04

[52] **U.S. Cl.** 800/200; 800/250; 800/DIG. 56; 435/240.4; 435/240.49; 435/240.5; 47/58; 47/DIG. 1

[58] **Field of Search** 800/200, 205, 800/235, 250, DIG. 56; 435/204.1, 240.4, 240.47, 240.49, 240.5; 47/58, 58.01, 58.03, DIG. 1

[56] **References Cited**

U.S. PATENT DOCUMENTS

5,276,265 1/1994 Fullerton 800/200

FOREIGN PATENT DOCUMENTS

1603390 11/1985 European Pat. Off. .

OTHER PUBLICATIONS

Conger, B.V., et al. (1987) "Somatic Embryogenesis From Cultured Leaf Segments of *Zea mays*", *Plant Cell Reports*, 6:345-347.

Duncan, D.R., et al. (1985) "The Production of Callus Capable of Plant Regeneration From Immature Embryos of Numerous *Zea mays* Genotypes", *Planta*, 165:322-332.

Edallo, et al. (1981) "Chromosomal Variation and Frequency of Spontaneous Mutation Associated with in Vitro Culture and Plant Regeneration in Maize", *Maydica*, XXVI: 39-56.

Green, et al., (1975) "Plant Regeneration From Tissue Cultures of Maize", *Crop Science*, vol. 15, pp. 417-421.

Green, C.E., et al. (1982) "Plant Regeneration in Tissue Cultures of Maize", *Maize for Biological Research*, pp. 367-372.

Hallauer, A.R. et al. (1988) "Corn Breeding" *Corn and Corn Improvement*, No. 18, pp. 463-481.

Meghji, M.R., et al. (1984) "Imbreeding Depression, Inbred & Hybrid Grain Yields, and Other Traits of Maize Genotypes Representing Three Eras", *Crop Science*, vol. 24, pp. 545-549.

Phillips, et al. (1988) "Cell/Tissue Culture and in Vitro Manipulation", *Corn & Corn Improvement*, 3rd Ed., ASA Publication, No. 18, pp. 345-387.

Poehlman et al., (1995) *Breeding Field Crop*, 4th Ed., Iowa State University Press, Ames, IA., pp. 132-155 and 321-344.

Rao, K.V., et al., (1986) "Somatic Embryogenesis in Glume Callus Cultures", *Maize Genetics Cooperative Newsletter*, No. 60, pp. 64-65.

Sass, John F. (1977) "Morphology", *Corn & Corn Improvement*, ASA Publication, Madison, Wisconsin, pp. 89-109.

Songstad, D.D. et al. (1988) "Effect of ACC (1-aminocyclopropane-1-carboxylic acid), Silver Nitrate & Norbonadiene on Plant Regeneration From Maize Callus Cultures", *Plant Cell Reports*, 7:262-265.

Tomes, et al. (1985) "The Effect of Parental Genotype on Initiation of Embryogenic Callus From Elite Maize (*Zea mays* L.) Germplasm", *Theor. Appl. Genet.*, vol. 70, pp. 505-509.

Troyer, et al. (1985) "Selection for Early Flowering in Corn: 10 Late Synthetics", *Crop Science*, vol. 25, pp. 695-697.

Umbeck, et al. (1983) "Reversion of Male-Sterile T-Cytoplasm Maize to Male Fertility in Tissue Culture", *Crop Science*, vol. 23, pp. 584-588.

Wright, Harold (1980) "Commercial Hybrid Seed Production", *Hybridization of Crop Plants*, Ch. 8: 161-176.

Wych, Robert D. (1988) "Production of Hybrid Seed", *Corn and Corn Improvement*, Ch. 9, pp. 565-607.

Lee, Michael (1994) "Inbred Lines of Maize and Their Molecular Markers", *The Maize Handbook* Ch. 65:423-432.

Boppenmaier, et al., "Comparisons Among Strains of Inbreds for RFLPs", *Maize Genetics Cooperative Newsletter*, 65:1991, p. 90.

Coe et al. In *Corn and Corn Improvement*. Third Edition. Sprague et al., eds. Ch. 3: 81-137 ASA-CSSA-SSSA Jan. 1988.

Hallauer et al. In *Corn and Corn Improvement*. Third Edition. Sprague et al., eds. Ch. 8:463-564 ASA-CSSA-SSSA Jan. 1988.

Primary Examiner—Gary Benzion

Attorney, Agent, or Firm—Zarley, McKee, Thomte, Voorhees & Sease

[57]

ABSTRACT

An inbred maize line, designated PH24E, the plants and seeds of inbred maize line PH24E, methods for producing a maize plant produced by crossing the inbred line PH24E with itself or with another maize plant, and hybrid maize seeds and plants produced by crossing the inbred line PH24E with another maize line or plant.

14 Claims, No Drawings

INBRED MAIZE LINE PH24E

FIELD OF THE INVENTION

This invention is in the field of maize breeding, specifically relating to an inbred maize line designated PH24E.

BACKGROUND OF THE INVENTION

The goal of plant breeding is to combine in a single variety or hybrid various desirable traits. For field crops, these traits may include resistance to diseases and insects, tolerance to heat and drought, reducing the time to crop maturity, greater yield, and better agronomic quality. With mechanical harvesting of many crops, uniformity of plant characteristics such as germination and stand establishment, growth rate, maturity, and plant and ear height, is important.

Field crops are bred through techniques that take advantage of the plant's method of pollination. A plant is self-pollinated if pollen from one flower is transferred to the same or another flower of the same plant. A plant is cross-pollinated if the pollen comes from a flower on a different plant.

Plants that have been self-pollinated and selected for type for many generations become homozygous at almost all gene loci and produce a uniform population of true breeding progeny. A cross between two different homozygous lines produces a uniform population of hybrid plants that may be heterozygous for many gene loci. A cross of two plants each heterozygous at a number of gene loci will produce a population of hybrid plants that differ genetically and will not be uniform.

Maize (*zea mays* L.), often referred to as corn in the United States, can be bred by both self-pollination and cross-pollination techniques. Maize has separate male and female flowers on the same plant, located on the tassel and the ear, respectively. Natural pollination occurs in maize when wind blows pollen from the tassels to the silks that protrude from the tops of the ears.

A reliable method of controlling male fertility in plants offers the opportunity for improved plant breeding. This is especially true for development of maize hybrids, which relies upon some sort of male sterility system. There are several options for controlling male fertility available to breeders, such as: manual or mechanical emasculation (or detasseling), cytoplasmic male sterility, genetic male sterility, gametocides and the like.

Hybrid maize seed is typically produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two inbred varieties of maize are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female). Providing that there is sufficient isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male), and the resulting seed is therefore hybrid and will form hybrid plants.

The laborious, and occasionally unreliable, detasseling process can be avoided by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of factors resulting from the cytoplasmic, as opposed to the nuclear, genome. Thus, this characteristic is inherited exclusively through the female parent in maize plants, since only the female provides cytoplasm to the fertilized seed. CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile. Usually seed from detasseled fertile maize and

CMS produced seed of the same hybrid are blended to insure that adequate pollen loads are available for fertilization when the hybrid plants are grown.

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Pat. Nos. 4,654,465 and 4,727,219 to Brar et al. and chromosomal translocations as described by Patterson in U.S. Pat. Nos. 3,861,709 and 3,710,511. These and all patents referred to are incorporated by reference. In addition to these methods, Albertsen et al., of Pioneer Hi-Bred, U.S. patent application Ser. No. 07/848,433, have developed a system of nuclear male sterility which includes: identifying a gene which is critical to male fertility; silencing this native gene which is critical to male fertility; removing the native promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered gene back into the plant; and thus creating a plant that is male sterile because the inducible promoter is not "on" resulting in the male fertility gene not being transcribed. Fertility is restored by inducing, or turning "on", the promoter, which in turn allows the gene that confers male fertility to be transcribed.

There are many other methods of conferring genetic male sterility in the art, each with its own benefits and drawbacks. These methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an antisense system in which a gene critical to fertility is identified and an antisense to that gene is inserted in the plant (see: Fabinjanski, et al. EPO 89/3010153.8 publication no. 329, 308 and PCT application PCT/CA90/00037 published as WO 90/08828).

Another system useful in controlling male sterility makes use of gametocides. Gametocides are not a genetic system, but rather a topical application of chemicals. These chemicals affect cells that are critical to male fertility. The application of these chemicals affects fertility in the plants only for the growing season in which the gametocide is applied (see Carlson, Glenn R., U.S. Pat. No. 4,936,904). Application of the gametocide, timing of the application and genotype specificity often limit the usefulness of the approach.

The use of male sterile inbreds is but one factor in the production of maize hybrids. The development of maize hybrids requires, in general, the development of homozygous inbred lines, the crossing of these lines; and the evaluation of the crosses. Pedigree breeding and recurrent selection breeding methods are used to develop inbred lines from breeding populations. Breeding programs combine the genetic backgrounds from two or more inbred lines or various other broad-based sources into breeding pools from which new inbred lines are developed by selfing and selection of desired phenotypes. The new inbreds are crossed with other inbred lines and the hybrids from these crosses are evaluated to determine which of those have commercial potential.

Pedigree breeding starts with the crossing of two genotypes, each of which may have one or more desirable characteristics that is lacking in the other or which complements the other. If the two original parents do not provide all the desired characteristics, other sources can be included in the breeding population. In the pedigree method, superior plants are selfed and selected in successive generations. In the succeeding generations the heterozygous condition gives way to homogeneous lines as a result of self-pollination and selection. Typically in the pedigree method of breeding five

or more generations of selfing and selection is practiced: $F_1 \rightarrow F_2$; $F_3 \rightarrow F_4$; $F_4 \rightarrow F_5$, etc.

Recurrent selection breeding, backcrossing for example, can be used to improve an inbred line. Backcrossing can be used to transfer a specific desirable trait from one inbred or source to an inbred that lacks that trait. This can be accomplished, for example, by first crossing a superior inbred (recurrent parent) to a donor inbred (non-recurrent parent), that carries the appropriate gene(s) for the trait in question. The progeny of this cross is then mated back to the superior recurrent parent followed by selection in the resultant progeny for the desired trait to be transferred from the non-recurrent parent. After five or more backcross generations with selection for the desired trait, the progeny will be heterozygous for loci controlling the characteristic being transferred, but will be like the superior parent for most or almost all other genes. The last backcross generation is then selfed to give pure breeding progeny for the gene(s) being transferred.

A single cross hybrid maize variety is the cross of two inbred lines, each of which has a genotype that complements the genotype of the other. The hybrid progeny of the first generation is designated F_1 . In the development of hybrids only the F_1 hybrid plants are sought. Preferred F_1 hybrids are more vigorous than their inbred parents. This hybrid vigor, or heterosis, can be manifested in many polygenic traits, including increased vegetative growth and increased yield.

The development of a hybrid maize variety involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, although different from each other, breed true and are highly uniform; and (3) crossing the selected inbred lines with different inbred lines to produce the hybrid progeny (F_1). During the inbreeding process in maize, the vigor of the lines decreases. Vigor is restored when two different inbred lines are crossed to produce the hybrid progeny (F_1). An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid between a defined pair of inbreds will always be the same. Once the inbreds that give a superior hybrid have been identified, the hybrid seed can be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained.

A single cross hybrid is produced when two inbred lines are crossed to produce the F_1 progeny. A double cross hybrid is produced from four inbred lines crossed in pairs ($A \times B$ and $C \times D$) and then the two F_1 hybrids are crossed again ($A \times B$) $\times$ ($C \times D$). Much of the hybrid vigor exhibited by F_1 hybrids is lost in the next generation (F_2). Consequently, seed from hybrid varieties is not used for planting stock.

Maize is an important and valuable field crop. Thus, a continuing goal of plant breeders is to develop high-yielding maize hybrids that are agronomically sound based on stable inbred lines. The reasons for this goal are obvious: to maximize the amount of grain produced with the inputs used and minimize susceptibility of the crop to environmental stresses. To accomplish this goal, the maize breeder must select and develop superior inbred parental lines for producing hybrids. This requires identification and selection of genetically unique individuals that occur in a segregating population. The segregating population is the result of a combination of crossover events plus the independent assortment of specific combinations of alleles at many gene loci that results in specific genotypes. Based on the number of segregating genes, the frequency of occurrence of an

individual with a specific genotype is less than 1 in 10,000. Thus, even if the entire genotype of the parents has been characterized and the desired genotype is known, only a few if any individuals having the desired genotype may be found in a large F_2 or S_0 population. Typically, however, the genotype of neither the parents nor the desired genotype is known in any detail.

In addition to the preceding problem, it is not known how the genotype will react with the environment. This genotype by environment interaction is an important, yet unpredictable, factor in plant breeding. A breeder of ordinary skill in the art cannot predict the genotype, how that genotype will interact with various environments or the resulting phenotypes of the developing lines, except perhaps in a very broad and general fashion. A breeder of ordinary skill in the art would also be unable to recreate the same line twice from the very same original parents as the breeder is unable to direct how the genomes combine or how they will interact with the environmental conditions. This unpredictability results in the expenditure of large amounts of research resources in the development of a superior new maize inbred line.

Pioneer research station staff propose about 400 to 500 new inbreds each year from over 2,000,000 pollinations. Of those proposed new inbreds, less than 50 and more commonly less than 30 are actually selected for commercial use.

SUMMARY OF THE INVENTION

According to the invention, there is provided a novel inbred maize line, designated PH24E. This invention thus relates to the seeds of inbred maize line PH24E, to the plants of inbred maize line PH24E, and to methods for producing a maize plant produced by crossing the inbred line PH24E with itself or another maize line. This invention further relates to hybrid maize seeds and plants produced by crossing the inbred line PH24E with another maize line.

Definitions

In the description and examples that follow, a number of terms are used herein. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided. NOTE: ABS is in absolute terms and %MN is percent of the mean for the experiments in which the inbred or hybrid was grown. These designators will follow the descriptors to denote how the values are to be interpreted. Below are the descriptors used in the data tables included herein.

ANT ROT=ANTHRACNOSE STALK ROT (*Colletotrichum graminicola*). A 1 to 9 visual rating indicating the resistance to Anthracnose Stalk Rot. A higher score indicates a higher resistance.

BAR PLT=BARREN PLANTS. The percent of plants per plot that were not barren (lack ears).

BRT STK=BRITTLE STALKS. This is a measure of the stalk breakage near the time of pollination, and is an indication of whether a hybrid or inbred would snap or break near the time of flowering under severe winds. Data are presented as percentage of plants that did not snap.

BU ACR=YIELD (BUSHEL/ACRE). Yield of the grain at harvest in bushels per acre adjusted to 15.5% moisture.

CLN=CORN LETHAL NECROSIS. Synergistic interaction of maize chlorotic mottle virus (MCMV) in combination with either maize dwarf mosaic virus (MDMV-A or MDMV-B) or wheat streak mosaic virus (WSMV). A 1 to 9

visual rating indicating the resistance to Corn Lethal Necrosis. A higher score indicates a higher resistance.

COM RST=COMMON RUST (*Puccinia sorghi*). A 1 to 9 visual rating indicating the resistance to Common Rust. A higher score indicates a higher resistance.

D/D=DRYDOWN. This represents the relative rate at which a hybrid will reach acceptable harvest moisture compared to other hybrids on a 1-9 rating scale. A high score indicates a hybrid that dries relatively fast while a low score indicates a hybrid that dries slowly.

DIP ERS=DIPLODIA EAR MOLD SCORES (*Diplodia maydis* and *Diplodia macrospora*). A 1 to 9 visual rating indicating the resistance to Diplodia Ear Mold. A higher score indicates a higher resistance.

DRP EAR=DROPPED EARS. A measure of the number of dropped ears per plot and represents the percentage of plants that did not drop ears prior to harvest.

D/T=DROUGHT TOLERANCE. This represents a 1-9 rating for drought tolerance, and is based on data obtained under Stress conditions. A high score indicates good drought tolerance and a low score indicates poor drought tolerance.

EAR HT=EAR HEIGHT. The ear height is a measure from the ground to the highest placed developed ear node attachment and is measured in inches.

EAR MLD=General Ear Mold. Visual rating (1-9 score) where a "1" is very susceptible and a "9" is very resistant. This is based on overall rating for ear mold of mature ears without determining the specific mold organism, and may not be predictive for a specific ear mold.

EAR SZ=EAR SIZE. A 1 to 9 visual rating of ear size. The higher the rating the larger the ear size.

ECB 1LF=EUROPEAN CORN BORER FIRST GENERATION LEAF FEEDING (*Ostrinia nubilalis*). A 1 to 9 visual rating indicating the resistance to preflowering leaf feeding by first generation European Corn Borer. A higher score indicates a higher resistance.

ECB 2IT=EUROPEAN CORN BORER SECOND GENERATION INCHES OF TUNNELING (*Ostrinia nubilalis*). Average inches of tunneling per plant in the stalk.

ECB 28C=EUROPEAN CORN BORER SECOND GENERATION (*Ostrinia nubilalis*). A 1 to 9 visual rating indicating post flowering degree of stalk breakage and other evidence of feeding by European Corn Borer, Second Generation. A higher score indicates a higher resistance.

ECB DPE=EUROPEAN CORN BORER DROPPED EARS (*Ostrinia nubilalis*). Dropped ears due to European Corn Borer. Percentage of plants that did not drop ears under second generation corn borer infestation.

EST CNT=EARLY STAND COUNT. This is a measure of the stand establishment in the spring and represents the number of plants that emerge on per plot basis for the inbred or hybrid.

EYE SPT=Eye Spot (*Kabatiella zeae* or *Aureobasidium zeae*). A 1 to 9 visual rating indicating the resistance to Eye Spot. A higher score indicates a higher resistance.

FUS ERS=FUSARIUM EAR ROT SCORE (*Fusarium moniliforme* or *Fusarium subglutinans*). A 1 to 9 visual rating indicating the resistance to Fusarium ear rot. A higher score indicates a higher resistance.

GDU=Growing Degree Units. Using the Barger Heat Unit Theory, which assumes that maize growth occurs in the temperature range 50° F.-86° F. and that temperatures outside this range slow down growth; the maximum daily heat unit accumulation is 36 and the minimum daily heat

unit accumulation is 0. The seasonal accumulation of GDU is a major factor in determining maturity zones.

GDU SHD=GDU TO SHED. The number of growing degree units (GDUs) or heat units required for an inbred line or hybrid to have approximately 50 percent of the plants shedding pollen and is measured from the time of planting. Growing degree units are calculated by the Barger Method, where the heat units for a 24-hour period are:

$$GDU = \frac{(\text{Max. temp.} + \text{Min. temp.})}{2} - 50$$

The highest maximum temperature used is 86° F. and the lowest minimum temperature used is 50° F. For each inbred or hybrid it takes a certain number of GDUs to reach various stages of plant development.

GDU SLK=GDU TO SILK. The number of growing degree units required for an inbred line or hybrid to have approximately 50 percent of the plants with silk emergence from time of planting. Growing degree units are calculated by the Barger Method as given in GDU SHD definition.

GIB ERS=GIBBERELLA EAR ROT (PINK MOLD) (*Gibberella zeae*). A 1 to 9 visual rating indicating the resistance to Gibberella Ear Rot. A higher score indicates a higher resistance.

GLF SPT=Gray Leaf Spot (*Cercospora zeae-maydis*). A 1 to 9 visual rating indicating the resistance to Gray Leaf Spot. A higher score indicates a higher resistance.

GOS WLT=Goss' Wilt (*Corynebacterium nebraskense*). A 1 to 9 visual rating indicating the resistance to Goss' Wilt. A higher score indicates a higher resistance.

GRN APP=GRAIN APPEARANCE. This is a 1 to 9 rating for the general appearance of the shelled grain as it is harvested based on such factors as the color of harvested grain, any mold on the grain, and any cracked grain. High scores indicate good grain quality.

H/POP=YIELD AT HIGH DENSITY. Yield ability at relatively high plant densities on 1-9 relative rating system with a higher number indicating the hybrid responds well to high plant densities for yield relative to other hybrids. A 1, 5, and 9 would represent very poor, average, and very good yield response, respectively, to increased plant density.

HC BLT=HELMINTHOSPORIUM CARBONUM LEAF BLIGHT (*Helminthosporium carbonum*). A 1 to 9 visual rating indicating the resistance to Helminthosporium infection. A higher score indicates a higher resistance.

HD SMT=HEAD SMUT (*Sphacelotheca reiliana*). This score indicates the percentage of plants not infected.

INC D/A=GROSS INCOME (DOLLARS PER ACRE). Relative income per acre assuming drying costs of two cents per point above 15.5 percent harvest moisture and current market price per bushel.

INCOME/ACRE. Income advantage of hybrid to be patented over other hybrid on per acre basis.

INC ADV=GROSS INCOME ADVANTAGE. GROSS INCOME advantage of variety #1 over variety #2.

L/POP=YIELD AT LOW DENSITY. Yield ability at relatively low plant densities on a 1-9 relative system with a higher number indicating the hybrid responds well to low plant densities for yield relative to other hybrids. A 1, 5, and 9 would represent very poor, average, and very good yield response, respectively, to low plant density.

MDM CPX=MAIZE DWARF MOSAIC COMPLEX (MDMV=Maize Dwarf Mosaic Virus and MCDV=Maize Chlorotic Dwarf Virus). A 1 to 9 visual rating indicating the

resistance to Maize Dwarf Mosaic Complex. A higher score indicates a higher resistance.

MST=HARVEST MOISTURE. The moisture is the actual percentage moisture of the grain at harvest.

MST ADV=MOISTURE ADVANTAGE. The moisture advantage of variety #1 over variety #2 as calculated by: MOISTURE of variety #2—MOISTURE of variety #1=MOISTURE ADVANTAGE of variety #1.

NLF BLT=Northern Leaf Blight (*Helminthosporium turcicum* or *Exserohilum turcicum*). A 1 to 9 visual rating indicating the resistance to Northern Leaf Blight. A higher score indicates a higher resistance.

PLT HT=PLANT HEIGHT. This is a measure of the height of the plant from the ground to the tip of the tassel in inches.

POL SC=POLLEN SCORE. A 1 to 9 visual rating indicating the amount of pollen shed. The higher the score the more pollen shed.

POL WT=POLLEN WEIGHT. This is calculated by dry weight of tassels collected as shedding commences minus dry weight from similar tassels harvested after shedding is complete.

It should be understood that the inbred can, through routine manipulation of cytoplasmic or other factors, be produced in a male-sterile form. Such embodiments are also contemplated within the scope of the present claims.

POP K/A=PLANT POPULATIONS. Measured as 1000s per acre.

POP ADV=PLANT POPULATION ADVANTAGE. The plant population advantage of variety #1 over variety #2 as calculated by PLANT POPULATION of variety #2—PLANT POPULATION of variety #1=PLANT POPULATION ADVANTAGE of variety #1.

PRM=PREDICTED RELATIVE MATURITY. This trait, predicted relative maturity, is based on the harvest moisture of the grain. The relative maturity rating is based on a known set of checks and utilizes standard linear regression analyses and is also referred to as the Comparative Relative Maturity Rating System that is similar to the Minnesota Relative Maturity Rating System.

PRM SHD=A relative measure of the growing degree units (GDU) required to reach 50% pollen shed. Relative values are predicted values from the linear regression of observed GDU's on relative maturity of commercial checks.

RTL LG=ROOT LODGING. Root lodging is the percentage of plants that do not root lodge; plants that lean from the vertical axis at an approximately 30° angle or greater would be counted as root lodged.

RTL ADV=ROOT LODGING ADVANTAGE. The root lodging advantage of variety #1 over variety #2.

SCT GRN=SCATTER GRAIN. A 1 to 9 visual rating indicating the amount of scatter grain (lack of pollination or kernel abortion) on the ear. The higher the score the less scatter grain.

SDG VGR=SEEDLING VIGOR. This is the visual rating (1 to 9) of the amount of vegetative growth after emergence at the seedling stage (approximately five leaves). A higher score indicates better vigor.

SEL IND=SELECTION INDEX. The selection index gives a single measure of the hybrid's worth based on information for up to five traits. A maize breeder may utilize his or her own set of traits for the selection index. One of the traits that is almost always included is yield. The selection index data presented in the tables represent the mean value averaged across testing stations.

SLF BLT=SOUTHERN LEAF BLIGHT (*Helminthosporium maydis* or *Bipolaris maydis*). A 1 to 9 visual rating indicating the resistance to Southern Leaf Blight. A higher score indicates a higher resistance.

SOU RST=SOUTHERN RUST (*Puccinia polysora*). A 1 to 9 visual rating indicating the resistance to Southern Rust. A higher score indicates a higher resistance.

STA GRN=STAY GREEN. Stay green is the measure of plant health near the time of black layer formation (physiological maturity). A high score indicates better late-season plant health.

STD ADV=STALK STANDING ADVANTAGE. The advantage of variety #1 over variety #2 for the trait STK CNT.

STK CNT=NUMBER OF PLANTS. This is the final stand or number of plants per plot.

STK LDG=STALK LODGING. This is the percentage of plants that did not stalk lodge (stalk breakage) as measured by either natural lodging or pushing the stalks and determining the percentage of plants that break below the ear.

STW WLT=Stewart's Wilt (*Erwinia stewartii*). A 1 to 9 visual rating indicating the resistance to Stewart's Wilt. A higher score indicates a higher resistance.

TAS BLS=TASSEL BLAST. A 1 to 9 visual rating was used to measure the degree of blasting (necrosis due to heat stress) of the tassel at the time of flowering. A 1 would indicate a very high level of blasting at time of flowering, while a 9 would have no tassel blasting.

TAS SZ=TASSEL SIZE. A 1 to 9 visual rating was used to indicate the relative size of the tassel. The higher the rating the larger the tassel.

TAS WT=TASSEL WEIGHT. This is the average weight of a tassel (grams) just prior to pollen shed.

TEX EAR=EAR TEXTURE. A 1 to 9 visual rating was used to indicate the relative hardness (smoothness of crown) of mature grain. A 1 would be very soft (extreme dent) while a 9 would be very hard (flinty or very smooth crown).

TILLER=TILLERS. A count of the number of tillers per plot that could possibly shed pollen was taken. Data is given as a percentage of tillers: number of tillers per plot divided by number of plants per plot.

TST WT=TEST WEIGHT (UNADJUSTED). The measure of the weight of the grain in pounds for a given volume (bushel).

TST WTA=TEST WEIGHT ADJUSTED. The measure of the weight of the grain in pounds for a given volume (bushel) adjusted for 15.5 percent moisture.

TSW ADV=TEST WEIGHT ADVANTAGE. The test weight advantage of variety #1 over variety #2.

WIN M%=PERCENT MOISTURE WINS.

WIN Y%=PERCENT YIELD WINS.

YLD=YIELD. It is the same as BU ACR ABS.

YLD ADV=YIELD ADVANTAGE. The yield advantage of variety #1 over variety #2 as calculated by: YIELD of variety #1—YIELD variety #2=yield advantage of variety #1.

YLD SC=YIELD SCORE. A 1 to 9 visual rating was used to give a relative rating for yield based on plot ear piles. The higher the rating the greater visual yield appearance.

DETAILED DESCRIPTION OF THE INVENTION

Inbred maize lines are typically developed for use in the production of hybrid maize lines. Inbred maize lines need to

be highly homogeneous, homozygous and reproducible to be useful as parents of commercial hybrids. There are many analytical methods available to determine the homozygotic and phenotypic stability of these inbred lines.

The oldest and most traditional method of analysis is the observation of phenotypic traits. The data is usually collected in field experiments over the life of the maize plants to be examined. Phenotypic characteristics most often observed are for traits associated with plant morphology, ear and kernel morphology, insect and disease resistance, maturity, and yield.

In addition to phenotypic observations, the genotype of a plant can also be examined. There are many laboratory-based techniques available for the analysis, comparison and characterization of plant genotype; among these are Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), and Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites.

The most widely used of these laboratory techniques are Isozyme Electrophoresis and RFLPs as discussed in Lee, M., "Inbred Lines of Maize and Their Molecular Markers," *The Maize Handbook*, (Springer-Verlag, New York, Inc. 1994, at 423-432) incorporated herein by reference. Isozyme Electrophoresis is a useful tool in determining genetic composition, although it has relatively low number of available markers and the low number of allelic variants among maize inbreds. RFLPs have the advantage of revealing an exceptionally high degree of allelic variation in maize and the number of available markers is almost limitless.

Maize RFLP linkage maps have been rapidly constructed and widely implemented in genetic studies. One such study is described in Boppenmaier, et al., "Comparisons among strains of inbreds for RFLPs", *Maize Genetics Cooperative Newsletter*, 65:1991, pg. 90, is incorporated herein by reference. This study used 101 RFLP markers to analyze the patterns of 2 to 3 different deposits each of five different inbred lines. The inbred lines had been selfed from 9 to 12 times before being adopted into 2 to 3 different breeding programs. It was results from these 2 to 3 different breeding programs that supplied the different deposits for analysis. These five lines were maintained in the separate breeding programs by selfing or sibbing and roguing off-type plants for an additional one to eight generations. After the RFLP analysis was completed, it was determined the five lines showed 0-2% residual heterozygosity. Although this was a relatively small study, it can be seen using RFLPs that the lines had been highly homozygous prior to the separate strain maintenance.

Inbred maize line PH24E is a yellow, dent maize inbred that is best suited as either a male or female in crosses for producing first generation F1 maize hybrids. Inbred maize line PH24E is best adapted to the Northcentral region of the United States and can be used to produce hybrids from approximately 106 relative maturity based on the Comparative Relative Maturity Rating System for harvest moisture of grain. As an inbred per se, maize line PH24E is a high yielding female with a resistance to barrenness. Inbred PH24E exhibits good cold test results, good stalks and dark green plants. As an inbred per se, PH24E exhibits excellent resistance to barrenness which makes it a reliable seed production parent. Inbred PH24E demonstrates excellent

general combining ability. When in hybrid combination, PH24E exhibits high yield for its maturity with a stability over different environments across the Northcentral Corn Belt. PH24E hybrids have excellent roots, good stalk strength and quality and short plant stature. PH24E hybrids' short plant stature and associated reduced crop residues, make them well adapted to conservation tillage cropping systems. PH24E hybrids perform well from the eastern to western edges of the Northcentral Corn Belt.

The inbred has shown uniformity and stability within the limits of environmental influence for all the traits as described in the Variety Description Information (Table 1) that follows. The inbred has been self-pollinated and earrowed a sufficient number of generations with careful attention paid to uniformity of plant type to ensure the homozygosity and phenotypic stability necessary to use in commercial production. The line has been increased both by hand and in isolated fields with continued observation for uniformity. No variant traits have been observed or are expected in PH24E.

Inbred maize line PH24E, being substantially homozygous, can be reproduced by planting seeds of the line, growing the resulting maize plants under self-pollinating or sib-pollinating conditions with adequate isolation, and harvesting the resulting seed, using techniques familiar to the agricultural arts.

TABLE 1

VARIETY DESCRIPTION INFORMATION
INBRED = PH24E
Type: Dent

A. Maturity: Average cross maturity zones. Zone: 0

GDU Shed: 1440

GDU Silk: 1490

No. Reps: 42

B. Plant Characteristics:

Plant height (to tassel tip): 209 cm

Length of top ear internode: 16 cm

Number of ears per stalk: Single

Ear height (to base of top ear): 80 cm

Number of tillers: None

Cytoplasm type: Normal

C. Leaf:

Color: (WF9) Medium Green*

Angle from Stalk: 30-60 degrees

Marginal Waves: (WF9) Few

Number of Leaves (mature plants): 18

Sheath Pubescence: (W22) Light

Longitudinal Creases: (PA11) Many

Length (Ear node leaf): 83 cm

Width (widest point, ear node leaf): 9 cm

D. Tassel:

Number lateral branches: 8

Branch Angle from central spike: >45 degrees

Peduncle Length (top leaf to basal branches): 18 cm

Anther Color: Yellow*

Glume Color: Green*

E. Ear (Husked Ear Data Except When Stated Otherwise):

Length: 17 cm

Weight: 141 gm

Mid-point Diameter: 43 mm

Silk Color: Red*

Husk Extension (Harvest stage): Medium (Barely Covering Ear)

Husk Leaf: Long (>15 cm)

Taper of Ear: Average

Position of Shank (dry husks): Upright

Kernel Rows: Distinct, Straight Number = 14

Husk Color (fresh): Light Green*

TABLE 1-continued

VARIETY DESCRIPTION INFORMATION	
INBRED = PH24E	
Type: Dent	
<u>Husk Color (dry): Buff*</u>	
<u>Shank Length: 13 cm</u>	
<u>Shank (No. of internodes): 9</u>	
<u>F. Kernel (Dried):</u>	
<u>Size (from ear mid-point)</u>	
Length: 11 mm	
Width: 9 mm	
Thick: 4 mm	
Shape Grade (% rounds) (35% medium round based on	
Parent Test Data) 20-40	
Pericarp Color: Colorless*	
Aleurone Color: Homozygous Yellow*	
Endosperm Color: Yellow*	
Endosperm Type: Normal Starch	
Gm Wt/100 Seeds (unsized): 33 gm	
<u>G. Cob:</u>	
Diameter at mid-point: 25 mm	
Strength: Strong	
Color: Red*	
<u>H. Diseases:</u>	
Corn Lethal Necrosis (MCMV = Maize Chlorotic Mottle Virus and	
MDMV = Maize Dwarf Mosaic Virus): Susceptible	
N. Leaf Blight (<i>E. turcicum</i>): Resistant	
Common Rust (<i>P. sorghi</i>): Resistant	
Eye Spot (<i>K. zeae</i>): Intermediate	
Gray Leaf Spot (<i>C. zeae</i>): Intermediate	
Fusarium Ear Mold (<i>F. moniliforme</i>): Resistant	
<u>I. Insects:</u>	
European Corn Borer-1 Leaf Damage (Pre-flowering): Intermediate	
European Corn Borer-2 (Post-flowering): Intermediate	
The above descriptions are based on a scale of 1-9, 1 being highly	
susceptible, 9 being highly resistant.	
S (Susceptible): Would generally represent a score of 1-3.	
I (Intermediate): Would generally represent a score of 4-5.	
R (Resistant): Would generally represent a score of 6-7.	
H (Highly Resistant) Would generally represent a score of 8-9. Highly	
resistant does not imply the inbred is immune.	
*In interpreting the foregoing color designations, reference may be had to the	
Munsell Glossy Book of Color, a standard color reference.	
Data for Items B, C, D, E, F, and G is based primarily on a maximum of 2	
reps from Johnston, Iowa grown in 1994, plus description information from	
the maintaining station.	
Data for Items B, C, D, E, F, and G is based primarily on a	
maximum of 2 reps from Johnston, Iowa grown in 1994,	
plus description information from the maintaining station.	

Industrial Applicability

This invention also is directed to methods for producing a maize plant by crossing a first parent maize plant with a second parent maize plant wherein the first or second parent maize plant is an inbred maize plant of the line PH24E. Further, both first and second parent maize plants can come from the inbred maize line PH24E. Thus, any such methods using the inbred maize line PH24E are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, and the like. All plants produced using inbred maize line PH24E as a parent are within the scope of this invention. Advantageously, the inbred maize line is used in crosses with other, different, maize inbreds to produce first generation (F_1) maize hybrid seeds and plants with superior characteristics.

As used herein, the term plant includes plant cells, plant protoplasts, plant cell tissue cultures from which maize plants can be regenerated, plant calli, plant clumps, and plant

cells that are intact in plants or parts of plants, such as embryos, pollen, ovules, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk and the like.

Duncan, Williams, Zehr, and Widholm, *Planta* (1985) 165:322-332 reflects that 97% of the plants cultured that produced callus were capable of plant regeneration. Subsequent experiments with both inbreds and hybrids produced 91% regenerable callus that produced plants. In a further study in 1988, Songstad, Duncan & Widholm in *Plant Cell Reports* (1988), 7:262-265 reports several media additions that enhance regenerability of callus of two inbred lines. Other published reports also indicated that "nontraditional" tissues are capable of producing somatic embryogenesis and plant regeneration. K. P. Rao, et al., *Maize Genetics Cooperation Newsletter*, 60:64-65 (1986), refers to somatic embryogenesis from glume callus cultures and B. V. Conger, et al., *Plant Cell Reports*, 6:345-347 (1987) indicates somatic embryogenesis from the tissue cultures of maize leaf segments. Thus, it is clear from the literature that the state of the art is such that these methods of obtaining plants are, and were, "conventional" in the sense that they are routinely used and have a very high rate of success.

Tissue culture of maize is described in European Patent Application, publication 160,390, incorporated herein by reference. Maize tissue culture procedures are also described in Green and Rhodes, "Plant Regeneration in Tissue Culture of Maize," *Maize for Biological Research* (Plant Molecular Biology Association, Charlottesville, Va. 1982, at 367-372) and in Duncan, et al., "The Production of Callus Capable of Plant Regeneration from Immature Embryos of Numerous *Zea Mays* Genotypes," 165 *Planta* 322-332 (1985). Thus, another aspect of this invention is to provide cells which upon growth and differentiation produce maize plants having the physiological and morphological characteristics of inbred line PH24E.

Maize is used as human food, livestock feed, and as raw material in industry. The food uses of maize, in addition to human consumption of maize kernels, include both products of dry- and wet-milling industries. The principal products of maize dry milling are grits, meal and flour. The maize wet-milling industry can provide maize starch, maize syrups, and dextrose for food use. Maize oil is recovered from maize germ, which is a by-product of both dry- and wet-milling industries.

Maize, including both grain and non-grain portions of the plant, is also used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry.

Industrial uses of maize include production of ethanol, maize starch in the wet-milling industry and maize flour in the dry-milling industry. The industrial applications of maize starch and flour are based on functional properties, such as viscosity, film formation, adhesive properties, and ability to suspend particles. The maize starch and flour have application in the paper and textile industries. Other industrial uses include applications in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications.

Plant parts other than the grain of maize are also used in industry: for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

The seed of inbred maize line PH24E, the plant produced from the inbred seed, the hybrid maize plant produced from the crossing of the inbred, hybrid seed, and various parts of the hybrid maize plant can be utilized for human food, livestock feed, and as a raw material in industry.

Performance Examples of PH24E

In the examples that follow, the traits and characteristics of inbred maize line PH24E are presented as an inbred per se and in hybrid combination. The data collected on inbred maize line PH24E is presented for key characteristics and traits.

The results in Table 2A compare PH24E to its parent, PHP02. The inbred per se results show PH24E is significantly higher yielding than PHP02. Inbred PH24E has higher pollen weight and staygreen than PHP02.

Table 2B compares PH24E to PHJ90, a line with similar characteristics. The inbred per se results show PH24E is significantly higher yielding with higher staygreen than PHJ90. Inbred PH24E exhibits higher resistance to stalk lodging and brittle stalk than PHJ90.

The results in Table 2C compare PH24E to PHAP9, another inbred line with similarly characteristics. Inbred PH24E exhibits greater resistance to stalk lodging and brittle stalk.

Table 2D compares PH24E to PHKW3, another inbred line with similar characteristics. Inbred PH24E is significantly higher yielding and is significantly shorter than inbred PHAP9. Inbred PH24E exhibits higher staygreen scores than inbred PHAP9.

Table 2E compares PH24E to its other parent, PHN82. The inbred per se results show PH24E is higher yielding with higher staygreen than PHN82. In addition, PH24E has a higher pollen score, tassel size and pollen weight than PHN82.

The results in Table 3A compare the general combining ability of PH24E to PHP02. The results show that the PH24E hybrids and the PHP02 hybrids have similar yields. The

PH24E hybrids have shorter stature than the PHP02 hybrids. Both sets of hybrids have above average resistance to stalk lodging and root lodging. The results in Table 3A are somewhat biased because they represent a limited number of experiments in which PH24E and PHP02 were crossed to the same tester. PH24E resembles inbred parent PHP02 much less than its other parent, PHN82. Inbreds PH24E and PHN82 have similar maturities of 1440-1490 and 1450-1500 heat units to shed and silk, respectively, and they have similar morphologies, yield levels, and harvest moistures. Inbred PHP02 is the older parent in terms of commercial production and is now in limited use in new commercial hybrids and will no longer be used in general combining ability testing. However, a professional plant breeder has expressed the expert opinion that PH24E is superior to PHP02, and that, if PH24E were to be compared to PHP02 using more appropriate testers than were used in these experiments, PH24E would have shown superior results to those of PHP02.

Table 3B compares the general combining ability of PH24E and PHJ90. The results show PH24E hybrids are significantly higher yielding with significantly higher staygreen than PHJ90 hybrids.

The results in Table 3C compares the general combining ability of PH24E and PHAP9. The results show PH24E hybrids are higher yielding with significantly higher staygreen than PHAP9 hybrids. Both PH24E hybrids and PHAP9 hybrids show above average resistance to root lodging and stalk lodging.

Table 3D compares the general combining ability of PH24E and PHKW3. The results show the PH24E hybrids are significantly higher yielding with superior staygreen than the PHKW3 hybrids. The PH24E hybrids exhibit higher resistance to root lodging, stock lodging and brittle stalk.

TABLE 2A

PAIRED INBRED COMPARISON DATA										
VARIETY #1 = PH24E										
VARIETY #2 = PHP02										
VAR #	BU ACR ABS	BU ACR % MN	MST ABS	TST WT ABS	SDG VGR ABS	EST CNT ABS	TIL LER ABS	GDU SHD ABS	GDU SLK ABS	
TOTAL SUM 1	91.7	135	18.3	56.7	3.5	36.2	2.5	137.8	140.9	
2	81.3	121	17.3	56.6	5.2	37.9	1.7	130.0	132.5	
LOCS	21	21	26	15	31	61	48	51	50	
REPS	40	40	45	29	45	95	64	56	55	
DIFF	10.5	14	1.0	0.1	1.8	1.7	0.8	7.8	8.4	
PROB	.001#	.005#	.002#	.746	.000#	.077*	.260	.000#	.000#	
VAR #	POL WT ABS	POL WT % MN	POL SC ABS	TAS BLS ABS	TAS SZ ABS	PLT HT ABS	EAR HT ABS	RT LDG ABS	STA GRN ABS	
TOTAL SUM 1	120.4	111	5.8	9.0	5.6	73.0	33.4	93.0	5.4	
2	83.8	77	6.4	6.3	6.4	72.1	30.0	93.5	4.0	
LOCS	11	11	15	2	16	28	26	9	20	
REPS	11	11	16	2	18	41	38	12	33	
DIFF	36.6	34	0.6	2.8	0.8	0.8	3.3	0.5	1.4	
PROB	.032+	.039+	.147	.500	.038+	.422	.001#	.836	.000#	
VAR #	STK LDG ABS	BRT STK ABS	GRN APP ABS	SCT GRN ABS	EAR SZ ABS	TEX EAR ABS	EAR MLD ABS	BAR PLT ABS	DRP EAR ABS	
TOTAL SUM 1	96.9	95.5	6.0	7.2	6.3	4.8	6.9	97.1	99.7	
2	97.6	91.0	5.3	7.8	6.2	4.0	7.4	97.9	99.7	
LOCS	11	16	4	9	6	5	11	36	4	
REPS	21	21	8	10	6	5	12	44	8	
DIFF	0.7	4.6	0.8	0.6	0.2	0.8	0.5	0.8	0.0	
PROB	.552	.271	.215	.347	.611	.016+	.096*	.368	.964	

TABLE 2A-continued

PAIRED INBRED COMPARISON DATA							
VARIETY #1 = PH24E							
VARIETY #2 = PHP02							
VAR		GLF	NLF	EYE	COM	ECB	ECB
#		SPT	BLT	SPT	RST	1LF	2SC
		ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	7.0	7.7	4.5	6.1	5.0	3.5
	2	2.0	5.3	5.5	5.8	5.6	4.5
	LOCS	1	3	2	6	7	3
	REPS	1	3	2	7	7	4
	DIFF	5.0	2.3	1.0	0.3	0.6	1.0
	PROB		.020+	.000#	.562	.231	.000#

* = 10% SIG

+ = 5% SIG

= 1% SIG

TABLE 2B

PAIRED INBRED COMPARISON DATA										
VARIETY #1 = PH24E										
VARIETY #2 = PHU90										
VAR		BU	BU	MST	TST	SDG	EST	TIL	GDU	GDU
#		ACR	ACR	ABS	WT	VGR	CNT	LER	SHD	SLK
		ABS	% MN	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	92.3	138	18.9	56.3	3.5	34.8	2.4	137.7	141.0
	2	73.8	107	16.9	56.3	5.1	36.5	3.4	133.8	137.3
	LOCS	19	19	22	13	28	51	41	43	42
	REPS	35	35	39	25	39	82	55	46	45
	DIFF	18.5	31	1.9	0.0	1.6	1.7	1.0	3.9	3.7
	PROB	.000#	.001#	.000#	.931	.000#	.114	.595	.000#	.000#
VAR		POL	POL	POL	TAS	TAS	PLT	EAR	RT	STA
#		WT	WT	SC	BLS	SZ	HT	HT	LDG	GRN
		ABS	% MN	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	109.5	82	5.5	9.0	5.5	72.5	33.4	92.3	5.4
	2	119.3	89	6.8	9.0	6.5	70.4	32.4	98.2	2.3
	LOCS	6	6	12	1	14	25	23	7	19
	REPS	6	6	13	1	16	36	33	10	31
	DIFF	9.8	6	1.3	0.0	1.0	2.0	1.1	6.0	3.1
	PROB	.250	.265	.007#		.003#	.079*	.238	.247	.000#
VAR		STK	BRT	GRN	SCT	EAR	TEX	EAR	BAR	DRP
#		LDG	STK	APP	GRN	SZ	EAR	MLD	FLT	EAR
		ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	98.4	96.6	6.3	7.4	6.2	4.8	7.2	96.7	99.5
	2	88.7	93.0	4.5	7.9	6.2	5.4	7.6	95.6	100.0
	LOCS	9	15	2	7	5	5	9	25	3
	REPS	17	20	4	8	5	5	10	31	6
	DIFF	9.6	3.7	1.8	0.5	0.0	0.6	0.3	1.2	0.5
	PROB	.055*	.083*	.258	.177	.000#	.070*	.195	.212	.423
VAR		GLF	NLF	EYE	COM	ECB	ECB			
#		SPT	BLT	SPT	RST	1LF	2SC			
		ABS	ABS	ABS	ABS	ABS	ABS			
TOTAL SUM	1	7.0	7.5	4.5	6.1	5.3	3.5			
	2	2.0	4.0	6.0	4.7	7.5	3.0			
	LOCS	1	2	2	5	6	3			
	REPS	1	2	2	6	6	4			
	DIFF	5.0	3.5	1.5	1.4	2.2	0.5			
	PROB		.258	.205	.052*	.015+	.225			

* = 10% SIG

+ = 5% SIG

= 1% SIG

TABLE 2C

PAIRED INBRED COMPARISON DATA										
VARIETY #1 = PH24E										
VARIETY #2 = PHAP9										
VAR #	BU ACR ABS	BU ACR % MN	MST ABS	SDG VGR ABS	EST CNT ABS	TIL LER ABS	GDU SHD ABS	GDU SLK ABS	POL WT ABS	
TOTAL SUM 1	32.0	69	10.4	3.1	22.8	3.2	137.4	140.7	106.0	
2	35.9	78	10.8	4.0	25.1	2.7	132.6	138.0	130.5	
LOCS	1	1	1	21	38	28	41	40	5	
REPS	1	1	1	21	53	29	42	41	5	
DIFF	3.8	8	0.4	0.9	2.4	0.5	4.8	2.7	24.4	
PROB				.052*	.026+	.743	.000#	.000#	.015+	
VAR #	POL WT % MN	POL SC ABS	TAS BLS ABS	TAS SZ ABS	PLT HT ABS	EAR HT ABS	RT LDG ABS	STA GRN ABS	STK LDG ABS	
TOTAL SUM 1	83	5.2	9.0	5.5	71.0	31.3	94.4	5.1	100.0	
2	104	6.2	9.0	6.4	77.8	32.1	100.0	4.6	96.0	
LOCS	5	11	1	15	16	15	4	7	1	
REPS	5	11	1	16	17	16	4	7	1	
DIFF	20	1.0	0.0	0.9	6.8	0.9	5.6	0.6	4.0	
PROB	.018+	.002#		.004#	.001#	.462	.061*	.280		
VAR #	BRT STK ABS	SCT GRN ABS	EAR SZ ABS	TEX EAR ABS	EAR MLD ABS	BAR PLT ABS	GLF SPT ABS	NLF BLT ABS	STW WLT ABS	
TOTAL SUM 1	97.1	7.2	6.3	5.0	6.9	95.3	4.0	7.7	7.0	
2	96.2	7.7	6.3	5.0	6.7	97.8	2.5	6.0	6.0	
LOCS	9	9	6	4	11	11	2	3	1	
REPS	9	10	6	4	12	11	2	3	1	
DIFF	0.9	0.4	0.0	0.0	0.2	2.4	1.5	1.7	1.0	
PROB	.722	.272	.000#	.000#	.676	.282	.656	.038+		
VAR #	EYE SPT ABS	COM RST ABS	ECB 1LF ABS	ECB 2SC ABS						
TOTAL SUM 1	4.5	5.8	5.0	3.5						
2	5.0	5.0	6.9	5.0						
LOCS	2	4	7	2						
REPS	2	4	7	2						
DIFF	0.5	0.8	1.9	1.5						
PROB	.500	.215	.032+	.205						

* = 10% SIG

+ = 5% SIG

= 1% SIG

TABLE 2D

PAIRED INBRED COMPARISON DATA										
VARIETY #1 = PH24E										
VARIETY #2 = PHKW3										
VAR #	BU ACR ABS	BU ACR % MN	MST ABS	TST WT ABS	SDG VGR ABS	EST CNT ABS	TIL LER ABS	GDU SHD ABS	GDU SLK ABS	
TOTAL SUM 1	95.5	142	18.6	56.3	3.4	37.2	2.7	138.8	142.0	
2	74.7	116	18.3	56.3	3.7	33.4	2.2	139.2	142.5	
LOCS	18	18	23	13	30	71	53	56	55	
REPS	33	33	38	23	41	110	74	67	66	
DIFF	20.9	26	0.4	0.1	0.3	3.7	0.4	0.4	0.4	
PROB	.000#	.003#	.238	.821	.366	.002#	.590	.368	.396	
VAR #	POL WT ABS	POL WT % MN	POL SC ABS	TAS BLS ABS	TAS SZ ABS	PLT HT ABS	EAR HT ABS	RT LDG ABS	STA GRN ABS	
TOTAL SUM 1	138.6	103	5.8	9.0	5.7	71.8	33.2	93.7	5.3	
2	157.9	117	5.9	7.8	5.0	77.4	33.5	89.3	4.0	
LOCS	20	20	22	2	15	25	23	9	20	
REPS	28	28	31	2	17	35	33	14	31	

TABLE 2D-continued

PAIRED INBRED COMPARISON DATA									
VARIETY #1 = PH24E									
VARIETY #2 = PHKW3									
DIFF	19.2	14	0.1	1.3	0.6	5.6	0.4	4.5	1.3
PROB	.030+	.029+	.627	.500	.112	.001#	.731	.502	.000#
VAR	STK	BRT	GRN	SCT	EAR	TEX	EAR	BAR	DRP
#	LDG	STK	APP	GRN	SZ	EAR	MLD	PLT	EAR
	ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	98.9	95.6	6.3	7.3	6.3	4.8	6.8	97.5
	2	96.3	97.7	7.0	7.4	4.8	6.6	7.1	94.1
LOCS	9	19	2	8	6	5	10	32	3
REPS	16	26	4	9	6	5	11	38	5
DIFF	2.7	2.1	0.8	0.1	1.5	1.8	0.3	3.4	0.5
PROB	.072*	.374	.830	.763	.017+	.001#	.496	.002#	.423
VAR	GLF	NLF	STW	EYE	COM	ECB	ECB		
#	SPT	BLT	WLT	SPT	RST	ILF	25C		
	ABS	ABS	ABS	ABS	ABS	ABS	ABS		
TOTAL SUM	1	4.0	7.7	7.0	4.5	6.1	5.0	3.7	
	2	2.5	6.0	5.0	6.5	5.7	6.6	3.3	
LOCS	2	3	1	2	6	7	3		
REPS	2	3	1	2	7	7	3		
DIFF	1.5	1.7	2.0	2.0	0.4	1.6	0.3		
PROB	.656	.130		.295	.486	.025+	.667		

* = 10% SIG

+ = 5% SIG

= 1% SIG

TABLE 2E

PAIRED INBRED COMPARISON DATA									
VARIETY #1 = PH24E									
VARIETY #2 = PHN82									
VAR	BU	BU	MST	TST	SDG	EST	TIL	GDU	GDU
#	ACR	ACR	ABS	WT	VGR	CNT	LER	SHD	SLK
	ABS	% MN	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	103.6	120	19.4	56.8	3.7	38.3	3.9	139.9
	2	94.2	112	21.7	55.9	4.0	41.5	3.7	144.0
LOCS	13	13	14	10	14	31	26	25	25
REPS	24	24	25	19	22	50	44	37	37
DIFF	9.4	9	2.3	0.9	0.3	3.2	0.2	4.1	4.8
PROB	.097*	.266	.000#	.000#	.326	.005#	.831	.000#	.000#
VAR	POL	POL	POL	TAS	PLT	EAR	RT	STA	STK
#	WT	WT	SC	SZ	HT	HT	LDG	GRN	LDG
	ABS	% MN	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	165.2	96	6.1	5.3	72.9	34.4	97.6	5.1
	2	109.5	63	4.9	3.9	72.3	28.3	100.0	3.8
LOCS	8	8	14	8	14	13	3	11	6
REPS	16	16	23	9	24	22	5	19	12
DIFF	55.6	34	1.2	1.4	0.6	6.2	2.4	1.3	1.0
PROB	.000#	.000#	.012+	.064*	.550	.000#	.423	.041+	.643
VAR	BRT	GRN	SCT	EAR	TEX	EAR	BAR	DRP	GLF
#	STK	APP	GRN	SZ	EAR	MLD	PLT	EAR	SPT
	ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS	ABS
TOTAL SUM	1	93.3	5.8	7.2	6.3	4.5	6.3	97.2	100.0
	2	95.1	6.4	6.8	5.3	4.8	7.0	98.2	100.0
LOCS	6	2	6	3	4	6	14	1	1
REPS	10	4	7	3	4	7	18	2	1
DIFF	1.7	0.7	0.3	1.0	0.3	0.7	1.0	0.0	1.0
PROB	.433	.758	.576	.225	.391	.102	.577		

TABLE 2E-continued

PAIRED INBRED COMPARISON DATA						
VARIETY #1 = PH24E						
VARIETY #2 = PHN82						
VAR #	NLF BLT ABS	EYE SPT ABS	COM RST ABS	ECB 1LF ABS	ECB 2SC ABS	
TOTAL SUM	1	8.0	4.0	5.8	4.0	3.5
	2	6.0	1.0	5.8	3.7	4.0
LOCS	1	1	1	4	3	2
REPS	1	1	4	3	2	
DIFF	2.0	3.0	0.0	0.3	0.5	
PROB			.000#	.667	.500	

* = 10% SIG

+ = 5% SIG

= 1% SIG

TABLE 3A

Average Inbred By Tester Performance Comparing PH24E To PHP02 Crossed To The Same Inbred Testers And Grown In The Same Experiments.											
	SEL IND % MN	PRM ABS	BU ACR ABS	BU ACR % MN	PRM SHD ABS	MST % MN	TST WT ABS	SDG VGR % MN	EST CNT % MN	GDU SHD % MN	
TOTAL SUM	REPS	16	15	68	68	12	78	52	17	50	28
	LDCS	16	15	62	62	12	68	48	17	48	25
	PH24E	100	101	160	101	100	96	54	83	101	99
	PHP02	102	101	162	103	99	96	54	98	107	98
	DIFF	2	0	2	2	1	0	0	16	6	2
	PR > T	0.37	0.91	0.45	0.18	0.02	0.89	0.44	0.02	0.00	0.00
	GDU SLK % MN	STK CNT % MN	FLT HT % MN	EAR HT % MN	RT LDG % MN	STA GRN % MN	STK LDG % MN	BRT STK % MN	GRH APP % MN	EYE SPT ABS	
TOTAL SUM	REPS	7	89	40	40	45	33	65	8	7	5
	LOCS	7	77	38	38	38	30	59	6	5	5
	PH24E	99	99	96	97	102	93	100	99	102	4
	PHP02	96	102	97	95	104	97	101	100	99	5
	DIFF	3	3	1	2	2	4	1	1	3	0
	PR > T	0.00	0.00	0.14	0.24	0.41	0.46	0.46	0.71	0.61	0.37
	ECB DPE ABS	ECB 1LF ABS	ECB 2SC ABS	DRP EAR % MN	NLF BLT ABS						
TOTAL SUM	REPS	12	2	4	45	4					
	LOCS	12	2	3	44	2					
	PH24E	100	7	7	100	7					
	PHP02	100	7	7	100	7					
	DIFF	1	1	0	0	0					
	PR > T	0.19	0.80	0.18	0.03	0.50					

*PR > T values are valid only for comparisons with LOCs >= 10.

TABLE 3B

Average Inbred By Tester Performance Comparing PH24E To PHJ90 Crossed To The Same Inbred Testers And Grown In The Same Experiments.										
	SEL IND % MN	PRM ABS	BU ACR ABS	BU ACR % MN	PRM SHD ABS	MST % MN	TST WT ABS	SDG VGR % MN	EST CNT % MN	GDU SHD % MN
TOTAL SUM	REPS	39	38	168	168	34	181	123	61	125
	LOCS	39	38	153	153	34	162	111	58	119
	PH24E	104	104	172	106	103	98	54	90	99
	PHJ90	97	103	162	99	103	100	54	100	99

TABLE 3B-continued

Average Inbred By Tester Performance Comparing PH24E To PHJ90 Crossed To The Same Inbred Testers And Grown In The Same Experiments.											
	DIFF PR > T	6 0.00	1 0.01	10 0.00	6 0.00	0 0.66	2 0.04	0 0.93	10 0.00	0 0.87	1 0.14
		GDU SLK % MN	STK CNT % MN	PLT HT % MN	EAR HT % MN	RT LDG % MN	STA GRN % MN	STK LDG % MN	BRT STK % MN	GRN APP % MN	EYE SPT ABS
TOTAL SUM	REPS	14	185	93	93	92	76	140	14	36	16
	LOCS	14	166	87	87	79	67	124	11	33	16
	PH24E	100	99	97	98	100	99	100	100	99	5
	PHJ90	101	100	97	99	104	82	100	102	103	4
	DIFF	0	1	0	1	4	18	0	2	4	1
	PR > T	0.64	0.26	0.65	0.48	0.27	0.00	0.96	0.25	0.34	0.01
		ECB DPE ABS	ECB 1LF ABS	ECB 2SC ABS	DRP EAR % MN	GLF SPT ABS	NLF BLT ABS				
TOTAL SUM	REPS	25	2	26	75	1	6				
	LOCS	25	2	23	70	1	3				
	PH24E	100	8	6	100	1	6				
	PHJ90	99	9	6	100	1	4				
	DIFF	0	1	1	0	0	1				
	PR > T	0.38		0.11	0.34		0.16				

*PR > T values are valid only for comparisons with Locs >= 10.

TABLE 3C

Average Inbred By Tester Performance Comparing PH24E To PHAP9 Crossed To The Same Inbred Testers And Grown In The Same Experiments.											
		SEL IND % MN	PRM ABS	BU ACR ABS	BU ACR % MN	PRM SHD ABS	MST % MN	TST WT ABS	SDG VGR % MN	EST CNT % MN	GDU SHD % MN
TOTAL SUM	REPS	64	65	256	256	56	259	179	121	230	94
	LOCS	64	65	246	246	56	249	173	113	220	90
	PH24E	102	105	173	105	104	98	55	89	98	100
	PHAP9	98	104	166	101	103	97	55	102	103	99
	DIFF	4	1	7	4	1	1	0	13	5	1
	PR > T	0.01	0.00	0.00	0.00	0.00	0.08	0.96	0.00	0.00	0.00
		GDU SLK % MN	STK CNT % MN	PLT HT % MN	EAR HT % MN	RT LDG % MN	STA GRN % MN	STK LDG % MN	BRT STK % MN	GRN APP % MN	EYE SPT ABS
TOTAL SUM	REPS	28	278	149	149	94	130	196	26	60	23
	LOCS	27	265	142	142	91	126	188	22	57	23
	PH24E	99	97	97	97	103	104	101	101	97	5
	PHAP9	100	100	104	105	102	98	102	98	91	5
	DIFF	1	3	6	8	1	6	1	4	6	0
	PR > T	0.03	0.00	0.00	0.00	0.70	0.04	0.13	0.34	0.07	0.26
		ECB DPE ABS	ECB 2SC ABS	DRP EAR % MN	GLF SPT ABS	NLF BLT ABS					
TOTAL SUM	REPS	57	30	124	3	7					
	LOCS	57	30	120	3	4					
	PH24E	100	7	100	3	7					
	PHAP9	99	6	100	3	6					
	DIFF	1	0	0	0	1					
	PR > T	0.02	0.18	0.03	0.42	0.04					

*PR > T values are valid only for comparisons with Locs >= 10.

TABLE 3D

Average Inbred By Tester Performance Comparing PH24E To PHKW3 Crossed To The Same Inbred Testers And Grown In The Same Experiments.											
		SEL IND % MN	PRH ABS	BU ACR ABS	BU ACR % MN	PRM SHD ABS	MST % MN	TST WT ABS	SDG VGR % MN	EST CNT % MN	GDU SHD % MN
TOTAL SUM	REPS	47	48	144	144	42	170	100	63	138	61
	LOCS	47	48	143	143	42	161	99	63	137	60
	PH24E	100	105	175	105	103	96	54	91	97	100
	PHKW3	95	102	168	101	103	92	56	94	97	100
	DIFF	5	2	7	4	0	4	2	3	0	0
	PR > T	0.05	0.00	0.00	0.00	0.61	0.00	0.00	0.40	0.86	0.88
		GDU SLK % MN	STK CNT % MN	PLT HT % MN	EAR HT % MN	RT LDG % MN	STA GRN % MN	STK LDG % MN	BRT STK % MN	GRN APP % MN	EYE SPT ABS
TOTAL SUM	REPS	13	180	99	99	92	95	135	8	38	20
	LOCS	13	171	98	98	85	91	129	6	37	20
	PH24E	99	98	98	98	104	103	100	102	104	6
	PHKW3	100	98	99	98	98	73	95	101	114	6
	DIFF	1	1	1	0	6	31	5	1	10	0
	PR > T	0.12	0.32	0.16	0.84	0.06	0.00	0.00	0.20	0.00	0.89
		ECB DPE ABS	ECB 1LF ABS	ECB 2SC ABS	DRP EAR % MN	NLF BLT ABS					
TOTAL SUM	REPS	21	1	12	81	6					
	LOCS	21	1	11	81	3					
	PH24E	100	8	5	100	6					
	PHKW3	100	7	4	100	3					
	DIFF	0	1	1	0	3					
	PR > T	0.33		0.02	0.20	0.01					

*PR > T values are valid only for comparisons with Locs ≥ 10 .

Deposits

A deposit of the seed of inbred PH24E is and has been maintained by Pioneer Hi-Bred International, Inc., 700 Capital Square, 400 Locust Street, Des Moines, Iowa 50309-2340, since prior to the filing date of this application. Access to this deposit will be available during the pendency of the application to the Commissioner of Patents and Trademarks and person determined by the Commissioner to be entitled thereto upon request. Upon allowance of any claims in the application, the Applicant(s) will make available to the public without restriction a deposit of at least 2500 seeds of inbred PH24E with the American Type Culture Collection (ATCC), Rockville, Md., 20852. The seeds deposited with the ATCC will be taken from the same deposit maintained at Pioneer Hi-Bred and described above. Additionally, Applicant(s) will meet all the requirements of 37 C.F.R. §1.801-1.809, including providing an indication of the viability of the sample when the deposit is made. This deposit of Inbred Maize Line PH24E will be maintained in the ATCC Depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the effective life of the patent, whichever is longer, and will be replaced if it ever becomes nonviable during that period. Applicant will impose no restrictions on the availability of the deposited material from the ATCC; however, Applicant has no authority to waive any restrictions imposed by law on the transfer of biological material or its transportation in commerce. Applicant does not waive any infringement of its rights granted under this patent or under the Plant Variety Protection Act (7 USC 2321 et seq.).

The foregoing invention has been described in detail by way of illustration and example for purposes of clarity and understanding. However, it will be obvious that certain

changes and modifications such as single gene modifications and mutations, somoclonal variants, variant individuals selected from large populations of the plants of the instant inbred and the like may be practiced within the scope of the invention, as limited only by the scope of the appended claims.

What is claimed is:

1. Seed of maize inbred line designated PH24E and having ATCC Accession No. 97925.

2. A maize plant and its parts produced by the seed of claim 1 and its plant parts.

3. Pollen of the plant of claim 2.

4. An ovule of the plant of claim 2.

5. A maize plant having all the physiological and morphological characteristics of the plant of claim 2.

6. A male sterile maize plant otherwise having all the physiological and morphological characteristics of the plant of claim 2.

7. A tissue culture of regenerable cells of an inbred maize plant PH24E, wherein the tissue regenerates plants capable of expressing all the morphological and physiological characteristics of the maize plant named PH24E, the seed of which have been deposited and have ATCC Accession No. 97925.

8. The tissue culture of claim 7, the tissue culture selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, silks, flowers, kernels, ears, cobs, husks, stalks, cells and protoplasts thereof.

9. A maize plant regenerated from the tissue culture of claim 7, capable of expressing all the morphological and physiological characteristics of inbred PH24E.

10. A method for producing first generation (F_1) hybrid maize seed comprising crossing a first inbred parent maize plant with a second inbred parent maize plant and harvesting

27

the resultant first generation (F_1) hybrid maize seed, wherein said first or second parent maize plant is the maize plant of claim 2.

11. The method of claim 10 wherein inbred maize plant PH24E is the female parent.

12. The method of claim 10 wherein inbred maize plant PH24E is the male parent.

28

13. An F_1 hybrid seed grown therefrom produced by crossing an inbred maize plant according to claim 2 with another, different maize plant.

14. An F_1 hybrid plant and its parts grown from the seed of claim 13.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5,689,034
DATED : November 18, 1997
INVENTOR(S) : Samuel Gregg Fullerton

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page.

Item [21], Appl. No., delete "523,660" and insert -- 08/523,660 --.

Column 25.


Delete rows 36-64 (the entire paragraph after "Deposits") and insert:

-- Applicant has made a deposit of at least 2500 seeds of inbred corn line PH24E with the American Type Culture Collection (ATCC), Rockville, MD 20852 USA, ATCC Deposit No. 97925. The seeds deposited with the ATCC on March 11, 1997 were taken from the deposit maintained by Pioneer Hi-Bred International, Inc., 700 Capital Square, 400 Locust Street, Des Moines, Iowa 50309-2340 since prior to the filing date of this application. This deposit of the Inbred Corn Line PH24E will be maintained in the ATCC depository, which is a public depository, for a period of thirty (30) years, or five years after the most recent request, or for the enforceable life of the patent, whichever is longer, and which will be replaced if it becomes nonviable during that period. Additionally, applicant has satisfied all the requirements of 37 C.F.R. §§1.801-1.809, including providing an indication of the viability of the sample. Applicant imposes no restrictions on the availability of the deposited material from the ATCC; however, applicant has no authority to waive any restrictions imposed by law on the transfer of biological material or its transportation in commerce. Applicant does not waive any infringement of its rights granted under this patent or under the Plant Variety Protection Act (7 U.S.C. § 2321 et seq.). Applicant has applied for U.S. Plant Variety Protection of PH24E under Application No. 9600204. --

Signed and Sealed this

Second Day of July, 2002

Attest:

A handwritten signature in black ink, appearing to read "James E. Rogan", with a horizontal line drawn underneath it.

Attesting Officer

JAMES E. ROGAN
Director of the United States Patent and Trademark Office