



US007820889B1

(12) **United States Patent**
Byrnes et al.

(10) **Patent No.:** **US 7,820,889 B1**
(45) **Date of Patent:** **Oct. 26, 2010**

(54) **INBRED CORN LINE LLD19BM**

OTHER PUBLICATIONS

(75) Inventors: **Kenneth J. Byrnes**, Bettendorf, IA (US);
Dale H. Storck, Sugar Grove, IL (US)

Australian Gov't 2008, The Biology of *Zea mays L. ssp mays* (maize or corn).*

(73) Assignee: **Agrigenetics, Inc.**, Indianapolis, IN (US)

Edwards et al 1981, Nucleic Acids Research 9(12): 2853-2869.*

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 469 days.

Allard, *In Principles of Plant Breeding*, John Wiley & Sons, Inc. pp 155-156, 1960.

Phillips, et al., *In Corn and Corn Improvement*, ASA Monograph No. 18, 3rd edition, p. 358, 1988.

Eshed, et al., Genetics (1996), vol. 143, pp. 1807-1817.

(21) Appl. No.: **11/823,977**

Kraft, et al., Theoretical Applied Genetics (2000), vol. 101, pp. 323-326.

(22) Filed: **Jun. 29, 2007**
(Under 37 CFR 1.47)

Murray, et al., Proceedings of the 43rd Annual Corn and Sorghum Industry Research Conference, vol. 43, p. 72-87, 1988.

* cited by examiner

Related U.S. Application Data

Primary Examiner—David H Kruse

(60) Provisional application No. 60/818,178, filed on Jun. 30, 2006.

(74) *Attorney, Agent, or Firm*—Eric J. Kraus

(51) **Int. Cl.**

(57) **ABSTRACT**

A01H 1/00 (2006.01)

An inbred corn line, designated LLD19BM, the plants and seeds of the inbred corn line LLD19BM, methods for producing a corn plant, either inbred or hybrid, produced by crossing the inbred corn line LLD19BM with itself or with another corn plant, and hybrid corn seeds and plants produced by crossing the inbred line LLD19BM with another corn line or plant and to methods for producing a corn plant containing in its genetic material one or more transgenes and to the transgenic corn plants produced by that method. This invention also relates to inbred corn lines derived from inbred corn line LLD19BM, to methods for producing other inbred corn lines derived from inbred corn line LLD19BM and to the inbred corn lines derived by the use of those methods.

A01H 4/00 (2006.01)

A01H 5/00 (2006.01)

A01H 5/10 (2006.01)

C12N 15/82 (2006.01)

(52) **U.S. Cl.** **800/320.1**; 435/412; 800/266; 800/269; 800/274; 800/275; 800/278; 800/298

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,491,296 A * 2/1996 Bergemann 800/275
5,523,520 A 6/1996 Hunsperger et al.
6,025,547 A 2/2000 Stucker
6,096,953 A 8/2000 Hoffbeck

51 Claims, No Drawings

1

INBRED CORN LINE LLD19BM

This application claims the benefit of U.S. Provisional Application No. 60/818,178, filed on Jun. 30, 2006.

FIELD OF THE INVENTION

This invention is in the field of corn breeding. In particular, the invention relates to an inbred corn line designated LLD19BM that includes plants and seeds of inbred corn line LLD19BM. Methods for producing corn plants, such as inbred corn plants, hybrid corn plants, or other corn plants, as by crossing inbred corn line LLD19BM with itself or any different corn plant are an integral part of this invention as are the resultant corn plants including the plant parts and seeds. This invention further relates to methods for producing LLD19BM-derived corn plants, to methods for producing male sterile LLD19BM corn plants and to methods for regenerating such plants from tissue cultures of regenerable cells as well as the plants obtained therefrom. Methods for producing a corn plant containing in its genetic material one or more transgenes and to the transgenic corn plants produced by that method are also a part of this invention.

BACKGROUND OF THE INVENTION

Corn (*Zea mays* L.) is the most important and abundant crop produced in the United States. Corn is used as human food, livestock feed, and as raw material in industry. The food uses of corn include kernels for human consumption, dry milling products such as grits, meal and flour, and wet milling products such as corn starch, corn syrups, and dextrose. Corn oil recovered from corn germ is a by-product of both dry and wet milling industries. Both grain and non-grain portions of corn plants are used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry.

Corn is used to produce ethanol while corn starch and flour are used in the paper and textile industries. Corn is also used in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications. Plant parts other than the grain of corn are also used in industry; for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

The goal of a corn breeder is to improve a corn plant's performance and therefore, its economic value by combining various desirable traits into a single plant. Improved performance is manifested in many ways. Higher yields of corn plants contribute to a more abundant food supply, a more profitable agriculture and a lower cost of food products for the consumer. Improved quality makes corn kernels more nutritious. Improved plant health increases the yield and quality of the plant and reduces the need for application of protective chemicals. Adapting corn plants to a wider range of production areas achieves improved yield and vegetative growth. Improved plant uniformity enhances the farmer's ability to mechanically harvest corn.

Corn is a monoecious plant, i.e., corn has imperfect flowers—male, pollen-producing flowers and separate female, pollen receiving flowers on the same plant. The male flowers are located at the top of the plant in the tassel, and the female flowers are located about midway up the stalk in the ear shoot. Each male flower has three anthers and each female flower includes a husk that envelops the cob and silks that emerge from the end of the cob and husks. Pollination is consummated by transfer of pollen from the tassels of the male flower to the silks of the female flowers.

2

Because corn has separate male and female flowers, corn breeding techniques take advantage of the plant's ability to be bred by both self-pollination and cross-pollination. Self-pollination occurs when pollen from the male flower is transferred to a female flower on the same plant. Cross-pollination occurs when pollen from the male flower is transferred to a female flower on a different plant.

A plant is sib-pollinated (a type of cross-pollination) when individuals within the same family or line are used for pollination (i.e. pollen from a family member plant is transferred to the silks of another family member plant). Self-pollination and sib-pollination techniques are traditional forms of inbreeding used to develop new inbred corn lines but other techniques exist to accomplish inbreeding. New inbred corn lines are developed by inbreeding heterozygous plants and practicing selection for superior plants for several generations until substantially homozygous plants are obtained. During the inbreeding process with corn, the vigor of the lines decreases and after a sufficient amount of inbreeding, additional inbreeding merely serves to increase seed of the developed inbred. Inbred corn lines are typically developed for use in the production of hybrid corn lines.

Natural, or open pollination, occurs in corn when wind blows pollen from the tassels to the silks that protrude from the tops of the ear shoot and may include both self- and cross-pollination. Vigor is restored when two different inbred lines are cross-pollinated to produce the first generation (F_1) progeny. A cross between two defined homozygous inbred corn plants always produces a uniform population of heterozygous hybrid corn plants and such hybrid corn plants are capable of being generated indefinitely from the corresponding inbred seed supply.

When two different, unrelated inbred corn parent plants are crossed to produce an F_1 hybrid, one inbred parent is designated as the male, or pollen parent, and the other inbred parent is designated as the female, or seed parent. Because corn plants are monoecious, hybrid seed production requires elimination or inactivation of pollen produced by the female parent to render the female parent plant male sterile. This serves to prevent the inbred corn plant designated as the female from self-pollinating. Different options exist for controlling male fertility in corn plants such as manual or mechanical emasculation (or detasseling), genetic male sterility, and application of gametocides. Incomplete removal or inactivation of the pollen in the female parent plant provides the potential for inbreeding which results in the unwanted production of self-pollinated or sib-pollinated seed. Typically, this seed is unintentionally harvested and packaged with hybrid seed.

The development of new inbred corn plants and hybrid corn plants is a slow, costly interrelated process that requires the expertise of breeders and many other specialists. The development of new hybrid corn varieties in a corn plant breeding program involves numerous steps, including: (1) selection of parent corn plants (germplasm) for initial breeding crosses; (2) inbreeding of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which individually breed true and are highly uniform; and, (3) crossing a selected inbred line with an unrelated line to produce the F_1 hybrid progeny having restored vigor.

Inbred corn plants and other sources of corn germplasm are the foundation material for all corn breeding programs. Despite the existence and availability of numerous inbred corn lines and other source germplasm, a continuing need still exists for the development of improved germplasm because existing inbred parent corn lines lose their commercial com-

petitiveness over time. The present invention addresses this need by providing a novel inbred parent corn line designated LLD19BM that contributes improved tonnage and digestibility relative to other hybrids containing the bm3 gene and in the same maturity group. To protect and to enhance yield production, trait technologies and seed treatment options provide additional crop plan flexibility and cost effective control against insects, weeds and diseases, thereby further enhancing the potential of hybrids with LLD19BM as a parent.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions of Plant Characteristics

In the description and examples that follow, a number of terms are used. To provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

Anther Color: Recorded at the time of pollen shed when anthers are actively dehiscing pollen as a standard color name and/or Munsell color code.

Anthocyanin in Brace Roots: This is a relative rating of the expression of anthocyanin in the brace roots (1=none, 2=faint, 3=50% purpling/moderate, and 4=dark) recorded two weeks after flowering.

Aphid Attractiveness: This represents the relative level of aphid numbers found on an emerging tassel up to and during the period of pollen shed rated as high, average or none.

Cob Color: Recorded as a standard color name and Munsell color code.

Digestibility: This represents the percent total dry matter digestibility of the whole plant.

DNDF: This is the percent of neutral detergent fiber digestibility measured in vitro.

Ear Mid Point Diameter: This represents the diameter of the unshelled ear in millimeters at the midpoint of the ear between its tip and its butt.

Ears Per Stalk: The total number of ears with seed set on each plant.

Ear Height: For inbred plants, this is the distance in centimeters from the ground to the highest placed ear node point of attachment of the ear shank of the uppermost developed ear on the stalk. For hybrid plants, this is the distance in centimeters from the ground to the base of the uppermost developed ear on the stalk.

Ear Node Leaf Width: Ear leaf node width at the top ear node at its maximum widest point measured in centimeters.

Ear Leaves: Ear leaves are defined as one or more distinct ear leaves on ear husks at flowering (usually >0.25 to 0.5") represented as present or absent. This characteristic may be difficult to determine, may be environmentally influenced, and is not recorded as present unless the ear leaves are present in sufficient size or on several plants in the middle of the row.

Ear Length: This is the length of an unhusked ear from the butt to the tip in centimeters.

Ear Node Leaf Length: This represents the length (cm) of the top ear node leaf from tip to collar.

Ear Position At Dry Husk: This represents the relative direction of the top ear observed 65 days after pollinating while still attached to the plant rated as 1=upright, 2=horizontal and 3=pending.

Ear Taper: This represents the relative taper of the unshelled ear rated as slight or nearly straight (1), average (2), or extreme or conical (3).

Endosperm Type: Endosperm is the material in the region of the kernel between the germ and the seed coat and is rated on the following scale: 1=sweet, 2=extra sweet (sh2), 3=nor-

mal starch, 4=high amylase starch, 5=waxy, 6=high protein, 7=high lysine, 8=supersweet (se), 9=high oil and 10=other—specify.

Environments: This is the number of different geographic locations where plants are grown together and in the same experiment.

50% Pollen (GDU from Planting): The number of GDUs after planting when 50% of the plants are shedding pollen.

50% Silk (GDU from Planting): The number of GDUs after emergence when 50% of the plants have extruded silk.

GDU: Growing degree unit(s) is a measure of the number of GDUs or heat units used in the tracking of flowering and maturation of inbred lines and hybrids. Growing degree units are calculated using the Barger Method, wherein the heat units for a 24-hour period are represented by the formula: $[(\text{Max. temp} + \text{Min. temp})/2] - 50$ and the highest maximum temperature used is 86° F. and the lowest minimum temperature used is 50° F.

Genotype: Refers to the genetic constitution of a corn plant or cell.

Glume Band: Recorded as absent or, if present, as a standard color name and Munsell color code.

Glume Color: Recorded after exposure to sunlight and just before extruding anthers as a standard color name and Munsell color code.

Hard Endosperm Color: This is the color of the region of the endosperm between the flourey endosperm and the aleurone layer in yellow dent corn recorded as a standard color name and a Munsell color code.

Husk Tightness: This represents the relative ability for husks to be removed either manually or through commercial production husking beds 65 days after flowering rated as loose (2), average (5) and 8 (tight).

Kernel Crown Color: This is the color of the portion of the kernel distal to the tip cap recorded as a standard color name and Munsell color code.

K Row Alignment: This is kernel row alignment and is scored as straight, slightly curved or spiral determined by standing the unshelled ear on its base and looking down at the tip.

Kernel Rows: The presence (distinct) or absence (indistinct) of defined kernel rows.

Lateral Tassel Branches: This represents the number of primary lateral tassel branches that originate from the central spike.

Leaf Color: Recorded as standard color name and Munsell color code.

Lignin Content: This represents the lignin content as a percent of the whole plant on a dry matter basis.

Moisture: Actual percentage moisture of the grain at harvest.

Munsell Code: This refers to a standard color reference, the Munsell Color Chart for Plant Tissues.

NDF: This is the neutral detergent fiber content as a percent of the whole plant on a dry matter basis.

Number of Kernel Rows: This is the average total number of kernel rows on the ear. If the rows are indistinct, then this value is an average number of kernels located around the circumference of the ear at the mid-point of its length.

Number of Leaves Above Ear: This represents the average number of leaves above the uppermost ear leaf.

Phenotype: Refers to the physical or external appearance of a corn plant.

Plant Height: For inbred plants, this is the plant height in centimeters from the ground to the tip of the tassel. For hybrid plants, this is the plant height in centimeters from the ground to the top leaf collar.

Root Lodging: This is the percentage of corn plants that root lodge, i.e., those plants that lean from the vertical axis at an approximate 30° angle or greater. For inbred plants, this measurement is taken just before anthesis whereas for hybrids, this measurement is taken at harvest.

Silk Color: Recorded 3 days after emergence using standard color name and Munsell color code.

Stalk Lodging: This is the percentage of corn plants that stalk lodge, i.e., those plants that are broken over, at or below the top ear node at harvest.

Standard Color Names: These color names include light green, medium green, dark green, very dark green, green-yellow, pale yellow, yellow, yellow orange, salmon, pink-orange, pink, light red, cherry red, red, red and white, pale purple, purple, colorless, white, white capped, buff, tan, brown, bronze and variegated.

Tassel Length: This is the length of the tassel from the top leaf collar to the tassel tip measured in centimeters.

Tons/Acre: This represents the amount in tons of all harvested above ground plant material per acre adjusted to 65% silage moisture.

Years: This refers to the number of calendar years included in a comparison.

II. Inbred Corn Line LLD19BM

A. Inbred Corn Plant LLD19BM

In accordance with one aspect of the present invention, provided is a new yellow dent inbred corn seed and plants thereof designated LLD19BM. The present invention further relates to a method for producing inbred corn seeds that includes, but is not limited to, the steps of planting seed of inbred corn LLD19BM in proximity to itself or to different seed from a same family or line, growing the resulting corn plants under self-pollinating or sib-pollinating conditions with adequate isolation, and harvesting resultant seed obtained from such inbred plants using techniques standard in the agricultural arts such as would be necessary to bulk-up seed such as for hybrid production. The present invention also relates to inbred seed produced by such a method.

In any cross between inbred corn plant LLD19BM and another inbred corn plant, LLD19BM may be designated as the male (pollen parent) or the female (seed parent). Optionally, the seed of inbred corn line LLD19BM may be pretreated to increase resistance of the seed and/or seedlings to stressed conditions, and further, the corn plants or surrounding soil may be treated with one or more agricultural chemicals before harvest. Such agricultural chemicals may include herbicides, insecticides, pesticides and the like. The present invention also relates to a corn plant that expresses substantially all of the physiological and morphological characteristics of inbred corn plant LLD19BM and to a substantially homogenous population of corn plants having all the physiological and morphological characteristics of inbred corn plant LLD19BM. Any corn plants produced from inbred corn plant LLD19BM are contemplated by the present invention and are, therefore, within the scope of this, invention. A description of physiological and morphological characteristics of corn plant LLD19BM is presented in Table 1.

TABLE 1

Inbred Trait	Value
50% Pollen (GDU from Planting)	1553
50% Silk (GDU from Planting)	1598
Anther Color (standard)	6 (Pale Yellow)

TABLE 1-continued

Inbred Trait	Value
5 Anther Color (Munsell)	5Y 8/6
Glume Color (standard)	01 & 12 (Light Green with faint Red)
Glume Color (Munsell)	2.5GY 7/4, 5R 4/6
Silk Color (standard)	5 (Pale Green-Yellow)
Silk Color (Munsell)	2.5GY 8/4
10 Glume Band (Pr/Abs)	Absent
Aphid Attractiveness (H, Av, N)	N
Plant Height (cm)	229
Ear Height (cm)	93
Ear Leaves (Pr/Abs)	Absent
Anthocyanin in brace roots	3.7
15 Leaf Color (standard)	3 (Dark Green)
Leaf Color (Munsell)	5GY 4/4
Number of Leaves Above Ear	5.5
Ear Node Leaf Length (cm)	74
Ear Node Leaf Width (cm)	10
Tassel Length (cm)	46
Lateral Tassel Branches	4
Ears Per Stalk	1
20 Ear Position at Dry Husk	n/a (upright)
Husk Tightness	5
Ear Length (cm)	18
Number of Kernel Rows	18
25 Kernel Rows (1= distinct, 2 = indistinct)	1
K Row Alignment	2
Ear Taper (1-slight, 2-average, 3 extreme)	2
30 Ear Mid Point Diameter (mm)	46
Endosperm Type	3
Cob Color (standard)	14
Cob Color (Munsell)	10R 3/6
Hard Endosperm Color (standard)	24 & 8 (Bronze and Yellow-Orange)
Hard Endosperm Color (Munsell)	10R 4/10
35 Kernel Crown Color (standard)	7 (Yellow)
Kernel Crown Color (Munsell)	2.5Y 8/10

Flag leaf commonly lays over tassel at the beginning of pollen shed LLD19BM has relatively weak roots compared to other inbreds of similar maturity

It should be appreciated by one having ordinary skill in the art that, for the quantitative characteristics identified in Table 1, the values presented are typical values. These values may vary due to the environment and accordingly, other values that are substantially equivalent are also within the scope of the invention.

Inbred corn line LLD19BM shows uniformity and stability within the limits of environmental influence for the traits described in Table 1. Inbred LLD19BM has been self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to ensure the homozygosity and phenotypic stability necessary to use in large scale, commercial production. The line has been increased both by hand and sib-pollinated in isolated fields with continued observations for uniformity. No variant traits have been observed or are expected in LLD19BM.

The present invention also relates to one or more corn plant parts of inbred corn plant LLD19BM. Corn plant parts include plant cells, plant protoplasts, plant cell tissue cultures from which corn plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, ovules, flowers, seeds, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, brace roots, lateral tassel branches, anthers, tassels, glumes, silks, tillers, and the like.

B. Inbred Corn Seed Designated LLD19BM

A corn kernel is composed of four structural parts: (1) the pericarp, which is a protective outer covering (also known as

bran or hull); (2) the germ (also known as an embryo); (3) the endosperm; and, (4) the tip cap, which is the point of attachment to the cob. Another aspect of the present invention is one or more parts of inbred corn seed LLD19BM, such as the pericarp of inbred corn seed LLD19BM or the germ and/or the endosperm of inbred corn seed LLD19BM which remain upon removal of the pericarp and adhering remnants of the seed coat.

Inbred corn seed designated LLD19BM may be provided as a substantially homogenous composition of inbred corn seed designated LLD19BM, that is, a composition that consists essentially of inbred corn seed LLD19BM. Such a substantially homogenous composition of inbred corn seed LLD19BM is substantially free from significant numbers of other inbred and/or hybrid seed so that the inbred seed forms from about 90% to about 100% of the total seed. Preferably, a substantially homogenous composition of the inbred corn seed contains from about 98.5%, 99%, or 99.5% to about 100% of the inbred seed, as measured by seed grow outs. The substantially homogenous composition of inbred corn seed of the invention may be separately grown to provide substantially homogenous populations of inbred corn plants. However, even if a population of inbred corn plants is present in a field with other different corn plants, such as in a commercial seed-production field of single-cross hybrid corn planted in a ratio of 1 male pollinator row to 4 female seed-parent rows, such a population would still be considered to be within the scope of the present invention.

Corn yield is affected by the conditions to which seeds and seedlings (young plants grown from seeds) are exposed. Seeds and seedlings may be exposed to one of, or a combination of, for example, cold, drought, salt, heat, pollutants, and disease, all of which are conditions that potentially retard or prevent the growth of crops therefrom. For example, temperature extremes are typical in the upper Midwest region of the United States. Furthermore, diseases evolved from pathogens and deterioration caused by fungi are potentially harmful to seeds and seedlings. Thus, it is desirable to treat seeds as by coating or impregnating the seeds with compositions that render the seeds and seedlings grown therefrom more hardy when exposed to such adverse conditions.

Accordingly, another aspect of the present invention relates to a coated and/or impregnated seed or corn inbred line designated LLD19BM and to coated and/or impregnated seed derived therefrom. Various agents have been used to treat seeds to increase resistance of the plants to stressed conditions, such as cold, drought, salt, and fungi. Such agents include, for example, sodium methylphenyl-pentadienate, trichloroacetic acid, polyoxyalkylene-organo-siloxane block copolymer, 5-aminolevulinic acid, salicylic acid, thiamethoxam, potassium chloride, and polyvinyl alcohol and are useful alone, or in combination in the present invention.

When pre-treating seeds according to the present invention such as before the seeds are planted, the seeds are contacted with the composition of interest, as by coating seeds, spraying seeds, and soaking seeds or a combination thereof, by methods well known to those skilled in the art.

C. Deposit Information

Applicants have made a deposit of at least 2,500 seeds of inbred corn plant LLD19BM with the American Type Culture Collection (ATCC), Manassas, Va. 20110 USA, under ATCC Accession No. PTA-9814. The seeds deposited with the ATCC on Feb. 10, 2009 were taken from a deposit maintained by Agrigenetics, Inc. d/b/a Mycogen Seeds since prior to the filing date of this application. Access to this deposit will be available during the pendency of the application to the Commissioner of Patents and Trademarks and persons determined

by the Commissioner to be entitled thereto upon request. Upon allowance of any claims in the application, the Applicant(s) will maintain and will make this deposit available to the public pursuant to the Budapest Treaty.

III. Processes of Preparing Novel Corn Plants

A. Novel Inbred Corn Plants Obtained from Inbred LLD19BM

Various breeding schemes may be used to produce new inbred corn lines from LLD19BM. In one method, generally referred to as the pedigree method, LLD19BM may be crossed with another different corn plant such as a second inbred parent corn plant, which either itself exhibits one or more selected desirable characteristic(s) or imparts selected desirable characteristic(s) to a hybrid combination. Examples of potentially desired characteristics include greater yield, better stalks, better roots, reduced time to crop maturity, better agronomic quality, higher nutritional value, higher starch extractability or starch fermentability, resistance and/or tolerance to insecticides, herbicides, pests, heat and drought, and disease, and uniformity in germination times, stand establishment, growth rate, maturity and kernel size. If the two original parents corn plants do not provide all the desired characteristics, then other sources can be included in the breeding population. Elite inbred lines, that is, pure breeding, homozygous inbred lines, can also be used as starting materials for breeding or source populations from which to develop inbred lines.

Thereafter, resulting seed is harvested and resulting superior progeny plants are selected and selfed or sib-mated in succeeding generations, such as for about 5 to about 7 or more generations, until a generation is produced that no longer segregates for substantially all factors for which the inbred parents differ, thereby providing a large number of distinct, pure-breeding inbred lines.

In another embodiment for generating new inbred corn plants, generally referred to as backcrossing, one or more desired traits may be introduced into inbred parent corn plant LLD19BM (the recurrent parent) by crossing the LLD19BM plants with another corn plant (referred to as the donor or non-recurrent parent) which carries the gene(s) encoding the particular trait(s) of interest to produce F_1 progeny plants. Both dominant and recessive alleles may be transferred by backcrossing. The donor plant may also be an inbred, but in the broadest sense can be a member of any plant variety or population cross-fertile with the recurrent parent. Next, F_1 progeny plants that have the desired trait are selected. Then, the selected progeny plants are crossed with LLD19BM to produce backcross progeny plants. Thereafter, backcross progeny plants comprising the desired trait and the physiological and morphological characteristics of corn inbred line LLD19BM are selected. This cycle is repeated for about one to about eight cycles, preferably for about 3 or more times in succession to produce selected higher backcross progeny plants that comprise the desired trait and all of the physiological and morphological characteristics of corn inbred line LLD19BM listed in Table 1 as determined at the 5% significance level when grown in the same environmental conditions. Exemplary desired trait(s) include insect resistance, cytoplasmic male sterility, enhanced nutritional quality, waxy starch, herbicide resistance, yield stability, yield enhancement and resistance to bacterial, fungal and viral disease. One of ordinary skill in the art of plant breeding would appreciate that a breeder uses various methods to help determine which plants should be selected from the segregating populations and ultimately which inbred lines will be used to develop

hybrids for commercialization. In addition to the knowledge of the germplasm and other skills the breeder uses, a part of the selection process is dependent on experimental design coupled with the use of statistical analysis. Experimental design and statistical analysis are used to help determine which plants, which family of plants, and finally which inbred lines and hybrid combinations are significantly better or different for one or more traits of interest. Experimental design methods are used to assess error so that differences between two inbred lines or two hybrid lines can be more accurately determined. Statistical analysis includes the calculation of mean values, determination of the statistical significance of the sources of variation, and the calculation of the appropriate variance components. Either a five or a one percent significance level is customarily used to determine whether a difference that occurs for a given trait is real or due to the environment or experimental error. One of ordinary skill in the art of plant breeding would know how to evaluate the traits of two plant varieties to determine if there is no significant difference between the two traits expressed by those varieties. For example, see Fehr, Walt, *Principles of Cultivar Development*, p. 261-286 (1987) which is incorporated herein by reference. Mean trait values may be used to determine whether trait differences are significant, and preferably the traits are measured on plants grown under the same environmental conditions.

This method results in the generation of inbred corn plants with substantially all of the desired morphological and physiological characteristics of the recurrent parent and the particular transferred trait(s) of interest. Because such inbred corn plants are heterozygous for loci controlling the transferred trait(s) of interest, the last backcross generation would subsequently be selfed to provide pure breeding progeny for the transferred trait(s).

Backcrossing may be accelerated by the use of genetic markers such as SSR, RFLP, SNP or AFLP markers to identify plants with the greatest genetic complement from the recurrent parent.

Direct selection may be applied where a single locus acts as a dominant trait, such as the herbicide resistance trait. For this selection process, the progeny of the initial cross are sprayed with the herbicide before the backcrossing. The spraying eliminates any plants which do not have the desired herbicide resistance characteristic, and only those plants which have the herbicide resistance gene are used in the subsequent backcross. In the instance where the characteristic being transferred is a recessive allele, it may be necessary to introduce a test of the progeny to determine if the desired characteristic has been successfully transferred. The process of selection, whether direct or indirect, is then repeated for all additional backcross generations.

It should be appreciated by those having ordinary skill in the art that backcrossing can be combined with pedigree breeding as where inbred LLD19BM is crossed with another corn plant, the resultant progeny are crossed back to inbred LLD19BM and thereafter, the resulting progeny of this single backcross are subsequently inbred to develop new inbred lines. This combination of backcrossing and pedigree breeding is useful as when recovery of fewer than all of the LLD19BM characteristics than would be obtained by a conventional backcross are desired.

In an additional embodiment of the present invention, new inbred corn plants can be developed by a method generally referred to as haploid breeding. In this methodology, haploid plants are generated from diploid, heterozygous corn plants that result from crossing inbred corn plant LLD19BM with another, different corn plant. Such haploid corn plants may be

generated by methods known to those skilled in the art such as by culturing haploid anthers or embryos from a diploid plant. Alternately, such haploid corn plant may be generated by crossing the diploid heterozygous corn plant with a corn plant that comprises a haploid inducing gene, such as the mutant gene "indeterminate gametophyte" (ig), which, when present in the female parent results in offspring with a greatly enhanced frequency of haploids of both maternal and paternal origin. Thereafter, homozygous diploid plants are produced by the doubling of a set of chromosomes (1N) from a haploid plant generated by self-pollination such as through use of a doubling agent, such as colchicine, nitrous oxide gas, heat treatment and trifluralin. See, e.g., Wan et al., "Efficient Production of Doubled Haploid Plants Through Colchicine Treatment of Anther-Derived Maize Callus", *Theoretical and Applied Genetics*, 77:889-892, 1989 and U.S. Patent Application No. 20030005479 the disclosure of which is expressly incorporated herein by reference. The technique of haploid breeding is advantageous because no subsequent inbreeding is required to obtain a homozygous plant from a heterozygous source. Thus, in another aspect of this invention a new inbred corn plant is developed by a method that includes the steps of crossing LLD19BM or a hybrid made with LLD19BM with another inbred corn plant having a propensity to generate haploids to produce haploid progeny plants, and selecting desirable inbred corn plants from the haploid progeny plants.

The present invention also relates to novel corn plants produced by a method generally referred to as mutation breeding whereby one or more new traits may be artificially introduced into inbred line LLD19BM. The goal of artificial mutagenesis is to increase the rate of mutation for a desired characteristic. Mutation rates can be increased by many different means including temperature, long-term seed storage, tissue culture conditions, radiation; such as X-rays, Gamma rays (e.g. cobalt 60 or cesium 137), neutrons, (product of nuclear fission by uranium 235 in an atomic reactor), Beta radiation (emitted from radioisotopes such as phosphorus 32 or carbon 14), or ultraviolet radiation (preferably from 2500 to 2900 nm), or chemical mutagens (such as base analogues (5-bromo-uracil), related compounds (8-ethoxy caffeine), antibiotics (streptonigrin), alkylating agents (sulfur mustard, nitrogen mustard, epoxides, ethylenamines, sulfates, sulfonates, sulfones, lactones), azide, hydroxylamine, nitrous acid, or acridines. Once a desired trait is observed through mutagenesis and selected, the trait may then be incorporated into existing germplasm by traditional breeding techniques. Details of mutation breeding can be found in "Principals of Cultivar Development" Fehr, 1993 Macmillan Publishing Company the disclosure of which is incorporated herein by reference.

The mutagenesis treatment may be applied to various stages of plant development, including but not limited to cell cultures, embryos, microspores and shoot apices as well as to corn kernels. By way of example, pollen may be mixed with a solution of 1 ml EMS and 100 mls Fisher paraffin oil (stock diluted by 1 ml and 15 mls oil solution) every minute for the first 5 minutes and then every five minutes for 45 minutes to keep the pollen suspended. Thereafter, the pollen/paraffin oil solution is brushed onto the silks of developing ears. A tassel bag is used to cover the ear to prevent contamination. The ear is picked at maturity and then resultant seeds or the plants therefrom are screened for the desired mutant trait(s).

Once inbred lines are created, the next step is to determine if the inbreds have any value. This is accomplished by techniques of measuring the combining ability of the new inbred plant, as well as the performance of the line itself. Combining ability refers to a line's contribution as a parent when crossed

11

with other lines to form hybrids. Specific combining ability (SCA) refers to the ability of a line to cross to another specific inbred to form a hybrid. General combining ability (GCA) refers to the ability of a line to cross to a wide range of lines to form hybrids. The methodology of forming hybrids to evaluate an inbred line's contribution as a parent for the purpose of selecting superior lines is interchangeably known as experimental, top or test crossing.

B. Novel Inbred Plants Obtained from a Hybrid Having Inbred LLD19BM as a Parent

In accordance with processes of the present invention, a hybrid plant having inbred LLD19BM as a parent is crossed with itself or any different corn plant such as an inbred corn plant or a hybrid corn plant to develop a novel inbred line. For example, a hybrid corn plant having inbred corn plant LLD19BM as a parent may be inbred, i.e., crossed to itself or sib-pollinated, and the resulting progeny each selfed for about 5 to about 7 or more generations, thereby providing a set of distinct, pure-breeding inbred lines wherein each of the lines received all of its alleles from the hybrid corn plant having inbred corn plant LLD19BM as a parent. Double haploid methods can also be used to obtain an inbred corn plant that is homozygous at essentially every locus, wherein the inbred corn plant received all of its alleles from the hybrid corn plant having inbred corn plant LLD19BM as a parent. In other embodiments, a hybrid corn plant having inbred corn plant LLD19BM as a parent is crossed with a different corn plant that may include any inbred corn plant that is not inbred corn plant LLD19BM, any hybrid corn plant that does not have LLD19BM as a parent, another germplasm source, a haploid or mutation inducing stock, or a trait donor plant, thereby providing a set of distinct, pure-breeding inbred lines. The resulting inbred lines could then be crossed with other inbred or non-inbred lines and the resulting inbred progeny analyzed for beneficial characteristics. In this way, novel inbred lines conferring desirable characteristics could be identified.

C. "Chasing Selfs"

Both female and male inbred seed may occasionally be found within a commercial bag of hybrid seed. Chasing the selfs involves identifying inbred plants within a stand of corn that has been grown from a bag of hybrid corn seed. Once the seed is planted, the inbred plants may be identified and selected due to their decreased vigor, i.e., by their short stature, narrower leaves, and smaller tassels relative to the hybrid plants that grow from the hybrid seed which predominates in a commercial bag of hybrid seed. By locating the inbred plants, isolating them from the rest of the plants, and self-pollinating them (i.e., "chasing selfs"), a breeder can obtain an inbred line that is identical to an inbred parent used to produce the hybrid.

Accordingly, another embodiment of the present invention is directed to a method for producing inbred corn plant LLD19BM comprising: (a) planting a collection of seed, such as a collection of seed comprising seed of a hybrid, one of whose parents is inbred corn plant LLD19BM, the collection also comprising seed of the inbred; (b) growing plants from said collection of seed; (c) identifying inbred parent plants; (d) controlling pollination in a manner which preserves homozygosity of the inbred parent plant; and, (e) harvesting resultant seed. Step (c) may further comprise identifying plants with decreased vigor, i.e., plants that appear less robust than the other plants, or identifying plants that have a genetic profile in accordance with the genetic profile of LLD19BM, such as an SSR genetic profile in accordance with Table 5 herein. Corn plants capable of expressing substantially all of the physiological and morphological characteristics of inbred

12

corn plant LLD19BM include corn plants obtained by chasing selfs from a bag of hybrid seed.

One having skill in the art will recognize that once a breeder has obtained inbred corn plant LLD19BM by chasing selfs from a bag of hybrid seed, the breeder can then produce new inbred plants such as by sib-pollinating, i.e., crossing the inbred corn plant LLD19BM with another inbred corn plant LLD19BM, or by crossing the inbred corn plant LLD19BM with a hybrid corn plant obtained by growing the collection of seed.

IV. Novel Hybrid Plants

A. Novel Hybrid Seeds and Plants

In yet another aspect of the invention, processes are provided for producing corn seeds or plants, which processes generally comprise crossing a first parent corn plant with a second parent corn plant wherein at least one of the first parent corn plant or the second parent corn plant is inbred parent corn plant LLD19BM. In some embodiments of the present invention, the first inbred corn plant is LLD19BM and is a female and in other embodiments the first inbred corn plant is LLD19BM and is a male. These processes may be further exemplified as processes for preparing hybrid corn seed or plants, wherein a first inbred corn plant is crossed with a second corn plant of a different, distinct variety to provide a hybrid that has, as one of its parents, the inbred corn plant variety LLD19BM. In this case, a second inbred variety is selected which confers desirable characteristics when in hybrid combination with the first inbred line. In these processes, crossing will result in the production of seed. The seed production occurs regardless whether the seed is collected.

Any time the inbred corn plant LLD19BM is crossed with another, different corn inbred, a first generation (F_1) corn hybrid plant is produced. As such, an F_1 hybrid corn plant may be produced by crossing LLD19BM with any second inbred corn plant. Therefore, any F_1 hybrid corn plant or corn seed which is produced with LLD19BM as a parent is part of the present invention.

When inbred corn plant LLD19BM is crossed with another inbred plant to yield a hybrid, the original inbred can serve as either the maternal or paternal plant with basically, the same characteristics in the hybrids. Occasionally, maternally inherited characteristics may express differently depending on the decision of which parent to use as the female. However, often one of the parental plants is preferred as the maternal plant because of increased seed yield and preferred production characteristics, such as optimal seed size and quality or ease of tassel removal. Some plants produce tighter ear husks leading to more loss, for example due to rot, or the ear husk may be so tight that the silk can not completely push out of the tip preventing complete pollination resulting in lower seed yields. There can be delays in silk formation which deleteriously affect timing of the reproductive cycle for a pair of parental inbreds. Seed coat characteristics can be preferable in one plant which may affect shelf life of the hybrid seed product. Pollen can shed better by one plant, thus rendering that plant as the preferred male parent. It is generally preferable to use LLD19BM as the male parent.

In embodiments of the present invention, the first step of "crossing" the first and the second parent corn plants comprises planting, preferably in pollinating proximity, seeds of a first inbred corn plant and a second, distinct inbred corn plant. As discussed herein, the seeds of the first inbred corn plant and/or the second inbred corn plant can be treated with compositions that render the seeds and seedlings grown therefrom more hardy when exposed to adverse conditions.

13

A further step comprises cultivating or growing the seeds of the first and second parent corn plants into plants that bear flowers. If the parental plants differ in timing of sexual maturity, techniques may be employed to obtain an appropriate nick, i.e., to ensure the availability of pollen from the parent corn plant designated the male during the time at which silks on the parent corn plant designated the female are receptive to the pollen. Methods that may be employed to obtain the desired nick include delaying the flowering of the faster maturing plant, such as, but not limited to delaying the planting of the faster maturing seed, cutting or burning the top leaves of the faster maturing plant (without killing the plant) or speeding up the flowering of the slower maturing plant, such as by covering the slower maturing plant with film designed to speed germination and growth or by cutting the tip of a young ear shoot to expose silk.

In a preferred embodiment, the corn plants are treated with one or more agricultural chemicals as considered appropriate by the grower.

A subsequent step comprises preventing self-pollination or sib-pollination of the plants, i.e., preventing the silks of a plant from being fertilized by any plant of the same variety, including the same plant. This is preferably done in large scale production by controlling the male fertility, e.g., treating the flowers so as to prevent pollen production or alternatively, using as the female parent a male sterile plant of the first or second parent corn plant (i.e., treating or manipulating the flowers so as to prevent pollen production, to produce an emasculated parent corn plant or using as a female, a cytoplasmic male sterile version of the corn plant). This control may also be accomplished in large scale production by physical removal of the tassel from the female plant, either by pulling the tassel by hand, cutting with a rotary cutter, or pulling with a mechanical tassel pulling machine. In small scale production, corn breeder's shoot bags, usually plastic or glassine, applied to cover the ear shoot prior to the extrusion of silks provide effective control of unwanted self-pollination or sib-pollination.

Yet another step comprises allowing cross-pollination to occur between the first and second parent corn plants. When the plants are not in pollinating proximity, this is done by placing a bag, usually paper, over the tassels of the first plant and another shoot bag over the ear shoot, prior to the extrusion of silk, of the incipient ear on the second plant. The bags are left in place usually overnight. Since pollen stops shedding each day and loses viability and new pollen is shed each morning, this assures that the silks are not pollinated from other pollen sources, that any stray pollen on the tassels of the first plant is dead, and that the only pollen transferred comes from the first plant. The pollen bag over the tassel of the first plant is then shaken vigorously to enhance release of pollen from the tassels and removed from the first plant. Finally, in one continuous motion, the shoot bag is removed from the silks of the incipient ear on the second plant, and the pollen bag containing the captured pollen is placed over the silks of the incipient ear of the second plant, shaken again to disperse the captured pollen, and left in place covering the developing ear to prevent contamination from any unwanted fresh airborne pollen. In large scale production, crossing is accomplished by isolated open-pollinated crossing fields whereby corn plants of the parent designated as the female, which are controlled for male fertility, are allowed to be pollinated by other plants of a different corn type where such plants are adjacent to the plants designated as the female parent.

A further step comprises harvesting the seeds, near or at maturity, from the ear of the plant that received the pollen. In a particular embodiment, seed is harvested from the female

14

parent plant, and when desired, the harvested seed can be grown to produce a first generation (F_1) hybrid corn plant.

Yet another step comprises drying and conditioning the seeds, including the treating, sizing (or grading) of seeds, and packaging for sale to growers for the production of grain or forage. As with inbred seed, it may be desirable to treat hybrid seeds with compositions that render the seeds and seedlings grown therefrom more hardy when exposed to adverse conditions. Mention should be made that resulting hybrid seed is sold to growers for the production of grain and forage and not for breeding or seed production.

Still further, the present invention provides a hybrid corn plant produced by growing the harvested seeds produced on the male-sterile plant as well as grain produced by the hybrid corn plant.

A single cross hybrid is produced when two different inbred parent corn plants are crossed to produce first generation F_1 hybrid progeny. Generally, each inbred parent corn plant has a genotype which complements the genotype of the other inbred parent. Typically, the F_1 progeny are more vigorous than the respective inbred parent corn plants. This hybrid vigor, or heterosis, is manifested in many polygenic traits, including markedly improved yields and improved stalks, roots, uniformity and insect and disease resistance. It is for this reason that single cross F_1 hybrids are generally the most sought after hybrid. A three-way, or modified single-cross hybrid is produced from three inbred lines (or synthetics) where two of the inbred lines are crossed ($A \times B$) and then the resulting F_1 hybrid is crossed with the third inbred ($A \times B$) $\times$ C, as where a modified female is used in the cross. A modified female provides an advantage of improved seed parent yield whereas a modified male improves pollen flow. A double cross hybrid is produced from four inbred lines crossed in pairs ($A \times B$ and $C \times D$), thereby resulting in two F_1 hybrids that are crossed again. Double cross hybrids are more common in countries wherein less demand exists for higher yielding single cross hybrids. Synthetic populations or crosses are developed by crossing two or more inbred lines (or hybrids, or germplasm sources) together and then employing one of many possible techniques to random mate the progeny. Random mating the progeny is any process used by plant breeders to make a series of crosses that will create a new germplasm pool from which new breeding lines can be derived. Much of the hybrid vigor exhibited by F_1 hybrids is lost in the next generation (F_2). Consequently, seed from hybrids are not typically used for planting stock.

The utility of the inbred plant LLD19BM also extends to crosses with species other than the mays species, such as *diploperennis*, *luxurians*, and *perennis*. Commonly, suitable species will be of the family Gramineae, and especially of the genera *Zea*, *Tripsacum*, *Coix*, *Schleachne*, *Polytoca*, *Chionachne*, and *Trilobachne*. Of these, *Zea* and *Tripsacum*, are most preferred. Varieties of the grain sorghum *Sorghum bicolor* (L.) Moench can be crossed with inbred corn line LLD19BM.

B. Physical Description of F_1 Hybrids and F_1 Hybrid Comparison

As mentioned above, testcross hybrids are progressively eliminated following detailed evaluations of their phenotype, including formal comparisons with other commercially successful hybrids. Research small-plot trials and commercial strip trials are used to compare the phenotypes of hybrids grown in as many environments as possible. They are performed in many environments to assess overall performance of the new hybrids and to select optimum growing conditions. Because the corn is grown in close proximity, differential effects of environmental factors that affect gene expression,

15

such as moisture, temperature, sunlight, and pests, are minimized. For a decision to be made to advance a hybrid, it is not necessary that the hybrid be better than all other hybrids. Rather, significant improvements must be shown in at least some traits that would create value for some applications or markets. Some testcross hybrids are eliminated despite being similarly competitive relative to the current commercial hybrids because of the cost to bring a new hybrid to market requires a new product to be a significant improvement over the existing product offering. Such hybrids may also be licensed to other parties who have a need in their commercial product portfolio.

The present invention provides F1 hybrid corn plants obtained from the corn plant LLD19BM. The physical characteristics of exemplary hybrids produced using LLD19BM as one inbred parent are set forth in Table 2-4. The results in Table 2 compare inbred LLD19BM and inbred D, when each inbred is crossed to the same tester line. Tables 3-4 also present a comparison of performance data for hybrids made with LLD19BM as one parent, versus selected commercial hybrids including Mycogen F2F797 and Mycogen F697.

TABLE 2

Characteristic	Tester A $\times$ LLD19BM	Tester A $\times$ Inbred D
Moisture	68.9	68.4
Tons/Acre	26.5	26.2
Stalk Lodging	0.0	0.0
Root Lodging	3.9	1.4
Plant Height	92.0	82.0
Ear Height	48.0	47.0
NDF	42.4	41.5
Digestibility	88.7	89.8
DNDF	72.3	73.7
Lignin Content	1.4	1.3
Environments	9	9
Years	1	1

TABLE 3

Characteristic	Tester A $\times$ LLD19BM	Mycogen F2F797
Moisture	68.9	69.2
Tons/Acre	26.5	25.2
Stalk Lodging	0.0	2.5
Root Lodging	3.9	1.7
Plant Height	92.0	86.0
Ear Height	48.0	48.0
NDF	42.4	42.4
Digestibility	88.7	89.7
DNDF	72.3	74.6
Lignin Content	1.4	1.3
Environments	9	9
Years	1	1

TABLE 4

Characteristic	Tester E $\times$ LLD19BM	Mycogen F697
Moisture	67.9	65.9
Tons/Acre	26.0	25.4
Stalk Lodging	0.0	0.0
Root Lodging	2.5	5.4
Plant Height	91.0	81.0
Ear Height	46.0	43.0
NDF	41.0	41.1
Digestibility	89.4	90.8
DNDF	73.4	73.4
Lignin Content	1.2	1.2
Environments	10	10
Years	1	1

16

V. Novel LLD19BM-Derived Plants

All plants produced using inbred corn plant LLD19BM as a parent are within the scope of this invention, including plants derived from inbred corn plant LLD19BM. This includes plants essentially derived from inbred LLD19BM with the term "essentially derived variety" having the meaning ascribed to such term in 7 U.S.C. §2104(a)(3) of the Plant Variety Protection Act, which definition is hereby incorporated by reference. This also includes progeny plant and parts thereof with at least one ancestor that is inbred corn plant LLD19BM and more specifically where the pedigree of this progeny includes 1, 2, 3, 4, and/or 5 or cross pollinations to inbred corn plant LLD19BM, or a plant that has LLD19BM as a progenitor. All breeders of ordinary skill in the art maintain pedigree records of their breeding programs. These pedigree records contain a detailed description of the breeding process, including a listing of all parental lines used in the breeding process and information on how such line was used. Thus, a breeder would know if LLD19BM were used in the development of a progeny line, and would also know how many breeding crosses to a line other than LLD19BM were made in the development of any progeny line. A progeny line so developed may then be used in crosses with other, different, corn inbreds to produce first generation F1 corn hybrid seeds and plants with superior characteristics.

Accordingly, another aspect of the present invention is methods for producing an inbred corn line LLD19BM-derived corn plant. This method for producing a LLD19BM-derived corn plant, comprises: (a) crossing inbred corn plant LLD19BM with a second corn plant to yield progeny corn seed; and, (b) growing the progeny corn seed, (under plant growth conditions), to yield the LLD19BM-derived corn plant. Such methods may further comprise the steps of: (c) crossing the LLD19BM-derived corn plant with itself or another corn plant to yield additional LLD19BM-derived progeny corn seed; (b) growing the progeny corn seed of step (d) (under plant growing conditions), to yield additional LLD19BM-derived corn plants; and (e) repeating the crossing and growing steps of (c) and (d) from 0 to 7 times to generate further LLD19BM-derived corn plants. Still further, this may comprise utilizing methods of haploid breeding and plant tissue culture methods to derive progeny of the LLD19BM-derived corn plant.

VI. Tissue Cultures and In Vitro Regeneration of Corn Plants

As is well known in this art, tissue culture of corn may be used for the in vitro regeneration of a corn plant. Accordingly, a further aspect of the invention relates to tissue cultures of the inbred corn plant designated LLD19BM, to tissue cultures of hybrid and derived corn plants obtained from LLD19BM, to plants obtained from such tissue cultures and to the use of tissue culture methodology in plant breeding. The term "tissue culture" includes a composition comprising isolated cells of the same type, isolated cells of a different type, or a collection of such cells organized into parts of a plant. Exemplary tissue cultures are protoplasts, calli and plant cells that are intact in plants or parts of plants, such as embryos, pollen, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk, and the like. In a preferred embodiment, the tissue culture comprises embryos, protoplasts, meristematic cells, pollen, leaves or anthers derived from immature tissues of these plant parts.

17

A. Immature Embryo Culture

To obtain immature embryos for callus culture initiation, ears are harvested from a corn plant, e.g., an inbred corn plant LLD19BM, a hybrid corn plant having LLD19BM as a parent or a LLD19BM-derived corn plant, approximately 9-10 days post-pollination. Initially, harvested ears are surface sterilized, as by scrubbing with Liqui-Nox® soap, immersion in 70% ethanol for 2-3 minutes, followed by immersion in 20% commercial bleach (0.1% sodium hypochlorite) for about 30 minutes.

Ears are rinsed in sterile, distilled water, and immature zygotic embryos are aseptically excised with the aid of a dissection microscope and cultured on a suitable initiation medium. One having ordinary skill in the art would understand that explants from other tissues, such as immature tassel tissue, intercalary meristems and leaf bases, apical meristems, and immature ears may also be the subject of callus culture initiation. Tissue culture media typically, contain amino acids, salts, sugars, hormones, and vitamins. Most of the media employed to regenerate inbred and hybrid plants have some similar components; the media differ in the composition and proportions of their ingredients depending on the particular application envisioned. Examples of media suitable for culture of plants cells include, but are not limited to an N6 medium, an MS media, or modifications thereof. A preferred media is a 15Ag10 medium (N6 Complete Medium, PhytoTechnology Laboratories C167), 1.0 mg/L 2,4-D, 20 g/L sucrose, 100 mg/L casein hydrolysate (enzymatic digest), 25 mM L-proline, 10 mg/L AgNO₃, 2.5 g/L Gelrite, pH 5.8, for about 2-3 weeks with the scutellum facing away from the medium. Tissue showing the proper morphology, Welter, et al., Plant Cell Rep. 14:725-729 (1995), is selectively transferred at biweekly intervals onto fresh 15Ag10 medium for about 6 weeks and then is transferred to "4" medium (N6 Complete Medium Phyto 167), 1.0 mg/L 2,4-D, 20 g/L sucrose, 100 mg/L casein hydrolysate (enzymatic digest), 6 mM L-proline, 2.5 g/L Gelrite, pH 5.8, at bi-weekly intervals for approximately 2 months. Hormones other than an auxin such as 2,4-D may be employed including dicamba, NAA, BAP, 2-NCA, ABA, and picloram. Modifications of these and other basic media may facilitate growth of recipient cells at specific developmental stages.

Regeneration is initiated by transferring callus tissue to a cytokinin-based induction medium, "28" (MS Salts, 30 g/L sucrose, 5 mg/L BAP, 0.025 mg/L 2,4-D at pH of 5.7.) Cells are allowed to grow in low light (13 $\mu\text{EM}^{-2}\text{s}^{-1}$) for one week then higher light (40 $\mu\text{EM}^{-2}\text{s}^{-1}$) for about another week. Regeneration of plants is completed by the transfer of mature and germinating embryos to a hormone-free medium, followed by the transfer of developed plantlets to soil and growth to maturity. In a preferred embodiment, the cells are transferred to a hormone-free regeneration medium, "28" lacking plant growth regulators. Thereafter, small (3-5 cm) plantlets are removed and placed into 150x25 mm culture tubes containing SHGA (Schenk and Hilldebrandt salts: 10 g/L sucrose, 1 g/L myo-inositol, 2 g/L Gelrite, pH 5.8) medium.

Once plantlets develop a sufficient root and shoot system, they are transplanted to 4-inch pots containing approximately 0.1 kg of METRO-MIX 360 soil in a growth room or greenhouse. They are grown with a 16-h photoperiod supplemented by a combination of high pressure sodium and metal halide lamps, and are watered as needed with a combination of three different Miracle Grow fertilizer formulations. At the 3-5 leaf stage, plants are transferred to five gallon pots containing approximately 4 kg METRO-MIX 360.

Primary regenerants (R1 plants) are self- or sib-pollinated after an additional 6-10 weeks in five gallon pots, and R1 seed

18

is collected at 40-45 days post-pollination. Alternately, when self- or sib-pollinations are not possible, plants may be outcrossed to elite inbreds.

It will be appreciated by those of ordinary skill in the art that regenerable cultures, including Type I and Type II cultures, may be initiated from immature embryos using other methods, such as those described in, for example, PCT Application WO 95/06128, the disclosure of which is incorporated herein by reference in its entirety.

B. Additional Tissue Cultures and Regeneration

Other means for preparing and maintaining plant tissue cultures are well known in the art. By way of example, a tissue culture comprising organs such as tassels or anthers and pollen (microspores) have been used to produce regenerated plants (U.S. Pat. Nos. 5,322,789 and 5,445,961). Also, meristematic cells (i.e., plant cells capable of continual cell division and characterized by an undifferentiated cytological appearance, normally found at growing points or tissues in plants such as root tips, stem apices, lateral buds, etc.) can be cultured to regenerate fertile plants (U.S. Pat. No. 5,736,369 the disclosure of which is specifically incorporated herein by reference).

VII. Male Sterility

Methods for controlling male fertility in corn plants offer the opportunity for improved plant breeding, particularly for the development of corn hybrids which require the implementation of a male sterility system to prevent the inbred parent plants from self-pollination.

Accordingly, another aspect of the present invention is male-sterile inbred corn plants designated LLD19BM and the production of hybrid corn seed using a male sterility system with such inbred female parent plants that are male sterile. In the event that inbred corn line LLD19BM is employed as the female parent, LLD19BM can be rendered male-sterile by, for example, removing the tassels or "detasseling" LLD19BM parental plants either manually or by machine. By way of example, alternate strips of two corn inbreds may be planted in a field followed by manual or mechanical removal of the pollen-bearing tassels from the designated female inbred. Provided that the female inbreds are sufficiently isolated from foreign corn pollen sources, the ears of the detasseled inbred will be fertilized only from the other male inbred, and the resulting seed will therefore be hybrid seed.

The laborious and occasionally unreliable detasseling process can be minimized by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of factors resulting from cytoplasmic as opposed to the nuclear genome. Thus, this characteristic is inherited exclusively through the female parent in corn plants since CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile. Seed from detasseled fertile corn and CMS produced seed of the same hybrid can be blended to insure that adequate pollen loads are available for fertilization when the hybrid plants are grown. Conventional backcrossing methodology may be used to introgress the CMS trait into inbred LLD19BM.

Alternatively, haploid breeding methods may also be employed to convert inbred LLD19BM to CMS sterility. Haploids are plants which contain only one-half of the chromosome number present in diploid somatic cells, which are cells other than haploid cells such as those found in the germ. There are a few stocks or genetic systems in corn which are known to generate haploids spontaneously. For example, plants are known which possess an indeterminate gameto-

phyte (ig) gene (Kermicle 1969 Science 166:1422-1424) which generate haploids. Additionally, a line known as Stock 6 (See, Birchler, J. A., "Practical Aspects of Haploid Production," The Corn Handbook, Freeling and Walbot (eds). pp. 386-388 (1996)) possesses a propensity to generate haploids. Moreover, RWS (Roeber and Geiger 2001, submitted to Crop Science), KEMS (Deimling, Roeber, and Geiger, 1997, Vortr. Pflanzenzuchtg 38:203-224), or KMS and ZMS (Chalyk, Bylich & Chebotar, 1994, MNL 68:47; Chalyk & Chebotar, 2000, Plant Breeding 119:363-364) also represent inducer lines which be used to produce haploid plants from any genotype. In a preferred aspect of the present invention, cytoplasmic male sterile female plants which possess the ig gene may be used to facilitate the generation of cytoplasmic male sterile inbred versions of LLD19BM of the present invention because the selection of haploids and subsequent doubling permits corn breeders to reach homogeneity more quickly and completely.

Detasseling can also be avoided by the use of chemically induced male sterility in the production of hybrid corn seed. Chemicals that induce male sterility include gametocides, pollen suppressants, and chemical hybridizing agents. The general procedure is to use a foliar spray before flowering, which inhibits production of viable pollen, but does not injure the pistillate reproductive organs or affect seed development. If the treatment is successful and all of the pollen killed, self-pollination will not occur in the treated plants, but the flowers will set seed freely from cross-pollination. In such a case, the parent plants used as the male may either not be treated with the chemical agent or may include a genetic factor which causes resistance to the sterilizing effects of the chemical agent. The use of chemically induced male sterility affects fertility in the plants only for the growing season in which the gametocide is applied (see Carlson, Glenn R., U.S. Pat. No. 4,963,904).

A further method for controlling male sterility includes the use of genes conferring male sterility, such as those disclosed in U.S. Pat. Nos. 3,861,709, 3,710,511, 4,654,465, 5,625,132, and 4,727,219, each of the disclosures of which are specifically incorporated herein by reference in their entirety. Both inducible and non-inducible male sterility genes can increase the efficiency with which hybrids are made, in that they eliminate the need to physically emasculate the corn plant used as a female in a given cross.

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Pat. Nos. 4,654,465 and 4,727,219 and chromosomal translocations as described by Patterson in U.S. Pat. Nos. 3,861,709 and 3,710,511. These and all patents referred to are incorporated by reference. In addition to these methods, U.S. Pat. No. 5,432,068, discloses a system of nuclear male sterility which includes: identifying a gene which is critical to male fertility; silencing this native gene which is critical to male fertility; removing the native promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered gene back into the plant; and thus creating a plant that is male sterile because the inducible promoter is not "on" resulting in the male fertility gene not being transcribed. Fertility is restored by inducing, or turning "on", the promoter, which in turn allows the gene which confers male fertility to be transcribed.

Other methods of conferring genetic male sterility exist in the art. These methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an

antisense system in which a gene critical to fertility is identified and an antisense to that gene is inserted in the plant (see Fabinjanski, et. al, EP089/3010153.8 publication no 329, 308 and PCT application PCT/CA90/00037 published as WO 90/08828).

The presence of a male-fertility restorer gene results in the production of a fully fertile F₁ hybrid progeny. If no restorer gene is present in the male parent, male-sterile hybrids are obtained. Such hybrids are useful where the vegetative tissue of the corn plant is used, e.g., for silage, but in most cases, the seeds will be deemed the most valuable portion of the crop, so fertility of the hybrids in these crops must be restored. Therefore, one aspect of the present invention concerns inbred corn plant LLD19BM comprising a single gene capable of restoring male fertility in an otherwise male-sterile inbred or hybrid plant. Examples of male-sterility genes and corresponding restorers which could be employed within the inbred of the invention are well known to those of skill in the art of plant breeding and are disclosed in, for example, U.S. Pat. Nos. 5,530,191, 5,689,041, 5,741,684, and 5,684,242, the disclosures of which are each specifically incorporated herein by reference in their entirety.

VIII. Corn Transformation

With the advent of molecular biological techniques that have allowed the isolation and characterization of genes that encode specific protein products, scientists in the field of plant biology developed a strong interest in engineering the genome of plants to contain and to express foreign genes, or additional, or modified versions of native or endogenous genes (perhaps driven by different promoters) to alter the traits of a plant in a specific manner. Such foreign, additional and/or modified genes are referred to herein collectively as "transgenes." The present invention, in particular embodiments, also relates to transformed versions of the claimed inbred corn line LLD19BM containing one or more transgenes.

Plant transformation involves the construction of an expression vector which will function in plant cells. Such a vector comprises DNA comprising a gene under control of or operatively linked to a regulatory element. The expression vector may contain one or more such operably linked gene/regulatory element combinations. The vector(s) may be in the form of a plasmid, and can be used, alone or in combination with other plasmids, to provide transformed corn plants, using transformation methods as described below to incorporate transgenes into the genetic material of the corn plant(s).

A. Expression Vectors for Corn Transformation/Marker Genes

Expression vectors include at least one genetic marker, operably linked to a regulatory element that allows transformed cells containing the marker to be either recovered by negative selection, i.e., inhibiting growth of cells that do not contain the selectable marker gene, or by positive selection, i.e., screening for the product encoded by the genetic marker. Many commonly used selectable marker genes for plant transformation are well known in the transformation arts, and include, for example, genes that code for enzymes that metabolically detoxify a selective chemical agent which may be an antibiotic or a herbicide, or genes that encode an altered target which is insensitive to the inhibitor. A few positive selection methods are also known in the art.

One commonly used selectable marker gene for plant transformation is the neomycin phosphotransferase II (nptII) gene, isolated from a bacterial source, which when placed under the

control of plant regulatory signals confers resistance to kanamycin. Fraley et al., *Proc. Natl. Acad. Sci. U.S.A.* 80: 4803 (1983). Another commonly used selectable marker gene is the hygromycin phosphotransferase gene which confers resistance to the antibiotic hygromycin. Vanden Elzen et al., *Plant Mol. Biol.* 5: 299 (1985).

Additional selectable marker genes of bacterial origin that confer resistance to antibiotics include gentamycin acetyl transferase, streptomycin phosphotransferase, aminoglycoside-3'-adenyl transferase, the bleomycin resistance determinant. Hayford et al., *Plant Physiol.* 86: 1216 (1988), Jones et al., *Mol. Gen. Genet.* 210: 86 (1987), Svab et al., *Plant Mol. Biol.* 14: 197 (1990), Hille et al., *Plant Mol. Biol.* 7: 171 (1986). Other selectable marker genes confer resistance to herbicides such as glyphosate, glufosinate or bromoxynil. Comai et al., *Nature* 317: 741-744 (1985), Gordon-Kamm et al., *Plant Cell* 2: 603-618 (1990) and Stalker et al., *Science* 242: 419-423 (1988).

Other selectable marker genes for plant transformation are not of bacterial origin. These genes include, for example, mouse dihydrofolate reductase, plant 5-enolpyruvylshikimate-3-phosphate synthase and plant acetolactate synthase. Eichholtz et al., *Somatic Cell Mol. Genet.* 13: 67 (1987), Shah et al., *Science* 233: 478 (1986), Charest et al., *Plant Cell Rep.* 8: 643 (1990).

Another class of marker genes for plant transformation require screening of presumptively transformed plant cells rather than direct genetic selection of transformed cells for resistance to a toxic substance such as an antibiotic. These genes are particularly useful to quantify or visualize the spatial pattern of expression of a gene in specific tissues and are frequently referred to as reporter genes because they can be fused to a gene or gene regulatory sequence for the investigation of gene expression. Commonly used genes for screening presumptively transformed cells include β -glucuronidase (GUS), β -galactosidase, luciferase and chloramphenicol acetyltransferase. Jefferson, R. A., *Plant Mol. Biol. Rep.* 5: 387 (1987), Teeri et al., *EMBO J.* 8: 343 (1989), Koncz et al., *Proc. Natl. Acad. Sci. U.S.A.* 84: 131 (1987), De Block et al., *EMBO J.* 3: 1681 (1984). Another approach to the identification of a relatively rare transformation events has been use of a gene that encodes a dominant constitutive regulator of the *Zea mays* anthocyanin pigmentation pathway. Ludwig et al., *Science* 247: 449 (1990).

Recently, in vivo methods for visualizing GUS activity that do not require destruction of plant tissue have been made available. Molecular Probes Publication 2908, Imagene Green™, p. 1-4 (1983) and Naleway et al., *J. Cell Biol.* 115: 151a (1991). However, these in vivo methods for visualizing GUS activity have not proven useful for recovery of transformed cells because of low sensitivity, high fluorescent backgrounds, and limitations associated with the use of luciferase genes as selectable markers.

More recently, a gene encoding Green Fluorescent Protein (GFP) has been utilized as a marker for gene expression in prokaryotic and eukaryotic cells. Chalfie et al., *Science* 263: 802 (1994). GFP and mutants of GFP may be used as screenable markers.

B. Promoters

Genes included in expression vectors must be driven by a nucleotide sequence comprising a regulatory element, for example, a promoter. Several types of promoters are now well known in the transformation arts, as are other regulatory elements that can be used alone or in combination with promoters.

As used herein "promoter" includes reference to a region of DNA upstream from the start of transcription and involved in

recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Examples of promoters under developmental control include promoters that preferentially initiate transcription in certain tissues, such as leaves, roots, seeds, fibers, xylem vessels, tracheids, or sclerenchyma. Such promoters are referred to as "tissue-preferred." Promoters which initiate transcription only in certain tissues are referred to as "tissue-specific." A "cell type" specific promoter primarily drives expression in certain cell types in one or more organs, for example, vascular cells in roots or leaves. An "inducible" promoter is a promoter which is under environmental control or is induced in response to chemical or hormonal stimuli. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions or the presence of light. Examples of chemicals that induce expression including salicylic acid and ABA. Tissue-specific, tissue-preferred, cell type specific, and inducible promoters constitute the class of "non-constitutive" promoters. A "constitutive" promoter is a promoter which is active under most environmental conditions and in all cells.

1. Inducible Promoters

An inducible promoter is operably linked to a gene for expression in corn. Optionally, the inducible promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in corn. With an inducible promoter the rate of transcription increases in response to an inducing agent. Any inducible promoter can be used in the instant invention. A particularly preferred inducible promoter is a promoter that responds to an inducing agent to which plants do not normally respond. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene, the transcriptional activity of which is induced by a glucocorticosteroid hormone.

2. Constitutive Promoters

A constitutive promoter is operably linked to a gene for expression in corn or the constitutive promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in corn. Many different constitutive promoters can be used in the present invention. Exemplary constitutive promoters include, but are not limited to, the promoters from plant viruses such as the 35S promoter from CaMV and the promoters from such genes as rice actin, maize ubiquitin, and corn H3 histone. Also, the ALS promoter, a XbaI/NcoI fragment 5' to the *Brassica napus* ALS3 structural gene (or a nucleotide sequence that has substantial sequence similarity to the XbaI/NcoI fragment) represents a particularly useful constitutive promoter.

3. Tissue-Specific or Tissue-Preferred Promoters

A tissue-specific promoter is operably linked to a gene for expression in corn. Optionally, the tissue-specific promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in corn. Plants transformed with a gene of interest operably linked to a tissue-specific promoter produce the protein product of the transgene exclusively, or preferentially, in a specific tissue. Any tissue-specific or tissue-preferred promoter can be utilized in the instant invention. Exemplary tissue-specific or tissue-preferred promoters include, but are not limited to, a seed-preferred promoter such as that from the phaseolin gene; a leaf-specific and light-induced promoter such as that from cab or rubisco; an anther-specific promoter such as that from LAT52; a pollen specific promoter such as that from Zm13 or a microspore-preferred promoter such as that from apg.

C. Signal Sequences for Targeting Proteins to Subcellular Compartments

Transport of protein produced by transgenes to a subcellular compartment such as the chloroplast, vacuole, peroxisome, glyoxysome, cell wall or mitochondrion, or for secretion into the apoplast, is accomplished by means of operably linking the nucleotide sequence encoding a signal sequence to the 5' and/or 3' region of a gene encoding the protein of interest. Targeting sequences at the 5' and/or 3' end of the structural gene may determine, during protein synthesis and processing, where the encoded protein is ultimately compartmentalized. The presence of a signal sequence directs a polypeptide to either an intracellular organelle or subcellular compartment or for secretion to the apoplast. Any signal sequence known in the art is contemplated by the present invention.

D. Foreign Protein Genes and Agronomic Genes

With transgenic plants according to the present invention, a foreign protein can be produced in commercial quantities. Thus, techniques for the selection and propagation of transformed plants, which are well understood in the art, yield a plurality of transgenic plants which are harvested in a conventional manner, and a foreign protein then can be extracted from a tissue of interest or from total biomass. Protein extraction from plant biomass can be accomplished by known methods.

According to a preferred embodiment, the transgenic plant provided for commercial production of foreign protein is corn. In another preferred embodiment, the biomass of interest is seed. For the relatively small number of transgenic plants that show higher levels of expression, a genetic map can be generated, primarily via conventional Restriction Fragment Length Polymorphisms (RFLP), Polymerase Chain Reaction (PCR) analysis, and Simple Sequence Repeats (SSR) which identifies the approximate chromosomal location of the integrated DNA molecule. For exemplary methodologies in this regard, see Glick and Thompson, METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY 269-284 (CRC Press, Boca Raton, 1993). Map information concerning chromosomal location is useful for proprietary protection of a subject transgenic plant. If unauthorized propagation is undertaken and crosses made with other germplasm, the map of the integration region can be compared to similar maps for suspect plants, to determine if the latter have a common parentage with the subject plant. Map comparisons would involve hybridizations, RFLP, PCR, SSR and sequencing, all of which are conventional techniques.

Likewise, by means of the present invention, agronomic genes can be expressed in transformed plants. More particularly, plants can be genetically engineered to express various phenotypes of agronomic interest. Exemplary genes implicated in this regard include, but are not limited to:

1. Genes that Confer Resistance to Pests or Disease and that Encode:

(a) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example Jones et al., Science 266: 789 (1994) (cloning of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*); Martin et al., Science 262: 1432 (1993) (tomato Pto gene for resistance to *Pseudomonas syringae* pv. tomato

encodes a protein kinase); Mindrinos et al., Cell 78: 1089 (1994) (*Arabidopsis* RSP2 gene for resistance to *Pseudomonas syringae*).

(b) A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser et al., Gene 48: 109 (1986), who disclose the cloning and nucleotide sequence of a Bt δ -endotoxin gene. Moreover, DNA molecules encoding δ -endotoxin genes can be purchased from American Type Culture Collection (Rockville, Md.), for example, under ATCC Accession Nos. 40098, 67136, 31995 and 31998.

(c) A lectin. See, for example, the disclosure by Van Damme et al., Plant Molec. Biol. 24: 25 (1994), who disclose the nucleotide sequences of several *Clivia miniata* mannose-binding lectin genes.

(d) A vitamin-binding protein such as avidin. See PCT application US93/06487 the contents of which are hereby incorporated by. The application teaches the use of avidin and avidin homologues as larvicides against insect pests.

(e) An enzyme inhibitor, for example, a protease inhibitor or an amylase inhibitor. See, for example, Abe et al., J. Biol. Chem. 262: 16793 (1987) (nucleotide sequence of rice cysteine proteinase inhibitor), Huub et al., Plant Molec. Biol. 21: 985 (1993) (nucleotide sequence of cDNA encoding tobacco proteinase inhibitor I), and Sumitani et al., Biosci. Biotech. Biochem. 57: 1243 (1993) (nucleotide sequence of *Streptomyces nitrosporeus* α -amylase inhibitor).

(f) An insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof. See, for example, the disclosure by Hammock et al., Nature 344: 458 (1990), of baculovirus expression of cloned juvenile hormone esterase, an inactivator of juvenile hormone.

(g) An insect-specific peptide or neuropeptide which, upon expression, disrupts the physiology of the affected pest. For example, see the disclosures of Regan, J. Biol. Chem. 269: 9 (1994) (expression cloning yields DNA coding for insect diuretic hormone receptor), and Pratt et al., Biochem. Biophys. Res. Comm. 163: 1243 (1989) (an allostatin is identified in *Diploptera punctata*). See also U.S. Pat. No. 5,266,317 to Tomalski et al., who disclose genes encoding insect-specific, paralytic neurotoxins.

(h) An insect-specific venom produced in nature by a snake, a wasp, etc. For example, see Pang et al., Gene 116: 165 (1992), for disclosure of heterologous expression in plants of a gene coding for a scorpion insectotoxic peptide.

(i) An enzyme responsible for an hyperaccumulation of a monoterpenes, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.

(j) An enzyme involved in the modification, including the post-translational modification, of a biologically active molecule, for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic. See PCT application WO 93/02197 in the name of Scott et al., which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under Accession Nos. 39637 and 67152. See also Kramer et al., Insect Biochem. Molec. Biol. 23: 691 (1993), who teach the nucleotide sequence of a cDNA encoding tobacco hookworm chitinase, and Kawalleck et al., Plant Molec. Biol. 21: 673 (1993), who provide the nucleotide sequence of the parsley ubi4-2 polyubiquitin gene.

(k) A molecule that stimulates signal transduction. For example, see the disclosure by Botella et al., *Plant Molec. Biol.* 24: 757 (1994), of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess et al., *Plant Physiol.* 104: 1467 (1994), who provide the nucleotide sequence of a corn calmodulin cDNA clone.

(l) A hydrophobic moment peptide. See PCT application WO95/16776 (disclosure of peptide derivatives of Tachyplexin which inhibit fungal plant pathogens) and PCT application WO95/18855 (teaches synthetic antimicrobial peptides that confer disease resistance), the respective contents of which are hereby incorporated by reference.

(m) A membrane permease, a channel former or a channel blocker. For example, see the disclosure by Jaynes et al., *Plant Sci.* 89: 43 (1993), of heterologous expression of a cecropin- β lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.

(n) A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See Beachy et al., *Ann. Rev. Phytopathol.* 28: 451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. *Id.*

(o) An insect-specific antibody or an immunotoxin derived therefrom. Thus, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor et al., Abstract #497, SEVENTH INT'L SYMPOSIUM ON MOLECULAR PLANT-MICROBE INTERACTIONS (Edinburgh, Scotland, 1994) (enzymatic inactivation in transgenic tobacco via production of single-chain antibody fragments).

(p) A virus-specific antibody. See, for example, Tavlaraki et al., *Nature* 366: 469 (1993), who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.

(q) A developmental-arrestive protein produced in nature by a pathogen or a parasite. Thus, fungal endo α -1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo- α -1,4-D-galacturonate. See Lamb et al., *Bio/Technology* 10: 1436 (1992). The cloning and characterization of a gene which encodes a bean endopolygalacturonase-inhibiting protein is described by Toubart et al., *Plant J.* 2: 367 (1992).

(r) A developmental-arrestive protein produced in nature by a plant. For example, Logemann et al., *Bio/Technology* 10: 305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

2. Genes that Confer Resistance to a Herbicide, for Example:

(a) A herbicide that inhibits the growing point or meristem, such as an imidazalinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee et al., *EMBO J.* 7: 1241 (1988), and Miki et al., *Theor. Appl. Genet.* 80: 449 (1990), respectively.

(b) Glyphosate (resistance imparted by mutant 5-enolpyruvyl-3-phosphokimate synthase (EPSP) and aroA genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin acetyl transferase (PAT) and *Streptomyces hygroscopicus* phosphinothricin acetyl transferase (bar) genes), and pyridinoyl or phenoxy propionic

acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. Pat. No. 4,940,835 to Shah et al., which discloses the nucleotide sequence of a form of EPSP which can confer glyphosate resistance. A DNA molecule encoding a mutant aroA gene can be obtained under ATCC accession No. 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. Pat. No. 4,769,061 to Comai. European patent application No. 0 333 033 to Kumada et al. and U.S. Pat. No. 4,975,374 to Goodman et al. disclose nucleotide sequences of glutamine synthetase genes which confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in European application No. 0 242 246 to Leemans et al. De Greef et al., *Bio/Technology* 7: 61 (1989), describe the production of transgenic plants that express chimeric bar genes coding for phosphinothricin acetyl transferase activity. Exemplary of genes conferring resistance to phenoxy propionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the Acc1-S1, Acc1-S2 and Acc1-S3 genes described by Marshall et al., *Theor. Appl. Genet.* 83: 435 (1992).

(c) A herbicide that inhibits photosynthesis, such as a triazine (psbA and gs+ genes) and a benzonitrile (nitrilase gene). Przibilla et al., *Plant Cell* 3: 169 (1991), describe the transformation of *Chlamydomonas* with plasmids encoding mutant psbA genes. Nucleotide sequences for nitrilase genes are disclosed in U.S. Pat. No. 4,810,648 to Stalker, and DNA molecules containing these genes are available under ATCC Accession Nos. 53435, 67441 and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes et al., *Biochem. J.* 285: 173 (1992).

3. Genes that Confer or Contribute to a Value-added Trait, such as:

(a) Modified fatty acid metabolism, for example, by transforming a plant with an antisense gene of stearoyl-ACP desaturase to increase stearic acid content of the plant. See Knultzon et al., *Proc. Natl. Acad. Sci. USA* 89: 2624 (1992).

(b) Decreased phytate content:

(i) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For example, see Van Hartingsveldt et al., *Gene* 127: 87 (1993), for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene.

(ii) A gene could be introduced that reduces phytate content. In corn, this, for example, could be accomplished, by cloning and then reintroducing DNA associated with the single allele which is responsible for corn mutants characterized by low levels of phytic acid. See Raboy et al., *Maydica* 35: 383 (1990).

(iii) Modified carbohydrate composition effected, for example, by transforming plants with a gene coding for an enzyme that alters the branching pattern of starch. See Shiroza et al., *J. Bacteriol.* 170: 810 (1988) (nucleotide sequence of *Streptococcus mutans* fructosyltransferase gene), Steinmetz et al., *Mol. Gen. Genet.* 200: 220 (1985) (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen et al., *Bio/Technology* 10: 292 (1992) (production of transgenic plants that express *Bacillus licheniformis* α -amylase), Elliot et al., *Plant Molec. Biol.* 21: 515 (1993) (nucleotide sequences of tomato invertase genes), Sogaard et al., *J. Biol. Chem.* 268: 22480 (1993) (site-directed mutagenesis of barley α -amylase gene), and Fisher et al., *Plant Physiol.* 102: 1045 (1993) (corn endosperm starch branching enzyme II).

E. Methods for Corn Transformation

Numerous methods for plant transformation have been developed, including biological and physical, plant transformation protocols. See, for example, Miki et al., "Procedures

for Introducing Foreign DNA into Plants" in Methods in Plant Molecular Biology and Biotechnology, Glick, B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 67-88. In addition, expression vectors and in vitro culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber et al., "Vectors for Plant Transformation" in Methods in Plant Molecular Biology and Biotechnology, Glick, B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 89-119.

1. *Agrobacterium*-Mediated Transformation

One method for introducing an expression vector into plants is based on the natural transformation system of *Agrobacterium*. See, for example, Horsch et al., Science 227: 1229 (1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria which genetically transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*, respectively, carry genes responsible for genetic transformation of the plant. See, for example, Kado, C. I., Crit. Rev. Plant. Sci. 10:1 (1991). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided by Gruber et al., supra, Miki et al., supra, and Moloney et al., Plant Cell Reports 8: 238 (1989). See also, U.S. Pat. No. 5,591,616, issued Jan. 7, 1997.

2. Direct Gene Transfer

Despite the fact the host range for *Agrobacterium*-mediated transformation is broad, some major cereal crop species and gymnosperms have generally been recalcitrant to this mode of gene transfer, even though some success has recently been achieved in rice and corn. Hiei et al., The Plant Journal 6: 271-282 (1994); U.S. Pat. No. 5,591,616, issued Jan. 7, 1997. Several methods of plant transformation, collectively referred to as direct gene transfer, have been developed as an alternative to *Agrobacterium*-mediated transformation.

A generally applicable method of plant transformation is microprojectile-mediated transformation wherein DNA is carried on the surface of microprojectiles measuring 1 to 4 μm (See e.g., U.S. Pat. No. 5,550,318; U.S. Pat. No. 5,736,369, U.S. Pat. No. 5,538,880; and PCT Publication WO 95/06128). The expression vector is introduced into plant tissues with a biolistic device that accelerates the microprojectiles to speeds of 300 to 600 m/s which is sufficient to penetrate plant cell walls and membranes. Sanford et al, Part. Sci. Technol. 5: 27 (1987), Sanford, J. C., Trends Biotech. 6: 299 (1988), Klein et al., Bio/Technology 6: 559-563 (1988), Sanford, J. C., Physiol Plant 79: 206 (1990), Klein et al., Biotechnology 10: 268 (1992). In corn, several target tissues can be bombarded with DNA-coated microprojectiles in order to produce transgenic plants, including, for example, callus (Type I or Type II), immature embryos, and meristematic tissue.

Another method for physical delivery of DNA to plants is sonication of target cells. Zhang et al., Bio/Technology 9: 996 (1991). Alternatively, liposome or spheroplast fusion have been used to introduce expression vectors into plants. Deshayes et al., EMBO J., 4: 2731 (1985), Christou et al., Proc Natl. Acad. Sci. U.S.A. 84: 3962 (1987). Direct uptake of DNA into protoplasts using CaCl_2 precipitation, polyvinyl alcohol or poly-L-ornithine have also been reported. Hain et al., Mol. Gen. Genet. 199: 161 (1985) and Draper et al., Plant Cell Physiol. 23: 451 (1982). Electroporation of protoplasts and whole cells and tissues have also been described. U.S. Pat. No. 5,384,253 and Donn et al. In Abstracts of VIIIth International Congress on Plant Cell and Tissue Culture IAPTC, A2-38, p 53 (1990); D'Halluin et al., Plant Cell 4: 1495-1505 (1992) and Spencer et al., Plant Mol. Biol. 24: 51-61 (1994).

Other methods which have been described for the genetic transformation of corn include, electroporation (U.S. Pat. No. 5,371,003) and silicon carbide fiber-mediated transformation (U.S. Pat. No. 5,302,532 and U.S. Pat. No. 5,464,765).

Following transformation of corn target tissues, expression of the above-described selectable marker genes allows for preferential selection of transformed cells, tissues and/or plants, using regeneration and selection methods now well known in the art. For example, transformed corn immature embryos.

The foregoing methods for transformation would typically be used for producing transgenic inbred lines. Transgenic inbred lines could then be crossed, with another (non-transformed or transformed) inbred line, in order to produce a transgenic hybrid corn plant. Alternatively, a genetic trait which has been engineered into a particular corn line using the foregoing transformation techniques could be moved into another line using traditional backcrossing techniques that are well known in the plant breeding arts. For example, a backcrossing approach could be used to move an engineered trait from a public, non-elite line into an elite line, or from a hybrid corn plant containing a foreign gene in its genome into a line or lines which do not contain that gene.

IX. Genetic Complements

In addition to phenotypic observations, a plant can also be described by its genotype. The genotype of a plant can be described through a genetic marker profile which can identify plants of the same variety, a related variety or be used to determine or to validate a pedigree. Genetic marker profiles can be obtained by techniques such as Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites, and Single Nucleotide Polymorphisms (SNPs), Isozyme Electrophoresis and Isoelectric Focusing. For example, see Berry, Don, et al., "Assessing Probability of Ancestry Using Simple Sequence Repeat Profiles: Applications to Corn Hybrids and Inbreds", Genetics, 2002, 161:813-824, which is incorporated by reference herein in its entirety.

Particular markers used for these purposes are not limited to the set of markers disclosed herewithin, but are envisioned to include any type of genetically stable marker and marker profile which provides a means of distinguishing varieties. In addition to being used for identification of inbred parents, a hybrid produced through the use of LLD19BM or its parents, and identification or verification of the pedigree of progeny plants produced through the use of LLD19BM, the genetic marker profile is also useful in breeding and developing back-cross conversions.

Means of performing genetic marker profiles using SSR polymorphisms are well known in the art. SSRs are genetic markers based on polymorphisms in repeated nucleotide sequences, such as microsatellites. The phrase "simple sequence repeats" or "SSR" refers to di-, tri- or tetra-nucleotide repeats within a genome. The repeat region may vary in length between genotypes while the DNA flanking the repeat is conserved such that the primers will work in a plurality of genotypes. A polymorphism between two genotypes represents repeats of different lengths between the two flanking conserved DNA sequences. A marker system based on SSRs

can be highly informative in linkage analysis relative to other marker systems in that multiple alleles may be present. Another advantage of this type of marker is that, through use of flanking primers, detection of SSRs can be achieved, for example, by the polymerase chain reaction (PCR). The PCR[®] detection is done by the use of two oligonucleotide primers flanking the polymorphic segment of repetitive DNA followed by DNA amplification. This step involves repeated cycles of heat denaturation of the DNA followed by annealing of the primers to their complementary sequences at low temperatures, and extension of the annealed primers with DNA polymerase. Size separation of DNA fragments on agarose or polyacrylamide gels following amplification, comprises the major part of the methodology.

DNA isolation and amplification were performed in the present invention as follows. DNA was extracted from plant leaf tissue using DNeasy 96 Plant Kit from Qiagen, Inc. (Valencia, Calif., U.S.A.) following an optimized September 2002 manufacturer's protocol. PCR amplifications were performed using a Quiagen HotStar[™] Taq master mix in an 8 μ l reaction format as follows: 2 μ l DNA (5 ng/ μ L+6 μ L of master mix). The PCR conditions were as follows: 12 mins. at 95° C., 40 cycles of 5 seconds at 94° C., 15 seconds at 55° C., 30 seconds at 72° C., 30 mins. at 72° C., followed by cooling to 4° C. Following isolation and amplification, markers can be scored by gel electrophoresis of the amplification products. Scoring of marker genotype is based on the size of the amplified fragment as measured by molecular weight (MW) rounded to the nearest integer. Multiple samples, comprised of fluorescently labeled DNA fragments were processed in an ABI 3700 capillary-based machine and precise allele sizing and locus genotyping were done by running GeneScan and Genotyper software (PE Applied Biosystems, Foster City, Calif.). When comparing lines, it is preferable if all SSR profiles are performed in the same lab. An SSR service is available to the public on a contractual basis by Paragen, Research Triangle Park, N.C. (formerly Celera AgGen of Davis, Calif.). Primers used for the SSRs suggested herein are publicly available and may be found in the Corn Genetics and Genomics Database at <http://www.maizegdb.org>, in Sharopova et al. (Plant Mol. Biol. 48(5-6):463-481), Lee et al. (Plant Mol. Biol. 48(5-6): 453-461). The chromosome locations on which such markers are located and the location on such chromosome are generally reported in the Database. SSR information is provided in TABLE 5.

A genetic marker profile of an inbred may be predictive of the agronomic traits of a hybrid produced using that inbred. For example, if an inbred of known genetic marker profile and phenotype is crossed with a second inbred of known genetic marker profile and phenotype it is possible to predict the phenotype of the F₁ hybrid based on the combined genetic marker profiles of the parent inbreds. Method for prediction of hybrid performance from genetic marker data are disclosed in U.S. Pat. No. 5,492,547, the disclosure of which is specifically incorporated herein by reference in its entirety. Such predictions may be made using any suitable laboratory-based techniques for the analysis, comparison and characterization of plant genotype.

The most widely used of these laboratory techniques are Isozyme Electrophoresis and RFLPs as discussed in Lee, M., "Inbred Lines of Corn and Their Molecular Markers," The Corn Handbook, (Springer-Verlag, New York, Inc. 1994, at 423-432) incorporated herein by reference. Isozyme Electrophoresis is a useful tool in determining genetic composition, although it has relatively low number of available markers and the low number of allelic variants among corn inbreds. RFLPs have the advantage of revealing an exceptionally high

degree of allelic variation in corn and the number of available markers is almost limitless. The present invention provides a genetic complement of the inbred corn plant variety designated LLD19BM. Further provided by the invention is a hybrid genetic complement, wherein the complement is formed by the combination of a haploid genetic complement from LLD19BM and another haploid genetic complement. Means for determining such a genetic complement are well-known in the art.

As used herein, the phrase "genetic complement" means an aggregate of nucleotide sequences, the expression of which defines the phenotype of a corn plant or a cell or a tissue of that plant. By way of example, a corn plant is genotyped to determine a representative sample of the inherited markers it possesses. Markers are alleles at a single locus. They are preferably inherited in codominant fashion so that the presence of both alleles at a diploid locus is readily detectable and they are free of environmental variation, i.e., their heritability is 1. This genotyping is preferably performed on at least one generation of the descendant plant for which the numerical value of the quantitative trait(s) of interest are also determined. The array of single locus genotypes is expressed as a profile of marker alleles, two at each locus. The marker allelic composition of each locus can be either homozygous or heterozygous. Homozygosity is a condition where both alleles at a locus are characterized by the same nucleotide sequence or size of a repeated sequence. Heterozygosity refers to different conditions of the gene at a locus.

The SSR genetic marker profile of inbred LLD19BM was determined. Because an inbred is essentially homozygous at all relevant loci, an inbred should, in almost all cases, have both the alleles of one size at each locus. In contrast, a diploid genetic marker profile of a hybrid should be the sum of those parents, e.g., if one inbred parent had the allele 168 (base pairs) at a particular locus, and the other inbred parent had 172, the hybrid is 168,172 by inference. Subsequent generations of progeny produced by selection and breeding are expected to be of genotype 168, 172, or 168,172 for that locus by inference. When the F₁ plant is used to produce an inbred, the locus should be either 168 or 172 for that position. Surprisingly, it has been observed that in certain instances, novel SSR genotypes arise during the breeding process. For example, a genotype of 170 may be observed at a particular locus positions from the cross of parental inbreds with 168 and 172 at that locus. Such a novel SSR genotype may further define an inbred plant from the parental inbreds from which it was derived. An SSR genetic marker profile of LLD19BM is presented in Table 5 wherein representative measured fragment lengths of alleles are given.

TABLE 5

Public Name Of Marker	LLD19BM Measured Fragment Length
bnlg1179	203.28
bnlg1614	180.06
phi109275	126.41
bnlg1866	101.59
bnlg1643	131.63
umc1128	153.14
umc1306	138.23
umc1064	143.25
umc1797	103
phi265454	230.52
phi402893	218.48
bnlg1338	167.03
umc1542	156.09

TABLE 5-continued

Public Name Of Marker	LLD19BM Measured Fragment Length
umc1776	134.56
phi083	125.87
bnlg1138	187.76
bnlg1633	161.06
bnlg1169	219.21
umc1525	132.21
umc1394	93.76
phi453121	212.75
umc1886	147.84
phi029	158.66
bnlg420	72.08
bnlg1160	131.97
phi046	64.63
umc1136	140.29
umc1008	155.54
phi072	142.13
umc1757	142.32
umc1509	133.12
nc004	138.58
umc1382	146.52
umc1964	135.38
umc1662	107.48
phi079	188.51
bnlg1189	216.54
phi066	154.22
umc1940	123.77
umc1101	151.63
umc1180	102.26
bnlg589	149.88
umc1050	95.06
umc1097	100.92
umc1478	135.97
phi113	129.93
bnlg1902	220.19
bnlg278	75.08
phi101	93.58
umc1225	130.19
umc1153	102.93
bnlg238	170.51
umc1186	213.9
umc1795	139.64
phi123	148.02
umc1127	157.55
umc1695	127.12
bnlg657	102.2
umc1944	140.6
dup013	150.18
phi082	130.19
umc1075	134.98
umc1327	78.19
phi119	158.78
phi125	109.33
umc1149	129.01
umc1724	135.08
bnlg1056	104.29
phi015	95.27
phi033	249.91
phi044	82.35
dup006	104.35
phi027	143
phi016	151.03
umc1357	169.25
umc1310	116.17
bnlg1129	195.55
umc1380	149.4
umc1152	159.7
umc1576	102.14
bnlg1079	175.67
phi084	154.38
umc1280	102.15
phi035	136.09
umc1506	114.53
umc1196	153.38

The present invention also provides a hybrid genetic complement formed by the combination of a haploid genetic complement of the corn plant LLD19BM with a haploid genetic complement of a second corn plant. Means for combining a haploid genetic complement from the foregoing inbred with another haploid genetic complement may comprise any method for producing a hybrid plant from LLD19BM. It is contemplated that such a hybrid genetic complement can be prepared using in vitro regeneration of a tissue culture of a hybrid plant of this invention.

In addition, plants and plant parts substantially benefiting from the use of LLD19BM in their development such as LLD19BM comprising a backcross conversion, or transgene, may be identified by having a molecular marker profile with a high percent identity to LLD19BM. Such a percent identity might be 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 99.9% identical to LLD19BM.

The SSR profile of LLD19BM also can be used to identify derived varieties and other progeny lines developed from the use of LLD19BM, as well as cells and other plant parts thereof. Progeny plants and plant parts produced using LLD19BM may be identified by having a molecular marker profile of at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 99.5% genetic contribution from corn plant LLD19BM.

All publications, patents and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All such publications, patents and patent applications are incorporated by reference herein to the same extent as if each was specifically and individually indicated to be incorporated by reference herein.

The foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding. However, it should be appreciated by those having ordinary skill in the art that certain changes and modifications such as single gene modifications and mutations, somoclonal variants, variant individuals selected from large populations of the plants of the instant inbred and the like may be practiced within the scope of the invention, as limited only by the scope of the appended claims, without departing from the true concept, spirit, and scope of the invention.

What is claimed is:

1. A seed of corn inbred line designated LLD19BM, or a part thereof, representative seed of the line having been deposited under ATCC Accession No. PTA-9814.
2. A part of the seed of claim 1 selected from the group consisting of pericarp, germ and endosperm.
3. The seed of claim 1, further comprising a coating.
4. A substantially homogenous composition of the corn seed of claim 1.
5. A corn plant produced by growing the seed of claim 1.
6. A part of the corn plant of claim 5, selected from the group consisting of an intact plant cell, a plant protoplast, an embryo, a pollen, an ovule, a flower, a kernel, a seed, an ear, a cob, a leaf, a husk, a stalk, a root, a root tip, a brace root, a lateral tassel branch, an anther, a tassel, a glume, a tiller and a silk.
7. Pollen of the plant of claim 5.
8. An ovule of the plant of claim 5.
9. A corn plant, or a part thereof, having all the physiological and morphological characteristics of the corn plant of claim 5.

33

10. A method for producing an inbred corn plant, comprising:

- (a) crossing inbred corn plant LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, with another different corn plant to yield progeny corn seed.

11. The method of claim 10, wherein the other, different corn plant is an inbred corn plant.

12. The method of claim 10, further comprising:

- (b) growing the progeny corn seed from step (a) under self-pollinating or sib-pollinating conditions for about 5 to about 7 generations; and
- (c) harvesting resultant seed.

13. The method of claim 10, further comprising selecting plants obtained from growing at least one generation of the progeny corn seed for a desirable trait.

14. A method of introducing a desired trait into corn inbred line LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, comprising:

- (a) crossing LLD19BM plants with plants of another corn line that comprise a desired trait to produce F1 progeny plants;
- (b) selecting F1 progeny plants that have the desired trait;
- (c) crossing selected progeny plants with LLD19BM plants to produce backcross progeny plants;
- (d) selecting for backcross progeny plants that comprise the desired trait and physiological and morphological characteristics of corn inbred line LLD19BM; and
- (e) performing steps (c) and (d) one or more times in succession to produce the selected or higher backcross progeny plants that comprise the desired trait and all of the physiological and morphological characteristics of corn inbred line LLD19BM listed in Table 1 as determined at the 5% significance level when grown in the same environmental conditions.

15. The method of claim 10, wherein the plants of the other corn line comprise a desired trait selected from the group consisting of male sterility, herbicide resistance, insect resistance, and resistance to bacterial, fungal and viral disease.

16. The method of claim 10, further comprising using genetic markers to compare a genetic complement of a progeny plant with a genetic complement of LLD19BM.

17. The method of claim 10, further comprising using direct or indirect selection to determine whether the desired trait is present in a progeny plant.

18. A method for producing a corn plant, comprising:

- (a) crossing the inbred corn plant of claim 5 with another different corn plant to produce a diploid progeny plant;
- (b) generating a haploid progeny plant from the diploid progeny plant;
- (c) generating a diploid plant from the haploid progeny plant; and,
- (d) selecting the diploid corn plant.

19. The method of claim 18, wherein the haploid progeny plant is generated by culturing a haploid explant from the diploid progeny plant.

20. The method of claim 18, wherein the haploid progeny plant is generated by crossing the progeny plant with another, different plant that induces haploid corn plants.

21. The method of claim 20, wherein the other, different plant is a corn plant that comprises a haploid-inducing gene.

22. The method of claim 18, wherein the diploid plant of step c) is generated by subjecting the haploid progeny plant to a treatment which induces chromosome doubling in the cultured explant.

34

23. The method of claim 18, wherein the diploid plant of step c) is generated by self pollinating the haploid progeny plant.

24. A method for producing a corn plant, comprising:

- (a) inducing a mutation in the corn plant, or a part thereof, of claim 9; and,
- (b) selecting mutated corn plants.

25. The method of claim 24, wherein the mutation is artificially induced by a method selected from the group consisting of elevated temperature, long-term seed storage, tissue culture conditions, radiation, and chemical mutagenesis.

26. A method for producing an inbred corn plant, comprising:

- (a) growing first generation hybrid corn plants having LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, as a parent corn plant;
- (b) inbreeding the first generation hybrid corn plants or crossing the first generation hybrid corn plants with different corn plants to yield progeny corn seed;
- (c) growing the progeny corn seed of step b) to yield further progeny corn seed;
- (d) repeating the inbreeding or the crossing and the growing steps of b) and c) from about 0 to about 7 times to generate inbred corn plants.

27. A method for producing inbred LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, comprising:

- (a) planting a collection of seed comprising seed of a hybrid, one of whose parents is an inbred LLD19BM, the collection also comprising seed of the inbred;
- (b) growing plants from the collection of seed;
- (c) identifying inbred parent plants;
- (d) controlling pollination in a manner which preserves the homozygosity of the inbred parent plant; and,
- (e) harvesting resultant seed.

28. The method of claim 27, wherein step (c) comprises identifying plants with decreased vigor.

29. A method for producing an inbred plant comprising: sib-pollinating plants obtained by growing the harvested resultant seed of step (e) of claim 27.

30. A method for producing an inbred plant comprising: crossing LLD19BM corn plants with corn plants obtained by growing the hybrid seed of step (a) of claim 27.

31. A method for producing a hybrid corn seed comprising crossing a first inbred parent corn plant with a second inbred parent corn plant and harvesting resultant hybrid corn seed, wherein the first inbred corn plant or the second inbred corn plant is the corn plant of claim 5.

32. A method for producing a hybrid corn seed comprising the steps of:

- (a) planting in pollinating proximity seeds of a first and a second inbred parent corn plants, wherein the first inbred corn plant or the second inbred corn plant is the corn plant of claim 9;
- (b) cultivating the seeds of the first and the second inbred corn plants into plants that bear flowers;
- (c) controlling the male fertility of the first or the second inbred corn plant to produce a male sterile corn plant;
- (d) allowing cross-pollination to occur between the first and second inbred corn plants; and,
- (e) harvesting seeds produced on the male sterile corn plant.

33. A hybrid corn seed produced by the method of claim 32.

34. A hybrid corn plant produced by growing the hybrid corn seed of claim 33.

35

35. A method for producing a LLD19BM-derived corn plant, comprising:

(a) crossing inbred corn line LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, with a second corn plant to yield progeny corn seed; and

(b) growing said progeny corn seed, under plant growth conditions, to yield said LLD19BM-derived corn plant.

36. The method of claim 35, further comprising:

(c) crossing the LLD19BM-derived corn plant with itself or another corn plant to yield additional LLD19BM-derived progeny corn seed;

(d) growing the progeny corn seed of step c) under plant growth conditions, to yield additional LLD19BM-derived corn plants; and

(e) repeating the crossing and growing steps of c) and d) from 0 to 7 times to generate further LLD19BM-derived corn plants.

37. The method of claim 35, still further comprising utilizing plant tissue culture methods and/or haploid breeding to derive progeny of said LLD19BM-derived corn plant.

38. A tissue culture of regenerable cells from the corn plant of claim 5.

39. A tissue culture according to claim 38, the cells or protoplasts of the tissue culture being from a tissue selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, silks, flowers, kernels, ears, cobs, husks, and stalks.

40. A corn plant regenerated from the tissue culture of claim 38, wherein the regenerated plant is capable of expressing all the morphological and physiological characteristics of inbred line LLD19BM.

41. A corn plant with all of the physiological and morphological characteristics of corn inbred LLD19BM, wherein the corn plant is produced by a tissue culture process using the corn plant of claim 5 as a starting material for the process.

42. A tissue culture of regenerable cells from the hybrid corn plant of claim 34.

43. A method for regenerating a corn plant comprising the steps of:

(a) culturing an explant comprising a tissue selected from the group consisting of a tissue obtained from inbred corn plant LLD19BM representative seed having been deposited under ATCC Accession No. PTA-9814, an immature tissue obtained from a hybrid corn plant having LLD19BM as a parent, and a LLD19BM-derived corn plant; and,

(b) initiating regeneration.

36

44. The process of claim 43, wherein the explant is an immature tissue.

45. A method for producing a male sterile LLD19BM corn plant, comprising:

(a) crossing the inbred corn plant of claim 5, with a cytoplasmic male sterile corn plant which generates haploids;

(b) identifying haploid plants; and,

(c) crossing the haploid plants with the inbred corn plant LLD19BM to produce male sterile LLD19BM corn plants.

46. A method for producing a corn plant that contains in its genetic material one or more transgenes, comprising crossing the corn plant of claim 5, wherein the plant has been transformed so that its genetic material contains one or more transgenes operably linked to one or more regulatory elements with either a second plant of another corn line, or a non-transformed corn plant of the line LLD19BM, so that the genetic material of the progeny that result from the cross contains the transgene(s) operably linked to a regulatory element.

47. A seed of a corn plant comprising one or more transgenes produced by transforming the plant according to claim 5 and producing seed from the transformed plant.

48. A corn plant produced by the method of claim 46.

49. A method for producing a population of LLD19BM, representative seed of the line having been deposited under ATCC Accession No. PTA-9814, progeny inbred corn plants comprising:

(a) obtaining a first generation progeny corn seed comprising the plant of claim 9 as a parent;

(b) growing the first generation progeny corn seed to produce F1 generation corn plants and obtaining self or sib pollinated seed from the F1 generation corn plants; and,

(c) producing successive filial generations to obtain a population of LLD19BM progeny inbred corn plants.

50. The population of LLD19BM progeny inbred corn plants produced by the method of claim 49, the population, on average, deriving 50% of its alleles from LLD19BM.

51. An F1 hybrid seed produced by crossing the inbred corn plant according to claim 5 with a variety of grain sorghum *Sorghum bicolor* (L.) Moench.

* * * * *