



US 20230389502A1

(19) **United States**

(12) **Patent Application Publication**
Frazier et al.

(10) **Pub. No.: US 2023/0389502 A1**

(43) **Pub. Date: Dec. 7, 2023**

(54) **HYBRID MAIZE VARIETY ARC-9913C**

(52) **U.S. Cl.**

(71) Applicant: **INARI AGRICULTURE
TECHNOLOGY, INC.**, Cambridge,
MA (US)

CPC **A01H 6/4684** (2018.05); **A01H 5/10**
(2013.01)

(72) Inventors: **Ryan Edward Frazier**, Omaha, NE
(US); **Scott Clarence Stelpflug**, West
Lafayette, IN (US)

(57)

ABSTRACT

(21) Appl. No.: **18/329,421**

(22) Filed: **Jun. 5, 2023**

Related U.S. Application Data

(60) Provisional application No. 63/365,910, filed on Jun.
6, 2022.

Publication Classification

(51) **Int. Cl.**
A01H 6/46 (2006.01)
A01H 5/10 (2006.01)

A novel maize variety designated ARC-9913C and seed, plants and plant parts thereof are produced by crossing inbred maize varieties. Methods for producing a maize plant by crossing hybrid maize variety ARC-9913C with another maize plant are disclosed. Methods for producing a maize plant containing in its genetic material one or more traits introgressed into ARC-9913C through backcrossing or genetic transformation, and to the maize seed, plant and plant part produced thereby are described. Maize variety ARC-9913C, the seed, the plant produced from the seed, and variants, mutants, and minor modifications of maize variety ARC-9913C are provided. Methods for producing maize varieties derived from maize variety ARC-9913C and methods of using maize variety ARC-9913C are disclosed.

HYBRID MAIZE VARIETY ARC-9913C**CROSS REFERENCE TO RELATED APPLICATION**

[0001] This application claims priority under 35 U.S.C. § 119 to provisional patent application U.S. Ser. No. 63/365,910, filed Jun. 6, 2022. The provisional patent application is herein incorporated by reference in its entirety, including without limitation, the specification, claims, and abstract, as well as any figures, tables, appendices, or drawings thereof.

BACKGROUND

[0002] The goal of hybrid development is to combine, in a single hybrid, various desirable traits. For field crops, these traits may include resistance to diseases and insects, resistance to heat and drought, reducing the time to crop maturity, greater yield, and better agronomic quality. With mechanical harvesting of many crops, uniformity of plant characteristics such as germination, stand establishment, growth rate, maturity, and plant and ear height are important. Traditional plant breeding is an important tool in developing new and improved commercial crops.

SUMMARY

[0003] Provided is a novel maize, *Zea mays* L., variety, seed, plant, cells and its parts designated as ARC-9913C, produced by crossing two maize inbred varieties. The hybrid maize variety ARC-9913C, the seed, the plant and its parts produced from the seed, and variants, mutants and minor modifications of maize ARC-9913C are provided. Processes are provided for making a maize plant containing in its genetic material one or more traits introgressed into ARC-9913C through locus conversion, backcrossing and/or transformation, and to the maize seed, plant and plant parts produced thereby. Methods for producing maize varieties derived from hybrid maize variety ARC-9913C are also provided. Also provided are maize plants having all the physiological and morphological characteristics of the hybrid maize variety ARC-9913C.

[0004] The hybrid maize plant may further comprise a cytoplasmic or nuclear factor capable of conferring male sterility or otherwise preventing self-pollination, such as by self-incompatibility. Parts of the disclosed maize plants are also provided, for example, pollen obtained from a hybrid plant and an ovule of the hybrid plant. Seed of the hybrid maize variety ARC-9913C is provided and may be provided as a population of maize seed of the variety designated ARC-9913C.

[0005] Compositions are provided comprising a seed of maize variety ARC-9913C comprised in plant seed growth media. In certain embodiments, the plant seed growth media is a soil or synthetic cultivation medium. In specific embodiments, the growth medium may be comprised in a container or may, for example, be soil in a field.

[0006] Hybrid maize variety ARC-9913C is provided comprising an added heritable trait. The heritable trait may be a genetic locus that is a dominant or recessive allele. In certain embodiments, the genetic locus confers traits such as, for example, male sterility, waxy starch, reduced lignin, herbicide tolerance or resistance, insect resistance, resistance to bacterial, fungal, nematode or viral disease, and altered or modified fatty acid, phytate, protein or carbohydrate metabolism. The genetic locus may be a naturally

occurring maize gene introduced into the genome of a parent of the variety by backcrossing, a natural or induced mutation, or a transgene introduced through genetic transformation techniques. When introduced through transformation, a genetic locus may comprise one or more transgenes integrated at a single chromosomal location.

[0007] A hybrid maize plant of the variety designated ARC-9913C is provided, wherein a cytoplasmically-inherited trait has been introduced into the hybrid plant. Such cytoplasmically-inherited traits are passed to progeny through the female parent in a particular cross. An exemplary cytoplasmically-inherited trait is the male sterility trait. Cytoplasmic-male sterility (CMS) is a pollen abortion phenomenon determined by the interaction between the genes in the cytoplasm and the nucleus. Alteration in the mitochondrial genome and the lack of restorer genes in the nucleus will lead to pollen abortion. With either a normal cytoplasm or the presence of restorer gene(s) in the nucleus, the plant will produce pollen normally. A CMS plant can be pollinated by a maintainer version of the same variety, which has a normal cytoplasm but lacks the restorer gene(s) in the nucleus, and continues to be male sterile in the next generation. The male fertility of a CMS plant can be restored by a restorer version of the same variety, which must have the restorer gene(s) in the nucleus. With the restorer gene(s) in the nucleus, the offspring of the male-sterile plant can produce normal pollen grains and propagate. A cytoplasmically inherited trait may be a naturally occurring maize trait or a trait introduced through genetic transformation techniques.

[0008] A tissue culture of regenerable cells of a plant of variety ARC-9913C is provided. The tissue culture can be capable of regenerating plants capable of expressing all of the physiological and morphological or phenotypic characteristics of the variety and of regenerating plants having substantially the same genotype as other plants of the variety. Examples of some of the physiological and morphological characteristics of the variety ARC-9913C that may be assessed include characteristics related to yield, maturity, and kernel quality. The regenerable cells in such tissue cultures can be derived, for example, from embryos, meristematic cells, immature tassels, microspores, pollen, leaves, anthers, roots, root tips, silk, flowers, kernels, ears, cobs, husks, or stalks, or from callus or protoplasts derived from those tissues. Maize plants regenerated from the tissue cultures and plants having all or essentially all of the physiological and morphological characteristics of variety ARC-9913C are also provided.

[0009] A method of producing hybrid maize seed comprising crossing a plant of variety GIQA9942 with a plant of variety GIP19629. In a cross, either parent may serve as the male or female. Processes are also provided for producing maize seeds or plants, which processes generally comprise crossing a first parent maize plant as a male or female parent with a second parent maize plant, wherein at least one of the first or second parent maize plants is a plant of the variety designated ARC-9913C. In such crossing, either parent may serve as the male or female parent. These processes may be further exemplified as processes for preparing hybrid maize seed or plants, wherein a first hybrid maize plant is crossed with a second maize plant of a different, distinct variety to provide a hybrid that has, as one of its parents, the hybrid maize plant variety ARC-9913C. In these processes, cross-

ing will result in the production of seed. The seed production occurs regardless of whether the seed is collected or not.

[0010] In some embodiments, the first step in “crossing” comprises planting, often in pollinating proximity, seeds of a first and second parent maize plant, and in many cases, seeds of a first maize plant and a second, distinct maize plant. Where the plants are not in pollinating proximity, pollination can nevertheless be accomplished by other means, such as by transferring a pollen or tassel bag from one plant to the other.

[0011] A second step comprises cultivating or growing the seeds of said first and second parent maize plants into plants that bear flowers (maize bears both male flowers (tassels) and female flowers (silks) in separate anatomical structures on the same plant).

[0012] A third step comprises preventing self-pollination of the plants, i.e., preventing the silks of a plant from being fertilized by any plant of the same variety, including the same plant. This can be done, for example, by emasculating the male flowers of the first or second parent maize plant, (i.e., treating or manipulating the flowers so as to prevent pollen production, to produce an emasculated parent maize plant). Self-incompatibility systems may also be used in some hybrid crops for the same purpose. Self-incompatible plants still shed viable pollen and can pollinate plants of other varieties but are incapable of pollinating themselves or other plants of the same variety.

[0013] A fourth step may comprise allowing cross-pollination to occur between the first and second parent maize plants. When the plants are not in pollinating proximity, this can be done by placing a bag, usually paper or glassine, over the tassels of the first plant and another bag over the silks of the incipient ear on the second plant. The bags are left in place for at least 24 hours. Since pollen is viable for less than 24 hours, this assures that the silks are not pollinated from other pollen sources, that any stray pollen on the tassels of the first plant is dead, and that the only pollen transferred comes from the first plant. The pollen bag over the tassel of the first plant is then shaken vigorously to enhance release of pollen from the tassels, and the shoot bag is removed from the silks of the incipient ear on the second plant. Finally, the pollen bag is removed from the tassel of the first plant and is placed over the silks of the incipient ear of the second plant, shaken again and left in place. Yet another step comprises harvesting the seeds from at least one of the parent maize plants. The harvested seed can be grown to produce a maize plant or hybrid maize plant.

[0014] Maize seed and plants are provided that are produced by a process that comprises crossing a first parent maize plant with a second parent maize plant, wherein at least one of the first or second parent maize plants is a plant of the variety designated ARC-9913C. Maize seed and plants produced by the process are first generation hybrid maize seed and plants produced by crossing an inbred with another, distinct inbred. Seed of an F1 hybrid maize plant, an F1 hybrid maize plant and seed thereof, specifically the hybrid variety designated ARC-9913C is provided.

[0015] The described plants may be analyzed by their genetic complement. Provided are maize plant cells that have a genetic complement in accordance with the disclosed maize plant cells, and plants, seeds and diploid plants containing such cells.

[0016] Plant genetic complements may be assessed by genetic marker profiles, and by the expression of phenotypic

traits that are characteristic of the expression of the genetic complement, e.g., isozyme typing profiles. It is understood that variety ARC-9913C could be identified by any of the many disclosed, well-known techniques used for genetic profiling.

DETAILED DESCRIPTION

[0017] A new and distinctive maize hybrid variety designated ARC-9913C is provided. Definitions

[0018] “Comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains,” “containing,” “characterized by” or any other variation thereof, are intended to cover a non-exclusive inclusion.

[0019] Unless expressly stated to the contrary, “or” is used as an inclusive term. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present). The indefinite articles “a” and “an” preceding an element or component are nonrestrictive regarding the number of instances (i.e., occurrences) of the element or component. Therefore “a” or “an” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

[0020] Maize, *Zea mays* L., can be referred to as maize or corn. Certain definitions used in the specification are provided below. To provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

[0021] “Allele” refers to any of one or more alternative forms of a genetic sequence. In a diploid cell or organism, the two alleles of a given sequence typically occupy corresponding loci on a pair of homologous chromosomes.

[0022] “Alter” refers to the utilization of up-regulation, down-regulation, or gene silencing in the context of genetic manipulation.

[0023] “Anthesis” refers to the time of a flower’s opening.

[0024] “Breeding cross” refers to the introduction of new genetic material into a plant for the development of a new variety. For example, one could cross plant A with plant B, wherein plant B would be genetically different from plant A. After the breeding cross, the resulting F1 plants could then be selfed or sibbed for one, two, three or more times (F1, F2, F3, etc.) until a new inbred variety is developed.

[0025] “Cell” refers to a plant cell, whether isolated, in tissue culture or incorporated in a plant or plant part.

[0026] “Cob color” refers to the color of the cob based on a visual assessment. Data values include deep red, red, pink and white.

[0027] “Cross pollination” refers to fertilization by the union of two gametes from different plants.

[0028] “Crossing” refers to the combination of genetic material by traditional methods such as a breeding cross or backcross, but also including protoplast fusion and other molecular biology methods of combining genetic material from two sources.

[0029] “Ear height” refers to the average height of the ear in inches when measured from the ground to the top developed ear node. The recorded number for a location is an average of 4 plants per plot.

[0030] “Early root lodging” refers to the number of plants leaning at about a 30-degree angle or more from the vertical

at the base of the plant are considered root lodged. The number of late root lodged plants is counted before anthesis and converted to a percent.

[0031] “Fall appearance” refers to a rating of total intactness of the plant in the fall. Scoring is on a 1 to 9 basis in which “1” is complete intactness and “9” is nothing left above the ear.

[0032] “F1 progeny” refers to a progeny plant produced by crossing a plant of one maize line with a plant of another maize line.

[0033] “Genetic complement” refers to the aggregate of nucleotide sequences, the expression of which defines the phenotype of, for example, a maize plant, or a cell or tissue of that plant. A genetic complement thus represents the genetic makeup of a cell, tissue or plant.

[0034] “Grain quality” refers to kernel/grain quality where 1=food grade quality w little dent and cap; 5=medium cap with medium grain texture; 9=high cap with rough grain texture.

[0035] “Growing degree units (GDUs)” refers to an estimation using the Barger Heat Unit Theory, which assumes that maize growth occurs in the temperature range 50° F. to 86° F. and that temperatures outside this range slow down growth; the maximum daily heat unit accumulation is 36 and the minimum daily heat unit accumulation is 0. The seasonal accumulation of GDU is a major factor in determining maturity zones.

[0036] “GDUs to black layer” refers to the number of growing degree units for a variety to have plants with corn kernels showing black layer formation (physiological maturity) GDUs to black layer is determined by summing the individual GDU daily values from the planting date to the date of 50% black layer.

[0037] “GDUs to pollen shed” refers to the number of growing degree units for a variety to have approximately 50% of the plants having pollen flowing from greater than or equal to 1.5 inches of main tassel spike as measured from the time of emergence. GDUs to pollen shed is determined by summing the individual GDU daily values from the emergence date to the date of 50% pollen shed.

[0038] “GDUs to silking” refers to the number of growing degree units for a variety to have approximately 50% of the plants having ears with exposed silks emergence as measured from the time of emergence. GDUs to silk is determined by summing the individual GDU daily values from the emergence date to the date of 50% silking.

[0039] “Gene silencing” refers to the interruption or suppression of the expression of a gene at the level of transcription or translation.

[0040] “Genotype” refers to the genetic makeup or profile of a cell or organism.

[0041] “Goss’ Wilt score” (*Clavibacter michiganensis* subspecies *nebraskensis*) refers to a 1 to 9 visual rating indicating the resistance to Goss’ Wilt. A lower score indicates a higher resistance. Data are collected only when sufficient selection pressure exists in the experiment measured.

[0042] “Grain yield” refers to the yield of the grain at harvest calculated in by weight per unit area, that is, bushels per acre, adjusted to 15% moisture.

[0043] “Greensnap” refers to the number of plants that snap below the ear as a result of brittleness following machine snapping or artificial selection pressure meant to

simulate high winds. The number of plants that snap is counted and converted to a percent.

[0044] “Haploid plant part” refers to a plant part or cell that has a haploid genotype.

[0045] “Harvest moisture” refers to the actual percentage moisture of the grain at harvest.

[0046] “Hybrid variety” refers to a substantially heterozygous hybrid line and minor genetic modifications thereof that retain the overall genetics of the hybrid line.

[0047] “Inbred” refers to a variety developed through inbreeding or doubled haploidy that preferably comprises homozygous alleles at about 95% or more of its loci. An inbred can be reproduced by selfing or growing in isolation so that the plants can only pollinate with the same inbred variety.

[0048] “Introgression” refers to the process of transferring genetic material from one genotype to another.

[0049] “Kernel-row average” refers to the average number of rows of kernels on a single ear.

[0050] “Late root lodging” refers to the number of plants leaning at a 30-degree angle or more from the vertical at the base of the plant are considered root lodged. The number of late root lodged plants is counted after anthesis and is converted to a percent.

[0051] “Late stalk lodging” refers to the number of plants that did stalk lodge. Plants are counted as stalk lodged if the plant is broken over or off below the ear. The number of late stalk lodged plants is counted just before harvest and converted to a percent.

[0052] “Linkage” refers to a phenomenon wherein alleles on the same chromosome tend to segregate together more often than expected by chance if their transmission was independent.

[0053] “Locus” refers to a specific location on a chromosome.

[0054] “Locus conversion” refers to plants within a variety that have been modified in a manner that retains the overall genetics of the variety and further comprises one or more loci with a specific desired trait, such as male sterility, insect resistance, disease resistance or herbicide tolerance or resistance. Examples of single locus conversions include mutant genes, transgenes and native traits finely mapped to a single locus. One or more locus conversion traits may be introduced into a single corn variety.

[0055] “Male sterility” refers to a male sterile plant is one which produces no viable pollen no (pollen that is able to fertilize the egg to produce a viable seed). Male sterility prevents self pollination. These male sterile plants are therefore useful in hybrid plant production.

[0056] “Nucleic acid” refers to an acidic, chainlike biological macromolecule consisting of multiple repeat units of phosphoric acid, sugar, and purine and pyrimidine bases.

[0057] “Percent identity” refers to the comparison of the alleles present in two varieties. For example, when comparing two inbred plants to each other, each inbred plant will have the same allele (and therefore be homozygous) at almost all of their loci. Percent identity is determined by comparing a statistically significant number of the homozygous alleles of two varieties. For example, a percent identity of 90% between ARC-9913C and other variety means that the two varieties have the same homozygous alleles at 90% of their loci.

[0058] “Plant” includes reference to an immature or mature whole plant, including a plant that has been detas-

seled or from which seed or grain has been removed. Seed or embryo that will produce the plant is also considered to be the plant.

[0059] “Plant Density” refers to a measurement of the number of corn plants per unit area after planting. Plant density is measured in thousands of plants per acre (PPA).

[0060] “Plant height” refers to the average height of the plant in inches when measured from the ground to the flag leaf node. The recorded number for a location is an average of 4 plants per plot.

[0061] “Plant part” includes leaves, stems, roots, seed, grain, embryo, pollen, ovules, flowers, ears, cobs, husks, stalks, root tips, anthers, pericarp, silk, tissue, cells and the like. In some embodiments, the plant part contains at least one cell of hybrid maize variety ARC-9913C or a locus conversion thereof.

[0062] “Relative maturity” refers to a predicted relative maturity based on the harvest moisture of the grain. The relative maturity rating is based on a known set of checks and utilizes standard linear regression analyses and is also referred to as the Comparative Relative Maturity Rating System that is similar to the Minnesota Relative Maturity Rating System.

[0063] “Resistance” is synonymous with tolerance and refers to the ability of a plant to withstand exposure to an insect, disease, herbicide or other condition. A resistant plant variety will have a level of resistance higher than a comparable wild-type variety.

[0064] “Seed” is synonymous with grain and refers to a fertilized and ripened ovule, consisting of the plant embryo, varying amounts of stored food material, and a protective outer seed coat.

[0065] “Self pollination” refers to the transfer of pollen from one flower to the same flower or another flower of the same plant.

[0066] “Sib pollination” refers to individual plants within the same family or variety are used for pollination.

[0067] “Silk color” refers to the predominant color of the corn silks after emergence from the ear, i.e., at the silking stage. The color is evaluated when silks are 1 -3 inches in length. Data values include yellow, green, red, pink, and purple.

[0068] “Simple-sequence repeats (SSRs)” refer to genetic markers based on polymorphisms in repeated nucleotide sequences, such as microsatellites. A marker system based on SSRs can be highly informative in linkage analysis relative to other marker systems in that multiple alleles may be present.

[0069] “Single-nucleotide polymorphism (SNP)” refers to a DNA sequence variation occurring when a single nucleotide in the genome differs between individual plant or plant varieties. The differences can be equated with different alleles, and indicate polymorphisms. A number of SNP markers can be used to determine a molecular profile of an individual plant or plant variety and can be used to compare similarities and differences among plants and plant varieties.

[0070] “Site specific integration” refers to genes that create a site for site specific DNA integration. This includes the introduction of FRT sites that may be used in the FLP/FRT system and/or Lox sites that may be used in the Cre/Loxp system.

[0071] “Southern Rust score” (*Puccinia polysora*) refers to a 1 to 9 visual rating indicating the resistance to Southern

Rust. A lower score indicates a higher resistance. Data are collected only when sufficient selection pressure exists in the experiment measured.

[0072] “Staygreen” is a measure of general plant health near the time of black layer formation (physiological maturity). Scoring is on a 1 to 9 basis in which “1” is the best score indicating better late-season plant health and “9” is the worst score.

[0073] “Test weight” refers to the weight of the grain in pounds for a given volume (bushel) adjusted for moisture less than or equal to 22%.

[0074] “Variety” refers to a maize line and minor genetic modifications thereof that retain the overall genetics of the line including but not limited to a locus conversion, a mutation, or a somaclonal variant.

[0075] All tables discussed in the Detailed Description section can be found at the end of the section.

[0076] Hybrid Maize Variety ARC-9913C is a single cross maize variety and can be made by crossing inbreds GIQA9942 and GIP19626. Locus conversions of Hybrid Maize Variety ARC-9913C can be made by crossing inbreds GIQA9942 and GIP19626 wherein GIQA9942 and/or GIP19626 comprise a locus conversion(s).

[0077] The maize variety has shown uniformity and stability within the limits of environmental influence for all the traits as described in the Variety Description Information (see Table 1, found at the end of the section). The inbred parents of this maize variety have been self-pollinated and ear-rowed a sufficient number of generations with careful attention paid to uniformity of plant type to ensure the homozygosity and phenotypic stability necessary for use in commercial hybrid seed production. The variety has been increased both by hand and in isolated fields with continued observation for uniformity. No variant traits have been observed or are expected in ARC-9913C.

[0078] Hybrid Maize Variety ARC-9913C can be reproduced by planting seeds of the inbred parent varieties, growing the resulting maize plants under cross pollinating conditions, and harvesting the resulting seed using techniques familiar to the agricultural arts.

[0079] In addition to phenotypic observations, a plant can also be described or identified by its genotype. The genotype of a plant can be characterized through a genetic marker profile. Genetic marker profiles can be obtained by techniques such as Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites, and Single Nucleotide Polymorphisms (SNPs).

[0080] Particular markers used for these purposes may include any type of marker and marker profile which provides a means of distinguishing varieties. A genetic marker profile can be used, for example, to identify plants of the same variety or related varieties or to determine or validate a pedigree. In addition to being used for identification of maize variety ARC-9913C and its plant parts, the genetic marker profile is also useful in developing a locus conversion of ARC-9913C.

[0081] Methods of isolating nucleic acids from maize plants and methods for performing genetic marker profiles

using SNP and SSR polymorphisms are well known in the art. SNPs are genetic markers based on a polymorphism in a single nucleotide. A marker system based on SNPs can be highly informative in linkage analysis relative to other marker systems in that multiple alleles may be present.

[0082] A method comprising isolating nucleic acids, such as DNA, from a plant, a plant part, plant cell or a seed of the disclosed maize plants is provided. The method can include mechanical, electrical and/or chemical disruption of the plant, plant part, plant cell or seed, contacting the disrupted plant, plant part, plant cell or seed with a buffer or solvent, to produce a solution or suspension comprising nucleic acids, optionally contacting the nucleic acids with a precipitating agent to precipitate the nucleic acids, optionally extracting the nucleic acids, and optionally separating the nucleic acids such as by centrifugation or by binding to beads or a column, with subsequent elution, or a combination thereof. If DNA is being isolated, an RNase can be included in one or more of the method steps. The nucleic acids isolated can comprise all or substantially all of the genomic DNA sequence, all or substantially all of the chromosomal DNA sequence or all or substantially all of the coding sequences (cDNA) of the plant, plant part, or plant cell from which they were isolated. The amount and type of nucleic acids isolated may be sufficient to permit whole genome sequencing of the plant from which they were isolated or chromosomal marker analysis of the plant from which they were isolated.

[0083] The methods can be used to produce nucleic acids from the plant, plant part, seed or cell, which nucleic acids can be, for example, analyzed to produce data. The data can be recorded. The nucleic acids from the disrupted cell, the disrupted plant, plant part, plant cell or seed or the nucleic acids following isolation or separation can be contacted with primers and nucleotide bases, and/or a polymerase to facilitate PCR sequencing or marker analysis of the nucleic acids. In some examples, the nucleic acids produced can be sequenced or contacted with markers to produce a genetic profile, a molecular profile, a marker profile, a haplotype, or any combination thereof. In some examples, the genetic profile or nucleotide sequence is recorded on a computer readable medium. In other examples, the methods may further comprise using the nucleic acids produced from plants, plant parts, plant cells or seeds in a plant breeding program, for example in making crosses, selection and/or advancement decisions in a breeding program. Crossing includes any type of plant breeding crossing method, including but not limited to crosses to produce hybrids, outcrossing, selfing, backcrossing, locus conversion, introgression and the like. Favorable genotypes and or marker profiles, optionally associated with a trait of interest, may be identified by one or more methodologies. In some examples one or more markers are used, including but not limited to AFLPs, RFLPs, ASH, SSRs, SNPs, indels, padlock probes, molecular inversion probes, microarrays, sequencing, and the like. In some methods, a target nucleic acid is amplified prior to hybridization with a probe. In other cases, the target nucleic acid is not amplified prior to hybridization, such as methods using molecular inversion probes. In some examples, the genotype related to a specific trait is monitored, while in other examples, a genome-wide evaluation including but not limited to one or more of marker panels, library screens, association studies, microarrays, gene chips, expression studies, or sequencing such as whole-genome

resequencing and genotyping-by-sequencing (GBS) may be used. In some examples, no target-specific probe is needed, for example by using sequencing technologies, including but not limited to next-generation sequencing methods such as sequencing by synthesis, sequencing by ligation, and single molecule sequencing (SMS or third-generation sequencing) which eliminate template amplification. Further technologies include optical sequencing systems, and nanopore sequencing. Each of these may be coupled with one or more enrichment strategies for organellar or nuclear genomes to reduce the complexity of the genome under investigation via PCR, hybridization, restriction enzyme, and expression methods. In some examples, no reference genome sequence is needed to complete the analysis. ARC-9913C and its plant parts can be identified through a molecular marker profile. Such plant parts may be either diploid or haploid. The plant part includes at least one cell of the plant from which it was obtained, such as a diploid cell, a haploid cell or a somatic cell. Also provided are plants and plant parts substantially benefiting from the use of variety ARC-9913C in their development, such as variety ARC-9913C comprising a locus conversion.

[0084] A breeder uses various methods to help determine which plants should be selected from segregating populations and ultimately which inbred varieties will be used to develop hybrids for commercialization. In addition to knowledge of the germplasm and plant genetics, a part of the hybrid selection process is dependent on experimental design coupled with the use of statistical analysis. Experimental design and statistical analysis are used to help determine which hybrid combinations are significantly better or different for one or more traits of interest. Experimental design methods are used to assess error so that differences between two hybrid varieties can be more accurately evaluated. Statistical analysis includes the calculation of mean values, determination of the statistical significance of the sources of variation, and the calculation of the appropriate variance components. One of ordinary skill in the art of plant breeding would know how to evaluate the traits of two plant varieties to determine if there is no significant difference between the two traits expressed by those varieties. Mean trait values may be used to determine whether trait differences are significant. Trait values should preferably be measured on plants grown under the same environmental conditions, and environmental conditions should be appropriate for the traits or traits being evaluated. Sufficient selection pressure should be present for optimum measurement of traits of interest such as herbicide tolerance or herbicide, insect or disease resistance. For example, a locus conversion of ARC-9913C for herbicide resistance or tolerance should be compared with an isogenic counterpart in the absence of the herbicide. In addition, a locus conversion for insect or disease resistance should be compared to the isogenic counterpart, in the absence of disease pressure or insect pressure.

[0085] BLUP, Best Linear Unbiased Prediction, values can be reported for maize hybrid ARC-9913C and/or maize hybrid ARC-9913C comprising locus conversions. BLUP values can also be reported for other hybrids adapted to the same growing region as maize hybrid ARC-9913C with corresponding locus conversions.

[0086] During the inbreeding process in maize, the vigor of the varieties decreases. However, vigor is restored when two different inbred varieties are crossed to produce the

hybrid progeny (F1). An important consequence of the homozygosity and homogeneity of the inbred varieties is that the hybrid between a defined pair of inbreds may be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained. Once the inbreds that create a superior hybrid have been identified, a continual supply of the hybrid seed can be produced using these inbred parents and the hybrid corn plants can then be generated from this hybrid seed supply.

[0087] ARC-9913C or its parents may also be used to produce a double cross hybrid or a three-way hybrid. A single cross hybrid is produced when two inbred varieties are crossed to produce the F1 progeny. A double cross hybrid is produced from four inbred varieties crossed in pairs (A×B and C×D) and then the two F1 hybrids are crossed again (A×B)×(C×D). A three-way cross hybrid is produced from three inbred varieties where two of the inbred varieties are crossed (A×B) and then the resulting F1 hybrid is crossed with the third inbred variety (A×B)×C. In each case, pericarp tissue from the female parent will be a part of and protect the hybrid seed.

[0088] Another form of commercial hybrid production involves the use of a mixture of male sterile hybrid seed and male pollinator seed. When planted, the resulting male sterile hybrid plants are pollinated by the pollinator plants. This method can be used to produce grain with enhanced quality grain traits, such as high oil, because desired quality grain traits expressed in the pollinator will also be expressed in the grain produced on the male sterile hybrid plant. In this method the desired quality grain trait does not have to be incorporated by lengthy procedures such as recurrent backcross selection into an inbred parent line.

[0089] Molecular data from ARC-9913C may be used in a plant breeding process. Nucleic acids may be isolated from a seed of ARC-9913C or from a plant, plant part, or cell produced by growing a seed of ARC-9913C, or from a seed of ARC-9913C with a locus conversion, or from a plant, plant part, or cell of ARC-9913C with a locus conversion. One or more polymorphisms may be isolated from the nucleic acids. A plant having one or more of the identified polymorphisms may be selected and used in a plant breeding method to produce another plant.

[0090] Hybrid variety ARC-9913C represents a new base genetic line into which a new locus or trait may be introduced or introgressed. Transformation and backcrossing represent two methods that can be used to accomplish such an introgression. "Locus conversion" is used to designate the product of such an introgression.

[0091] To select and develop a superior hybrid, it is necessary to identify and select genetically unique individuals that occur in a segregating population. The segregating population is the result of a combination of crossover events plus the independent assortment of specific combinations of alleles at many gene loci that results in specific and unique genotypes. Once such a variety is developed its value to society is substantial since it is important to advance the germ plasm base as a whole to maintain or improve traits such as yield, disease resistance, pest resistance and plant performance in extreme weather conditions. Locus conversions are routinely used to add or modify one or a few traits of such a line and this further enhances its value and usefulness to society.

[0092] Backcrossing can be used to improve inbred varieties and a hybrid variety which is made using those inbreds.

Backcrossing can be used to transfer a specific desirable trait from one variety, the donor parent, to an inbred called the recurrent parent which has overall good agronomic characteristics yet that lacks the desirable trait. This transfer of the desirable trait into an inbred with overall good agronomic characteristics can be accomplished by first crossing a recurrent parent to a donor parent (non-recurrent parent). The progeny of this cross is then mated back to the recurrent parent followed by selection in the resultant progeny for the desired trait to be transferred from the non-recurrent parent.

[0093] Traits may be used by those of ordinary skill in the art to characterize progeny. Traits are commonly evaluated at a significance level, such as a 1%, 5% or 10% significance level, when measured in plants grown in the same environmental conditions. For example, a locus conversion of ARC-9913C may be characterized as having essentially the same or essentially all of the phenotypic traits or physiological and morphological traits or characteristics as ARC-9913C. By essentially all of the phenotypic characteristics or morphological and physiological characteristics, it is meant that all of the characteristics of a plant are recovered that are otherwise present when compared in the same environment, other than an occasional variant trait that might arise during backcrossing or direct introduction of a transgene or genetic modification. The traits used for comparison may be those traits shown in Table 1 as determined at the 5% significance level when grown under the same environmental conditions. Molecular markers can also be used during the breeding process for the selection of qualitative traits. For example, markers can be used to select plants that contain the alleles of interest during a backcrossing breeding program. The markers can also be used to select for the genome of the recurrent parent and against the genome of the donor parent. Using this procedure can minimize the amount of genome from the donor parent that remains in the selected plants.

[0094] A backcross or locus conversion of ARC-9913C can be developed when DNA sequences are introduced through backcrossing, with a parent of ARC-9913C utilized as the recurrent parent. Naturally occurring, modified and transgenic DNA sequences may be introduced through backcrossing techniques. A backcross or locus conversion may produce a plant with a trait or locus conversion in at least one or more backcrosses, including at least 2 backcrosses, at least 3 backcrosses, at least 4 backcrosses, at least 5 backcrosses and the like. Molecular marker assisted breeding or selection may be utilized to reduce the number of backcrosses necessary to achieve the backcross conversion. A backcross locus conversion can be made in as few as two backcrosses.

[0095] The complexity of the backcross conversion method depends on the type of trait being transferred (a single gene or closely linked genes compared to unlinked genes), the level of expression of the trait, the type of inheritance (cytoplasmic or nuclear), dominant or recessive trait expression, and the types of parents included in the cross. It is understood by those of ordinary skill in the art that for single locus or gene traits that are relatively easy to classify, the backcross method is effective and relatively easy to manage. Desired traits that may be transferred through backcross conversion include, but are not limited to, waxy starch, sterility (nuclear and cytoplasmic), fertility restoration, grain color (white), nutritional enhancements, drought tolerance, nitrogen utilization, altered fatty acid profile, increased digestibility, low phytate, industrial

enhancements, disease resistance (bacterial, fungal, or viral), insect resistance, and herbicide tolerance or resistance. A locus conversion, also called a trait conversion, can be a native trait or a transgenic trait. In addition, a recombination site itself, such as an FRT site, Lox site or other site specific integration site, may be inserted by backcrossing and utilized for direct insertion of one or more genes of interest into a specific plant variety. The trait of interest is transferred from the donor parent to the recurrent parent, in this case, an inbred parent of the disclosed maize variety.

[0096] A single locus may contain several transgenes, such as a transgene for disease resistance that, in the same expression vector, also contains a transgene for herbicide tolerance or resistance. The gene for herbicide tolerance or resistance may be used as a selectable marker and/or as a phenotypic trait. A single locus conversion of a site specific integration system allows for the integration of multiple genes at a known recombination site in the genome. At least one, at least two or at least three and less than ten, less than nine, less than eight, less than seven, less than six, less than five or less than four locus conversions may be introduced into the plant by backcrossing, introgression or transformation to express the desired trait, while the plant, or a plant grown from the seed, plant part or plant cell, otherwise retains the phenotypic characteristics of the deposited seed when grown under the same environmental conditions.

[0097] The backcross or locus conversion may result from either the transfer of a dominant allele or a recessive allele. Selection of progeny containing the trait of interest can be accomplished by direct selection for a trait associated with a dominant allele. Transgenes transferred via backcrossing typically function as a dominant single gene trait and are relatively easy to classify. Selection of progeny for a trait that is transferred via a recessive allele, such as the waxy starch characteristic, requires growing and selfing the first backcross generation to determine which plants carry the recessive alleles. Recessive traits may require additional progeny testing in successive backcross generations to determine the presence of the locus of interest. The last backcross generation is usually selfed to give pure breeding progeny for the gene(s) being transferred, although a backcross conversion with a stably introgressed trait may also be maintained by further backcrossing to the recurrent parent with selection for the converted trait.

[0098] Along with selection for the trait of interest, progeny are selected for the phenotype and/or genotype of the recurrent parent. While occasionally additional polynucleotide sequences or genes may be transferred along with the backcross conversion, the backcross conversion variety fits into the same hybrid combination as the recurrent parent inbred variety and contributes the effect of the additional locus added through the backcross.

[0099] When one or more traits are introgressed into the variety a difference in quantitative agronomic traits, such as yield or dry down, between the variety and an introgressed version of the variety in some environments may occur. For example, the introgressed version, may provide a net yield increase in environments where the trait provides a benefit, such as when a variety with an introgressed trait for insect resistance is grown in an environment where insect pressure exists, or when a variety with herbicide tolerance is grown in an environment where the herbicide is used.

[0100] The modified ARC-9913C may be further characterized as having essentially the same phenotypic charac-

teristics of maize variety ARC-9913C such as are listed in Table 1 when grown under the same or similar environmental conditions and/or may be characterized by percent identity to ARC-9913C as determined by molecular markers, such as SSR markers or SNP markers. Examples of percent identity determined using markers include at least 95%, 96%, 97%, 98%, 99% or 99.5%.

[0101] Traits can be used by those of ordinary skill in the art to characterize progeny. Traits are commonly evaluated at a significance level, such as a 1%, 5% or 10% significance level, when measured in plants grown in the same environmental conditions.

[0102] Hybrid seed production requires elimination or inactivation of pollen produced by the female inbred parent. Incomplete removal or inactivation of the pollen provides the potential for self-pollination. A reliable method of controlling male fertility in plants offers the opportunity for improved seed production. There are several ways in which a maize plant can be manipulated so that it is male sterile. These include use of manual or mechanical emasculation (or detasseling), use of one or more genetic factors that confer male sterility, including cytoplasmic genetic and/or nuclear genetic male sterility, use of gametocides and the like. A male sterile variety designated ARC-9913C may include one or more genetic factors, which result in cytoplasmic genetic and/or nuclear genetic male sterility. The male sterility may be either partial or complete male sterility.

[0103] Hybrid maize seed is often produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two inbred varieties of maize are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female). Provided that there is sufficient isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male), and the resulting seed is therefore hybrid and will form hybrid plants.

[0104] Large scale commercial maize hybrid production, as it is practiced today, requires the use of some form of male sterility system which controls or inactivates male fertility. A reliable method of controlling male fertility in plants also offers the opportunity for improved plant breeding. This is especially true for development of maize hybrids, which relies upon some sort of male sterility system. There are several ways in which a maize plant can be manipulated so that it is male sterile. These include use of manual or mechanical emasculation (or detasseling), cytoplasmic genetic male sterility, nuclear genetic male sterility, gametocides and the like.

[0105] The laborious detasseling process can be avoided by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of genetic factors in the cytoplasm, as opposed to the nucleus, and so nuclear linked genes are not transferred during backcrossing. Thus, this characteristic is inherited exclusively through the female parent in maize plants, since only the female provides cytoplasm to the fertilized seed. CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile, and either option may be preferred depending on the intended use of the hybrid. The same hybrid seed, a portion produced from detasseled fertile maize and a portion produced using the CMS system can be blended to insure that adequate pollen loads are available for fertilization when the hybrid plants

are grown. CMS systems have been successfully used since the 1950's, and the male sterility trait is routinely backcrossed into inbred varieties.

[0106] There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, and chromosomal translocations. In addition to these methods, a system of nuclear male sterility which includes: identifying a gene which is needed for male fertility; silencing this native gene which is needed for male fertility; removing the native promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered gene back into the plant; and thus creating a plant that is male sterile because the inducible promoter is not "on" resulting in the male fertility gene not being transcribed is known to those skilled in the corn breeding art. Fertility is restored by inducing, or turning "on", the promoter, which in turn allows the gene that confers male fertility to be transcribed.

[0107] These, and the other methods of conferring genetic male sterility in the art, each possess their own benefits and drawbacks. Some other methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an antisense system in which a gene needed for fertility is identified and an antisense to that gene is inserted in the plant.

[0108] Another system for controlling male sterility makes use of gametocides. Gametocides are not a genetic system, but rather a topical application of chemicals. These chemicals affect cells that are needed for male fertility. The application of these chemicals affects fertility in the plants only for the growing season in which the gametocide is applied. Application of the gametocide, timing of the application and genotype specificity often limit the usefulness of the approach and it is not appropriate in all situations.

[0109] Transgenes and transformation methods facilitate engineering of the genome of plants to contain and express heterologous genetic elements, such as foreign genetic elements, or additional copies of endogenous elements, or modified versions of native or endogenous genetic elements to alter at least one trait of a plant in a specific manner. Any sequences, such as DNA, whether from a different species or from the same species, which have been stably inserted into a genome using transformation are referred to collectively as "transgenes" and/or "transgenic events". Transgenes can be moved from one genome to another using breeding techniques which may include, for example, crossing, backcrossing or double haploid production. In some embodiments, a transformed variant of ARC-9913C may comprise at least one transgene but could contain at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 and/or no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, or 2. Transformed versions of the claimed maize variety ARC-9913C containing and inheriting the transgene thereof are provided.

[0110] Numerous methods for plant transformation have been developed, including biological and physical plant transformation protocols. In addition, expression vectors and in vitro culture methods for plant cell or tissue transformation and regeneration of plants are available.

[0111] In general, methods to transform, modify, edit or alter plant endogenous genomic DNA include altering the plant native DNA sequence or a pre-existing transgenic sequence including regulatory elements, coding and non-

coding sequences. These methods can be used, for example, to target nucleic acids to pre-engineered target recognition sequences in the genome. Such pre-engineered target sequences may be introduced by genome editing or modification. As an example, a genetically modified plant variety is generated using "custom" or engineered endonucleases such as meganucleases produced to modify plant genomes. Another site-directed engineering method is through the use of zinc finger domain recognition coupled with the restriction properties of restriction enzyme. A transcription activator-like (TAL) effector-DNA modifying enzyme (TALE or TALEN) is also used to engineer changes in plant genome. Site-specific modification of plant genomes can also be performed using the bacterial type II CRISPR (clustered regularly interspaced short palindromic repeats)/Cas (CRISPR-associated) system. The Cas9/guide RNA-based system allows targeted cleavage of genomic DNA guided by a customizable small noncoding RNA in plants.

[0112] Plant transformation methods may involve the construction of an expression vector. Such a vector comprises a DNA sequence that contains a gene under the control of or operatively linked to a regulatory element, for example a promoter. The vector may contain one or more genes and one or more regulatory elements.

[0113] A transgenic event which has been stably engineered into the germ cell line of a particular maize plant using transformation techniques, could be moved into the germ cell line of another variety using traditional breeding techniques that are well known in the plant breeding arts. These varieties can then be crossed to generate a hybrid maize variety plant such as maize variety plant ARC-9913C which comprises a transgenic event. For example, a backcrossing approach is commonly used to move a transgenic event from a transformed maize plant to another variety, and the resulting progeny would then comprise the transgenic event(s). Also, if an inbred variety was used for the transformation then the transgenic plants could be crossed to a different inbred to produce a transgenic hybrid maize plant.

[0114] Various genetic elements can be introduced into the plant genome using transformation. These elements include, but are not limited to genes; coding sequences; inducible, constitutive, and tissue specific promoters; enhancing sequences; and signal and targeting sequences. In addition, transformability of a variety can be increased by introgressing the trait of high transformability from another variety known to have high transformability, such as Hi-II.

[0115] With transgenic or genetically modified plants, a foreign protein can be produced in commercial quantities. Thus, techniques for the selection and propagation of transformed plants, which are well understood in the art, yield a plurality of transgenic or genetically modified plants that are harvested in a conventional manner, and a foreign protein then can be extracted from a tissue of interest or from total biomass. Protein extraction from plant biomass can be accomplished by known methods.

[0116] Transgenic events can be mapped by one of ordinary skill in the art and such techniques are well known to those of ordinary skill in the art.

[0117] Plants can be genetically engineered or modified to express various phenotypes of agronomic interest. Through the transformation or modification of maize the expression of genes can be altered to enhance disease resistance, insect resistance, herbicide tolerance, agronomic traits, grain quality and other traits. Transformation can also be used to insert

DNA sequences which control or help control male-sterility. DNA sequences native to maize as well as non-native DNA sequences can be transformed into maize and used to alter levels of native or non-native proteins. Various promoters, targeting sequences, enhancing sequences, and other DNA sequences can be inserted into the maize genome for the purpose of altering the expression of proteins. Reduction of the activity of specific genes (also known as gene silencing, or gene suppression) is desirable for several aspects of genetic engineering in plants.

[0118] Many techniques for gene silencing are well known to one of skill in the art, including but not limited to knock-outs (such as by insertion of a transposable element such as mu or other genetic elements such as a FRT, Lox or other site specific integration site, antisense technology; co-suppression, virus-induced gene silencing; target-RNA-specific ribozymes; hairpin structures; MicroRNA; ribozymes; oligonucleotide mediated targeted modification; Zn-finger targeted molecules; and other methods or combinations of the above methods known to those of skill in the art.

[0119] Exemplary nucleotide sequences that may be altered by genetic engineering include, but are not limited to, those categorized below.

[0120] 1. Transgenes That Confer Resistance to Insects or Disease and That Encode:

[0121] (A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. A plant resistant to a disease is one that is more resistant to a pathogen as compared to the wild type plant.

[0122] (B) A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. DNA molecules encoding delta-endotoxin genes can be purchased from American Type Culture Collection (Manassas, Va.), for example, under ATCC Accession Nos. 40098, 67136, 31995 and 31998.

[0123] (C) An insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof

[0124] (D) An insect-specific peptide which, upon expression, disrupts the physiology of the affected pest. For example, an insect diuretic hormone receptor or an allostatin.

[0125] (E) An enzyme responsible for a hyperaccumulation of a monoterpene, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.

[0126] (F) An enzyme involved in the modification, including the post-translational modification, of a biologically active molecule; for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic.

[0127] (G) A molecule that stimulates signal transduction. For example, calmodulin cDNA clones.

[0128] (H) A hydrophobic moment peptide.

[0129] (I) A membrane permease, a channel former or a channel blocker.

[0130] (J) A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. Coat protein-mediated resistance may be conferred upon transformed plants against, for example, alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus.

[0131] (K) An insect-specific antibody or an immunotoxin derived therefrom. For example, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect.

[0132] (L) A virus-specific antibody. Plants expressing recombinant antibody genes may be protected from virus attack.

[0133] (M) A developmental-arrestive protein produced in nature by a pathogen or a parasite. For example, fungal endo alpha-1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo-alpha-1,4-D-galacturonase.

[0134] (N) A developmental-arrestive protein produced in nature by a plant. For example, plants expressing the barley ribosome-inactivating gene may have an increased resistance to fungal disease.

[0135] (O) Genes involved in the Systemic Acquired Resistance (SAR) Response and/or the pathogenesis related genes

[0136] (P) Antifungal genes.

[0137] (Q) Detoxification genes, such as for fumonisin, beauvericin, moniliformin and zearalenone and their structurally related derivatives.

[0138] (R) Cystatin and cysteine proteinase inhibitors.

[0139] (S) Defensin genes.

[0140] (T) Genes conferring resistance to nematodes.

[0141] (U) Genes that confer resistance to *Phytophthora* Root Rot, such as the Rps 1, Rps 1-a, Rps 1-b, Rps 1-c, Rps 1-d, Rps 1-e, Rps 1-k, Rps 2, Rps 3-a, Rps 3-b, Rps 3-c, Rps 4, Rps 5, Rps 6, Rps 7 and other Rps genes.

[0142] (V) Genes that confer resistance to Brown Stem Rot.

[0143] (W) Genes that confer resistance to *Colletotrichum*. This includes the Rcg locus that may be utilized as a single locus conversion.

[0144] 2. Transgenes That Confer Tolerance to An Herbicide, For Example:

[0145] (A) An herbicide that inhibits the growing point or meristem, such as an imidazolinone or a sulfonyleurea. Exemplary genes in this category code for mutant acetolactate synthase (ALS) and acetohydroxyacid synthase (AHAS) enzyme.

[0146] (B) Glyphosate (tolerance imparted by mutant 5-enolpyruvyl-3-phosphokimate synthase (EPSP) and aroA genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin acetyl transferase (PAT) and *Streptomyces hygroscopicus* phosphinothricin acetyl transferase (bar) genes), and pyridinoxy or phenoxy propionic acids and cyclohexones (ACCase inhibitor-encoding

genes). Glyphosate tolerance is also imparted to plants that express a gene that encodes a glyphosate oxido-reductase enzyme. In addition, glyphosate tolerance can be imparted to plants by the over expression of genes encoding glyphosate N-acetyltransferase. A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC accession No. 39256. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is known to those skilled in the art as are exemplary genes conferring resistance to phenoxy propionic acids, cyclohexanediones and cyclohexones, such as sethoxydim and haloxyfop, are the *Accl-S1*, *Accl-S2* and *Accl-S3* genes.

[0147] (C) An herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes), glutathione S-transferase and a benzonitrile (nitrilase gene) such as bromoxynil. Nucleotide sequences for nitrilase genes are known, and DNA molecules containing these genes are available under ATCC Accession Nos. 53435, 67441 and 67442.

[0148] (D) Other genes that confer tolerance to herbicides include: a gene encoding a chimeric protein of rat cytochrome P4507A1 and yeast NADPH-cytochrome P450 oxidoreductase, genes for glutathione reductase and superoxide dismutase, and genes for various phosphotransferases.

[0149] (E) An herbicide that inhibits protoporphyrinogen oxidase (protox or PPO) is necessary for the production of chlorophyll, which is necessary for all plant survival. The protox enzyme serves as the target for a variety of herbicidal compounds. PPO-inhibitor herbicides can inhibit growth of all the different species of plants present, causing their total destruction. The development of plants containing altered protox activity which are tolerant to these herbicides are known to those skilled in the art.

[0150] (F) Dicamba (3,6-dichloro-2-methoxybenzoic acid) is an organochloride derivative of benzoic acid which functions by increasing plant growth rate such that the plant dies.

[0151] 3. Transgenes That Confer or Contribute to an Altered Grain Characteristic, Such as:

[0152] (A) Altered fatty acids.

[0153] (1) Down-regulation of stearoyl-ACP desaturase to increase stearic acid content of the plant.

[0154] (2) Elevating oleic acid via FAD-2 gene modification and/or decreasing linolenic acid via FAD-3 gene modification.

[0155] (3) Altering conjugated linolenic or linoleic acid content.

[0156] (4) Altering LEC1, AGP, Dek1, Superalt milps, various *Ipa* genes such as *Ipa1*, *Ipa3*, *hpt* or *hgt*.

[0157] (B) Altered phosphate content.

[0158] (1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant.

[0159] (2) Modulating a gene that reduces phytate content. In maize, this, for example, could be accomplished, by cloning and then re-introducing DNA associated with one or more of the alleles, such as the LPA alleles, identified in maize mutants characterized by low levels of phytic acid.

[0160] (C) Altered carbohydrates affected, for example, by altering a gene for an enzyme that affects the branching pattern of starch or, a gene altering thio-

doxin such as NTR and/or TRX and/or a gamma zein knock out or mutant such as *cs27* or *TUSC27* or *en27*. The fatty acid modification genes may also be used to affect starch content and/or composition through the interrelationship of the starch and oil pathways.

[0161] (D) Altered antioxidant content or composition, such as alteration of tocopherol or tocotrienols.

[0162] (E) Altered essential seed amino acids.

[0163] 4. Genes that Control Male-sterility

[0164] There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility.

[0165] (A) Introduction of a deacetylase gene under the control of a tapetum-specific promoter and with the application of the chemical N-Ac-PPT.

[0166] (B) Introduction of various stamen-specific promoters.

[0167] (C) Introduction of the barnase and the barstar gene.

[0168] For additional examples of nuclear male and female sterility systems and genes.

[0169] 5. Genes that create a site for site specific DNA integration. This includes the introduction of FRT sites that may be used in the FLP/FRT system and/or Lox sites that may be used in the Cre/Lox system. Other systems that may be used include the *Gin* recombinase of phage Mu, the *Pin* recombinase of *E. coli*, and the R/RS system of the pSR1 plasmid.

[0170] 6. Genes that affect abiotic stress resistance (including but not limited to flowering, ear and seed development, enhancement of nitrogen utilization efficiency, altered nitrogen responsiveness, drought resistance or tolerance, cold resistance or tolerance, and salt resistance or tolerance) and increased yield under stress.

[0171] Other genes and transcription factors that affect plant growth and agronomic traits such as yield, flowering, plant growth and/or plant structure, can be introduced or introgressed into plants.

[0172] The development of maize hybrids in a maize plant breeding program requires, in general, the development of homozygous inbred lines, the crossing of these lines, and the evaluation of the crosses. Maize plant breeding programs combine the genetic backgrounds from two or more inbred varieties or various other germplasm sources into breeding populations from which new inbred varieties are developed by selfing and selection of desired phenotypes. Hybrids also can be used as a source of plant breeding material or as source populations from which to develop or derive new maize varieties.

[0173] Plant breeding techniques known in the art and used in a maize plant breeding program include, but are not limited to, recurrent selection, mass selection, bulk selection, backcrossing, making double haploids, pedigree breeding, open pollination breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation. Often combinations of these techniques are used. The inbred varieties derived from hybrids can be developed using plant breeding techniques as described above. New inbreds are crossed with other inbred varieties and the hybrids from these crosses are evaluated to determine which of those have commercial potential. The

oldest and most traditional method of analysis is the observation of phenotypic traits but genotypic analysis may also be used.

[0174] Methods for producing a maize plant by crossing a first parent maize plant with a second parent maize plant wherein either the first or second parent maize plant is a maize plant of the variety ARC-9913C are provided. The other parent may be any other maize plant, such as another inbred variety or a plant that is part of a synthetic or natural population. Any such methods using the maize variety ARC-9913C in crossing or breeding are provided, such as, for example: selfing, sibbing, backcrosses, mass selection, pedigree breeding, bulk selection, hybrid production, crosses to populations, and the like.

[0175] Recurrent selection is a method used in a plant breeding program to improve a population of plants. ARC-9913C is suitable for use in a recurrent selection program. The method entails individual plants cross pollinating with each other to form progeny. The progeny are grown and the superior progeny selected by any number of selection methods, which include individual plant, half-sib progeny, full-sib progeny, selfed progeny and topcrossing. The selected progeny are cross pollinated with each other to form progeny for another population. This population is planted and again superior plants are selected to cross pollinate with each other. Recurrent selection is a cyclical process and therefore can be repeated as many times as desired. The objective of recurrent selection is to improve the traits of a population. The improved population can then be used as a source of breeding material to obtain inbred varieties to be used in hybrids or used as parents for a synthetic cultivar. A synthetic cultivar is the resultant progeny formed by the intercrossing of several selected inbreds.

[0176] ARC-9913C is suitable for use in mass selection. Mass selection is a useful technique when used in conjunction with molecular marker enhanced selection. In mass selection seeds from individuals are selected based on phenotype and/or genotype. These selected seeds are then bulked and used to grow the next generation. Bulk selection requires growing a population of plants in a bulk plot, allowing the plants to self-pollinate, harvesting the seed in bulk and then using a sample of the seed harvested in bulk to plant the next generation. Instead of self-pollination, directed pollination could be used as part of the breeding program.

[0177] The production of double haploids from ARC-9913C can also be used for the development of inbreds. Double haploids are produced by the doubling of a set of chromosomes (1N) from a heterozygous plant to produce a completely homozygous individual. For example, a method is provided of obtaining a substantially homozygous ARC-9913C progeny plant by obtaining a seed from the cross of ARC-9913C and another maize plant and applying double haploid methods to the F1 seed or F1 plant or to any successive filial generation. Methods for producing plants by doubling haploid seed generated by a cross of the disclosed plants, or parts thereof, with a different maize plant are provided. The use of double haploids substantially decreases the number of generations required to produce an inbred with similar genetics or characteristics to ARC-9913C. This can be advantageous because the process omits the generations of selfing needed to obtain a homozygous plant from a heterozygous source.

[0178] Haploid induction systems have been developed for various plants to produce haploid tissues, plants and seeds. The haploid induction system can produce haploid plants from any genotype by crossing a selected variety (as female) with an inducer variety. Such inducer varieties for maize include Stock 6, RWS, KEMS, or KMS and ZMS, and indeterminate gametophyte (ig) mutation.

[0179] In particular, a process of making seed substantially retaining the molecular marker profile of maize variety ARC-9913C is provided. Obtaining a seed of hybrid maize variety ARC-9913C further comprising a locus conversion, wherein representative seed is produced by crossing a first plant of variety GIQA9942 or a locus conversion thereof with a second plant of variety GIP19626 or a locus conversion thereof, and wherein representative seed of said varieties GIQA9942 and GIP19626 have been deposited and wherein said maize variety ARC-9913C further comprising a locus conversion has 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% A of the same polymorphisms for molecular markers as the plant or plant part of maize variety ARC-9913C. Sequences for the public markers can be found, for example, in the Panzea database which is available online from Panzea. The type of molecular marker used in the molecular profile can be but is not limited to Single Nucleotide Polymorphisms, SNPs. A process of making seed retaining essentially the same phenotypic, physiological, morphological or any combination thereof characteristics of maize variety ARC-9913C is also contemplated. Obtaining a seed of hybrid maize variety ARC-9913C further comprising a locus conversion, wherein representative seed is produced by crossing a first plant of variety GIQA9942 or a locus conversion thereof with a second plant of variety GIP19626 or a locus conversion thereof, and wherein representative seed of said varieties GIQA9942 and GIP19626 have been deposited and wherein said maize variety ARC-9913C further comprising a locus conversion has essentially the same morphological characteristics as maize variety ARC-9913C when grown in the same environmental conditions. The same environmental conditions may be, but is not limited to, a side-by-side comparison. The characteristics can be or include, for example, those listed in Table 1. The comparison can be made using any number of professionally accepted experimental designs and statistical analysis.

[0180] Methods of tissue culturing cells of ARC-9913C and a tissue culture of ARC-9913C is provided. "Tissue culture" includes plant protoplasts, plant cell tissue culture, cultured microspores, plant calli, plant clumps, and the like. In certain embodiments, the tissue culture comprises embryos, protoplasts, meristematic cells, pollen, leaves or anthers derived from immature tissues of pollen, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk, and the like. Phrases such as "growing the seed" or "grown from the seed" include embryo rescue, isolation of cells from seed for use in tissue culture, as well as traditional growing methods.

[0181] Methods for preparing and maintaining plant tissue cultures are well known in the art. In certain embodiments, cells are provided which upon growth and differentiation produce maize plants having the genotype and/or phenotypic characteristics of variety ARC-9913C.

[0182] Methods of harvesting the grain of the F1 plant of variety ARC-9913C and using the F2 grain as seed for planting are provided. Also provided are methods of using

the seed of variety ARC-9913C, or selfed grain harvested from variety ARC-9913C, as seed for planting.

[0183] Embodiments include cleaning the seed, treating the seed, and/or conditioning the seed and seed produced by such cleaning, conditioning, treating or any combination thereof. Cleaning the seed is understood in the art to include removal of one or more of foreign debris such as weed seed, chaff, and non-seed plant matter from the seed. Conditioning the seed is understood in the art to include controlling the temperature and rate of dry down of the seed and storing the seed in a controlled temperature environment. Seed treatment is the application of a composition to the seed such as a coating or powder. Methods for producing a treated seed include the step of applying a composition to the seed or seed surface. Seeds are provided which have on the surface a composition. Biological active components such as bacteria can also be used as a seed treatment. Some examples of compositions include active components such as insecticides, fungicides, pesticides, antimicrobials, germination inhibitors, germination promoters, cytokinins, and nutrients. Biological active components, such as bacteria, can also be used as a seed treatment. Carriers such as polymers can be used to increase binding of the active component to the seed.

[0184] To protect and to enhance yield production and trait technologies, seed treatment options can provide additional crop plan flexibility and cost effective control against insects, weeds and diseases, thereby further enhancing the described inventions. Seed material can be treated, typically surface treated, with a composition comprising combinations of chemical or biological herbicides, herbicide safeners, insecticides, fungicides, germination inhibitors and enhancers, nutrients, plant growth regulators and activators, bactericides, nematocides, avicides and/or molluscicides. These compounds are typically formulated together with further carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. The coatings may be applied by impregnating propagation material with a liquid formulation or by coating with a combined wet or dry formulation.

[0185] Some seed treatments that may be used on crop seed include, but are not limited to, one or more of abscisic acid, acibenzolar-S-methyl, avermectin, amitrol, azaconazole, azospirillum, azadirachtin, azoxystrobin, *Bacillus* spp. (including one or more of *cereus*, *firmus*, *megaterium*, *pumilis*, *sphaericus*, *subtilis* and/or *thuringiensis*), *Bradyrhizobium* spp. (including one or more of *betae*, *canariense*, *elkanii*, *iriomotense*, *japonicum*, *liaonigense*, *pachyrhizi* and/or *yuanmingense*), captan, carboxin, chitosan, clothianidin, copper, cyazypyr, difenoconazole, etidiazole, fipronil, fludioxonil, fluoxastrobin, fluquinconazole, flurazole, fluxofenim, harpin protein, imazalil, imidacloprid, ipconazole, isoflavenoids, lipo-chitooligosaccharide, mancozeb, manganese, maneb, mefenoxam, metalaxyl, metconazole, myclobutanil, PCNB, penflufen, *penicillium*, pen thiopyrad, permethrine, picoxystrobin, prothioconazole, pyraclostrobin, rynaxypyr, S-metolachlor, saponin, sedaxane, TCMTB, tebuconazole, thiabendazole, thiamethoxam, thiocarb, thiram, tolclofos-methyl, triadimenol, *trichoderma*, trifloxystrobin, triticonazole and/or zinc. PCNB seed coat refers to EPA registration number 00293500419, containing quitozen and terrazole. TCMTB refers to 2-(thiocyanomethylthio) benzothiazole.

[0186] Seed varieties and seeds with specific transgenic traits may be tested to determine which seed treatment

options and application rates may complement such varieties and transgenic traits to enhance yield. For example, a variety with good yield potential but head smut susceptibility may benefit from the use of a seed treatment that provides protection against head smut, a variety with good yield potential but cyst nematode susceptibility may benefit from the use of a seed treatment that provides protection against cyst nematode, and so on. Likewise, a variety encompassing a transgenic trait conferring insect resistance may benefit from the second mode of action conferred by the seed treatment, a variety encompassing a transgenic trait conferring herbicide resistance may benefit from a seed treatment with a safener that enhances the plants resistance to that herbicide, etc. Further, the good root establishment and early emergence that results from the proper use of a seed treatment may result in more efficient nitrogen use, a better ability to withstand drought and an overall increase in yield potential of a variety or varieties containing a certain trait when combined with a seed treatment.

[0187] Another embodiment is a method of harvesting the grain or plant material of the F1 plant of variety ARC-9913C and using the grain or plant material in a commodity. Methods of producing a commodity plant product are also provided. Examples of maize grain or plant material as a commodity plant product include, but are not limited to, oils, meals, flour, starches, syrups, proteins, cellulose, silage, and sugars. Maize grain is used as human food, livestock feed, and as raw material in industry. The food uses of maize, in addition to human consumption of maize kernels, include both products of dry- and wet-milling industries. The principal products of maize dry milling are grits, meal and flour. The maize wet-milling industry can provide maize starch, maize syrups, and dextrose for food use. Maize oil is recovered from maize germ, which is a by-product of both dry- and wet-milling industries. Processing the grain can include one or more of cleaning to remove foreign material and debris from the grain, conditioning, such as addition of moisture to the grain, steeping the grain, wet milling, dry milling and sifting.

[0188] Maize, including both grain and non-grain portions of the plant, is also used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry.

[0189] Industrial uses of maize include production of ethanol, maize starch in the wet-milling industry and maize flour in the dry-milling industry. The industrial applications of maize starch and flour are based on functional properties, such as viscosity, film formation, adhesive properties, and ability to suspend particles. The maize starch and flour have application in the paper and textile industries. Other industrial uses include applications in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications.

[0190] Plant parts other than the grain of maize are also used in industry: for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

[0191] The seed of the maize variety, the plant produced from the seed, a plant produced from crossing of maize variety ARC-9913C and various parts of the maize plant and transgenic versions of the foregoing, can be utilized for human food, livestock feed, and as a raw material in industry.

[0192] All publications, patents, and patent applications mentioned in the specification are incorporated by reference

for the purpose cited to the same extent as if each was specifically and individually indicated to be incorporated by reference.

[0193] Applicant has made a deposit of at least 625 seeds of parental maize inbred varieties GIQA9942 and GIP19626 with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110-2209 USA, with ATCC Deposit Nos. PTA-_____ and PTA-_____, respectively. The seeds deposited with the ATCC on _____ for PTA-_____ and on _____ for PTA-_____, were obtained from the seed of the variety maintained by Inari Agriculture, Inc., 3400 Kent Avenue, West Lafayette, Indiana since before the filing date of this application. Access to this seed will be available during the pendency of the application to the Commissioner of Patents and Trademarks and persons determined by the Commissioner to be entitled thereto upon request. Upon issuance of any claims in the application, the Applicant will make available to the public, pursuant to 37 C.F.R. § 1.808, a sample(s) of the deposit of at least 625 seeds of parental maize inbred varieties GIQA9942 and GIP19626 with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110-2209. The deposits of the seed of parental maize inbred varieties for Hybrid Maize Variety ARC-9913C will be maintained in the ATCC depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the enforceable life of the patent, whichever is longer, and will be replaced if it becomes nonviable during that period. Additionally, Applicant has or will satisfy all of the requirements of 37 C.F.R. §§ 1.801-1.809, including providing an indication of the viability of the sample upon deposit. Applicant has no authority to waive any restrictions imposed by law on the transfer of biological material or its transportation in commerce. Applicant does not waive any infringement of the rights granted under this patent or rights applicable to Hybrid Maize Variety ARC-9913C and/or its parental maize inbred varieties GIQA9942 and GIP19626 under either the patent laws or the Plant Variety Protection Act (7 USC 2321 et seq.). Unauthorized seed multiplication is prohibited.

TABLE 1

Variety Description Information ARC-9913C		
Characteristic	Value	Number of Locations
Relative Maturity (RM)	113	—
Plant Density	28,863	26
Grain Yield (Yield)	176.9	26
Harvest Moisture (PCTHOH)	17.5	25
Test Weight (TWT)	60.8	25
Average Plant Height (PHT_AVE)	97.3	2
Average Ear Height (EHT_AVE)	45.4	2
Percent Total Late Root Lodging (TRTLP)	5.91	3
Percent Late Stalk Lodging (STLP)	8.15	5
Percent Greensnap (GSPPE)	4.8	3
Percent Early Root Lodging (ERTLP)	13.6	2
Fall Appearance (Fap)	4.4	7
Stay green (StyGrn)	5.2	6
Goss' Wilt (GW)	1	1
GDUs to Pollen Shed (GDU_P50)	1290	2
GDUs to Silking (GDU_S50)	1330	2
GDUs to Black Layer (GDU_R6)	2558	24

TABLE 1-continued

Variety Description Information ARC-9913C		
Characteristic	Value	Number of Locations
Kernel-Row Average (KRN_AVE)	17	9
Cob Color (Cob-COLA)	Deep red	1
Grain Quality (Qual_K)	1	1
Southern Rust (SR)	5	1
Silk color	Green	3

We claim:

1. A seed of hybrid maize variety ARC-9913C, representative seed produced by crossing a first plant of variety GIQA9942 with a second plant of variety GIP19626, wherein representative seed of the varieties GIQA9942 and GIP19626 have been deposited under ATCC Accession Numbers PTA-_____ and _____, respectively.

2. The seed of claim 1, that is free from a transgenic event.

3. The hybrid maize variety ARC-9913C seed of claim 1, further comprising a seed treatment on the surface of the seed.

4. A plant or plant part of hybrid maize variety ARC-9913C grown from the seed of claim 1, wherein the plant part comprises at least one cell of hybrid maize variety ARC-9913C.

5. A method of producing the seed of claim 1, the method comprising crossing a plant of variety GIQA9942 with a plant of variety GIP19626.

6. A method for producing nucleic acids, the method comprising isolating nucleic acids from the hybrid maize variety ARC-9913C seed of claim 5.

7. A method of producing a commodity plant product comprising starch, syrup, silage, fat or protein, the method comprising producing the commodity plant product from the plant or plant part of claim 4.

8. A method for producing a second maize plant, the method comprising applying plant breeding techniques to the plant or plant part of claim 4 to produce the second maize plant.

9. A method of producing a modified corn hybrid plant, where the method comprises genome editing the plant of claim 4.

10. A modified corn plant produced by the method of claim 9, where the plant comprises the genome edit and otherwise comprises all of the physiological and morphological characteristics of hybrid maize variety ARC-9913C.

11. A seed of hybrid maize variety ARC-9913C, representative seed produced by crossing a first plant of variety GIQA9942 with a second plant of variety 7SSDE1042, wherein representative seed of the varieties GIQA9942 and GIP19626 have been deposited under ATCC Accession Numbers PTA-_____ and PTA-_____, respectively, further comprising a transgene, wherein the transgene is introduced by backcrossing or genetic transformation into the variety GIQA9942, the variety GIP19626, or both varieties GIQA9942 and GIP19626.

12. A plant or plant part grown from the hybrid maize variety ARC-9913C seed of claim 11, the plant part comprising at least one cell of hybrid maize variety ARC-9913C further comprising the transgene.

13. A method for producing the hybrid maize variety ARC-9913C seed of claim 11, the method comprising cross-

ing a first plant of variety GIQA9942 with a second plant of variety GIP19626, representative seed of the varieties GIQA9942 and GIP19626 having been deposited under ATCC Accession Numbers PTA-_____ and PTA-_____, respectively, wherein at least one of the varieties GIQA9942 and GIP19626 further comprises the transgene.

14. The seed of claim 11, wherein the transgene confers a property selected from the group consisting of male sterility, herbicide tolerance, insect resistance, disease resistance, waxy starch, modified fatty acid metabolism, modified phytic acid metabolism, modified carbohydrate metabolism and modified protein metabolism.

15. A method for producing nucleic acids, the method comprising isolating nucleic acids from the seed of claim 11.

16. A seed of hybrid maize variety ARC-9913C further comprising a locus conversion, wherein a plant grown from the seed comprises a trait conferred by the locus conversion, and wherein the seed is produced by crossing a first plant of variety GIQA9942 with a second plant of variety GIP19626, wherein the first plant, the second plant or both further comprise the locus conversion, and wherein representative seed of the varieties GIQA9942 and GIP19626 have been deposited under ATCC Accession Numbers PTA-_____ and PTA-_____, respectively and wherein the seed produces a plant having otherwise all the physiological and morphological characteristics as maize variety ARC-9913C when grown under the same environmental conditions.

* * * * *