

MOLECULAR MECHANISM OF HEMOLYMPH CLOTTING SYSTEM IN *LIMULUS*

Sadaaki Iwanaga,¹ Toshiyuki Miyata,² Fuminori Tokunaga,³ and Tatsushi Muta⁴

Department of Biology, Faculty of Science, Kyushu University 33, Fukuoka 812, Japan

(Received 11.6.92 by Editor-in-Chief B. Blombäck)

ABSTRACT *Limulus* (horseshoe crab) hemolymph is known to be very sensitive to bacterial endotoxin (LPS), which causes a rapid coagulation response. Hemolymph contains a single type of hemocyte that undergoes aggregation, adhesion, and degranulation in response to LPS. The granule contents are released into the hemolymph, where they form an insoluble gel. We have characterized four components involved in this coagulation response that comprise a cascade of three serine protease zymogens (factor C, factor B, and proclotting enzyme) and one clottable protein (coagulogen). Of these components, factor C sensitive to LPS is a protein composed of five complement-related domains (“Sushi” or SCR), an EGF-like domain, and a C-type lectinlike domain as well as a putative amino-terminal LPS-binding domain. This domain structure is very similar to that of selectin family of cell adhesion molecules, suggesting that it might also function as a cell adhesion molecule after the release into the hemolymph. Factor B and the proclotting enzyme share a common Cys-rich motif (“cliplike” domain) in the amino-terminal portions. This domain is also found in a putative serine protease zymogen (“easter”) in *Drosophila*, which is essential for normal embryonic development. All four of the components of the cascade and an antibacterial protein (anti-LPS factor) are localized to a specific type of the hemocyte granule. Another antibacterial peptide (tachyplesins I and II) is localized in a distinct granule population. The contents of both granule populations are released into the hemolymph in response to LPS, where they cooperate in immobilization and killing of Gram-negative bacteria.

INTRODUCTION

Invertebrates have characteristic host defense systems that differ from mammalian immune systems (1–6). In *Limulus* (horseshoe crab), an *Arthropoda* species, this defense system is carried by the hemolymph, which contains a type of cell called an amebocyte. The amebocytes

¹To whom correspondence should be addressed.

²Present address: Laboratory of Thrombosis Research, National Cardiovascular Center Research Institute, Fujishirodai 5, Suita-565, Japan.

³Present address: Department of Life Science, Faculty of Science, Himeji Institute of Technology, Harima Science Park City, Kamigori, Hyogo 678-12, Japan.

⁴Present address: Cardiovascular Biology Research Program, Oklahoma Medical Research Foundation, 825 N.E. 13th Street, Oklahoma City, Oklahoma 73104-5073, USA.

are extremely sensitive to endotoxin (LPS), which is a major component of the cell surface of Gram-negative bacteria (7-11). When hemolymph comes into contact with Gram-negative bacteria or LPS, the amebocytes begin to degranulate, and hemolymph coagulation is initiated by the granule components (12-17). This response is thought to be very important for limulus not only in preventing the leakage of hemolymph, but also in engulfing invading microorganisms (18-20).

During the past decade, we have studied the molecular mechanism of hemolymph coagulation in limulus and have established a protease cascade, as shown in Figure 1. This cascade is based on three kinds of serine protease zymogens, factor C, factor B, proclotting enzyme, and one clottable protein, coagulogen. LPS activates the zymogen factor C to the active factor $\bar{C}$. Factor $\bar{C}$ then activates factor B to factor $\bar{B}$, which in turn converts the proclotting enzyme to the clotting enzyme. Each activation proceeds by limited proteolysis. The resulting clotting enzyme cleaves two bonds in coagulogen, which is a fibrinogen-like molecule in arthropods, to yield an insoluble coagulin gel. The coagulation cascade is also activated by (1,3)- β -D-glucan, a serine protease that is tentatively named factor G is activated, leading to activation of the proclotting enzyme.

In this review, we will focus on the molecular structures and functions of limulus clotting factors and antimicrobial substances so far studied in our laboratory.

Limulus Hemocytes

The limulus hemolymph contains two types of cells, granular hemocytes and cyanocytes (12,14-17,20). Based on cell morphology, there appears to be only one type of hemocyte in the systemic circulation of the adult intermolt animal, the so-called granulocyte or amebocyte (4). This cell is an oval, plate-shaped structure, 15-20 μm in its longest dimension. Figure 2 shows an electron micrograph of the hemocyte separated from Japanese horseshoe crab, *Tachypleus (T) tridentatus* (20). The cell contains numerous dense granules classed into two major types: large (L) and small (S) granules. The L-granules are larger (up to 1.5 μm in diameter) and less electron dense than the S-granules (<0.6 μm in diameter). The L-granules contain at least 20 protein components including, 4 clotting factors and 1 antimicrobial factor (anti-LPS factor), whereas the S-granules exclusively contain the other antimicrobial substance, named tachyplesin peptide, in addition to 6 major protein components. After




FIG. 1.

Hemolymph coagulation cascade system in *Limulus (Tachypleus tridentatus)*




FIG. 2.

A. A scanning electron micrograph of hemocytes; **B.** Granular hemocytes (G) and non-granular hemocytes (NG); **C.** A cross-section of a granular hemocyte showing the distribution of various organelles. C. golgi vacuole with condensed dense materials; CT. centriole; D. dense (small) granules; G. golgi complex; L. large granules; P. golgi vacuole with dispersed dense materials (P-granule). Published by permission of Springer-Verlag, Heidelberg. Original in Ref. 20.

treatment with endotoxin or LPS, the L-granules are released more rapidly than the S-granules, although almost all granules are finally exocytosed (20). This exocytosis is associated with clot formation, the process being complete within 90 seconds. The clot is more soft than mammalian fibrin clot and contains coagulin gel generated from its precursor, named coagulogen.

Coagulogen

The clottable protein, coagulogen, has a functional similarity with vertebrate fibrinogen (21–23) and is known to play a central role in the limulus clotting system (Fig. 1) (24–29). In the




FIG. 3.

The gross structure of *Limulus* coagulogen. The arrows indicate sites cleaved by *Limulus* clotting enzyme in the conversion of coagulogen to coagulin gel.

intact animal, soluble precursor of this protein is absent from the plasma, but is instead sequestered in the cytoplasmic L-granules of the hemocytes (14,20). This granule exocytosis releases coagulogen into the external milieu (12,15). The extracellular clot is composed of fibers of the protein, named coagulin (11).

Figure 3 shows a gross structure of coagulogen and its structural change by limulus clotting enzyme (30–33). Coagulogen is an invertebrate fibrinogen-like substance, and it is converted from a soluble form to an insoluble gel, coagulin. The conversion to a gel is mediated by a clotting enzyme in the hemocytes (25,26). Coagulogen consists of a single basic polypeptide chain and has a calculated molecular weight of 19,700 (± 50). This protein contains three segments, A chain, peptide C, and B chain (34). The clotting enzyme cleaves the Arg-Gly and Arg-Thr linkages, both located at the NH₂-terminal region (24,25). The Arg-Gly linkage cleaved by the clotting enzyme is the same type as that cleaved by α -thrombin in the transformation of mammalian fibrinogen to fibrin. Moreover, the COOH-terminal octapeptide sequences of A chain and peptide C exhibit great homology to one another, and their sequences are very similar to that of primate fibrinopeptide B (25,32). The amino acid sequences of four coagulogens isolated from the hemocytes of American (*Limulus (L) polyphemus*), Japanese (*T. tridentatus*), and two Southeast Asian (*T. gigas* and *Carcinoscorpius (C) rotundicauda*) horseshoe crabs are shown in Fig. 4 (34–40). Upon gelation of all the coagulogens by a limulus clotting enzyme, a large peptide, named peptide C, is released in common from the inner portion of the parent molecules. The resulting gel consists of two chains of A (18 residues) and B (129 residues), bridged by two disulfide linkages. The overall sequences of the four coagulogens have considerable homology, particularly in the A and B chain regions, consisting of 147 residues (40). Surprisingly, 73%–92% of the sequences in these regions contain the same amino acid residues in identical positions, and homologies are found along the entire molecules. In addition, 16 half-cystine residues of the four are in the same positions in the sequences, suggesting that they are paired in the same fashion and, thus, that the molecules have very similar conformations. In contrast, the sequence homology of the peptide C region is less than that of the A and B chain regions. The homologies amount to approximately 43%–57% of the structure (41,42). It is of interest that coagulogen appears to contain an α -helical (14.9%) region exclusively around the peptide C segment released by the clotting enzyme (39), suggesting that there is a marked conformational change in the transformation of coagulo-




FIG. 4.

Amino acid sequence identities among Asian (*T.t.*, *T.g.*, *C.r.*) and American (*L.p.*) horseshoe crabs. Chemically identical residues in the sequences are framed. The bonds cleaved by horseshoe crab clotting enzyme are indicated by arrows. *L.p.*, *Limulus polyphemus*; *T.t.*, *Tachypileus tridentatus*; *T.g.*, *Tachypileus gigas*; *C.r.*, *Carcinoscorpius rotundicauda*. Published by permission of the Japanese Biochemical Society, Tokyo. Original in Ref. 39.

gen to coagulin gel. On the other hand, the β -sheet (24.6%), reverse-turn (29.7%), and coiled (30.9%) structure are commonly distributed in the B chain segment, suggesting that it is a β -type protein (39).

We also identified a cloned DNA fragment for coagulogen from a cDNA library of the limulus hemocyte mRNA (37). Figure 5 shows the nucleotide sequences of two pre-coagulogen cDNA clones and their predicted amino acid sequences. The cDNA of 946 bp encoded for pre-coagulogen II is found to code for a 20-bp noncoding region at the 5' end, a leader sequence of 20 amino acids, followed by the entire coding region of coagulogen, and a 341-bp noncoding region at the 3' end. The predicted amino acid sequence of this clone is in good agreement with the sequence determined by the protein-based study, except for the residue at positions 82 and 86, the sequence of which is identical to that found in coagulogen isolated from *T. gigas*. On the other hand, the sequence of cDNA encoded for pre-coagulogen I agreed completely with that previously established (38), indicating that the hemocyte coagulogens are encoded by two very similar but distinct mRNAs.




FIG. 5.

Nucleotide sequences of two pre-coagulogen cDNA clones, p4i and p3C21, and their predicted amino acid sequences. The numbers above and below the sequences show the nucleotide positions and the amino acid positions respectively. The positive numbers of the amino acids start from the NH₂-terminus of coagulogen. The putative poly (A) additional signal, AATAAA, is underlined. The clone p4i was 811 bp in length starting from nucleotide position 137 encoded for the amino acid residues 20-175 in coagulogen type I. The clone p3C21 was 946 bp in length starting from nucleotide position 1 encoded for coagulogen type II, (G)_n and (A)_n are the oligo(G) and oligo(A) residues; oc is the ochre translation stop codon TAA. Published by permission of the Japanese Biochemical Society, Tokyo. Original in Ref. 37.




FIG. 6.

Phylogenetic relationship of living horseshoe crab species based on mutation distances calculated from the amino acid sequences of coagulogens (6). Published by permission of the Japanese Biochemical Society, Tokyo. Original in Ref. 39.

As well known, protein sequences have been used to estimate the evolutionary rate and phylogenetic relationships among different species at the molecular level (43,44). Based on the data on amino acid differences and minimum base substitutions among the sequences of coagulogens from the four species, the mutation distance between Limulinae and Tachypleinae is approximately 67.3; that between *T. gigas* and the other Asian species is 25 and that between *T. tridentatus* and *C. rotundicauda* is 20. A cladogram (based on the mutation distances among coagulogen sequences) indicating the phylogeny for the living horseshoe crab species is illustrated in Figure 6. These data agree with previous reports (32) that *T. tridentatus* and *C. rotundicauda* are phylogenetically more closely related than the other species of horseshoe crab, and the fact that *Limulus* is evolutionarily distant from the other three species is supported by the results of immunochemical studies on coagulogen (33) and sialic acid binding lectin (32). According to Shuster (43), divergence from the common ancestor of Limulinae and Tachypleinae occurred in the late Jurassic or early Cretaceous period, corresponding to 110–160 million years ago. The calculated divergence time between *T. gigas* and the other Asian species is 52.5 million years and that between *T. tridentatus* and *C. rotundicauda* is 36.3 million years. These values differ from those suggested by Shuster (44) and also from those estimated previously based on the amino acid sequences of peptide C (residues 19–46) of coagulogen (32). The approximate mutation rate per amino acid site in coagulogen per year (R) can be calculated by the method of Kimura using the following equation; $R = A / 13.5 \times 10^7 \times B \times 2$. A is the average number of amino acid replacements separating Limulinae and Tachypleinae, and B is the number of amino acid residues in coagulogen. The value of 13.5×10^7 is the average time in years from the late Jurassic and early Cretaceous periods to the present. We can calculate A and B as 54.7 and 175, respectively, and, thus, R for coagulogen is 1.16×10^{-9} . This value is approximately a half of that obtained for peptide C, 2.03×10^{-9} (39,42).

Limulus Clotting Factors

Proclotting Enzyme. *Limulus* proclotting enzyme is activated by both factor B and factor G (Fig. 1) and it is a single chain glycoprotein with an apparent molecular mass of 54 kDa. Upon activation by factor B (Fig. 1), it is converted to a two chain active form, clotting enzyme

(31,49,50), composed of a light (L, 25 kDa) and a heavy (H, 31 kDa) chains (48,52). The clotting enzyme cleaves two bonds in coagulogen to yield a coagulin gel. Thus, proclotting enzyme/clotting enzyme seems to be a prothrombin/ α -thrombin counterpart to invertebrates, because both catalyze a final step for gelation. Interestingly, however, the clotting enzyme has a substrate specificity similar to that of mammalian coagulation factor Xa. It catalyzes the conversion of bovine prothrombin to α -thrombin at a reasonable rate (48). As the activity of clotting enzyme is inhibited by several serine protease inhibitors (45–48), this protein is thought to be a serine protease zymogen (51), and it was proved, afterwards, by its primary structure (53).

Figure 7 shows cDNA and the entire amino acid sequences of proclotting enzyme (53). The sequence from the first ATG codon shows characteristics of a signal sequence. The 29 amino acid residues (amino acid –29 to –1) preceding the amino-terminal glutamine residue of mature protein seems to constitute a prepro-sequence. The hydropathy profile calculated by the method of Kyte and Doolittle indicates that the presequence region is highly hydrophobic, suggesting that this region acts as a signal peptide. The sequence linked between the mature protein and the signal peptide is assumed to consist of 12 (or 5) residues and appears to represent a propeptide region of the immature protein. In fact, the carboxyl-terminal sequence of this region shows a sequence, -Arg-X-Arg-Arg (amino acid –4 to –1) similar to those found in serine protease zymogens associated in mammalian coagulation and complement systems (Fig. 8). The presence of this sequence strongly suggests that an enzyme, which proteolytically processes the dibasic cleavage site, might also exist in the invertebrate hemocyte. The propeptide region might function so as to prevent an unexpected activation of the proclotting enzyme, the result being clot formation within the cell.

The mature protein is composed of 346 amino acid residues with a molecular mass of 38,194 (Fig. 7). The amino-terminal residue of the intact proclotting enzyme is masked by a pyroglutamyl residue. All three potential glycosylation sites, one in the L chain and two in the H chain, for *N*-linked carbohydrate chains (Asn-X-Ser/Thr) are actually glycosylated. Furthermore, at least six *O*-linked carbohydrate chains are present in the L chain. The cleavage site associated with the zymogen activation is Arg₉₈-Ile₉₉, the sequence of which connects the L and H chains. The sequence preceding this site has a unique structure, -Thr-Thr-Thr-Thr-Arg. This sequence agrees well with the substrate specificity of the limulus factor B̄ (54). The H chain of the proclotting enzyme is composed of a typical structure of serine protease. The locations of four disulfide linkages in the H chain are the same as those in prothrombin and factor IX. As previously mentioned, clotting enzyme shows a substrate specificity similar to that of factor Xa. In fact, the entire sequence of the H chain corresponding to the serine protease region shows a close similarity with that of human factor Xa (34.1%). It contains also the His₁₄₃-Asp₁₉₉-Ser₂₉₇ triad known to be the catalytic triad of serine proteases. The substrate binding site corresponding to Ser₁₈₉ in chymotrypsin numbering is aspartic acid, thereby indicating that this clotting enzyme has a typical trypsinlike specificity.

The amino-terminal portion of the L chain consists of a novel disulfide-knotted structure. This structure forms a “cliplike” shape consisting of a discrete domain. Following this domain, the serine and threonine-rich sequence are highly glycosylated with *O*-linked carbohydrate chains. After we published the entire sequence of proclotting enzyme, Dr. A. Day (Dept. of Biochemistry, Univ. of Oxford) informed us that the proclotting enzyme has a sequence similarity in its amino-terminal light chain portion with *Drosophila* protein, named serine protease easter precursor. Figure 9 shows a similarity of parts of these proteins, indicating that all cysteine residues are aligned. Although the significance is still unknown, these results may

	GCA	3
ATACAGGCTACAAACATCTCATCAGACGCATTACCTGGTTGTTAATTCCGCGAAAGCTTCACGAGAACAAGTCA		78
AACCTAGCTTGTGGTCCCGGACTGACTGCCTGGAAAGTAGTTCCTCAAAATTGCGGCTTTATAAATCAACCTAAA		153
CAAAACCGTCTGAACTTTACCGTTCACATCCACCGAGGTAGCCGGTCTGCTCTCAGCAGTATGTTGGTGAAT		228
	M L V N	-26
AACGTGTTTTCTACTGTGTTTCCACTCTTGATGTCTGTGGTTAGATGCAGTACTCTCAGCAGACGCGTAGA		303
N V F S L L C F P L L H S V V R C S T L S R Q R R		-1
CAGTTTGTTCCTGACGAGGAAGAACTTTGCTCAAACCGATTTACTGAAGAAGGAACATGCAAAATGTCTTG		378
Q F V F P D E E E L C S N R F T E E G T C K N V L		25
I		
GATTGTAGAATACTTTTACAAAAAATGATTATAATTTACTCAAAGAATCAATATGCGGCTTTGAAGGCATAACA		453
D C R I L L Q K N D Y N L L K E S I C G F E G I T		50
CCCAAAGTTTGTGTCGGAATCAAGCCATGTAAATTTCAAGTACACAGGCACCTCCAGAAACCACTACGACTGAA		528
P K V C C P K S S H V I S S T Q A P P E T T T T E		75
CGCCCAACCAACAGATCAACCCCAATCTTCTGAAGTGTGTGGAATTCACAATACCACTACCGATTATT		603
R P P K Q I P P N L P E V C G I H N T T T T R I I		100
GGAGTTCGGAAGCACCTATTGGAGCCTGGCGTGGATGACTGCTGTCTACATAAAACAGGAGGAATCAGAAGT		678
G G R E A P I G A W P W M T A V Y I K Q G G I R S		125
GTTCAGTGTGGTGGCGCACTTGTCTACTAACAGGCACGTGATTACAGCTTCGCACTGTGTGTAAACAGTGCAGGA		753
V Q C G G A L V T N R H V I T A S (II) C V V N S A G		150
ACAGATGTGATGCCAGCTGATGATTCTCGGTTCTGCTGGGTGAACACAATTTATACAGTACCGATGACGATTTCG		828
T D V H P A D V F S V R L G E H N L Y S T D D D S		175
AATCCAATAGATTTTGCAGTTACGTGGTGAACATCAGCAACACTTTGTACTCGGACGTATTTGAATGACATC		903
N P I D F A V T S V K H H E H F V L A T Y L N (D) I		200
GCAATTTCTAACGTTAAATGACACAGTTACGTTTACAGACAGAATTCGACCCATTGTCTACCTTATCGTAAGTTG		978
A I L T L N D T V T F T D R I R P I C L P Y R K L		225
AGATACGATGATCTAGCAATGAGAAACCGTTTATCACTGGATGGGGAACACAGCATTTAAGCGCCCATCTAGT		1053
R Y D D L A H R K P F I T G W G T T A F N G P S S		250
GCAGTGTTCAGAGAAGTACAGTTACCAATATGGGAACACGAGGCTGTAGACAGCCCTACGAGAAGGATTTAAAT		1128
A V L R E V Q L P I W E H E A C R Q A Y E K D L N		275
ATTACAAACGTGTATATGTGTCTGGCTTTCAGATGGCGGAAGGATGCTTGCAGGGTGATTTCTGGAGGTCCA		1203
I T N V Y M C A G F A D G G K D A C Q G D (S) G G P		300
ATGATGTTGCCTGTTAAAAACCGAGAGTTTTATCTCATTGGAATTGTGTCTTTCCGAAAGAAATGCGCATTCGCT		1278
H M L P V K T G E F Y L I G I V S F G K K C A L P		325
GGATTTCCTGGGTTTACAAAAAGTGACAGAGTTTTAGATTGGAATTCAGAACATATGGTGTAGACTAAAGCT		1353
G F P G V Y T K V T E F L D M J A E H M V *		346
GTGAATGACTCCATTTCGAATATTAAACCATACTCCGCTGTAGCAAAACGACTGAAGAATCAAGCAGGGGACAA		1428
AAACCACTTTTTTTTTTTTTTTCTTCAATAGTTCTACCATGTATTGAAAAACCAACATAAGACAATCAATATA		1501

FIG. 7.

Nucleotide and deduced amino acid sequences of proclotting enzyme. Nucleotide (*upper*) and deduced amino acid sequences (*lower*) are shown. The amino acid sequences are numbered from the amino-terminus of the mature protein. The *underlined* amino acid residues were confirmed by protein sequencing. The cleavage sites accompanied by maturation and activation are shown by *arrowheads*. The potential attachment sites for *N*-linked carbohydrate chains are indicated by *closed diamonds*. The active triads of the serine protease domain are *circled*. Published by permission of the American Society for Biochemistry and Molecular Biology, Inc. (ASBMB). Original in Ref. 53.




FIG. 8.

Sequence comparisons of processing sites of limulus immature clotting factors with those of immature mammalian vitamin K dependent clotting factors and others. An arrow indicates the sites cleaved by a microsomal processing enzyme (105).

suggest that the “cliplike” domain each found in limulus proclotting enzyme and factor B is one of the common structural elements in the serine protease zymogen of invertebrate animals.

An immunohistochemical study utilizing anticlotting enzyme antibody demonstrates that the proclotting enzyme is present in the L-granules in hemocytes (20).

Clotting Factor B. Factor B is activated by factor C and then, in turn, activates proclotting enzyme (Fig. 1) (55,56). Factor B is a single-chain glycoprotein with a molecular mass of 64,000 (nonreduced SDS-PAGE). The purified factor B, however, gives three bands with apparent molecular masses of 64,000, 40,000 (heavy chain), and 25,000 (light chain) after reduction (54). Therefore, the purified preparation contains two molecular species; single-chain factor B and two-chain factor B bridged by a disulfide linkage. The two-chain species appears




FIG. 9.

Sequence similarity of “Clip-like” domain of limulus proclotting enzyme and factor B with that of serine protease zymogen easter from *Drosophila* (106).

most likely to be a factor B zymogen cleaved in a disulfide loop that links the heavy and the light chains. Upon activation of the zymogen factor B by active factor $\bar{C}$, it is converted into factor $\bar{B}$ with the heavy ($M_r = 32,000$) and light chains ($M_r = 25,000$), releasing an activation peptide. When the purified factor $\bar{B}$ is treated with [^3H]DFP, one major peak with radioactivity before reduction is found at the position corresponding to the major protein band of factor $\bar{B}$. On reduction, the major radioactive peak migrates to the position corresponding to the mobility of the heavy chain ($M_r = 32,000$) derived from factor $\bar{B}$, indicating that the DFP-reactive site is located in the heavy chain portion of factor $\bar{B}$.

Recently, we succeeded to determine the cDNA sequence for zymogen factor B (57). It consists of 400 amino acid residues with 23 residues of signal sequence. The mature protein is composed of 377 amino acids with a calculated molecular mass of 40,570. Three potential glycosylation sites for *N*-linked carbohydrate chain are found in the mature polypeptide sequence. The entire amino acid sequence has a similarity with that of limulus proclotting enzyme. Particularly, the sequence identity of the carboxyl-terminal serine protease domains between factor B and proclotting enzyme is 43.9%. Moreover, the amino-terminal regions up to 60th residue of both proteins share a similar structure with six cysteines (Fig. 9), suggesting that these proteins are arisen from a gene duplication.

Factor $\bar{B}$ most efficiently hydrolyzes Boc-Met-Thr-Arg-4-methylcoumaryl-7-amide-(MCA) and shows a relatively weak activity towards Boc-Leu-Gly-Arg-MCA, which is the best substrate for limulus clotting enzyme (54). The good substrate (Boc-Val-Pro-Arg-MCA) for factor $\bar{C}$ is also weakly hydrolyzed by factor $\bar{B}$. Factor $\bar{B}$ shows an apparent affinity for hydroxyl amino acid (Thr or Ser) at the P_2 site (nomenclature of Schechter and Berger). As regards to natural substrates, it does not activate any mammalian plasma coagulation factors, such as factor IX, factor X, prothrombin, plasminogen, protein C, and prekallikrein. It also does not convert fibrinogen to fibrin gel. Moreover, factor $\bar{B}$ does not catalyze the activation of the zymogen factor C or the conversion of coagulogen to coagulin. The amidase activity of factor $\bar{B}$ is strongly inhibited by α_2 -plasmin inhibitor, but is not inhibited by antithrombin III, which is a potent inhibitor of factor $\bar{C}$ and limulus clotting enzyme. Among four synthetic inhibitors, DFP has the strongest inhibitory effect. Benzamidine, leupeptin, and PCMB partially inhibit the factor $\bar{B}$ activity (54).

Clotting Factor C. As shown in Fig. 1, factor C is an initial activator of the clotting cascade triggered by LPS. Factor C has a molecular weight (M_r) of 123,000 glycoprotein consisting of a heavy chain of $M_r = 80,000$ and a light chain of $M_r = 43,000$ (58–60). Factor C is converted to an activated form in the presence of LPS with a $M_r = 123,000$, designated factor $\bar{C}$ (62–64). Upon activation, a single cleavage of Phe-Ile bond in the light chain occurs, resulting in the accumulation of two new fragments of $M_r = 34,000$ (B chain) and 8,500 (A chain) (62). A diisopropylfluorophosphate-sensitive active site is localized in the B chain of factor $\bar{C}$. We determined the entire amino acid sequence of factor C using recombinant DNA technique (65). The zymogen consists of 994 residues with a calculated molecular mass of 109,648 kDa (Fig. 10). In the H and A chains, there are several interesting amino acid sequences. First, factor C has five repeating units ("Sushi domain or short consensus repeat) of about 60 amino acid residues each, which are found in many proteins participating in the mammalian complement system. Factor C is the first example of a protein that has the complement-like structure in invertebrates. Second, an epidermal growth factor (EGF)-like domain is present in the H chain. This EGF-like domain is most homologous with that of laminin B₂ chain. Proteins containing this type of domain involve several blood coagulation factors, complement factors,

Seikagaku Corporation Exhibit 2067 - Page 12 of 32
Charles River Laboratories, Inc. v. Seikagaku Corporation
PGR2025-00023

some receptors including low density lipoprotein (LDL) receptor. Third, a lectinlike domain is unexpectedly found in between the third and fourth "Sushi" domains. This region is composed of about 120 amino acids and shows homology with so-called C-type lectins (65). This type of domain is also found in proteins whose functions are not considered only to bind the carbohydrate moiety. Any sugar binding abilities in factor C have not been demonstrated, thus far.

Two other characteristic sequences are also found in the H chain (Fig. 12). One is a Cys-rich region, which is located in the amino-terminal portion of the H chain. Eleven residues of cysteines out of 56 residues are clustered in this region (Fig. 10). The other is in the carboxyl-terminal portion of the H chain, in which 13 out of 48 residues are proline. This region has homology with the connecting region of mammalian coagulation factor XII (65). Its significance, however, remains unknown.

The B chain portion of factor C is composed of a typical serine protease, with an active site triad. The B chain shows the strongest homology with human α -thrombin (36.7%). This is consistent with the result that factor C shows a similar specificity toward synthetic fluorogenic peptide substrates comparable with α -thrombin (58). The substrate binding site of aspartic acid is conserved, which indicates that factor C is a trypsin type enzyme, although cleavage site associated with the zymogen activation is the Phe-Ile (60). The proline residue at the P₂ site might be important for the cleavage. In fact, the peptide bond cleaved during conversion to the two-chain form, Arg-Ser, is also preceded by a proline residue. The mosaic structure of factor C is summarized in Fig. 12. Immunohistochemical observation with electron microscopy utilizing anti-factor C antibody demonstrates that factor C is localized in the L-granules in the cell (20).

To our knowledge, proteases with Sushi domains are only mammalian complement C1r, C1s, C2, factor B, and limulus factor C, as shown in Fig. 11. Moreover, proteases that have both Sushi and EGF domains include only C1r, C1s, and factor C, all of which are associated with the initiation of cascade reaction. "Selectins" including three membrane proteins, endothelial leukocyte adhesion molecule, lymph node homing receptor, and granule membrane protein 140, have been reported to contain a lectinlike, an EGF-like domain and several "Sushi" domains (Fig. 12). The similarity in domain composition between factor C and those proteins is of interest, because these three proteins all seem to be associated with biologic defense systems. Another example that contain all these domains is a proteoglycan core protein

FIG. 10.

The composite nucleotide sequence (*upper*) and the deduced amino acid sequence (*lower*) of factor C. A polyadenylation signal (AATAAA) is *double-underlined*. The amino acid sequence is *numbered* beginning at the amino terminus of the mature factor C molecule. The amino acid residues that have been confirmed by sequencing of purified peptides are *underlined* and include data from Tokunaga *et al.* (60). Amino-terminal amino acid sequencing of each chain identified the three cleavage sites marked by *arrow-heads*. The potential carbohydrate attachment sites for (Asn-Xaa-Ser/Thr) are indicated with *closed diamonds* and the putative attachment site confirmed not to carry carbohydrate is shown as an *open diamond*. The amino acid residues suspected to form the catalytic triad of the serine protease domain are *circled*. Published by permission of the American Society for Biochemistry and Molecular Biology, Inc. (ASBMB). Original in Ref. 65.




FIG. 11.

Comparison of the structures of serine proteases containing Sushi domains. The domain structures of serine proteases with Sushi domains including limulus factor C, C1r, C1s, C2, and complement factor B are illustrated. Published by permission of the American Society for Biochemistry and Molecular Biology, Inc. (ASBMB). Original in Ref. 65.





FIG. 12.

The domain structure of limulus zymogen factor C sensitive to LPS. Factor C has a mosaic structure of several domains including a signal peptide (-25 to -1), one Cys-rich region (1-76), one EGF-like domain (77-111), five Sushi domains (117-170, 174-229, 235-296, 551-609, and 618-665, located within the fifth Sushi domain) and one serine protease domain (738-994). The H, A, and B chain correspond to each domain as illustrated. These domain structures are compared with those of the so-called "selectin" family. Published by permission of the American Society for Biochemistry and Molecular Biology, Inc. (ASBMB). Original in Ref. 65.

that is present extracellular matrix (65). On the other hand, the fact that factor C is structurally related to mammalian complement factors indicates evolutionary relationship between two systems. Factor C is a newly discovered type of serine protease zymogen, a "coagulation-complement factor," which may play an important role both in hemostasis and defense mechanisms. In considering the structural similarity of initiators of the two systems, the limulus coagulation and mammalian complement systems may have evolved from a common origin. Concerning the evolution of host defense systems in the animal kingdom, this is consistent with a possible common evolutionary origin of the coagulation and complement cascades.

The zymogen factor C is rapidly activated not only by bacterial endotoxins (58,61) but also by acylated (β 1-6)-D-glucosamine disaccharide bisphosphate (synthetic *Escherichia coli*-type lipid A), and the corresponding 4'-monophosphate analogues, as shown in Tables I and II (63,64). However, the corresponding nonphosphorylated lipid A does not activate factor C, indicating that a phosphate ester group linked with the (β 1-6)-D-glucosamine disaccharide back bone is required for the zymogen activation. In addition, the zymogen factor C is significantly activated by acidic phospholipids, such as phosphatidylinositol, phosphatidylglycerol, and cardiolipin, but not at all by neutral phospholipids (Table III). The rate of this activation, however, is affected markedly by ionic strength in the reaction mixture, although such an effect is not observed in the lipid-A-mediated activation of factor C. A variety of negatively charged surfaces, such as sulfatide, dextran sulfate, and ellagic acid, which are known

TABLE I
Structure of chemically synthesized lipid A analogues. Published by permission of Elsevier Science Publishers BV, Amsterdam, Holland. Original in Ref. 63.



The chemical structure shows a disaccharide backbone consisting of a glucose unit (left) and a glucosamine unit (right) linked by a β 1-6 glycosidic bond. The glucose unit has a hydroxyl group at C2 and a phosphate group at C6 (R2O). The glucosamine unit has a hydroxyl group at C2 and a phosphate group at C6 (OR1). Both units have fatty acid chains at C1 and C3. The substituents R1, R2, R3, and R4 are defined in the table below.

Number	Substituent			
	R ₁	R ₂	R ₃	R ₄
403	-H	-H		
404	-H	-PO(OH) ₂		
405	-PO(OH) ₂	-H		
406	-PO(OH) ₂	-PO(OH) ₂		
503	-H	-H		
504	-H	-PO(OH) ₂		
505	-PO(OH) ₂	-H	-CO(CH ₂) ₁₀ CH ₃	-CO(CH ₂) ₁₂ CH ₃
506	-PO(OH) ₂	-PO(OH) ₂		

TABLE II

Activation of factor C by lipid A analogues. Synthetic lipid A analogues were dissolved in 0.025% triethylamine (1 mg/ml), sonicated at room temperature for 10 min. and diluted with 0.05 M Tris/HCl (pH 8.0) containing 0.5 mg/ml human serum albumin. As the compounds 403 and 503 were not soluble in the above buffer, their sonicated emulsions were used alone for assay. The activation of factor C (7.7 nM) at various concentrations of lipid A analogues were examined, and the minimal concentration of each lipid A analogue for 50% activation of factor C (ED_{50}) was estimated. Published by permission of Elsevier Science Publishers BV, Amsterdam, Holland. Original in Ref. 63.

Compound	Activation	Concentration of lipid A to activate 50% of factor C (ED_{50})
		nM
403	—	
404	+	16
405	+	143
406	+	6
503	—	
504	+	66
505	+	220
506	+	55

as typical initiators for activation of the mammalian intrinsic clotting system, do not show any effect on the zymogen factor C activation. These results suggest that lipid A is the most effective trigger to initiate the activation of the horseshoe crab hemolymph clotting system. In addition to nonenzymatic activators described previously, the zymogen factor C is rapidly ac-

TABLE III

Activation of factor C by phospholipids. Phospholipids were dissolved in water (1 mg/ml), sonicated at room temperature for 10 min and diluted with 0.05 M Tris/HCl (pH 8.0) containing 0.5 mg/ml human serum albumin. Then the activation of factor C (7.7 nM) at various concentrations of phospholipids was examined and the minimal concentration of each phospholipid for 50% activation of factor C was estimated, as shown in Table I. Published by permission of Elsevier Science Publishers BV, Amsterdam, Holland. Original in Ref. 63.

Phospholipids	Activation	Concentration of phospholipid to activate 50% of factor C (ED_{50})
		nM
Phosphatidic acid	+	>10000
Phosphatidylglycerol	+	22
Phosphatidylserine	+	45
Phosphatidylinositol	+	8
Phosphatidylethanolamine	—	
Phosphatidylcholine	—	
Diphosphatidylglycerol (cardiolipin)	+	24

tivated by α -chymotrypsin and rat mast cell chymase (63,66), but not by trypsin, which activates many serine protease zymogens associated with mammalian coagulation, fibrinolysis, and complement systems. Thus, limulus factor C seems to be a novel type of serine protease zymogen susceptible to α -chymotrypsin. The amidase activity of factor C toward Boc-Val-Pro-Arg-MCA is strongly inhibited by antithrombin III, but not by α_2 -antiplasmin. Soybean trypsin inhibitor and hirudin has no inhibitory effect on factor C activity. DFP and D-Phe-Pro-Arg-chloromethyl ketone have a potent inhibitory effect. Benzamidine and leupeptin also have inhibitory activity at high concentrations (5 mM and 0.23 mM, respectively), while *p*-chloromercuribenzoate has no apparent inhibitory effect (58).

Figure 13 summarizes the gross structures of limulus clotting factors so far described. Like mammalian clotting factors, factor C, factor B and proclotting enzyme are typical ser-




FIG. 13.

The domain structures of limulus clotting factors and antimicrobial substances. The arrowheads indicate the cleavage sites for zymogen activation. The potential carbohydrate attachment sites are indicated with closed diamonds.

is closely associated with its biologic function. In charge distribution along the polypeptide chains, the region in the disulfide loop is highly basic. Especially, the region from arginine-41 to lysine-49 has basic amino acids at every second residues, and the region from arginine-61 to arginine-76 is also rich in basic residues, in which they have positive charges almost at every third residues. Hydropathic profile indicates that amino-terminal region of ALF is highly hydrophobic, which render this molecule amphipathic (70).

ALF inhibits the LPS-mediated activation of factor C (69). Because the activation of factor C induced by synthetic lipid A is also inhibited by ALF, it appears to interact with lipid A portion of LPS. ALF also shows the hemolytic activity against red blood cells (RBC) sensitized with LPS (78). Not only human RBC but also horse and chicken RBC are susceptible to the hemolysis. The minimal concentration of ALF required for hemolysis is dependent on the concentration of LPS used for the sensitization. Under optimal conditions, the minimal hemolytic concentration of ALF is 0.48 $\mu\text{g/ml}$. The hemolysis mediated with ALF completes within 1 minute at 37°C, and even at 0°C this reaction completes within 5 minutes. The LPS sensitization is essential for these phenomenon, because ALF does not show any activities against intact cells. This activity of ALF can be neutralized by free LPS but not by polysaccharide portion of LPS. ALF shows cell lysis activity against human polymorphonuclear leukocytes, mononuclear cells, and leukemia cells (K562), which are exposed with LPS (78).

Furthermore, ALF exhibits growth inhibitory activity against some, not all, strains of Gram-negative bacteria (77,82,83). Among the strains tested Re type strains, *Salmonella minnesota* R595 and *S. typhimurium* 1102, are most susceptible to the antibacterial action of ALF. Because ALF binds to lipid A portion of LPS, it seems likely that ALF neutralizes biologic activities of LPS. In fact, ALF suppresses the pyrogenicity of LPS in rabbits (92-94). Thus, it would be expected that ALF might be clinically useful to remove LPS and to suppress LPS pyrogenicity. As described earlier, ALF contains two positively charged regions and an amino-terminal hydrophobic region. If the two positively charged cluster regions form β -sheet and α -helix structure, respectively, the positive charges would cluster at the same side of the molecule. These two positively charged clusters might provide interaction site(s) with phosphate groups in the lipid A portion of LPS. Once ALF binds with LPS on cell membrane, the membrane structure seems to be perturbed by insertion of the hydrophobic amino-terminal region, including approximately up to 27th residues. The region appears to have a sufficient length to go across a lipid bilayer-like transmembrane α -helices of bacteriorhodopsin molecule (Fig. 15). The facts that ALF is localized in the L-granules and has strong antimicrobial activities do suggest that ALF is secreted during degranulation of the hemocyte induced by LPS and functions as antibactericidal substance following engulfment of the invaders by coagulation system (20).

Tachyplesins and Their Analogs. Tachyplesin I is a cationic peptide first isolated from acid extracts of Japanese horseshoe crab (*T. tridentatus*) hemocyte debris. This peptide inhibits the LPS-mediated activation of factor C in a similar manner to that of anti-LPS factor (74). Afterwards, two peptide analogs of tachyplesin I—tachyplesins II and III—have been isolated from hemocytes and two Southeast Asian horseshoe crabs, *T. gigas* and *C. rotundicauda* (75,79). Moreover, in the hemocyte debris of American *L. polyphemus*, two tachyplesin analogs, polyphemusins I and II, have also been found (75,90).

Tachyplesin is highly stable at low pH or high temperature, because the LPS-binding ability of tachyplesin is not affected even in 0.1% trifluoroacetic acid used for HPLC and by heat treatment in neutral pH buffer at 100°C for 30 minutes. This stability seems to be due to the




FIG. 15.

A speculative scheme of anti-LPS factor (ALF) which interacts with LPS at the positively charged regions and perturbs a cell membrane structure at the NH_2 -terminal hydrophobic region.

rigid structure imposed by the two disulfide linkages. The primary structures of tachyplesin family are shown in Figure 16. They are highly basic polypeptides composed of 17 or 18 residues with arginine α -amide as the carboxyl-terminal end. Naturally occurring peptides containing arginine α -amide at the COOH termini have been reported in a scorpion polypeptide toxin and sarcotoxins from *Sarcophaga peregrina* (108). Tachyplesin shows a characteristic structure with three tandem repeats of a tetrapeptide sequence, i.e., hydrophobic amino acid-Cys-hydrophobic amino acid-Arg, indicating that its amphipathic nature is closely associated with biological activity (74,91). Figure 17 shows the conformational structure of tachyplesin I (80,85). The $^1\text{H-NMR}$ spectrum of tachyplesin I in aqueous solution could be completely assigned, and the secondary structure is substantiated by interpretations of the nuclear Overhauser effect, coupling constant, amide exchange rate, and temperature dependence of the amide chemical shift. Tachyplesin I takes on a fairly rigid conformation constrained by two disulfide bridges and adopts a conformation consisting of an antiparallel β -sheet (residues 3-8 and 11-16) connected by a β -turn (residues 8-11). In this planar conformation, five bulky hydrophobic side groups are localized in one side of the plane, and six cationic side groups are




FIG. 16.

Primary structures of six kinds of tachyplesins and its analogs isolated from four species of horseshoe crabs. Tachyplesin III was isolated from *T. gigas*, I from *T. tridentatus*, *T. gigas* and *C. rotundicauda*, and II from *T. tridentatus* respectively. Polyphemusins I and II were isolated from *L. polyphemus*.

distributed at the “tail” part of the molecule (residues 1–5 and 14–17). This is presumed to be closely associated with the bactericidal activity (95).

As summarized in Table IV, tachyplesin displays potent antimicrobial activity against several strains of microorganisms (74,82). It is effective against not only rough and smooth strains




FIG. 17.

Schematic representation of the antiparallel β -pleated sheet structure of tachyplesin I.

TABLE IV
Antimicrobial activity of native and synthetic tachypleins and polyphemusins (75). Published by permission of the Japanese Biochemical Society, Tokyo. Original in Ref. 75.

Strain	Minimal inhibitory concentration ($\mu\text{g/ml}$)						
	Tachyplein I		Tachyplein II		Polyphemusin I		Polyphemusin II
	Native	Synthetic	Native	Synthetic	Native	Synthetic	
Gram (-)							
<i>Salmonella typhimurium</i> LT2	3.1	1.6-3.1	3.1	1.6-3.1	3.1	3.1-6.3	6.3
<i>Salmonella typhimurium</i> 1102	0.8-1.6	0.8-1.6	1.6	1.6-3.1	3.1	3.1	3.1
<i>Escherichia coli</i> K12	1.6-3.1	1.6-3.1	3.1	1.6-3.1	6.3	6.3	12.5
<i>Salmonella minnesota</i> 1114W	3.1	6.3	3.1	6.3	6.3	12.5	12.5
<i>Salmonella minnesota</i> R595	1.6	1.6	1.6	1.6	3.1	3.1	3.1
<i>Pseudomonas aeruginosa</i>	12.5	—	—	—	—	—	—
Gram (+)							
<i>Staphylococcus aureus</i> 209P	3.1	3.1	1.6-3.1	6.3	6.3	6.3	6.3
<i>Staphylococcus aureus</i> ATCC 25923	6.3	12.5	6.3	12.5	6.3	6.3	12.5
<i>Bacillus subtilis</i>	3.13	—	—	—	—	—	—
Fungus							
<i>Candida albicans</i> M9	3.1	3.1	3.1	3.1	6.3	6.3	6.3
<i>Cryptococcus neoformans</i> IMF40040	1.56	—	—	—	—	—	—

of Gram-negative but also Gram-positive cells, such as *Staphylococcus* species. In the presence of tachyplein at 3.5 $\mu\text{g/ml}$, *Salmonella* strains irreversibly lose viability. This antimicrobial potency is comparable to that of anti-LPS factor previously described. Therefore, tachyplein seems very likely to act as an antimicrobial peptide for the defense of horseshoe crabs against microbial infections (84,86).

Protein Precursors of Tachypleins I and II. The cDNA encoding for tachyplein precursors has been cloned and sequenced (Fig. 18). The tachyplein precursors consist of 77 amino acids with 23 residues in a presegment, and there are two types of mRNAs corresponding to the iso-peptides of tachypleins I and II (81). Both precursors contain a putative signal peptide, a processing peptide sequence, and a carboxyl-terminal amidation signal "Gly-Lys-Arg" connected to the mature tachyplein peptide. Moreover, an unusual acidic amino acid cluster with Asp-Glu-Asp-Glu-Asp-Asp-Glu-Glu-COOH is present in the carboxyl-terminal portions of both precursors. This acidic region might interact with a cationic part of the tachyplein peptide to stabilize a conformational structure of the precursor for proteolytic degradation. The amino acid sequence of precursor molecule suggests that the mature tachyplein might be generated through several steps of the processing. First, the amino-terminal signal sequence of the tachyplein precursor is processed in a stepwise fashion by a signal peptidase followed by a dipeptidyl aminopeptidase, because the Glu-Ala-Glu-Ala sequence preceding the mature tachyplein peptide is fit for the substrate specificity of a dipeptidyl aminopeptidase. Second, the cleavage of Arg-Asn bond at positions 20 and 21 is occurred by a processing enzyme that recognizes the dibasic sequence of Lys-Arg at positions 19 and 20. Third, release of Lys-19 and Arg-20 is taken place by a carboxypeptidase B-like enzyme, and finally an oxidative amidation is catalyzed by an enzyme recognizing the carboxyl-terminal glycine residue.

Northern blot analyses of total RNA prepared from various tissues indicate that the tachyplein precursor is expressed exclusively in the hemocytes, in accordance with the high




FIG. 18.

Amino acid sequence of tachyplesin I precursor. The residues circled thickly constitute a mature tachyplesin. The charged residues are indicated by + and -.

levels of tachyplesin peptides recovered therefrom (81). The mRNAs from heart and brain tissues also contain the transcript identical in size (0.7 kb) to that found in the hemocytes, but the signals are relatively weak. The amount of the tachyplesin precursor mRNA in other tissues, including those of the hepatopancreas, stomach, intestine, muscle, and coxal gland, is negligible. Although little is known about the hematopoietic tissue for production of hemocytes in horseshoe crab, our observations suggest that cardiac tissues may be the site of formation of circulating hemocytes. The subcellular localization of tachyplesin in the hemocytes proves to be in the S-granules, as determined immunohistochemically (20).

CONCLUSION

From all the results described previously, we speculate that bacterial endotoxin first contacts with an LPS receptor or LPS-binding proteins probably present in plasma membrane of the hemocyte and activates factor C located in the granules, and the resulting active factor C induces the activation of the intracellular clotting system (Fig. 19). The β -glucan-mediated pathway (67,68) also could not be disregarded for activation of this system. The clot generated during the activation encapsulates and immobilizes the Gram-negative bacteria, and the released peptide tachyplesin and anti-LPS factor act as the bactericidal substances. Therefore, this intracellular clotting system may have a crucial role in host defense against invading microorganisms.

Figure 20 shows an outline of the cellular and humoral defense systems in limulus. Although we have not mentioned about plasma components in this review, there are many humoral factors, such as proteinase inhibitors (87,96), α_2 -macroglobulin (88,89), lectins (102), C-reactive protein (101), and polyphemins (4), all of which are important in antimicrobial defense (104). Furthermore, Liu and his colleagues (107) have also reported several endotoxin-




FIG. 19.

The hypothetical processes of coagulogen-based clotting system and containment of invading Gram-negative bacteria.




FIG. 20.

Cellular and humoral defense systems in *Limulus*.

binding proteins and a cell adhesion protein found in the limulus hemocytes (97-99,105,106). Although the exact functions of these substances are not known at present, they may act in concert with other components to serve as the biological defense of this animal (100,103). However, we have still less knowledge about cell biology and physiology of horseshoe crab hemocytes in comparison with mammalian blood cells.

Acknowledgements

This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan. The authors wish to express their thanks to Drs. T. Morita, T. Nakamura, M. Niwa, Y. Miura, T. Shigenaga, R. Hashimoto, N. Seki, and T. Oda for their collaborations of this work, and to I. Edamitsu for expert secretarial assistance.

REFERENCES

1. MacFARLAN, R.G. The hemostatic mechanism in man and other animals. In: *Symp. Zool Soc. London, Vol. 27*, New York: Academic Press, 1970, pp. 1-6.
2. ARCHER, R.K. Blood coagulation in vertebrate animals other than man. In: *Recent Advances in Blood Coagulation*, L. Poller (Ed.). J. & A. Churchill Ltd., 1969, pp. 29-37.
3. RAVINDRANATH, M.H. Hemocytes in hemolymph coagulation of arthropods. *Biol. Rev.* 55, 139-170, 1980.
4. ARMSTRONG, P.B. Cellular and humoral immunity in the horseshoe crab. In: *Limulus polyphemus. Immunology of Insects and Other Arthropods*, A.P. Gupta (Ed.). Boca Raton, FL: CRC Press, 1991, pp. 3-17.
5. IWANAGA, S., MORITA, T., MIYATA, T., NAKAMURA, T., HIRANAGA, M. and OHTSUBO, S. The limulus coagulation system sensitive to bacterial endotoxins. In: *Bacterial Endotoxin: Recent Analytical, Synthetic and Biochemical Approaches*, Y.J. Homma et al. (Eds.). Heidelberg: Verlag Chemie, 1983, pp. 365-382.
6. MORITA, T., NAKAMURA, T., OHTSUBO, S., TANAKA, S., MIYATA, T., HIRANAGA, M. and IWANAGA, S. Intracellular proteinases and inhibitors associated with the hemolymph coagulation system of the horseshoe crabs (*Tachypleus tridentatus*, *Limulus polyphemus*). In: *Proteinase Inhibitors: Medical and Biological Aspects*, N. Katunuma, H. Umezawa, and H. Holzer (Eds.). Berlin: Springer-Verlag, 1983, pp. 229-241.
7. BANG, F.B. A bacterial disease of *Limulus polyphemus*. *Bull. Johns Hopkins Hosp.* 98, 325-350, 1956.
8. LEVIN, J. and BANG, F.B. The role of endotoxin in the extracellular coagulation of *Limulus* blood. *Bull. Johns Hopkins Hosp.* 115, 265-274, 1964.
9. LIU, T.-Y., SEID, R.C.JR., TAI, J.Y., LIANG, S.-M., SAKMAR, T.P. and ROBBINS, J.B. Studies on *Limulus* lysate coagulating system. In: *Biomedical Applications of the Horseshoe Crab (Limulidae)*. E. Cohen (Ed.). New York: Alan R. Liss, Inc., 1979, pp. 147-158.
10. MORITA, T., NAKAMURA, T., MIYATA, T. and IWANAGA, S. Biochemical characterization of limulus clotting factors and inhibitors which interact with bacterial endotoxins. *Prog. Clin. Biol. Res.* 189, 53-64, 1985.
11. IWANAGA, S., MORITA, T., MIYATA, T., NAKAMURA, T. and AKETAGAWA, J. The hemolymph coagulation system in invertebrate animals. *J. Protein Chem.* 5, 225-268, 1986.

12. DUMONT, J.N., ANDERSON, E. and WINNER, G. Some cytologic characteristics of the hemocytes of *Limulus* during clotting. *J. Morphol.* 119, 181-208, 1966.
13. ARMSTRONG, P.B. Interaction of the motile blood cells of the horseshoe crab, *Limulus*. Studies on contact paralysis of pseudopodial activity and cellular overlapping *in vitro*. *Exp. Cell Res.* 107, 127-138, 1977.
14. MÜRER, E.H., LEVIN, J. and HOLM, R. Isolation and studies of the granules of the amebocytes of *Limulus polyphemus*, the horseshoe crab. *J. Cell Physiol.* 86, 533-542, 1975.
15. ORNBERG, R.L. and REESE, T.S. Beginning of exocytosis captured by rapid-freezing of *Limulus* amebocytes. *J. Cell Biol.* 90, 40-54, 1981.
16. ARMSTRONG, P.B. and RICKLES, F.R. Endotoxin-induced degranulation of the *Limulus* amebocyte. *Exp. Cell Res.* 140, 15-24, 1982.
17. ORNBERG, R.L., Exocytosis in *Limulus* amebocytes. In: *Blood Cells of Marine Invertebrates: Experimental Systems in Cell Biology and Comparative Physiology, MBL Lectures in Biology vol 6*, W.D. Cohen (Ed.). New York: Alan R. Liss, Inc., 1985, pp. 127-142.
18. BRANDIN, E.R. and PISTOLE, T.G. Presence of microorganisms in the hemolymph of the horseshoe crab, *Limulus polyphemus*. *Appl. Environ. Microbiol.* 49, 718-723, 1985.
19. LEVIN, J. The role of amebocytes in the blood coagulation mechanism of the horseshoe crab *Limulus polyphemus*. In: *Blood Cells of Marine Invertebrates. Experimental Systems in Cell Biology and Comparative Physiology, MBL Lectures in Biology vol 6*, W.D. Cohen (Ed.). New York: Alan R. Liss, Inc., 1985, pp. 145-163.
20. TOH, Y., MIZUTANI, A., TOKUNAGA, F., MUTA, T. and IWANAGA, S. Structure of hemocytes of the Japanese horseshoe crab *Tachypleus tridentatus*: Fine structure, morphological changes during coagulation and localization of clotting factors and antimicrobial substances. *Cell Tissue Res.* 266, 137-147, 1991.
21. SOLUM, N.O. Some characteristics of the clottable protein of *Limulus polyphemus* blood cells. *Thromb. Diath. Haemorrh.* 23, 170-181, 1970.
22. SOLUM, N.O. The coagulogen of *Limulus polyphemus* hemocytes. A comparison of the clotted and non-clotted forms of the molecule. *Thromb. Res.* 2, 55-70, 1973.
23. NAKAMURA, S., IWANAGA, S., HARADA, T. and NIWA, M. A clottable protein (coagulogen) from amebocyte lysate of Japanese horseshoe crab (*Tachypleus tridentatus*). *J. Biochem. (Tokyo)* 80, 1011-1021, 1976.
24. NAKAMURA, S., TAKAGI, T., IWANAGA, S., NIWA, M. and TAKAHASHI, K. A clottable protein (coagulogen) of horseshoe crab hemocytes: Structural change of its polypeptide chain during gel formation. *J. Biochem. (Tokyo)* 80, 649-652, 1976.
25. NAKAMURA, S., TAKAGI, T., IWANAGA, S., NIWA, M. and TAKAHASHI, K. Amino acid sequence studies on the fragments produced from horseshoe crab coagulogen during gel formation: Homologies with primate fibrinopeptide B. *Biochem. Biophys. Res. Commun.* 72, 902-908, 1976.
26. TAI, J.Y., SEID, R.C.JR., HUHN, R.D. and LIU, T.-Y. Studies on *Limulus* amebocyte lysate II: Purification of the coagulogen and the mechanism of clotting. *J. Biol. Chem.* 252, 4773-4776, 1977.
27. NAKAMURA, S., MORITA, T., IWANAGA, S., NIWA, M. and TAKAHASHI, K. A sensitive substrate for the clotting enzyme in horseshoe crab hemocytes. *J. Biochem. (Tokyo)* 81, 1567-1577, 1977.
28. SHISHIKURA, F. and SEKIGUCHI, K. Comparative studies on horseshoe crab coagulogens. *J. Exp. Zool.* 206, 241-246, 1978.

29. MOSESSON, M.W., WOLFENSTEIN-TODEL, C., LEVIN, J. and BERTRAND, O. Characterization of amebocyte coagulogen from the horseshoe crab (*Limulus polyphemus*). *Thromb. Res.* 14, 765-779, 1979.
30. TAKAGI, T., HOKAMA, Y., MORITA, T., IWANAGA, S., NAKAMURA, S. and NIWA, M. Amino acid sequence studies on horseshoe crab (*Tachypleus tridentatus*) coagulogen and the mechanism of gel formation. In: *Biomedical Applications of the Horseshoe Crab (Limulidae)*. E. Cohen (Ed.). New York: Alan R. Liss, Inc., 1979, pp. 69-184.
31. NAKAMURA, S., SHISHIKURA, F., IWANAGA, S., TAKAGI, T., TAKAHASHI, K., NIWA, M. and SEKIGUCHI, K. Horseshoe crab coagulogens. Their structure and gelation mechanism. In: *Frontiers in Protein Chemistry*, T.-Y. Liu, G. Mamiya, and T. Yasunobu (Eds.). New York: Elsevier North Holland, Inc., 1980, pp. 495-514.
32. SHISHIKURA, F., NAKAMURA, S., TAKAHASHI, K. and SEKIGUCHI, K. Horseshoe crab phylogeny based on amino acid sequence of the fibrino-peptide-like peptide C. *J. Exp. Zool.* 223, 89-91, 1982.
33. SHISHIKURA, F., NAKAMURA, S., TAKAHASHI, K. and SEKIGUCHI, K. Coagulogens from the four living species of horseshoe crab (Limulidae): Comparison of their biochemical and immunochemical properties. *J. Biochem. (Tokyo)* 94, 1279-1287, 1983.
34. MIYATA, T., HIRANAGA, M., UMEZU, M. and IWANAGA, S. The complete amino acid sequence of coagulogen isolated from *Limulus polyphemus* amebocytes. *Ann. New York Acad. Sci.* 408, 651-654, 1983.
35. MIYATA, T., USUI, K. and IWANAGA, S. The amino acid sequence of coagulogen isolated from Southeast Asian horseshoe crab *Tachypleus gigas*. *J. Biochem. (Tokyo)* 95, 1793-1801, 1984.
36. IWANAGA, S., MORITA, T., MIYATA, T. and NAKAMURA, T. The hemolymph coagulation system in *Limulus*. In: *Microbiology—1985*. American Society for Microbiology, 1985, pp. 29-32.
37. MIYATA, T., MATSUMOTO, H., HATTORI, M., SAKAKI, Y. and IWANAGA, S. Two types of coagulogen mRNAs found in horseshoe crab (*Tachypleus tridentatus*) hemocytes: Molecular cloning and nucleotide sequences. *J. Biochem. (Tokyo)* 100, 213-220, 1986.
38. TAKAGI, T., HOKAMA, Y., MIYATA, T., MORITA, T. and IWANAGA, S. Amino acid sequence of Japanese horseshoe crab (*Tachypleus tridentatus*) coagulogen B chain: Completion of the coagulogen sequence. *J. Biochem. (Tokyo)* 95, 1445-1457, 1984.
39. SRIMAL, S., MIYATA, T., KAWABATA, S. and IWANAGA, S. The complete amino acid sequence of coagulogen isolated from Southeast Asian horseshoe crab, *Carcinoscorpius rotundicauda*. *J. Biochem. (Tokyo)* 98, 305-318, 1985.
40. MIYATA, T., HIRANAGA, M., UMEZU, M. and IWANAGA, S. Amino acid sequence of the coagulogen from *Limulus polyphemus* hemocytes. *J. Biol. Chem.* 259, 8924-8933, 1984.
41. CHENG, S.M., SUZUKI, A., ZON, G. and LIU, T.-Y. Characterization of a complementary deoxyribonucleic acid for the coagulogen of *Limulus polyphemus*. *Biochim. Biophys. Acta* 868, 1-8, 1986.
42. MIYATA, T. and IWANAGA, S. Limulus coagulogen (invertebrate fibrinogen-like substance): Its structure, function and molecular evolution. In: *Fibrinogen 4. Current Basic and Clinical Aspects*, M. Matsuda, S. Iwanaga, A. Takada and A. Henschen (Eds.). Amsterdam: Elsevier Science Publishers B.V. (Biomedical Division), 1990, pp. 29-33.
43. SHUSTER, C.N.JR. Serological correspondence among horseshoe "crabs" (Limulidae). *Zoologica. N.Y.* 47, 1-7, 1962.

44. SHUSTER, C.N.JR. Xiphosurida. In: *Encyclopedia of Science and Technology*, vol. 14. New York: McGraw-Hill, Book. Co. Inc., 1982, pp. 766-770.
45. SULLIVAN, J.D.JR. and WATSON, S.W. Purification and properties of the clotting enzyme from *Limulus* lysate. *Biochem. Biophys. Res. Commun.* 66, 845-855, 1975.
46. TAI, J.Y. and LIU, T.-Y. Studies on *Limulus* amebocyte lysate. Isolation of pro-clotting enzyme. *J. Biol. Chem.* 252, 2178-2181, 1977.
47. YOUNG, N.S., LEVIN, J. and PRENDERGAST, R.A. An invertebrate coagulation system activated by endotoxin: Evidence for enzymatic mediation. *J. Clin. Invest.* 51, 1790-1797, 1972.
48. NAKAMURA, S., MORITA, T., HARADA-SUZUKI, T., IWANAGA, S., TAKAHASHI, K. and NIWA, M. A clotting enzyme associated with the hemolymph coagulation system of horseshoe crab (*Tachypleus tridentatus*): Its purification and characterization. *J. Biochem. (Tokyo)* 92, 781-792, 1982.
49. NAKAMURA, S. and LEVIN, J. Endotoxin-mediated *Limulus* proclotting enzyme activator and detection of a previously undescribed protease (protease N). *Biochem. Biophys. Res. Commun.* 108, 1619-1623, 1982.
50. NAKAMURA, S. and LEVIN, J. Fractionation of *Limulus* amebocyte lysate. Characterization of activation of the proclotting enzyme by an endotoxin-mediated activator. *Biochim. Biophys. Acta.* 707, 217-225, 1982.
51. HARADA-SUZUKI, T., MORITA, T., IWANAGA, S., NAKAMURA, S. and NIWA, M. Further studies on the chromogenic substrate assay method for bacterial endotoxins using horseshoe crab (*Tachypleus tridentatus*) hemocyte lysate. *J. Biochem. (Tokyo)* 92, 793-800, 1982.
52. NAKAMURA, T., MORITA, T. and IWANAGA, S. Intracellular proclotting enzyme in limulus (*Tachypleus tridentatus*) hemocytes: Its purification and properties. *J. Biochem. (Tokyo)* 97, 1561-1574, 1985.
53. MUTA, T., HASHIMOTO, R., MIYATA, T., NISHIMURA, H., TOH, Y. and IWANAGA, S. Proclotting enzyme from horseshoe crab hemocytes: cDNA cloning, disulfide locations, and subcellular localization. *J. Biol. Chem.* 265, 22426-22433, 1990.
54. NAKAMURA, T., HORIUCHI, T., MORITA, T. and IWANAGA, S. Purification and properties of intracellular clotting factor, factor B, from horseshoe crab (*Tachypleus tridentatus*) hemocytes. *J. Biochem. (Tokyo)* 99, 847-857, 1986.
55. OHKI, M., NAKAMURA, T., MORITA, T. and IWANAGA, S. A new endotoxin sensitive factor associated with hemolymph coagulation system of horseshoe crab (Limulidae). *FEBS Lett.* 120, 217-220, 1980.
56. TORANO, A.E., NAKAMURA, S. and LEVIN, J. Properties of the clotting enzyme responsible for endotoxin-mediated *Limulus* coagulation. *Thromb. Res.* 34, 407, 1984.
57. MUTA, T., HASHIMOTO, R., ODA, T., MIYATA, T. and IWANAGA, S. *Limulus* clotting enzyme and factor B associated with endotoxin-sensitive coagulation cascade: Novel serine protease zymogens with a new type of "disulfide-knotted domain." *Thrombos. Haemostas.* 65 (abstract), 935, 1991.
58. NAKAMURA, T., MORITA, T. and IWANAGA, S. Lipopolysaccharide-sensitive serine-protease zymogen (factor C) found in *Limulus* hemocytes: Isolation and characterization. *Eur. J. Biochem.* 154, 511-521, 1986.
59. HARADA, T., MORITA, T., IWANAGA, S. and NIWA, M. A new chromogenic substrate method for assay of bacterial endotoxins using *Limulus* hemocyte lysate. *Prog. Clin. Biol. Res.* 29, 202-220, 1979.
60. TOKUNAGA, F., MIYATA, T., NAKAMURA, T., MORITA, T., KUMA, K., MIYATA,

- T. and IWANAGA, S. Lipopolysaccharide-sensitive serine-protease zymogen (factor C) of horseshoe crab hemocytes: Identification and alignment of proteolytic fragment produced during the activation show that it is a novel type of serine protease. *Eur. J. Biochem.* 167, 405–416, 1987.
61. URAGOH, K., SUEISHI, K., NAKAMURA, T. and IWANAGA, S. A novel immunohistochemical method for *in vivo* detection of endotoxin using horseshoe crab factor C. *J. Histochem. Cytochem.* 36, 1275–1283, 1988.
62. TOKUNAGA, F., NAKAMURA, T., MUTA, T., MIYATA, T. and IWANAGA, S. Intracellular serine protease zymogen, factor C, from limulus hemocytes: Its structure and function. In: *Proceeding of International Symposium on Intracellular Protein Catabolism*, N. Katunuma and E. Kominami (Eds.). Tokyo: Japan Scientific Societies Press, 1989, pp. 120–128.
63. NAKAMURA, T., TOKUNAGA, F., MORITA, T., IWANAGA, S., KUSUMOTO, S., SHIBA, T., KOBAYASHI, T. and INOUE, K. Intracellular serine-protease zymogen, factor C, from horseshoe crab hemocytes: Its activation by synthetic lipid A analogues and acidic phospholipids. *Eur. J. Biochem.* 176, 89–94, 1988.
64. NAKAMURA, T., TOKUNAGA, F., MORITA, T. and IWANAGA, S. Interaction between lipopolysaccharide and intracellular serine protease zymogen, factor C, from horseshoe crab (*Tachypleus tridentatus*) hemocytes. *J. Biochem. (Tokyo)* 103, 370–374, 1988.
65. MUTA, T., MIYATA, T., MISUMI, Y., TOKUNAGA, F., NAKAMURA, T., TOH, Y., IKEHARA, Y. and IWANAGA, S. Limulus factor C: An endotoxin sensitive serine protease zymogen with a mosaic structure of complement-like, epidermal growth factor-like, and lectin-like domains. *J. Biol. Chem.* 266, 6554–6561, 1991.
66. TOKUNAGA, F., NAKAJIMA, H. and IWANAGA, S. Purification and characterization of lipopolysaccharide-sensitive serine protease zymogen (factor C) isolated from *Limulus polyphemus* hemocytes: A newly identified intracellular zymogen activated by α -chymotrypsin, not by trypsin. *J. Biochem. (Tokyo)* 109, 150–157, 1991.
67. MORITA, T., TANAKA, S., NAKAMURA, T. and IWANAGA, S. A new (1-3)- β -D-glucan-mediated coagulation pathway found in *Limulus* amebocytes. *FEBS Lett.* 129, 318–321, 1981.
68. KAKINUMA, A., ASANO, T., TORII, H. and SUGINO, Y. Gelation of *Limulus* amebocyte lysate by an antitumor (1-3)- β -D-glucan. *Biochem. Biophys. Res. Commun.* 101, 433–439, 1981.
69. TANAKA, S., NAKAMURA, T., MORITA, T. and IWANAGA, S. *Limulus* anti-LPS factor: An anticoagulant which inhibits the endotoxin-mediated activation of *Limulus* coagulation system. *Biochem. Biophys. Res. Commun.* 105, 717–723, 1982.
70. AKETAGAWA, J., MIYATA, T., OHTSUBO, S., NAKAMURA, T., HAYASHIDA, H., MIYATA, T. and IWANAGA, S. Primary structure of *Limulus* anticoagulant anti-lipopolysaccharide factor. *J. Biol. Chem.* 261, 7375–7365, 1986.
71. NACHUM, R., WATSON, S.W. and SIEGEL, S.E. Antimicrobial defense mechanisms in the horseshoe crab, *Limulus polyphemus*: Effect of sodium chloride on bactericidal activity. *J. Invertebr. Pathol.* 36, 382, 1980.
72. SMITH, R.H. and PISTOLE, T.G. Bactericidal activity of granules isolated from amebocytes of the horseshoe crab, *Limulus polyphemus*. *J. Invertebr. Pathol.* 45, 272, 1985.
73. PISTOLE, T.G. and GRAF, S.A. Antibacterial activity in *Limulus*. In: *Hemolytic and Humoral Immunity in Arthropoda*, A.P. Gupta (Ed.). New York: John Wiley & Sons, 1986, p. 331.
74. NAKAMURA, T., FURUNAKA, H., MIYATA, T., TOKUNAGA, F., MUTA, T.,

- IWANAGA, S., NIWA, M., TAKAO, T. and SHIMONISHI, Y. Tachyplesin, a class of antimicrobial peptide from hemocytes of the horseshoe crab (*Tachypleus tridentatus*). *J. Biol. Chem.* 263, 16709-16713, 1988.
75. MIYATA, T., TOKUNAGA, F., YONEYA, Y., YOSHIKAWA, K., IWANAGA, S., NIWA, M., TAKAO, T., and SHIMONISHI, Y. Antimicrobial peptides isolated from horseshoe crab hemocytes, Tachyplesin II, and Polyphemusins I and II: Chemical structures and biological activity. *J. Biochem. (Tokyo)* 106, 663-668, 1989.
76. MUTA, T., MIYATA, T., TOKUNAGA, F., NAKAMURA, T. and IWANAGA, S. Primary structure of anti-lipopolysaccharide factor from American horseshoe crab, *Limulus polyphemus*. *J. Biochem. (Tokyo)* 101, 1321-1330, 1987.
77. MORITA, T., OHTSUBO, S., NAKAMURA, T., TANAKA, S., IWANAGA, S., OHASHI, K. and NIWA, M. Isolation and biological activities of *Limulus* anticoagulant (anti-LPS factor) which interacts with lipopolysaccharide (LPS). *J. Biochem. (Tokyo)* 97, 1611-1620, 1985.
78. OHASHI, K., NIWA, M., NAKAMURA, T., MORITA, T. and IWANAGA, S. Anti-LPS factor in the horseshoe crab, *Tachypleus tridentatus*: Its hemolytic activity on the red blood cell sensitized with lipopolysaccharide. *FEBS Lett.* 176, 207-210, 1984.
79. MUTA, T., FUJIMOTO, T., NAKAJIMA, H. and IWANAGA, S. Tachyplesins isolated from hemocytes of Southeast Asian horseshoe crab (*Carcinoscorpius rotundicauda* and *Tachypleus gigas*): Identification of a new Tachyplesin, Tachyplesin III, and a processing intermediate of its precursor. *J. Biochem. (Tokyo)* 108, 261-266, 1990.
80. KAWANO, K., YONEYA, T., MIYATA, T., YOSHIKAWA, K., TOKUNAGA, F., TERADA, Y. and IWANAGA, S. Antimicrobial peptide, tachyplesin I, isolated from hemocytes of the horseshoe crab (*Tachypleus tridentatus*). *J. Biol. Chem.* 265, 15365-15367, 1990.
81. SHIGENAGA, T., MUTA, T., TOH, Y., TOKUNAGA, F. and IWANAGA, S. Antimicrobial tachyplesin peptide precursor: cDNA cloning and cellular localization in horseshoe crab (*Tachypleus tridentatus*). *J. Biol. Chem.* 265, 21350-21354, 1990.
82. NIWA, M., HUA, H.E., IWANAGA, S., MORITA, T., MIYATA, T., NAKAMURA, T., AKETAGAWA, J., MUTA, T., TOKUNAGA, F. and OHASHI, K. Biological activities of anti-LPS factor and LPS binding peptide from horseshoe crab amebocytes. In: *Endotoxin*, H. Friedman et al. (Eds.). *Adv. Exp. Med. Biol.* 256, New York: Plenum Publishing Co., 1990, pp. 257-271.
83. MUTA, T., NAKAMURA, T., FURUNAKA, H., TOKUNAGA, F., MIYATA, T., NIWA, M. and IWANAGA, S. Primary structures and functions of anti-lipopolysaccharide factor and tachyplesin peptide found in horseshoe crab hemocytes. In: *Endotoxin*. H. Friedman et al. (Eds.). *Adv. Exp. Med. Biol.* 256, New York: Plenum Publishing Co., 1990, pp. 273-285.
84. MORIMOTO, M., MORI, H., OTAKE, T., UEBA, N., KUNITA, N., NIWA, M., MURAKAMI, T. and IWANAGA, S. Inhibitory effect of tachyplesin I on the proliferation of human immunodeficiency virus *in vitro*. *Chemotherapy* 37, 206-211, 1991.
85. KAWANO, K., YONEYA, T., MIYATA, T., YOSHIKAWA, K., TOKUNAGA, F., TERADA, Y. and IWANAGA, S. Conformational analysis of tachyplesin I, LPS binding antimicrobial peptide, isolated from hemocytes of the horseshoe crab by ¹H NMR. *Peptide Chemistry 1990*. Osaka: Protein Research Foundation, 1991, pp. 385-388.
86. MURAKAMI, T., NIWA, M., TOKUNAGA, F., MIYATA, T. and IWANAGA, S. Direct virus inactivation of tachyplesin I and its isopeptides from horseshoe crab hemocytes. *Chemotherapy* 37, 327-334, 1991.

87. NAKAMURA, T., HIRAI, T., TOKUNAGA, F., KAWABATA, S. and IWANAGA, S. Purification and amino acid sequence of Kunitz-type protease inhibitor found in the hemocytes of the horseshoe crab (*Tachypleus tridentatus*). *J. Biochem. (Tokyo)* 101, 1297–1306, 1987.
88. ARMSTRONG, P.B., QUIGLEY, J.P. and RICKLES, F.R. The *Limulus* blood cell secretes α_2 -macroglobulin when activated. *Biol. Bull. (Woods Hole, MA)* 178, 137–143, 1990.
89. ARMSTRONG, P.B. and QUIGLEY, J.P. α_2 -Macroglobulin: A recently-discovered defense system in arthropods. In: *Immunology of Insects and Other Arthropods*, A.P. Gupta (Ed.). Boca Raton, FL, 1991, in press.
90. AKAJI, K., FUJII, N., TOKUNAGA, F., MIYATA, T., IWANAGA, S. and YAJIMA, H. Studies on peptides CLXVIII: Syntheses of three peptides isolated from horseshoe crab hemocytes, tachyplesin I, tachyplesin II, and polyphemusin I. *Chem. Pharm. Bull.* 37, 2661–2664, 1989.
91. SHIEH, T.C., KOHZUMA, T., PARK, N.G., OHNO, M., NAKAMURA, T., IWANAGA, S. and YAMAMOTO, T. Synthesis and properties of tachyplesin I, a lipopolysaccharide-binding peptide, from *Tachypleus tridentatus*. *FEBS Lett.* 252, 121–124, 1989.
92. ROSLANSKY, P.F. and NOVITSKY, T.J. Sensitivity of *Limulus* amebocyte lysate (LAL) to LAL-reactive glucans. *J. Clin. Microbiol.* 29, 2477–2483, 1991.
93. ALPERT, G., BALDWIN, G., THOMPSON, C., WAINWRIGHT, N., NOVITSKY, T.J., GILLS, Z., PARSONNET, J., FLEISHER, G.R. and SIBER, G.R. *Limulus* anti-lipopolysaccharide factor protects rabbit from *Meningococcal* endotoxin shock. *J. Infect. Dis.* 165, 494–500, 1992.
94. WAINWRIGHT, N.R., MILLER, R.J., PAUS, E., NOVITSKY, T.J., FLETCHER, M.A., McKENNA, T.M. and WILLIAMS, T. Endotoxin binding and neutralizing activity by a protein from *Limulus polyphemus*. In: *Cellular and Molecular Aspects of Endotoxin Reaction*, A. Nowotny, J.J. Spitzer, and E.J. Ziegler (Eds.). Amsterdam: Elsevier Science Publishers (Biomedical Division), 1991, pp. 315–325.
95. YONEZAWA, A., KUWAHARA, J., FUJII, N. and SUGIURA, Y. Binding of tachyplesin I to DNA revealed by footprinting analysis: Significant contribution of secondary structure to DNA binding and implication for biological action. *Biochemistry* 31, 2998–3004, 1992.
96. ARMSTRONG, P.B. and QUIGLEY, J.P. Protease inhibitory activity released from the horseshoe crab blood cell during exocytosis. *Biochim. Biophys. Acta* 827, 453–459, 1985.
97. LIANG, S.-M., SAKMAR, T.P. and LIU, T.-Y. Studies on *Limulus* amebocyte lysate III: Purification of an endotoxin-binding protein from *Limulus* amebocyte membranes. *J. Biol. Chem.* 255, 5586–5590, 1980.
98. LIANG, S.-M., LIANG, C.-M. and LIU, T.-Y. Studies on *Limulus* amebocytes. Isolation and identification of a membrane-bound protein activator of cyclic nucleotide phosphodiesterase from *Limulus* amebocytes. *J. Biol. Chem.* 256, 4968–4972, 1981.
99. BRSEY, C.R. Histological response to injury in the horseshoe crab, *Limulus polyphemus*. *Can. J. Zool.* 57, 1158–1165, 1977.
100. WEINBERG, J.B., SMITH, P.F. and KAHANE, I. Bacterial lipopolysaccharides and Mycoplasmal lipoglycans: A comparison between their abilities to induce macrophage-mediated tumor killing and *Limulus* amebocyte lysate clotting. *Biochem. Biophys. Res. Commun.* 97, 493–499, 1980.

101. NGUYEN, N.Y., SUZUKI, A., CHENG, S.-M., ZON, G. and LIU, T.-Y. The amino acid sequence of *Limulus* C-reactive protein. *J. Biol. Chem.* 261, 10450, 10455, 1986.
102. PISTOLE, T.G. Bacterial agglutinins from *Limulus polyphemus*—an overview. *Prog. Clin. Biol. Res.* 29, 547–553, 1979.
103. COPELAND, D.E. and LEVIN, J. The fine structure of the amebocyte in the blood of *Limulus polyphemus*. I. Morphology of the normal cell. *Biol. Bull.* 169, 449–457, 1985.
104. LEVIN, J. The horseshoe crab: A model for Gram-negative sepsis in marine organs and humans. In: *Bacterial Endotoxins: Pathophysiological Effects, Clinical Significance, and Pharmacological Control*. Alan R. Liss, Inc., 1988, pp. 3–15.
105. MINETTI, C.A.S.A., LIN, Y., CISLO, T. and LIU, T.-Y. Purification and characterization of an endotoxin-binding protein with protease inhibitory activity from *Limulus* amebocytes. *J. Biol. Chem.* 266, 20773–20780, 1991.
106. SAITO, H., YOSHIOKA, Y., UEHARA, N., AKETAGAWA, J., TANAKA, S. and SHIBATA, Y. Relationship between conformation and biological response for (1-3)- β -D-glucans in the activation of coagulation factor G from limulus amebocyte lysate and host-mediated antitumor activity: Demonstration of single helix conformation as a stimulant. *Carbohydr. Res.* 212, 81–190, 1991.
107. LIU, T., LIN, Y., CISLO, T., MINETTI, C.A.S.A., BABA, J.M.K. and LIU, T.-Y. Limunectin: A phosphocholine-binding protein from *Limulus* amebocytes with adhesion-promoting properties. *J. Biol. Chem.* 266, 14813–14821, 1991.
108. OKADA, M. and NATORI, S. Primary structure of sarcotoxin I, an antibacterial protein induced in the hemolymph of *Sarcophaga peregrina* (Flesh fly) larvae. *J. Biol. Chem.* 260, 7174–7177, 1985.