

Note

Synthesis and the structure of mixed carboxylatoplatinum(IV) complexes involving one trifluoroacetate ligand

Hae Jin Kim, Kwan Mook Kim, Rita Song ^{*}, Youn Soo Sohn ^{*}*Department of Chemistry, Division of Nano Science, Ewha Womans University, Seoul 120-750, South Korea*

Received 29 April 2004; accepted 24 July 2004

Available online 28 August 2004

Abstract

Mono(trifluoroacetato)platinum(IV) complexes of the formula $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{TFAc})]$ (dach = *trans*-(±)-1,2-diaminocyclohexane, TFAc = trifluoroacetate, L = acetate or propionate) could be prepared from the reaction of $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{OH})]$ with trifluoroacetic anhydride in the presence of a base and the crystal structure of compound **4** was determined by X-ray crystallography. In vitro anti-tumor activity of complex **4** ($\text{ED}_{50} = 3.1 \mu\text{M}$) was found to be much higher than carboplatin ($\text{ED}_{50} = 10.2 \mu\text{M}$).

© 2004 Elsevier B.V. All rights reserved.

Keywords: Platinum(IV); Antitumor drug; Electrophilic substitution**1. Introduction**

Platinum(IV) complexes have been extensively studied for the past two decades to improve the toxicity and severe side effects of the platinum(II)-based anticancer drugs [1–3]. Platinum(IV) complexes have shown the possibility as an orally active anticancer drug, which may be greatly improved both in toxicity and the acquired drug resistance caused by cisplatin or carboplatin [4–7]. The lipophilicity of the Pt(IV) complexes has been regarded as one of the most important factors in structure–activity relationship [8,9]. We reported previously a new series of platinum(IV) complexes of the formula $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{OH})]$ (dach = *trans*-(±)-1,2-diaminocyclohexane, L = acetate or propionate) obtained by selective nucleophilic substitution of tetrahydroxoplatinum(IV) complexes by carboxylic acids with appropriate pK_a values ($\text{pK}_a = 4\text{--}5$) [10]. These Pt(IV) complexes were fur-

ther derivatized by the well-known electrophilic substitution of the remaining hydroxo ligand with different carboxylic anhydrides to tune further their lipophilicity and water solubility [11]. This synthetic method afforded various mixed carboxylatoplatinum(IV) complexes, $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3\text{L}']$ (L' = acetate, propionate, valerate, and pivalate), but resulted in unexpected products involving two trifluoroacetate (TFAc) ligands instead of one TFAc of the formula $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_2(\text{TFAc})_2]$, when the reaction was carried out with trifluoroacetic anhydride. Here, we report a new method to obtain pure complexes involving only one TFAc ligand of the formula $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{TFAc})]$, and the crystal structure and in vitro antitumor activity of the complexes will be briefly discussed.

2. Experimental**2.1. Materials and measurements**

Potassium tetrachloroplatinate(II) purchased from Kojima, *trans*-(±)-1,2-diaminocyclohexane (dach),

^{*} Corresponding authors. Tel : +82 2 3277 2345; fax : +82 2 3277 2384 (Y.S. Sohn), Tel.: +82 2 3277 3589; fax: +82 2 3277 2384 (R. Song).

E-mail addresses: rsong@ewha.ac.kr (R. Song), yssohn@ewha.ac.kr (Y.S. Sohn).

acetic anhydride (Ac_2O), propionic anhydride (Pr_2O) and trifluoroacetic anhydride (TFAC_2O) from Aldrich were used without further purification. Elemental analyses were carried out in the Organic Chemistry Research Center of Sogang University. IR spectra were recorded as KBr pellets on a Perkin Elmer 16FPC FT-IR spectrophotometer. ^1H NMR spectra were measured with Bruker 250 MHz spectrometer. ESI-Mass measurements were performed by HP5989A equipped with HP59987A as an electron-spray source. A mixture of methanol and water (80:20) containing 1% formic acid was used as solvent for the mass analysis. The starting platinum(IV) complexes, $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OAc})_3(\text{OH})]$ (**1**) and $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OPr})_3(\text{OH})]$ (**2**) were prepared according to the previous method as depicted in Scheme 1 [10].

2.2. Synthesis

Trifluoroacetic anhydride (0.5 mmol) and triethylamine (0.75 mmol) were added to a solution of tris(carboxylato)monohydroxoplatinum(IV) complexes (0.2 mmol) in CH_2Cl_2 (4 ml). After the reaction mixture was stirred for 4 h at room temperature, the solution was evaporated to dryness. The crude solid was dissolved in methanol and precipitated with ether, and purified by column chromatography (silica gel, CH_2Cl_2 : methanol = 10:1, v/v). The crystals suitable for X-ray crystallography were obtained by slow evaporation of the methanolic solution.

2.2.1. $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OAc})_3(\text{OTFAc})]$ (**3**)

Yield, 59%. *Anal.* Calc. for $\text{C}_{14}\text{H}_{23}\text{F}_3\text{N}_2\text{O}_8\text{Pt}$: C, 28.05; H, 3.87; N, 4.67. Found: C, 28.03; H, 3.61; N, 4.68%. ^1H NMR (D_2O): 2.72 (m, 2H, NCH), 2.25 (m, 2H, CH_2), 2.07 (s, 6H, eq- CH_3CO_2), 2.04 (s, 3H, ax- CH_3CO_2), 1.55 (m, 4H, CH_2), 1.19 (m, 2H, CH_2). IR (KBr, cm^{-1}): ν_{max} 3052, 2968, 1721, 1642, 1360, 1306, 1192. MS (ESI): m/z 598.8 $[\text{M}^+]$.

2.2.2. $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OPr})_3(\text{OTFAc})]$ (**4**)


Yield, 68%. *Anal.* Calc. for $\text{C}_{17}\text{H}_{29}\text{F}_3\text{N}_2\text{O}_8\text{Pt}$: C, 31.83; H, 4.56; N, 4.37. Found: C, 31.83; H, 4.56; N, 4.37; Found: C, 31.84; H, 4.59; N, 4.34%. ^1H NMR (D_2O , δ): 2.80 (m, 2H, NCH), 2.32 (m, 8H, OPr- CH_2 and CH_2), 1.61 (m, 4H, CH_2), 1.23 (m, 2H, CH_2), 1.03 (q, 9H, OPr- CH_3). IR (KBr, cm^{-1}): ν_{max} 2984, 2942, 1722, 1632, 1462, 1350, 1210. MS (ESI): m/z 664.0 $[\text{M} + \text{Na}^+]$.

2.3. Assay of *in vitro* cytotoxicity

The new present complexes **3** and **4** were subjected to bioassay according to the previous method [12].

2.4. X-ray structure determination

All the X-ray data were collected on an Enraf-Nonius CAD4 automated diffractometer equipped with a Mo X-ray tube and a graphite crystal monochromator. The orientation matrix and unit cell dimensions were determined from 25 machine centered reflections in the 2θ range of 15–25°. The variation of intensities was monitored by repeated check of intensities of three reflections every 1 h during the data collection period. Absorption corrections were applied by empirical psi scan on three reflection planes with a χ value of near 90°. A direct or Patterson method (SHELXS-97) [13] was employed to locate the platinum atom. Subsequent cycles of Fourier map and least square refinements located other atoms (SHELXL-97) [14]. All the nonhydrogen atoms were refined anisotropically. Hydrogen atoms were included in the structure factor calculation using a riding model. All the calculations were carried out using VAX and PC computers. The crystallographic data for complex **4** are listed in Table 1.



Scheme 1. The synthetic route to Pt(IV) complexes of mixed carboxylates.

Table 1
Crystallographic data for $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OPr})_3(\text{TFAc})]$

Empirical formula	$\text{C}_{17}\text{H}_{29}\text{F}_3\text{N}_2\text{O}_8\text{Pt}$
Formula weight	641.51
Crystal system	monoclinic
Space group	$P2_1/c$ (No. 14)
<i>Unit cell dimensions</i>	
a (Å)	10.891(2)
b (Å)	17.343(4)
c (Å)	12.143(4)
α (°)	90
β (°)	95.33(2)
γ (°)	90
Volume (Å ³)	2283.5(9)
Z	4
D_{calc} (mg/m ³)	1.866
Absorption coefficient (mm ⁻¹)	6.213
$F(000)$	1256
Crystal size (mm)	$0.12 \times 0.20 \times 0.22$
θ range for data collection	$1.68\text{--}25.16^\circ$
Reflections collected/unique	4313/4080 [$R_{\text{int}} = 0.0582$]
Completeness to $\theta = 25.16$	99.3%
Refinement method	full-matrix least-squares on F^2
Data/restraints/parameters	4080/10/280
Goodness-of-fit on F^2	1.055
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0662$, $wR_2 = 0.1645$
R indices (all data)	$R_1 = 0.1522$, $wR_2 = 0.2015$
Largest difference peak and hole (e Å ⁻³)	1.526 and -1.661

$R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|$, $wR_2 = \{\sum (F_o^2 - F_c^2)^2 / \sum wF_o^4\}^{1/2}$, where $w = 1 / \{\sum^2 F_o^2 + (0.0197P)^2 + 0.00P\}$, and where $P = \{\max(F_o^2, 0) + 2F_c^2\} / 3$.

3. Results and discussion

The reaction of the tetrahydroxoplatinum(IV) complex with acetic or propionic acid afforded tricarboxylato-monohydroxoplatinum(IV) complexes by proton mediated nucleophilic substitution as reported previously [10]. The remaining hydroxo ligand was further derivatized by electrophilic substitution with various carboxylic anhydrides. Generally, the electrophilic substitution reaction of the axial hydroxo ligand of (diamine)platinum(IV) complexes was easily carried out without side reaction [15], particularly in the case of the complexes bearing the chelating equatorial dicarboxylate ligand. The alternative electrophilic substitution reaction with acyl chloride needs the presence of a base, such as triethylamine or pyridine, to remove hydrochloric acid formed [16]. When $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{OH})]$ was reacted with acetic or propionic anhydride, the reaction proceeded cleanly, leading to $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3\text{L}']$ ($\text{L}' = \text{acetate}$, propionate) because the generated acid is weak acid ($\text{p}K_{\text{a}} = 4\text{--}5$) and their conjugate base is not so strong to replace the carboxylate ligand coordinated to the platinum(IV) atom [17]. However, in the reaction with trifluoroacetic anhydride, bis(trifluoroacetato)Pt(IV) complexes of the formula $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_2(\text{TFAc})_2]$ were obtained probably because the trifluoroacetyl group (CF_3CO^+) was presumed to attack the hydroxo ligand,

forming a trifluoroacetate ligand, but the strong trifluoroacetic acid resultant from the anhydride was subjected to nucleophilic substitution to replace one of the three carboxylate ligands coordinated to the platinum(IV) atom. In other words, the strong trifluoroacetic acid seems to weaken and labilize the Pt–O bond of other carboxylate ligands so that the nucleophilic substitution by the trifluoroacetate ion may easily take place. Therefore, the elimination of the resultant trifluoroacetic acid may be necessary to avoid the formation of the unexpected product substituted with the second trifluoroacetate. In fact, the addition of an equivalent amount of a base afforded the pure reaction product involving only one trifluoroacetate and three intact carboxylate ligands.

The electrophilic substitution of the axial hydroxo ligand by trifluoroacetic anhydride did not change considerably the ^1H NMR spectra of the complexes compared with that of the starting $[\text{Pt}^{\text{IV}}(\text{dach})\text{L}_3(\text{OH})]$. The ^1H NMR spectra taken in D_2O showed characteristic multiplet pattern of the two protons near the amine group of dach, probably due to the asymmetrically substituted trifluoroacetate ligand at axial position. Three acetate ligands in equatorial and axial position gave the distinct signals at 2.08 and 2.05 ppm, respectively, although the difference of the chemical shift of propionate in $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OPr})_3(\text{OH})]$ is less evident.

The ORTEP drawing of complex **4** is drawn in Fig. 1 and its relevant distances and angles is listed in Table 2. The crystal structure of complex **4** is almost the same as that found in other platinum(IV) complexes reported previously [17]. All compounds derived from tetrahydroxoplatinum(IV) complexes exhibited similar structural characteristics. The local geometry around the




Fig. 1. ORTEP drawing with an atomic labeling scheme for $[\text{Pt}^{\text{IV}}(\text{dach})(\text{OPr})_3(\text{TFAc})]$.

Table 2

Selected bond lengths (Å), intramolecular hydrogen bond length (Å) and angles (°) for [Pt^{IV}(dach)(OPr)₃(TFAc)]

Pt(1)–O(7)	2.004(14)	Pt(1)–O(5)	2.006(13)
Pt(1)–O(3)	2.008(14)	Pt(1)–N(1)	2.016(14)
Pt(1)–O(1)	2.033(13)	Pt(1)–N(2)	2.056(16)
O(1)–C(7)	1.29(2)	C(8)–F(1)	1.482(13)
N(1)–O(2)	2.701	N(2)–O(6)	2.788
N(2)–O(4)	2.724	N(1)–O(8)	2.738
O(7)–Pt(1)–O(5)	80.7(6)	O(7)–Pt(1)–O(3)	90.8(6)
O(7)–Pt(1)–N(1)	97.4(6)	O(5)–Pt(1)–O(1)	87.7(6)
O(3)–Pt(1)–O(1)	174.4(6)	N(1)–Pt(1)–O(1)	95.7(7)
O(5)–Pt(1)–N(2)	100.0(6)	N(1)–Pt(1)–N(2)	82.4(6)

Table 3

In vitro antitumor activity of platinum(IV) complexes of mixed carboxylates

Complex	Cytotoxicity
	ED ₅₀ (μM)
[Pt ^{IV} (dach)(OAc) ₃ (OH)] ¹⁰ (1)	>40
[Pt ^{IV} (dach)(OPr) ₃ (OH)] ¹⁰ (2)	27.6
[Pt ^{IV} (dach)(OAc) ₂ (OTFAc) ₂] ¹⁰	7.2
[Pt ^{IV} (dach)(OPr) ₂ (OTFAc) ₂] ¹⁰	1.3
[Pt ^{IV} (dach)(OAc) ₃ (OTFAc)](3)	9.5
[Pt ^{IV} (dach)(OPr) ₃ (OTFAc)](4)	3.1
JM216	1.2
Carboplatin	10.2

platinum(IV) is a distorted octahedral arrangement with three propionate and one trifluoroacetate groups (O(5)–Pt–O(7) = 80.7° and O(1)–Pt–O(3) = 174.4(6)°). The four carboxylate ligands occupy their positions with intramolecular hydrogen bonds by the carbonyl groups with two amine groups (N(1)–O(2) = 2.701, N(2)–O(6) = 2.788, N(2)–O(4) = 2.724 and N(1)–O(8) = 2.738 Å).

The in vitro antitumor activity of the present Pt(IV) complexes are listed along with reference complexes in Table 3. The present complexes were designed to have the optimal range of partition coefficient (log*P*) between those of [Pt^{IV}(dach)(OPr)₄] (log*P* = 0.18) and of [Pt^{IV}(dach)(OBU)₄] (OBU = butanoyl, log*P* = 1.54) [8,11]. The in vitro cytotoxicity against the murine leukemia L1210 cell line shows that the present complexes exhibit higher activity than carboplatin and comparable activity to JM216.

In conclusion, we have prepared mono(trifluoroacetato)platinum(IV) complexes in the presence of a base, and the structure was confirmed by various physico-chemical analyses including X-ray crystallography. Their in vitro antitumor activity was found to be higher than carboplatin and their in vivo activity is to be assayed. The methodology to obtain a variety of the platinum(IV) complexes with a wide range of lipophilicity or water solubility was further extended and conformed to

the previously proposed mechanism involving proton assisted nucleophilic substitution of the platinum(IV) complexes.

Acknowledgement

This research was supported by Korea Research Foundation Grant (KRF-2001-042-D0054).

Appendix A. Supporting information

Tables of crystallographic details, non-hydrogen positional parameters, bond distances and angles, anisotropic thermal parameters of non-hydrogen atoms, and hydrogen coordinates and isotropic thermal parameters for complex **4** (8 pages). Ordering information is given on any current masthead page. Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.ica.2004.07.037](https://doi.org/10.1016/j.ica.2004.07.037).

References

- [1] B.K. Keppler, *Metal Complexes in Cancer Therapy*, VCH, Weinheim, 1993.
- [2] D.D. Von Hoff, R. Schilsky, C.M. Reichert, R.L. Reddick, M. Rozenzweig, R.C. Young, F.M. Muggia, *Cancer Treat. Rep.* 63 (1979) 1527.
- [3] W.B. Pratt, R.W. Ruddon, W.D. Ensminger, J. Maybaum, *The Anticancer Drugs*, second ed., Oxford University Press, New York, 1994, pp. 133–168.
- [4] Z.H. Siddik, G.S. Hagopian, G. Thai, Tomisaki, S. Toyomasu, T.A.R. Khokhar, *J. Inorg. Chem.* 77 (1999) 65.
- [5] L. Cai, K. Lim, S. Ren, R.S. Cadena, W.T. Beck, *J. Med. Chem.* 44 (2001) 2959.
- [6] M.D. Hall, T.W. Hambley, *Coord. Chem. Rev.* 232 (2002) 49.
- [7] C.F.J. Barnard, J.F. Vollano, P.A. Chaloner, S.Z. Dewa, *Inorg. Chem.* 35 (1996) 3280.
- [8] Y.A. Lee, S.S. Lee, K.M. Kim, C.O. Lee, Y.S. Sohn, *J. Med. Chem.* 43 (7) (2000) 1409.
- [9] R. Kizu, T. Nakanishi, M. Miyazaki, T. Tashiro, M. Noji, A. Matsuzawa, M. Eriguchi, Y. Takeda, N. Akiyama, Y. Kidani, *Anti-Cancer Drugs* 7 (1996) 248.
- [10] R. Song, K.M. Kim, Y.S. Sohn, *Inorg. Chimica. Acta* 338 (2002).
- [11] R. Song, S.Y. Park, Y.-S. Kim, Y.M. Kim, S.J. Kim, B.T. Ahn, Y.S. Sohn, *J. Inorg. Biochem.* 96 (2003) 339.
- [12] Y.S. Sohn, H. Baek, Y.H. Cho, Y.A. Lee, O.S. Jung, C.O. Lee, Y.S. Kim, *Int. J. Pharm.* 153 (1997) 79.
- [13] G.M. Sheldrick, *SHELXS-97: A Program for Structure Determination*, University of Göttingen, Germany, 1997.
- [14] G.M. Sheldrick, *SHELXL-97: A Program for Structure Refinement*, University of Göttingen, Germany, 1997.
- [15] S. Shamsuddin, C.C. Santillan, J.L. Stark, K.H. Whitmire, Z.H. Siddik, A.R. Khokhar, *J. Inorg. Biochem.* 71 (1998) 29.
- [16] M. Galanski, B.K. Keppler, *Inorg. Chim. Acta* 265 (1997) 271.
- [17] K.M. Kim, Y.A. Lee, S.S. Lee, Y.S. Sohn, *Inorg. Chim. Acta* 292 (1999) 52.

```
#####
#
# This file contains crystal structure data downloaded from the
# Cambridge Structural Database (CSD) hosted by the Cambridge
# Crystallographic Data Centre (CCDC).
#
# Full information about CCDC data access policies and citation
# guidelines are available at http://www.ccdc.cam.ac.uk/access/V1
#
# Audit and citation data items may have been added by the CCDC.
# Please retain this information to preserve the provenance of
# this file and to allow appropriate attribution of the data.
#
#####

data_complex_4
_audit_block_doi          10.5517/cc8zpsp
_database_code_depnum_ccdc_archive 'CCDC 267833'
loop_
_citation_id
_citation_doi
_citation_year
1 10.1016/j.ica.2004.07.037 2005
_audit_update_record
;
2005-04-04 deposited with the CCDC.      2025-07-11 downloaded from the
CCDC.
;

_audit_creation_method      'Manually at CCDC'

_chemical_name_systematic
;
?
;
_chemical_name_common        ?
_chemical_formula_sum        'C17 H29 F3 N2 O8 Pt'
_chemical_formula_moiety      '? '

_chemical_formula_weight      641.51

_symmetry_cell_setting        monoclinic
_symmetry_space_group_name_H-M P2(1)/c
_symmetry_Int_Tables_number    14

loop_
_symmetry_equiv_pos_as_xyz
?
?
?

_cell_length_a                10.891(2)
_cell_length_b                17.343(4)
_cell_length_c                12.143(4)
```

```

_cell_angle_alpha          90.00
_cell_angle_beta           95.33(2)
_cell_angle_gamma          90.00
_cell_volume               2283.5(9)
_cell_formula_units_Z      4

_exptl_crystal_density_diffn  1.866

_diffn_ambient_temperature  293(2)

_diffn_source              ?

_diffn_radiation_probe     x-ray

_chemical_melting_point    ?

_exptl_crystal_description ?
_exptl_crystal_colour      ' ? '
_exptl_special_details
;
#include here any other information not included in the CIF
;

_refine_ls_R_factor_gt     0.0662

loop_
_atom_site_label
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
Pt(1) 0.0434(1) 0.0452(1) 0.7565(1) 0.049(1)
N(1) 0.0471(16) -0.0537(8) 0.8440(13) 0.051(4)
N(2) -0.0873(15) -0.0138(10) 0.6563(13) 0.052(4)
O(1) 0.1689(13) 0.0112(8) 0.6524(11) 0.057(3)
O(2) 0.2748(16) -0.0681(11) 0.7752(16) 0.091(6)
O(3) -0.0756(15) 0.0894(9) 0.8564(12) 0.066(4)
O(4) -0.2387(15) 0.0943(13) 0.7344(16) 0.108(7)
O(5) 0.0374(14) 0.1475(8) 0.6792(12) 0.062(4)
O(6) -0.0437(19) 0.1070(9) 0.5136(13) 0.086(5)
O(7) 0.1847(13) 0.0975(8) 0.8441(11) 0.061(4)
O(8) 0.1666(16) 0.0393(9) 1.0017(14) 0.078(5)
C(1) -0.0200(20) -0.1139(15) 0.7810(20) 0.082(8)
C(2) -0.0460(30) -0.1876(14) 0.8420(20) 0.080(7)
C(3) -0.1180(40) -0.2435(17) 0.7710(30) 0.119(12)
C(4) -0.2210(30) -0.2110(17) 0.7110(30) 0.120(12)
C(5) -0.2010(20) -0.1403(15) 0.6450(20) 0.083(7)
C(6) -0.1210(20) -0.0850(16) 0.7210(20) 0.087(8)
C(7) 0.2579(18) -0.0348(13) 0.6854(17) 0.079(8)
C(8) 0.3583(16) -0.0414(12) 0.6090(15) 0.122(13)
C(9) -0.1870(20) 0.1090(12) 0.8200(20) 0.069(7)
C(10) -0.2410(30) 0.1535(17) 0.9190(30) 0.108(11)
C(11) -0.3530(30) 0.1900(30) 0.8780(20) 0.145(18)

```

C(12) -0.0110(30) 0.1586(15) 0.5806(18) 0.077(7)
 C(13) -0.0300(30) 0.2435(15) 0.5530(20) 0.091(9)
 C(14) -0.1190(40) 0.2750(19) 0.6150(30) 0.157(18)
 C(15) 0.2215(19) 0.0806(13) 0.9414(19) 0.058(5)
 C(16) 0.3470(20) 0.1170(20) 0.9830(20) 0.133(15)
 C(17) 0.4220(30) 0.0830(30) 1.0770(30) 0.152(16)
 F(1) 0.3910(40) -0.1234(13) 0.5930(30) 0.370(30)
 F(2) 0.4690(20) 0.0000(30) 0.6580(20) 0.370(30)
 F(3) 0.3190(30) -0.0076(19) 0.4993(17) 0.310(30)
 H(1A) 0.1258 -0.0687 0.8606 0.062
 H(1B) 0.0131 -0.0458 0.9078 0.062
 H(2A) -0.1543 0.0159 0.6401 0.063
 H(2B) -0.0568 -0.0275 0.5928 0.063
 H(1) 0.0351 -0.1299 0.7261 0.099
 H(2C) 0.0314 -0.2109 0.8706 0.096
 H(2D) -0.0914 -0.1750 0.9052 0.096
 H(3A) -0.0647 -0.2658 0.7199 0.143
 H(3B) -0.1449 -0.2849 0.8171 0.143
 H(4A) -0.2813 -0.1988 0.7629 0.144
 H(4B) -0.2579 -0.2500 0.6611 0.144
 H(5A) -0.2795 -0.1162 0.6215 0.099
 H(5B) -0.1600 -0.1534 0.5804 0.099
 H(6) -0.1745 -0.0663 0.7750 0.104
 H(10A) -0.2560 0.1177 0.9780 0.129
 H(10B) -0.1820 0.1919 0.9496 0.129
 H(11A) -0.3861 0.2186 0.9357 0.218
 H(11B) -0.4115 0.1514 0.8503 0.218
 H(11C) -0.3374 0.2243 0.8184 0.218
 H(13A) 0.0470 0.2711 0.5680 0.109
 H(13B) -0.0563 0.2489 0.4743 0.109
 H(14A) -0.1337 0.3278 0.5937 0.235
 H(14B) -0.0910 0.2724 0.6919 0.235
 H(14C) -0.1947 0.2463 0.6014 0.235
 H(16A) 0.3966 0.1182 0.9206 0.159
 H(16B) 0.3310 0.1703 1.0018 0.159
 H(17A) 0.4982 0.1104 1.0893 0.228
 H(17B) 0.4387 0.0298 1.0606 0.228
 H(17C) 0.3781 0.0855 1.1416 0.228

loop_
 _geom_bond_atom_site_label_1
 _geom_bond_atom_site_label_2
 _geom_bond_distance
 Pt(1) O(7) 2.004(14)
 Pt(1) O(5) 2.006(13)
 Pt(1) O(3) 2.008(14)
 Pt(1) N(1) 2.016(14)
 Pt(1) O(1) 2.033(13)
 Pt(1) N(2) 2.056(16)
 N(1) C(1) 1.45(3)
 N(2) C(6) 1.52(3)
 O(1) C(7) 1.29(2)
 O(2) C(7) 1.23(3)
 O(3) C(9) 1.30(2)

O(4) C(9) 1.17(3)
 O(5) C(12) 1.28(3)
 O(6) C(12) 1.24(3)
 O(7) C(15) 1.25(2)
 O(8) C(15) 1.22(3)
 C(1) C(6) 1.35(3)
 C(1) C(2) 1.52(3)
 C(2) C(3) 1.47(3)
 C(3) C(4) 1.40(4)
 C(4) C(5) 1.49(4)
 C(5) C(6) 1.54(3)
 C(7) C(8) 1.5021(13)
 C(8) F(3) 1.4819(13)
 C(8) F(1) 1.4820(13)
 C(8) F(2) 1.4821(13)
 C(9) C(10) 1.59(3)
 C(10) C(11) 1.43(4)
 C(12) C(13) 1.52(3)
 C(13) C(14) 1.40(4)
 C(15) C(16) 1.54(3)
 C(16) C(17) 1.47(4)

loop_
 _atom_site_aniso_label
 _atom_site_aniso_U_11
 _atom_site_aniso_U_22
 _atom_site_aniso_U_33
 _atom_site_aniso_U_23
 _atom_site_aniso_U_13
 _atom_site_aniso_U_12
 #U11 U22 U33 U23 U13 U12
 Pt(1) 0.059(1) 0.042(1) 0.043(1) 0.000(1) -0.003(1) 0.001(1)
 N(1) 0.065(10) 0.036(9) 0.049(9) 0.006(7) -0.020(8) 0.004(8)
 N(2) 0.056(10) 0.056(10) 0.043(9) 0.001(8) -0.004(8) 0.001(8)
 O(1) 0.060(8) 0.054(8) 0.059(9) 0.002(7) 0.019(7) 0.008(7)
 O(2) 0.074(11) 0.107(15) 0.095(14) 0.041(11) 0.019(10) 0.025(10)
 O(3) 0.087(11) 0.056(9) 0.055(9) -0.020(7) 0.010(8) 0.005(8)
 O(4) 0.053(9) 0.170(20) 0.091(13) -0.044(14) -0.037(9) 0.037(11)
 O(5) 0.080(9) 0.040(8) 0.065(9) 0.013(7) 0.000(8) 0.004(7)
 O(6) 0.134(16) 0.064(11) 0.058(10) 0.004(9) -0.010(10) -0.017(10)
 O(7) 0.070(9) 0.064(10) 0.051(9) 0.012(7) 0.006(7) -0.015(8)
 O(8) 0.082(11) 0.080(12) 0.067(10) 0.010(9) -0.015(9) -0.036(10)
 C(1) 0.090(17) 0.077(18) 0.078(16) 0.017(14) -0.003(13) -0.051(15)
 C(2) 0.110(20) 0.057(14) 0.071(16) 0.009(12) -0.012(14) -0.009(14)
 C(3) 0.170(30) 0.080(20) 0.100(20) -0.008(16) -0.060(20) -0.030(20)
 C(4) 0.130(30) 0.100(20) 0.120(20) 0.020(20) -0.050(20) -0.070(20)
 C(5) 0.089(17) 0.076(17) 0.080(17) -0.025(14) -0.004(14) -0.029(15)
 C(6) 0.073(16) 0.099(19) 0.078(16) 0.017(16) -0.047(14) -0.009(15)
 C(7) 0.110(20) 0.073(17) 0.059(14) -0.015(13) 0.022(14) 0.033(15)
 C(8) 0.090(20) 0.140(30) 0.140(30) 0.060(30) 0.040(20) 0.030(20)
 C(9) 0.049(12) 0.039(12) 0.110(20) 0.007(12) -0.012(13) 0.003(10)
 C(10) 0.120(20) 0.090(20) 0.120(20) -0.006(18) 0.050(20) 0.058(19)
 C(11) 0.090(20) 0.260(50) 0.075(18) -0.060(20) -0.011(16) 0.100(30)
 C(12) 0.110(20) 0.075(17) 0.044(13) 0.018(12) 0.001(13) 0.016(15)

C(13) 0.110(20) 0.059(16) 0.100(20) 0.017(15) -0.029(17) 0.023(15)
 C(14) 0.250(50) 0.070(20) 0.140(30) 0.020(20) -0.020(30) 0.060(30)
 C(15) 0.057(13) 0.055(12) 0.060(14) -0.005(11) -0.007(11) -0.018(11)
 C(16) 0.088(19) 0.180(30) 0.120(20) 0.070(20) -0.081(19) -0.070(20)
 C(17) 0.120(30) 0.210(40) 0.120(30) -0.010(30) -0.050(20) -0.030(30)
 F(1) 0.410(70) 0.370(60) 0.380(60) 0.030(50) 0.300(60) 0.090(50)
 F(2) 0.110(20) 0.800(110) 0.220(40) 0.180(60) 0.050(20) -0.020(40)
 F(3) 0.330(50) 0.310(40) 0.320(40) 0.220(40) 0.240(40) 0.190(40)

loop_
 _geom_angle_atom_site_label_1
 _geom_angle_atom_site_label_2
 _geom_angle_atom_site_label_3
 _geom_angle
 O(7) Pt(1) O(3) 90.8(6)
 O(5) Pt(1) O(3) 87.2(6)
 O(7) Pt(1) N(1) 97.4(6)
 O(7) Pt(1) O(5) 80.7(6)
 O(5) Pt(1) N(1) 176.0(6)
 O(3) Pt(1) N(1) 89.3(7)
 O(7) Pt(1) O(1) 86.2(6)
 O(5) Pt(1) O(1) 87.7(6)
 O(3) Pt(1) O(1) 174.4(6)
 N(1) Pt(1) O(1) 95.7(7)
 O(7) Pt(1) N(2) 173.7(6)
 O(5) Pt(1) N(2) 100.0(6)
 O(3) Pt(1) N(2) 95.6(7)
 N(1) Pt(1) N(2) 82.4(6)
 O(1) Pt(1) N(2) 87.5(6)
 C(1) N(1) Pt(1) 110.5(13)
 C(6) N(2) Pt(1) 106.6(11)
 C(7) O(1) Pt(1) 121.1(12)
 C(9) O(3) Pt(1) 122.2(16)
 C(12) O(5) Pt(1) 124.4(16)
 C(15) O(7) Pt(1) 124.1(13)
 C(6) C(1) N(1) 111(2)
 C(6) C(1) C(2) 114(2)
 N(1) C(1) C(2) 117.0(19)
 C(3) C(2) C(1) 112(2)
 C(4) C(3) C(2) 113(3)
 C(3) C(4) C(5) 117(3)
 C(4) C(5) C(6) 107(2)
 C(1) C(6) N(2) 111(2)
 C(1) C(6) C(5) 119(3)
 N(2) C(6) C(5) 110.5(19)
 O(2) C(7) O(1) 127.6(16)
 O(2) C(7) C(8) 117.0(17)
 O(1) C(7) C(8) 115.1(16)
 F(3) C(8) F(1) 108.5(14)
 F(3) C(8) F(2) 108.8(14)
 F(1) C(8) F(2) 109.1(14)
 F(3) C(8) C(7) 111.2(12)

F(1) C(8) C(7) 110.4(12)
 F(2) C(8) C(7) 108.8(13)
 O(4) C(9) O(3) 127(2)
 O(4) C(9) C(10) 127(2)
 O(3) C(9) C(10) 106(2)
 C(11) C(10) C(9) 108(2)
 O(6) C(12) O(5) 125(2)
 O(6) C(12) C(13) 122(2)
 O(5) C(12) C(13) 113(2)
 C(14) C(13) C(12) 110(3)
 O(8) C(15) O(7) 125.0(19)
 O(8) C(15) C(16) 121(2)
 O(7) C(15) C(16) 114(2)
 C(17) C(16) C(15) 120(3)

loop_
 _geom_torsion_atom_site_label_1
 _geom_torsion_atom_site_label_2
 _geom_torsion_atom_site_label_3
 _geom_torsion_atom_site_label_4
 _geom_torsion
 _geom_torsion_site_symmetry_1
 _geom_torsion_site_symmetry_2
 _geom_torsion_site_symmetry_3
 _geom_torsion_site_symmetry_4
 _geom_torsion_publ_flag
 ? ? ? ? ? ? ? ? ? ?

#END
 #Attach the next CIF data_number_n+1 below