DIAGNOSTIC AND STATISTICAL MANUAL OF MENTAL DISORDERS

FIFTH EDITION

DSM-5™

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00001

American Psychiatric Association

Officers 2012–2013

PRESIDENT DILIP V. JESTE, M.D.
PRESIDENT-ELECT JEFFREY A. LIEBERMAN, M.D.
TREASURER DAVID FASSLER, M.D.
SECRETARY ROGER PEELE, M.D.

Assembly

SPEAKER R. SCOTT BENSON, M.D.
SPEAKER-ELECT MELINDA L. YOUNG, M.D.

Board of Trustees

JEFFREY AKAKA, M.D.
CAROL A. BERNSTEIN, M.D.
BRIAN CROWLEY, M.D.
ANITA S. EVERETT, M.D.
JEFFREY GELLER, M.D., M.P.H.
MARC DAVID GRAFF, M.D.
JAMES A. GREENE, M.D.
JUDITH F. KASHTAN, M.D.
MOLLY K. MCVOY, M.D.
JAMES E. NININGER, M.D.
JOHN M. OLDHAM, M.D.
ALAN F. SCHATZBERG, M.D.
ALIK S. WIDGE, M.D., PH.D.

ERIK R. VANDERLIP, M.D.,
MEMBER-IN-TRAINING TRUSTEE-ELECT

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00002

DIAGNOSTIC AND STATISTICAL MANUAL OF MENTAL DISORDERS,

FIFTH EDITION

DSM-5™

New School Library





Washington, DC
London, England

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00003

Copyright © 2013 American Psychiatric Association

DSM and DSM-5 are trademarks of the American Psychiatric Association. Use of these terms is prohibited without permission of the American Psychiatric Association.

ALL RIGHTS RESERVED. Unless authorized in writing by the APA, no part of this book may be reproduced or used in a manner inconsistent with the APA's copyright. This prohibition applies to unauthorized uses or reproductions in any form, including electronic applications.

Correspondence regarding copyright permissions should be directed to DSM Permissions, American Psychiatric Publishing, 1000 Wilson Boulevard, Suite 1825, Arlington, VA 22209-3901.

Manufactured in the United States of America on acid-free paper.

ISBN 978-0-89042-554-1 (Hardcover)

ISBN 978-0-89042-555-8 (Paperback)

American Psychiatric Association
1000 Wilson Boulevard
Arlington, VA 22209-3901
www.psych.org

The correct citation for this book is American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA, American Psychiatric Association, 2013.

Library of Congress Cataloging-in-Publication Data
Diagnostic and statistical manual of mental disorders : DSM-5. — 5th ed.
p. ; cm.
DSM-5
DSM-V
Includes index.
ISBN 978-0-89042-554-1 (hardcover : alk. paper) — ISBN 978-0-89042-555-8 (pbk. : alk. paper)
I. American Psychiatric Association. II. American Psychiatric Association. DSM-5 Task Force.
III. Title: DSM-5. IV. Title: DSM-V.
[DNLM: 1. Diagnostic and statistical manual of mental disorders. 5th ed. 2. Mental Disorders—classification. 3. Mental Disorders—diagnosis. WM 15]
RC455.2.C4
616.89'075—dc23

2013011061

British Library Cataloguing in Publication Data
A CIP record is available from the British Library.

Text Design—Tammy J. Cordova

Manufacturing—Edwards Brothers Malloy

RC
455
.2
.C4
D5A
2013

Contents

DSM-5 Classification xiii
Preface xli

Section I
DSM-5 Basics

Introduction 5
Use of the Manual 19
Cautionary Statement for Forensic Use of DSM-5 25

Section II
Diagnostic Criteria and Codes

Neurodevelopmental Disorders 31
Schizophrenia Spectrum and Other Psychotic Disorders 87
Bipolar and Related Disorders 123
Depressive Disorders 155
Anxiety Disorders. 189
Obsessive-Compulsive and Related Disorders 235
Trauma- and Stressor-Related Disorders 265
Dissociative Disorders 291
Somatic Symptom and Related Disorders 309
Feeding and Eating Disorders 329
Elimination Disorders 355
Sleep-Wake Disorders. 361
Sexual Dysfunctions 423
Gender Dysphoria 451

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00005

Disruptive, Impulse-Control, and Conduct Disorders 461
Substance-Related and Addictive Disorders 481
Neurocognitive Disorders. 591
Personality Disorders 645
Paraphilic Disorders 685
Other Mental Disorders 707
Medication-Induced Movement Disorders and Other Adverse Effects of Medication 709
Other Conditions That May Be a Focus of Clinical Attention . . 715

Section III
Emerging Measures and Models

Assessment Measures 733
Cultural Formulation 749
Alternative DSM-5 Model for Personality Disorders 761
Conditions for Further Study 783

Appendix

Highlights of Changes From DSM-IV to DSM-5 809
Glossary of Technical Terms 817
Glossary of Cultural Concepts of Distress 833
Alphabetical Listing of DSM-5 Diagnoses and Codes (ICD-9-CM and ICD-10-CM) 839
Numerical Listing of DSM-5 Diagnoses and Codes (ICD-9-CM) 863
Numerical Listing of DSM-5 Diagnoses and Codes (ICD-10-CM) 877
DSM-5 Advisors and Other Contributors 897

Index 917

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00006

Bipolar and Related Disorders

Bipolar and related disorders are separated from the depressive disorders in DSM-5 and placed between the chapters on schizophrenia spectrum and other psychotic disorders and depressive disorders in recognition of their place as a bridge between the two diagnostic classes in terms of symptomatology, family history, and genetics. The diagnoses included in this chapter are bipolar I disorder, bipolar II disorder, cyclothymic disorder, substance/medication-induced bipolar and related disorder, bipolar and related disorder due to another medical condition, other specified bipolar and related disorder, and unspecified bipolar and related disorder.

The bipolar I disorder criteria represent the modern understanding of the classic manic-depressive disorder or affective psychosis described in the nineteenth century, differing from that classic description only to the extent that neither psychosis nor the lifetime experience of a major depressive episode is a requirement. However, the vast majority of individuals whose symptoms meet the criteria for a fully syndromal manic episode also experience major depressive episodes during the course of their lives.

Bipolar II disorder, requiring the lifetime experience of at least one episode of major depression and at least one hypomanic episode, is no longer thought to be a "milder" condition than bipolar I disorder, largely because of the amount of time individuals with this condition spend in depression and because the instability of mood experienced by individuals with bipolar II disorder is typically accompanied by serious impairment in work and social functioning.

The diagnosis of cyclothymic disorder is given to adults who experience at least 2 years (for children, a full year) of both hypomanic and depressive periods without ever fulfilling the criteria for an episode of mania, hypomania, or major depression.

A large number of substances of abuse, some prescribed medications, and several medical conditions can be associated with manic-like phenomena. This fact is recognized in the diagnoses of substance/medication-induced bipolar and related disorder and bipolar and related disorder due to another medical condition.

The recognition that many individuals, particularly children and, to a lesser extent, adolescents, experience bipolar-like phenomena that do not meet the criteria for bipolar I, bipolar II, or cyclothymic disorder is reflected in the availability of the other specified bipolar and related disorder category. Indeed, specific criteria for a disorder involving short-duration hypomania are provided in Section III in the hope of encouraging further study of this disorder.

Bipolar I Disorder

Diagnostic Criteria

For a diagnosis of bipolar I disorder, it is necessary to meet the following criteria for a manic episode. The manic episode may have been preceded by and may be followed by hypomanic or major depressive episodes.

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00007

Manic Episode

A. A distinct period of abnormally and persistently elevated, expansive, or irritable mood and abnormally and persistently increased goal-directed activity or energy, lasting at least 1 week and present most of the day, nearly every day (or any duration if hospitalization is necessary).

B. During the period of mood disturbance and increased energy or activity, three (or more) of the following symptoms (four if the mood is only irritable) are present to a significant degree and represent a noticeable change from usual behavior:

1. Inflated self-esteem or grandiosity.
2. Decreased need for sleep (e.g., feels rested after only 3 hours of sleep).
3. More talkative than usual or pressure to keep talking.
4. Flight of ideas or subjective experience that thoughts are racing.
5. Distractibility (i.e., attention too easily drawn to unimportant or irrelevant external stimuli), as reported or observed.
6. Increase in goal-directed activity (either socially, at work or school, or sexually) or psychomotor agitation (i.e., purposeless non-goal-directed activity).
7. Excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments).

C. The mood disturbance is sufficiently severe to cause marked impairment in social or occupational functioning or to necessitate hospitalization to prevent harm to self or others, or there are psychotic features.

D. The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment) or to another medical condition.

Note: A full manic episode that emerges during antidepressant treatment (e.g., medication, electroconvulsive therapy) but persists at a fully syndromal level beyond the physiological effect of that treatment is sufficient evidence for a manic episode and, therefore, a bipolar I diagnosis.

Note: Criteria A–D constitute a manic episode. At least one lifetime manic episode is required for the diagnosis of bipolar I disorder.

Hypomanic Episode

A. A distinct period of abnormally and persistently elevated, expansive, or irritable mood and abnormally and persistently increased activity or energy, lasting at least 4 consecutive days and present most of the day, nearly every day.

B. During the period of mood disturbance and increased energy and activity, three (or more) of the following symptoms (four if the mood is only irritable) have persisted, represent a noticeable change from usual behavior, and have been present to a significant degree:

1. Inflated self-esteem or grandiosity.
2. Decreased need for sleep (e.g., feels rested after only 3 hours of sleep).
3. More talkative than usual or pressure to keep talking.
4. Flight of ideas or subjective experience that thoughts are racing.
5. Distractibility (i.e., attention too easily drawn to unimportant or irrelevant external stimuli), as reported or observed.
6. Increase in goal-directed activity (either socially, at work or school, or sexually) or psychomotor agitation.
7. Excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments).

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00008

C. The episode is associated with an unequivocal change in functioning that is uncharacteristic of the individual when not symptomatic.
D. The disturbance in mood and the change in functioning are observable by others.
E. The episode is not severe enough to cause marked impairment in social or occupational functioning or to necessitate hospitalization. If there are psychotic features, the episode is, by definition, manic.
F. The episode is not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment).

Note: A full hypomanic episode that emerges during antidepressant treatment (e.g., medication, electroconvulsive therapy) but persists at a fully syndromal level beyond the physiological effect of that treatment is sufficient evidence for a hypomanic episode diagnosis. However, caution is indicated so that one or two symptoms (particularly increased irritability, edginess, or agitation following antidepressant use) are not taken as sufficient for diagnosis of a hypomanic episode, nor necessarily indicative of a bipolar diathesis.

Note: Criteria A–F constitute a hypomanic episode. Hypomanic episodes are common in bipolar I disorder but are not required for the diagnosis of bipolar I disorder.

Major Depressive Episode

A. Five (or more) of the following symptoms have been present during the same 2-week period and represent a change from previous functioning; at least one of the symptoms is either (1) depressed mood or (2) loss of interest or pleasure.

   Note: Do not include symptoms that are clearly attributable to another medical condition.

   1. Depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, or hopeless) or observation made by others (e.g., appears tearful). (**Note:** In children and adolescents, can be irritable mood.)
   2. Markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated by either subjective account or observation).
   3. Significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month), or decrease or increase in appetite nearly every day. (**Note:** In children, consider failure to make expected weight gain.)
   4. Insomnia or hypersomnia nearly every day.
   5. Psychomotor agitation or retardation nearly every day (observable by others; not merely subjective feelings of restlessness or being slowed down).
   6. Fatigue or loss of energy nearly every day.
   7. Feelings of worthlessness or excessive or inappropriate guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick).
   8. Diminished ability to think or concentrate, or indecisiveness, nearly every day (either by subjective account or as observed by others).
   9. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.

B. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.
C. The episode is not attributable to the physiological effects of a substance or another medical condition.

Note: Criteria A–C constitute a major depressive episode. Major depressive episodes are common in bipolar I disorder but are not required for the diagnosis of bipolar I disorder.

Note: Responses to a significant loss (e.g., bereavement, financial ruin, losses from a natural disaster, a serious medical illness or disability) may include the feelings of intense

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00009

sadness, rumination about the loss, insomnia, poor appetite, and weight loss noted in Criterion A, which may resemble a depressive episode. Although such symptoms may be understandable or considered appropriate to the loss, the presence of a major depressive episode in addition to the normal response to a significant loss should also be carefully considered. This decision inevitably requires the exercise of clinical judgment based on the individual's history and the cultural norms for the expression of distress in the context of loss.[1]

Bipolar I Disorder

A. Criteria have been met for at least one manic episode (Criteria A–D under "Manic Episode" above).

B. The occurrence of the manic and major depressive episode(s) is not better explained by schizoaffective disorder, schizophrenia, schizophreniform disorder, delusional disorder, or other specified or unspecified schizophrenia spectrum and other psychotic disorder.

Coding and Recording Procedures

The diagnostic code for bipolar I disorder is based on type of current or most recent episode and its status with respect to current severity, presence of psychotic features, and remission status. Current severity and psychotic features are only indicated if full criteria are currently met for a manic or major depressive episode. Remission specifiers are only indicated if the full criteria are not currently met for a manic, hypomanic, or major depressive episode. Codes are as follows:

Bipolar I disorder	Current or most recent episode manic	Current or most recent episode hypomanic*	Current or most recent episode depressed	Current or most recent episode unspecified**
Mild (p. 154)	296.41 (F31.11)	NA	296.51 (F31.31)	NA
Moderate (p. 154)	296.42 (F31.12)	NA	296.52 (F31.32)	NA
Severe (p. 154)	296.43 (F31.13)	NA	296.53 (F31.4)	NA

[1] In distinguishing grief from a major depressive episode (MDE), it is useful to consider that in grief the predominant affect is feelings of emptiness and loss, while in MDE it is persistent depressed mood and the inability to anticipate happiness or pleasure. The dysphoria in grief is likely to decrease in intensity over days to weeks and occurs in waves, the so-called pangs of grief. These waves tend to be associated with thoughts or reminders of the deceased. The depressed mood of a MDE is more persistent and not tied to specific thoughts or preoccupations. The pain of grief may be accompanied by positive emotions and humor that are uncharacteristic of the pervasive unhappiness and misery characteristic of a major depressive episode. The thought content associated with grief generally features a preoccupation with thoughts and memories of the deceased, rather than the self-critical or pessimistic ruminations seen in a MDE. In grief, self-esteem is generally preserved, whereas in a MDE, feelings of worthlessness and self-loathing are common. If self-derogatory ideation is present in grief, it typically involves perceived failings vis-à-vis the deceased (e.g., not visiting frequently enough, not telling the deceased how much he or she was loved). If a bereaved individual thinks about death and dying, such thoughts are generally focused on the deceased and possibly about "joining" the deceased, whereas in a major depressive episode such thoughts are focused on ending one's own life because of feeling worthless, undeserving of life, or unable to cope with the pain of depression.

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00010

Bipolar I disorder	Current or most recent episode manic	Current or most recent episode hypomanic*	Current or most recent episode depressed	Current or most recent episode unspecified**
With psychotic features*** (p. 152)	296.44 (F31.2)	NA	296.54 (F31.5)	NA
In partial remission (p. 154)	296.45 (F31.73)	296.45 (F31.73)	296.55 (F31.75)	NA
In full remission (p. 154)	296.46 (F31.74)	296.46 (F31.74)	296.56 (F31.76)	NA
Unspecified	296.40 (F31.9)	296.40 (F31.9)	296.50 (F31.9)	NA

*Severity and psychotic specifiers do not apply; code 296.40 (F31.0) for cases not in remission.
**Severity, psychotic, and remission specifiers do not apply. Code 296.7 (F31.9).
***If psychotic features are present, code the "with psychotic features" specifier irrespective of episode severity.

In recording the name of a diagnosis, terms should be listed in the following order: bipolar I disorder, type of current or most recent episode, severity/psychotic/remission specifiers, followed by as many specifiers without codes as apply to the current or most recent episode.

Specify:

With anxious distress (p. 149)
With mixed features (pp. 149–150)
With rapid cycling (pp. 150–151)
With melancholic features (p. 151)
With atypical features (pp. 151–152)
With mood-congruent psychotic features (p. 152)
With mood-incongruent psychotic features (p. 152)
With catatonia (p. 152). **Coding note:** Use additional code 293.89 (F06.1).
With peripartum onset (pp. 152–153)
With seasonal pattern (pp. 153–154)

Diagnostic Features

The essential feature of a manic episode is a distinct period during which there is an abnormally, persistently elevated, expansive, or irritable mood and persistently increased activity or energy that is present for most of the day, nearly every day, for a period of at least 1 week (or any duration if hospitalization is necessary), accompanied by at least three additional symptoms from Criterion B. If the mood is irritable rather than elevated or expansive, at least four Criterion B symptoms must be present.

Mood in a manic episode is often described as euphoric, excessively cheerful, high, or "feeling on top of the world." In some cases, the mood is of such a highly infectious quality that it is easily recognized as excessive and may be characterized by unlimited and haphazard enthusiasm for interpersonal, sexual, or occupational interactions. For example, the individual may spontaneously start extensive conversations with strangers in public. Often the predominant mood is irritable rather than elevated, particularly when the individual's wishes are denied or if the individual has been using substances. Rapid shifts in mood over brief periods of time may occur and are referred to as lability (i.e., the alterna-

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00011

Unspecified Bipolar and Related Disorder

296.80 (F31.9)

This category applies to presentations in which symptoms characteristic of a bipolar and related disorder that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning predominate but do not meet the full criteria for any of the disorders in the bipolar and related disorders diagnostic class. The unspecified bipolar and related disorder category is used in situations in which the clinician chooses *not* to specify the reason that the criteria are not met for a specific bipolar and related disorder, and includes presentations in which there is insufficient information to make a more specific diagnosis (e.g., in emergency room settings).

Specifiers for Bipolar and Related Disorders

Specify if:

With anxious distress: The presence of at least two of the following symptoms during the majority of days of the current or most recent episode of mania, hypomania, or depression:

1. Feeling keyed up or tense.
2. Feeling unusually restless.
3. Difficulty concentrating because of worry.
4. Fear that something awful may happen.
5. Feeling that the individual might lose control of himself or herself.

Specify current severity:

Mild: Two symptoms.

Moderate: Three symptoms.

Moderate-severe: Four or five symptoms.

Severe: Four or five symptoms with motor agitation.

Note: Anxious distress has been noted as a prominent feature of both bipolar and major depressive disorder in both primary care and specialty mental health settings. High levels of anxiety have been associated with higher suicide risk, longer duration of illness, and greater likelihood of treatment nonresponse. As a result, it is clinically useful to specify accurately the presence and severity levels of anxious distress for treatment planning and monitoring of response to treatment.

With mixed features: The mixed features specifier can apply to the current manic, hypomanic, or depressive episode in bipolar I or bipolar II disorder:

Manic or hypomanic episode, with mixed features:

A. Full criteria are met for a manic episode or hypomanic episode, and at least three of the following symptoms are present during the majority of days of the current or most recent episode of mania or hypomania:

   1. Prominent dysphoria or depressed mood as indicated by either subjective report (e.g., feels sad or empty) or observation made by others (e.g., appears tearful).
   2. Diminished interest or pleasure in all, or almost all, activities (as indicated by either subjective account or observation made by others).
   3. Psychomotor retardation nearly every day (observable by others; not merely subjective feelings of being slowed down).

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00012

4. Fatigue or loss of energy.
5. Feelings of worthlessness or excessive or inappropriate guilt (not merely self-reproach or guilt about being sick).
6. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.

B. Mixed symptoms are observable by others and represent a change from the person's usual behavior.

C. For individuals whose symptoms meet full episode criteria for both mania and depression simultaneously, the diagnosis should be manic episode, with mixed features, due to the marked impairment and clinical severity of full mania.

D. The mixed symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, other treatment).

Depressive episode, with mixed features:

A. Full criteria are met for a major depressive episode, and at least three of the following manic/hypomanic symptoms are present during the majority of days of the current or most recent episode of depression:

1. Elevated, expansive mood.
2. Inflated self-esteem or grandiosity.
3. More talkative than usual or pressure to keep talking.
4. Flight of ideas or subjective experience that thoughts are racing.
5. Increase in energy or goal-directed activity (either socially, at work or school, or sexually).
6. Increased or excessive involvement in activities that have a high potential for painful consequences (e.g., engaging in unrestrained buying sprees, sexual indiscretions, or foolish business investments).
7. Decreased need for sleep (feeling rested despite sleeping less than usual; to be contrasted with insomnia).

B. Mixed symptoms are observable by others and represent a change from the person's usual behavior.

C. For individuals whose symptoms meet full episode criteria for both mania and depression simultaneously, the diagnosis should be manic episode, with mixed features.

D. The mixed symptoms are not attributable to the physiological effects of a substance (e.g., a drug of abuse, a medication, or other treatment).

Note: Mixed features associated with a major depressive episode have been found to be a significant risk factor for the development of bipolar I or bipolar II disorder. As a result, it is clinically useful to note the presence of this specifier for treatment planning and monitoring of response to treatment.

With rapid cycling (can be applied to bipolar I or bipolar II disorder): Presence of at least four mood episodes in the previous 12 months that meet the criteria for manic, hypomanic, or major depressive episode.

Note: Episodes are demarcated by either partial or full remissions of at least 2 months or a switch to an episode of the opposite polarity (e.g., major depressive episode to manic episode).

Note: The essential feature of a rapid-cycling bipolar disorder is the occurrence of at least four mood episodes during the previous 12 months. These episodes can occur in any combination and order. The episodes must meet both the duration and

Atossa Therapeutics, Inc. Exhibit 1009
Intas Pharms Ltd. v. Atossa Therapeutics, Inc.
PGR2026-00015
Page 00013