

A Glycosylation-Deficient Endothelial Cell Mutant with Modified Responses to Transforming Growth Factor- β and Other Growth Inhibitory Cytokines: Evidence for Multiple Growth Inhibitory Signal Transduction Pathways

Véronique Fafeur,* Bryan O'Hara, and Peter Böhlen

Medical Research Division, American Cyanamid, Pearl River, New York 10965

Submitted September 2, 1992; Accepted December 1, 1992

An endothelial cell line (M40) resistant to growth inhibition by transforming growth factor-beta type 1 (TGF β 1) was isolated by chemical mutagenesis and growth in the presence of TGF β 1. Like normal endothelial cells, this mutant is characterized by high expression of type II TGF β receptor and low expression of type I TGF β receptor. However, the mutant cells display a type II TGF β receptor of reduced molecular weight as a result of a general defect in N-glycosylation of proteins. The alteration does not impair TGF β 1 binding to cell surface receptors or the ability of TGF β 1 to induce fibronectin or plasminogen activator inhibitor-type I production. M40 cells were also resistant to growth inhibition by tumor necrosis factor alpha (TNF α) and interleukin-1 alpha (IL-1 α) but were inhibited by interferon- γ (IFN γ) and heparin. These results imply that TGF β 1, TNF α , and IL-1 α act through signal transducing pathways that are separate from pathways for IFN γ and heparin. Basic fibroblast growth factor was still mitogenic for M40, further suggesting that TGF β 1, TNF α , and IL-1 α act by direct inhibition of cell growth rather than by interfering with growth stimulatory pathways.

INTRODUCTION

The regulation of new capillary blood vessel formation is thought to be under the control of growth-stimulatory and -inhibitory factors (Folkman and Klagsbrun, 1987). In vitro, these factors can modulate endothelial cell proliferation, migration, and synthesis of specific proteins. Although considerable progress has been made in understanding the mechanism of action of growth-stimulatory factors, most notably basic fibroblast growth factor (bFGF),¹ the mechanisms underlying growth inhibition remain more elusive.

* Present address: Laboratory of Molecular Oncology, Pasteur Institute of Lille, Lille, France

¹ Abbreviations used: bFGF, basic fibroblast growth factor; DMEM, Dulbecco's modified Eagle's medium; DSS, disuccinimidyl suberate; ECM, extracellular matrix; EMS, ethyl methanesulfonate; HEPES, N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid; IFN γ , interferon-gamma; IL-1 α , interleukin-1 alpha; PAI-1, plasminogen activator inhibitor-type I; PBS, phosphate-buffered saline; PMSF, phenylmethylsulfonyl fluoride; SDS, sodium dodecyl sulfate; TGF β 1, transforming

The best known growth inhibitor, transforming growth factor-beta type 1 (TGF β 1), inhibits basal and bFGF-induced endothelial cell proliferation (Baird and Durkin, 1986; Fräter-Schroeder *et al.*, 1986) and cell migration (Heimark *et al.*, 1986; Müller *et al.*, 1987). TGF β 1 also stimulates extracellular matrix (ECM) formation in endothelial cells by inducing the synthesis of ECM proteins, such as fibronectin and types IV and V collagens (Müller *et al.*, 1987; Madri *et al.*, 1988). TGF β 1 further contributes to ECM formation by means of inhibiting the degradation of ECM proteins by specific proteases (increased production of plasminogen activator inhibitor-type 1 [PAI-1] and decreased plasminogen activator activity) (Saksela *et al.*, 1987; Pepper *et al.*, 1990). In most cells, TGF β 1 binds with high affinity to cell surface receptor glycoproteins of 53 kDa (type I) and 70–85 kDa (type II) and with lower affinity to 200–

growth factor-beta type 1; TNF α , tumor necrosis factor alpha; Tris, tris(hydroxymethyl)aminomethane.

400 kDa (type III) receptors (Cheifetz *et al.*, 1988; Segarini *et al.*, 1989). Endothelial cells exhibit only the type I and type II receptors, with the latter being predominantly expressed (Fafeur *et al.*, 1990). Although the mechanisms by which TGF β 1 exerts its effects on endothelial cells are still poorly understood, available evidence suggests that the growth inhibitory effect is not caused by interference with known mitogenic signaling pathways (reviewed in Massagué, 1990; Sporn and Roberts, 1990). For example, TGF β 1 blocks the proliferation of endothelial cells without altering the signaling capacity of bFGF receptors (Fafeur *et al.*, 1991). Recent data suggest that growth arrest occurs in late G1 and that earliest effects of TGF β 1 involve regulation of genes controlling cell adhesion and enhanced formation of ECM (Massagué, 1990) and that TGF β -induced growth inhibition may be linked to suppression of phosphorylation of the retinoblastoma gene product (Laiho *et al.*, 1990a). More recently, the type II TGF β receptor of a porcine renal epithelial cell line was cloned and characterized as a serine/threonine kinase (Lin *et al.*, 1992), thus lending support to the notion that this receptor is involved in signaling through a novel pathway.

Other cytokines, including tumor necrosis factor- α (TNF α) (Fräter-Schroeder *et al.*, 1987), interleukin-1 α (IL-1 α) (Cozzolino *et al.*, 1990), interleukin-6 (Böhlen *et al.*, 1987), and interferons (INFs) (Böhlen *et al.*, 1987; Friesel *et al.*, 1987) were also reported to inhibit endothelial cell proliferation. Although all these factors inhibit the growth of endothelial cells in a manner similar to that of TGF β , their signal transducing mechanisms are most likely different, as evidenced by the considerable structural differences in the cytoplasmic domains of their respective receptors (Aguet *et al.*, 1988; Sims *et al.*, 1988; Smith *et al.*, 1990; Lin *et al.*, 1992).

We found it intriguing that several cytokines with apparently different mechanisms of action would all inhibit potently the growth of endothelial cells. In an attempt to understand better the mechanisms by which growth inhibition of endothelial cells is effected, we tried to produce a cell mutant that is resistant to the growth inhibitory activity of TGF β 1. Here we report the isolation of a glycosylation-deficient endothelial cell mutant that is resistant to TGF β 1-induced growth inhibition while still being responsive to some other actions normally mediated by this cytokine. This mutant also proved to be resistant to growth inhibitory activities of TNF α and IL-1 α but not to that of IFN γ nor that of heparin, a well-documented but rather ill-characterized inhibitor of bovine endothelial cell growth.

MATERIALS AND METHODS

Chemicals

Porcine platelet TGF β 1 and human recombinant-IL-1 α and -TNF α were purchased from R&D Systems (Minneapolis, MN). Bovine brain

bFGF was isolated as described (Gospodarowicz *et al.*, 1984). A recombinant human bFGF analogue, Glu³⁵-Ser^{78,96}-bFGF, with biological activity being indistinguishable from native bFGF, was expressed in *Escherichia coli* and purified as described (Seddon *et al.*, 1991). Recombinant human IFN γ was a gift from Biogen (Geneva, Switzerland). Heparin was purchased from Hepar Industries (Franklin, OH). Ethyl methanesulfonate was purchased from Sigma (St. Louis, MO). Iodinated TGF β 1 was a gift of Dr. Anita Roberts (National Cancer Institute, Bethesda, MD) or purchased from New England Nuclear (Boston, MA) (both preparations had specific activities of 1.9–2.2 μ Ci/pmol). ³⁵S-protein labeling mix (77% ³⁵S-methionine, 18% ³⁵S-cysteine), D-[6-³H(N)]-glucosamine hydrochloride, and D-(2-³H)-mannose were purchased from New England Nuclear. Rabbit antiserum against human PAI-1 was kindly provided by Dr. Daniel Rifkin (New York University, NY), and rabbit antiserum against bovine fibronectin was purchased from Calbiochem (La Jolla, CA). Protein G-Agarose beads were from Genex (Gaithersburg, MD), and tunicamycin was purchased from Sigma.

Production of TGF β 1-Resistant Endothelial Cell Mutants

Endothelial cells from adult bovine aortic arch were prepared and grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% calf serum (Hyclone, Logan, UT) as previously described (Gospodarowicz *et al.*, 1984). Preconfluent endothelial cells (passage 5) maintained in DMEM/10% calf serum (10 ml/100-mm plates) were treated for 24 h with 0.4 μ l/ml ethyl methanesulfonate (EMS). The treatment resulted in the death of 80% of the cells. The remaining cells were washed and fed with fresh medium. Four days later, a total of 14×10^6 cells were split into 100-mm plates at 3×10^5 cells per plate in fresh medium. The following day, 2 ng/ml of TGF β 1 was added. The cells were fed fresh medium containing TGF β 1 every 5 d. After 2–3 wk of treatment, clonal cell lines were isolated by dilution cloning of the colony-derived cultures and expanded in the presence of 1 ng/ml bFGF and in the absence of TGF β 1. Cell lines were subjected to two additional rounds of selection for 7–10 d in TGF β 1, with 7- to 10-d periods of culture in TGF β 1-free/bFGF-containing medium between rounds of TGF β 1 selection. Resultant cell lines were periodically tested for resistance to TGF β 1-induced cell growth. Normal (or wild-type) endothelial cells that were not exposed to EMS and TGF β 1 selection but were grown in parallel with the mutant cells and were used at the same passage number (between passage 5 and 20) served as control cells. Additional control cells that were exposed or not exposed to EMS, and subjected to alternative treatments with TGF β 1 and bFGF as described above, were also cultured in parallel and used in most experiments. They gave comparable results to wild-type cells.

Cell Growth Assay

Cells were seeded in 24-multiwell plates at a density of 10 000 cells/0.5 ml of DMEM/10% calf serum per well. The following day, cells were fed growth factors or growth inhibitors. At different time intervals, cells were detached from the culture dishes with trypsin and counted in a Coulter particle counter (Coulter, Hialeah, FL).

Assays of Fibronectin and PAI-1 Production

Subconfluent cell cultures in 35-mm dishes were fed serum-free DMEM supplemented with or without 5 ng/ml of TGF β 1. After 15 h, the medium was replaced with cysteine- and methionine-free DMEM containing 30 μ Ci/ml ³⁵S-methionine/cysteine with or without 5 ng/ml of TGF β 1. After 5 h of labeling, the medium was removed and retained for further analysis. Cell monolayers were washed with ice-cold phosphate-buffered saline (PBS) and were extracted with 0.5 ml of a solution containing 1 mM urea, 1 mM dithiothreitol, 10 mM tris(hydroxymethyl)aminomethane (Tris)-HCl, pH 7.4, 10 mM EDTA, and 2 mM phenylmethylsulfonyl fluoride (PMSF) (Boehringer Mann-

heim, Indianapolis, IN) according to a published procedure (Ignatz and Massagué, 1986). The supernatants of the cell extracts and samples of the media were lyophilized and diluted with 0.5 ml of 0.5% Triton X-100. A portion of the samples was retained for further analysis by sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis. Fibronectin and PAI-1 were immunoprecipitated from the diluted extracts with anti-fibronectin and anti-PAI-1 antibodies, respectively, as follows. Protein G-Agarose beads (10 μ l) were incubated with an excess of anti-fibronectin or of anti-PAI-1 antibody solution (5 μ l) at 4°C for 1 h. The beads were then washed once in 0.5% Triton X-100, and the medium or cell samples were added. Incubation was allowed to take place for 2 h at 4°C. The beads were then washed four times in 0.5% Triton X-100, 0.15 M NaCl, 25 mM Tris-HCl, pH 7.4. Fibronectin or PAI-1 were released by heating at 100°C for 2 min in electrophoresis sample buffer and electrophoresed on a 6%-SDS polyacrylamide gel. Gels were fixed and treated with an autoradiographic enhancer (Amplify; Amersham, Arlington Heights, IL) according to the manufacturer's suggestion, dried, and exposed for autoradiography.

Receptor Binding Assay

Binding of ^{125} I-TGF β 1 to cell monolayers was performed as described previously (Fafeur *et al.*, 1990). Subconfluent monolayers of endothelial cells in 24-multiwell plates were incubated in binding buffer (DMEM, pH 7.4, containing 25 mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid [HEPES] and 0.1% bovine serum albumin) for 2 h at 37°C. Various concentrations of ^{125} I-TGF β 1, with or without 5 nM unlabeled TGF β 1, were added and incubation allowed for another 2 h at room temperature. Cultures were then washed four times with binding buffer and solubilized with a solution containing 1% Triton X-100, 10% glycerol, HEPES 25 mM, pH 7.4. Nonspecific binding measured in the presence of 5 nM unlabeled TGF β 1 was subtracted from total binding to determine specific binding. The data were analyzed by the method of Scatchard (1949).

Affinity-Labeling Assay

Affinity-labeling of subconfluent endothelial cell monolayers with ^{125} I-TGF β 1 and disuccinimidyl suberate (DSS) was performed as previously described (Fafeur *et al.*, 1990). Briefly, cultures were incubated with 100 pM ^{125} I-TGF β 1, in the presence or absence of 5 nM unlabeled TGF β 1, for 3–5 h at 4°C. Cultures were washed and cross-linked using 0.5 mM DSS. The cells were then detached from the plates with a cell scraper in ice-cold sucrose buffer containing 250 mM sucrose, 10 mM Tris-HCl, pH 7.4, 1 mM EDTA, 10 μ g/ml leupeptin (Boehringer Mannheim), and 1 mM PMSF. After centrifugation at 13 000 rpm, the pellets were resuspended in 50 mM Tris-HCl, pH 6.9, 2% SDS, 1% glycerol, and 10% 2-mercaptoethanol. The samples were boiled for 2 min and electrophoresed on 6% SDS polyacrylamide gels. Gels were stained for protein with Coomassie Blue, dried, and exposed to film with enhancing screens.

Metabolic Labeling With Radioactive Sugars

Subconfluent cell cultures in 35-mm wells were fed serum-free DMEM containing 50 μ Ci/ml of ^{3} H-mannose or 50 μ Ci/ml of ^{3} H-glucosamine. After 18 h at 37°C, cells were washed twice with PBS, extracted in sucrose buffer, and analyzed by 6% SDS polyacrylamide gel electrophoresis (see above, Affinity-Labeling Assays).

RESULTS

Isolation of an Endothelial Cell Mutant Resistant to TGF β 1-Induced Growth Inhibition

Previous studies have described the isolation of TGF β -resistant clones from a mink lung epithelial cell line

(Chinkers, 1987). We used the same strategy, which consisted in maintaining sparse cell cultures in the presence of 2 ng/ml TGF β 1 after chemical mutagenesis of the cells and isolating cell colonies that grew under these conditions. Although cells treated with TGF β 1 for 2–3 wk were usually able to resume proliferation after replating in fresh TGF β 1-free media, a significant portion of colonies failed to grow after cloning. We found that the rate of survival could be improved when isolated TGF β 1-resistant colonies were subsequently grown in the presence of bFGF (1 ng/ml). After treatment with EMS, TGF β 1-resistant clones of endothelial cells were obtained and taken through two additional cycles of TGF β 1-resistance selection interrupted by bFGF-growth stimulation periods. Of 35 TGF β 1-resistant colonies originally isolated from 14×10^6 mutagenized endothelial cells, only one clone (M40) remained completely resistant to TGF β 1 for over 6 mo in culture and was further characterized. Another TGF β 1-resistant clone had characteristics similar to those of the M40 clone (a modified TGF β type II receptor, see below) but had very poor growth properties and did not survive after several months in culture.

Exposure of sparsely cultured nonmutagenized control endothelial cells (4.2×10^6 cells tested) to 2 ng/ml TGF β 1 for 2–3 wk resulted in complete growth arrest of all cells, and no cells resistant to TGF β 1-induced growth inhibition emerged.

Defect in Growth-Inhibitory Response

The growth-inhibitory response of normal endothelial cells to TGF β 1 was half-maximal at 0.05 ng/ml and maximal at 1 ng/ml (Figure 1). The mutant M40 cells were unresponsive to TGF β 1 concentrations of up to 10 ng/ml (Figure 1), and the rate of proliferation was




Figure 1. Inhibition of cell proliferation by TGF β 1 in normal (○) and M40 mutant (●) endothelial cells. Duplicate samples were counted on day 6 after plating. Individual values deviated <10% from averages.

essentially the same in the presence or absence of 1 ng/ml of TGF β 1 (Figure 2). M40 cells, like normal endothelial cells, were responsive to the growth-promoting activity of bFGF (4 ng/ml) (Figure 2). TGF β 1, which is a potent inhibitor of bFGF-induced endothelial cell proliferation (Baird and Durkin, 1986; Fräter-Schroeder *et al.*, 1986), had no effect on bFGF-stimulated M40 cells (Figure 2).

Resistance to Other Endothelial Growth Inhibitors

In control experiments, we established that near maximal growth-inhibitory responses were obtained when endothelial cells were incubated with the inhibitors TNF α (1 ng/ml), IL-1 α (10 ng/ml), IFN γ (10 ng/ml), and heparin (10 μ g/ml) (Figure 3). At those concentrations, M40 cells were not inhibited by TNF α and IL-1 α (Figure 3, left), but normal growth inhibition occurred with IFN γ and heparin (Figure 3, right). This selective resistance to the various growth inhibitors was also found when M40 cells were growth stimulated by bFGF.

Morphological Changes in Response to TGF β 1

Normal endothelial cells show dramatic morphological changes in response to TGF β 1, involving an enlarged and flattened shape (Chinkers, 1987) (Figure 4). In contrast, M40 cells did not change shape in response to TGF β 1 (Figure 4). In the absence of TGF β 1, both cell types had a cobblestone appearance at confluence, characteristic of endothelial cells.

TGF β 1-Induced Synthesis of Extracellular Matrix Proteins

To determine whether the observed lack of growth inhibition by TGF β 1 is the result of a general impairment of the signal transducing mechanism, we tested whether other cellular responses to TGF β 1 were altered in the mutant endothelial cells. In this context, TGF β 1 can affect cell adhesion by enhancing synthesis and deposition of extracellular matrix components, such as fibronectin (Ignatz and Massagué, 1986; Müller *et al.*, 1987). We found that when biosynthetically labeled normal and mutant endothelial cells were treated with TGF β 1, increased secretion of a 220-kDa protein, identified as fibronectin by immunoprecipitation, was observed with both types of cells (Figure 5). Furthermore, in a similar experiment, the secretion of PAI-1, another TGF β 1-inducible protein thought to be involved in the regulation of extracellular matrix production (Sprengers and Kluft, 1987), was clearly stimulated in TGF β 1-treated normal and mutant cells (Figure 5). Comparable results were obtained when fibronectin and PAI-1 were determined in cell extracts rather than in conditioned media, suggesting that TGF β 1 stimulated the synthesis of these proteins both in normal and mutant cells. These data indicate that at least some TGF β 1-associated signaling




Figure 2. Growth curves of normal (top) and M40 (bottom) endothelial cells without added factors (○), in the presence of 1 ng/ml TGF β 1 (□), in the presence of 4 ng/ml bFGF (●), and with a combination of 1 ng/ml TGF β 1 and 4 ng/ml bFGF (■). Points represent duplicate samples. Individual values deviated <10% from averages. Although the results shown indicate that TGF β 1 alone slightly inhibited growth of M40 cells, this was not confirmed in two other experiments (e.g., see Figure 3; TGF β 1 has a slight stimulatory effect).

pathways remained intact in the M40 cell mutant. However, levels of fibronectin and PAI-1 were reproducibly lower in mutant cells, as compared with normal endothelial cells.

Characterization of TGF β Receptors

In previous experiments (Fafeur *et al.*, 1990), two types of TGF β receptors of ~65 and 85 kDa (type I and II receptors, respectively) were identified by affinity labeling techniques in bovine endothelial cells. Similar experiments performed in this study with normal and M40 cells showed that both cell types contained the same receptor types, although the type II receptor in normal endothelial cells appeared to be closer to 90 kDa (Figure 6). As previously observed, that 85–90 kDa type




Figure 3. Growth curves of normal (top) and mutant M40 (bottom) endothelial cells. Left, growth in the absence of added factor (○), TGF β 1 (1 ng/ml) (▲), TNF α (1 ng/ml) (■), and IL-1 α (10 ng/ml) (●) are shown. Right, growth curves in the absence of added factors (○), IFN γ (10 ng/ml) (□), and heparin (10 μ g/ml) (●). Cells were counted in duplicate after various times as indicated. Individual values deviated <10% from averages.

II receptor is the predominant form in normal cells, with the 65-kDa type I receptor sometimes being barely detectable (Fafeur *et al.*, 1990) (Figure 6). M40 shows the same receptor distribution; however, the type II receptor is clearly of lower molecular weight than that of normal cells (85 vs. 90 kDa). In addition, this type II receptor migrates as a much broader species, suggesting that there is heterogeneity in the glycosylation pattern of this receptor. The type I receptor also appears to be slightly smaller than the corresponding normal receptor (Figures 6 and 7).

Modification of TGF β Receptor Glycosylation

Exposure of cells to increasing concentrations of tunicamycin, a metabolic inhibitor of N-linked glycosylation (Fuhrmann *et al.*, 1985), resulted in a dose-dependent loss of the type II receptor in normal (90-kDa band)

and M40 (85-kDa band) cells. Instead, a band corresponding to a smaller cross-linked protein of $\sim$ 75 kDa appeared in tunicamycin-treated normal and mutant endothelial cells (Figure 7). These data indicate that a structural alteration of the type II receptor in mutant cells is located within the glycosylated part of this receptor. We were not able to conclude from our gel data whether tunicamycin treatment also caused a glycosylation defect in the type I receptor. Differences in migration of this receptor from tunicamycin-treated and control cells are expected to be small (5 kDa) (Cheifetz *et al.*, 1988) and, in our hands, were too small to be assessed reliably.

Given this evidence for a structural alteration of the predominant type II receptor in mutant endothelial cells, we investigated the binding characteristics of TGF β 1 by Scatchard analysis. TGF β 1 bound about equally to endothelial cell surface receptors of both normal and M40




Figure 4. Effect of TGF β 1 on the morphology of normal and mutant M40 endothelial cells. Cells were seeded at a low density (50 000 cells per 35-mm dish) and fed the following day with TGF β 1 (1 ng/ml) or control buffer (C). Phase-contrast photomicrographs of the cultures were taken after 5 d of growth. Magnification $\times 160$.

cells. Scatchard analysis of the data revealed that M40 cells expressed one high-affinity TGF β -binding site (25 200 sites/cell, $K_d = 7.0$ pM). These results are comparable with those obtained with normal endothelial cells (19 300 sites/cell, $K_d = 6.3$ pM) (Figure 8).

Evidence for a General Glycosylation Defect

We examined the ability of normal and M40 cells to incorporate radioactive sugars into the carbohydrate chains of proteins. When cells were labeled with ^3H -glucosamine, cellular proteins in M40 cells had a significantly higher mobility in SDS gel electrophoresis than proteins from normal cells (Figure 9). When cells were labeled with ^3H -mannose, labeled proteins were detectable at much lower levels in M40 cells than in normal cells (Figure 9). These results indicate that the M40 cells have a general glycosylation defect. The results, obtained when cells were incubated for 24 h with tunicamycin (Figure 7), further suggest that this gly-

cosylation defect affects at least the processing of N-linked carbohydrate chains of proteins.

DISCUSSION

The data presented here are discussed in the context of two main issues. First, it is intriguing that responses to TGF β 1 appear to be differentially impaired in M40 mutant endothelial cells. We observed that these cells cannot respond any longer to the normally growth inhibitory signal of TGF β 1 but remain able to signal the induction of the synthesis of specific proteins, fibronectin, and PAI-1, both well known for their TGF β 1-responsiveness.

Increasing evidence suggests that the type I and/or type II receptors, but not the type III receptor, are responsible for transducing the growth regulatory effect of TGF β 1. Some cell types, such as bovine endothelial cells, express only the type I and type II receptors and are responsive to various actions of TGF β 1, such as




Figure 5. Effect of TGF β 1 on the amounts of fibronectin and PAI-1 in the conditioned media of normal (WT) and mutant (M40) endothelial cells. Cells were incubated with (+) or without (-) 5 ng/ml of TGF β 1 for 15 h and labeled for the last 5 h with ^{35} S-methionine/cysteine mixture. Conditioned media were directly analyzed by 6% SDS-polyacrylamide gel electrophoresis, either before or after immunoprecipitation with antibodies against PAI-1 or fibronectin. Equal numbers of counts from control and TGF β -treated cultures ($\sim 9 \times 10^6$ cpm) were loaded into each lane. Note that the molecular weight of fibronectin from mutant cells is slightly lower than that of fibronectin synthesized by normal endothelial cells. An additional PAI-1-like immunoreactive protein of ~ 40 kDa was consistently detected in M40 cells but not in normal endothelial cells.

growth inhibition or induction of proteins involved in controlling extracellular matrix modeling. Previous experiments with mink lung epithelial cell mutants that lack either type I or type I and type II receptors showed that those cells were refractory to growth inhibition by TGF β 1 and other TGF β -mediated cellular responses, such as elevated production of fibronectin (Boyd and

Figure 6. Autoradiograms of ^{125} I-TGF β 1 cross-linked to normal (WT) and mutant (M40) endothelial cell surface proteins. Subconfluent cells were affinity-labeled by incubation with 100 pM ^{125} I-TGF β 1 alone (-) or in combination (+) with 5 nM unlabeled TGF β 1, followed by treatment with disuccinimidyl suberate. Membrane extracts were analyzed by electrophoresis on 6% SDS-polyacrylamide gels. The positions corresponding to molecular size markers are indicated. Arrows indicate bands corresponding to the 90- and 65-kDa labeled cross-linked proteins (the molecular size refers to the receptor-ligand complexes).





Figure 7. Effect of tunicamycin on TGF β receptor subtypes in normal (WT) and mutant (M40) endothelial cells. Cells were incubated for 24 h with various concentrations of tunicamycin (0.1, 0.3, 1 μ g/ml). Samples were then affinity-labeled with ^{125} I-TGF β 1 and processed as indicated in Figure 6.

Massagué, 1989; Laiho *et al.*, 1990b,c). It was concluded that the type I and type II receptors are necessary for those responses to TGF β 1 and it was suggested that both receptors may be coordinately expressed as part of a common signaling TGF β 1 receptor complex (Laiho *et al.*, 1990b,c). However, mutants lacking only the type




Figure 8. Scatchard plots of the binding of TGF β 1 to normal (O) and M40 (●) endothelial cells. Subconfluent cells (50 000 cells per well) were incubated with various concentrations of ^{125} I-TGF β 1 for 2 h at room temperature. Nonspecific binding was determined in the presence of 5 nM unlabeled TGF β 1.




Figure 9. Incorporation of radioactive sugars into proteins in normal (WT) and mutant (M40) endothelial cells. Cells were labeled for 18 h with either ^3H -glucosamine (^3H -Gluc) or with ^3H -mannose (^3H -Man). Samples were analyzed by 6% SDS-polyacrylamide gel electrophoresis. The volumes of cell extracts loaded were adjusted so that the two lanes would contain a similar amount of tritium counts (2×10^5 and 2.5×10^4 cpm per lane for ^3H -glucosamine and ^3H -mannose, respectively). Coomassie blue staining of gels indicated that normal and mutant cell lines contained similar amounts of protein.

II receptor have not been found in these studies, and it is therefore difficult to assess the precise role of the type II receptor in TGF β 1 signal transduction. Other evidence implicates more specifically the type II receptor in transmitting TGF β 1 growth inhibitory effects. By investigating the mechanism by which bFGF counteracts the ability of TGF β 1 to inhibit DNA synthesis in bovine endothelial cells, we found that expression of the type II receptor (which is the major TGF β 1 receptor on endothelial cells) is selectively down-regulated by bFGF (Fafeur *et al.*, 1990). Furthermore, a recent study showed that in a hybrid cell line obtained by fusion of two human carcinoma cell lines that are not growth-inhibited by TGF β 1, growth inhibitory response to this cytokine was restored (Geiser *et al.*, 1992). In contrast, TGF β 1-mediated activation of fibronectin, collagenase, and PAI-1 gene expression occurred in both hybrid and parental tumor cells. The hybrid cells expressed large quantities of the type II receptor as compared with the parental carcinoma cells. From these experiments, it was concluded that the type II receptor is necessary for mediating the effects of TGF β 1 on inhibition of growth but not on gene activation of the hybrid cells (Geiser *et al.*, 1992).

In M40 mutant cells, the type II receptor is clearly structurally altered by aberrant glycosylation. At present, we cannot rule out the possibility that the type I receptor has a similar glycosylation defect (see RESULTS). Regardless, the fact that by Scatchard analysis we found no reduction of TGF β 1 binding to the mutant cells suggests that TGF β 1 binds to glycosylation-defective TGF β receptors, as well as to their normal counterparts. Assuming that normal ligand-receptor binding is required for initiation of a normal signaling cascade, it is therefore likely that growth-inhibitory signaling is impaired downstream from receptor-ligand binding.

The glycosylation defect in M40 cells may interfere with other signaling mechanisms that are not directly related to the functionality of the TGF β receptor(s). The finding that synthesis of fibronectin and PAI-1 is still

induced in M40 cells shows that at least part of the TGF β 1-mediated signal transduction system is intact in M40 mutants. The differential effects could be the result of either separate transduction mechanisms mediated by type I and type II receptors, triggering either growth inhibition or specific protein synthesis, or of a common initial signal pathway mediated by these receptors, split downstream and leading to multiple actions. Whatever the mechanism, it is of interest that the results of the experiments with the hybrid carcinoma cell lines mentioned above (Geiser *et al.*, 1992) also suggest the existence of independent processes leading to growth inhibition and gene activation by TGF β 1. Furthermore, in Mv1Lu epithelial cells transfected to express the large T antigen, TGF β 1 is no longer growth inhibitory, whereas its effects on protooncogene c-jun induction and PAI-1, fibronectin, and thrombospondin synthesis are not altered as compared with parental cells (Laiho *et al.*, 1991).

The second issue concerns the differential response of mutant endothelial cells to various growth inhibitors, of which modes of action are largely unknown. The cytokines TGF β 1, TNF α , IL-1 α , and IFN γ all potentially inhibit growth of normal endothelial cells. In contrast, with the exception of IFN γ , these cytokines are without growth-inhibitory effects on M40 cells. The TNF, IL-1, and IFN γ receptors, like the TGF β receptors, are glycoproteins, and their structures are unrelated (Aguet *et al.*, 1988; Sims *et al.*, 1988; Smith *et al.*, 1990; Lin *et al.*, 1992). Furthermore, their cytoplasmic domains do not contain sequences typical of protein tyrosine kinases, and this suggests that these cytokines may act on cell growth through novel signal transducing pathways. A possible interpretation of our data is that separate transducing pathways are triggered by these cytokines and are all impaired in the mutant cells. Alternatively, the similar loss of growth inhibitory responses to TGF β 1, TNF α , and IL-1 α by M40 cells, both in the presence or absence of bFGF, suggests that the common growth inhibitory actions of these cytokines, triggered by unrelated ligand-specific receptors, may nevertheless be mediated by shared elements of their signaling pathways. In both cases, the general N-glycosylation defect in M40 cells suggests that protein(s) that must be N-glycosylated to function normally may be involved.

The fact that M40 cells remain growth inhibited by IFN γ suggests a separate inhibitory mechanism. Although very little is known about the signaling mechanisms of the various inhibitory factors, some other data support the concept of multiple inhibitory pathways. For example, increasing evidence suggests that IFN γ may interfere directly with growth stimulatory mechanisms induced by growth factors, including FGF (Friesel *et al.*, 1987; Hosang, 1988; Mundschauf and Faller, 1991), in contrast to TGF β 1, which does not appear to inhibit growth factor-induced events (see Sporn and Roberts, 1990).

Our data regarding the action of heparin on bovine endothelial cells also support the concept of multiple inhibitory mechanisms. Although the growth inhibitory activity of heparin on some cell types, e.g., smooth muscle cells and bovine endothelial cells, is well documented, its mechanism of action is unknown and generally not thought to be related to those of the cytokines. However, heparin action also may be mediated, at least in part, by typical signaling transduction pathways (Pukac *et al.*, 1992).

ACKNOWLEDGMENTS

We are very grateful to Dr. Anita Roberts, National Institutes of Health (Bethesda, MD), for providing ¹²⁵I-TGF β 1 and to Dr. Daniel Rifkin, New York University, for providing anti-PAI-1 antibodies. We thank Jean Coll, Vincent Laudet, Simon Saule, and Bernard Vandenbunder (Institut Pasteur, Lille, France) for helpful discussions.

REFERENCES

- Aguet, M., Dembic, Z., and Merlin, G. (1988). Molecular cloning and expression of the human interferon- γ receptor. *Cell* 55, 273–280.
- Baird, A., and Durkin, T. (1986). Inhibition of endothelial cell proliferation by type beta-transforming growth factor: interactions with acidic and basic fibroblast growth factor. *Biochem. Biophys. Res. Commun.* 138, 476–482.
- Böhlen, P., Fräter-Schroeder, M., Michel, T., and Jiang, Z.P. (1987). Inhibitors of endothelial cell proliferation. In: *Angiogenesis. Current Communications in Molecular Biology*, ed. D. Rifkin and M. Klagsbrun, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory, 119–124.
- Boyd, F.T., and Massagué, J. (1989). Transforming growth factor-beta inhibition of epithelial cell proliferation linked to the expression of a 53-kDa membrane receptor. *J. Biol. Chem.* 264, 2272–2278.
- Cheifetz, S., Andres, J.L., and Massagué, J. (1988). The transforming growth factor-beta receptor type III is a membrane proteoglycan. *J. Biol. Chem.* 263, 1689–1691.
- Chinkers, M. (1987). Isolation and characterization of mink lung epithelial cell mutants resistant to transforming growth factor beta. *J. Cell. Physiol.* 130, 1–5.
- Cozzolino, F., Torcia, M., Aldinucci, D., Ziche, M., Almerigogna, F., Bani, D., and Stern, D.M. (1990). Interleukin-1 is an autocrine regulator of human endothelial cell growth. *Proc. Natl. Acad. Sci. USA* 87, 6487–6491.
- Fafeur, V., Jiang, Z.P., and Böhlen, P. (1991). Signal transduction by bFGF, but not TGF β 1, involves arachidonic acid metabolism in endothelial cells. *J. Cell. Physiol.* 149, 277–283.
- Fafeur, V., Terman, B.I., Blum, J., and Böhlen, P. (1990). Basic FGF treatment of endothelial cells down-regulates the 85-kDa TGF β receptor subtype and decreases the growth inhibitory response to TGF β 1. *Growth Factors* 3, 237–245.
- Folkman, J., and Klagsbrun, M. (1987). Angiogenic factors. *Science* 235, 442–447.
- Fräter-Schroeder, M., Müller, G., Birchmeier, W., and Böhlen, P. (1986). Transforming growth factor-beta inhibits endothelial cell proliferation. *Biochem. Biophys. Res. Commun.* 137, 295–302.
- Fräter-Schroeder, M., Risau, W., Hallmann, R., Gautschi-Sova, P., and Böhlen, P. (1987). Tumor necrosis factor-alpha, a potent inhibitor of endothelial cell proliferation in vitro, is angiogenic in vivo. *Proc. Natl. Acad. Sci. USA* 84, 5277–5281.
- Friesel, R., Komoriya, A., and Maciag, T. (1987). Inhibition of endothelial cell proliferation by gamma-interferon. *J. Cell. Biol.* 104, 689–696.
- Fuhrmann, U., Bause, E., and Ploegh, H. (1985). Inhibitors of oligosaccharide processing. *Biochem. Biophys. Acta* 825, 95–110.
- Geiser, A.G., Burmester, J.K., Webbing, R., Roberts, A.B., and Sporn, M.B. (1992). Inhibition of growth by transforming growth factor-beta following fusion of two nonresponsive human carcinoma cell lines. *J. Biol. Chem.* 267, 2588–2593.
- Gospodarowicz, D., Cheng, J., Lui, G.M., Baird, A., and Böhlen, P. (1984). Isolation of brain fibroblast growth factor by heparin-sepharose affinity chromatography: identity with pituitary fibroblast growth factor. *Proc. Natl. Acad. Sci. USA* 81, 6963–6967.
- Heimark, R.L., Twardzik, D.R., and Schwartz, S.M. (1986). Inhibition of endothelial regeneration by type-beta transforming growth factor from platelets. *Science* 233, 1078–1080.
- Hosang, M. (1988). Recombinant interferon- γ inhibits the mitogenic effect of platelet-derived growth factor at a level distal to the growth factor receptor. *J. Cell. Physiol.* 134, 396–404.
- Ignatz, R.A., and Massagué, J. (1986). Transforming growth factor-beta stimulates the expression of fibronectin and collagen and their incorporation into the extracellular matrix. *J. Biol. Chem.* 261, 4337–4345.
- Laiho, M., DeCaprio, J.A., Ludlow, J.W., Livingston, D.M., and Massagué, J. (1990a). Growth inhibition of TGF β linked to suppression of retinoblastoma protein phosphorylation. *Proc. Natl. Acad. Sci. USA* 87, 820–823.
- Laiho, M., Ronnstrand, L., Heino, J., Decaprio, J.A., Ludlow, J.W., Livingston, D.M., and Massagué, J. (1991). Control of junB and extracellular matrix protein expression by transforming growth factor β 1 is independent of simian virus 40 T antigen-sensitive growth-inhibitory events. *Mol. Cell. Biol.* 11, 972–978.
- Laiho, M., Weis, M.B.W., Boyd, F.T., Ignatz, R.A., and Massagué, J. (1990b). Responsiveness to transforming growth factor-beta (TGF- β) restored by genetic complementation between cells defective in TGF- β receptors I and II. *J. Biol. Chem.* 266, 9108–9112.
- Laiho, M., Weis, F.M.B., and Massagué, J. (1990c). Concomitant loss of transforming growth factor TGF β receptor types I and II in TGF β -resistant cell mutants implicates both receptor types in signal transduction. *J. Biol. Chem.* 265, 18518–18524.
- Lin, H.Y., Wang, X.F., Ng-Eaton, E., Weinberg, R.A., and Lodish, H.F. (1992). Expression cloning of the TGF β type II receptor, a functional transmembrane serine/threonine kinase. *Cell* 68, 775–785.
- Madri, J.A., Pratt, B.M., and Tucker, A.M. (1988). Phenotypic modulation of endothelial cells by transforming growth factor-beta depends upon the composition and organization of the extracellular matrix. *J. Cell Biol.* 106, 1375–1384.
- Massagué, J. (1990). The transforming growth factor-beta family. *Annu. Rev. Cell Biol.* 6, 597–641.
- Müller, G., Behrens, J., Nussbaumer, U., Böhlen, P., and Birchmeier, W. (1987). Inhibitory action of transforming growth factor beta on endothelial cells. *Proc. Natl. Acad. Sci. USA* 84, 5600–5604.
- Mundschauf, L.J., and Faller, D.S. (1991). Balb/c-3T3 fibroblasts resistant to growth inhibition by beta interferon exhibit aberrant platelet derived-growth factor, epidermal growth factor, and fibroblast growth factor signal transduction. *Mol. Cell. Biol.* 11, 3148–3154.
- Pepper, M.S., Belin, D., Montesano, R., Orci, L., and Vassalli, J.D. (1990). Transforming growth factor-beta 1 modulates basic fibroblast

growth factor-induced proteolytic and angiogenic properties of endothelial cells in vitro. *J. Cell Biol.* 111, 743–755.

Pukac, L.A., Ottlinger, M.E., and Karnovsky, M.J. (1992). Heparin suppresses specific 2nd messenger pathways for proto-oncogene expression in rat vascular smooth muscle cells. *J. Biol. Chem.* 267, 3707–3711.

Saksela, O., Moscatelli, D., and Rifkin, D.B. (1987). The opposing effects of basic fibroblast growth factor and transforming growth factor beta on the regulation of plasminogen activator activity in capillary endothelial cells. *J. Cell Biol.* 105, 957–963.

Scatchard, G. (1949). The attractions of proteins for small molecules and ions. *Ann. NY Acad. Sci.* 51, 660–672.

Seddon, A., Decker, M., Müller, T., Armellino, D., Kovesdi, I., Gluzman, Y., and Böhlen, P. (1991). Structure activity relationships in basic FGF. *Ann. NY Acad. Sci.* 638, 98–108.

Segarini, P.R., Rosen, D.M., and Seyedin, S.M. (1989). Binding of transforming growth factor-beta to cell surface proteins varies with cell type. *Mol. Endocrinol.* 3, 262–272.

Sims, J.M., March, C.J., Cosman, D., Widmer, M.B., MacDonald, H.R., MacMahan, C., Grubin, C.E., Wignall, J.M., Jackson, J.L., Call, S.M., Friend, D., Alpert, A.L., Gillis, S., Urdal, D.L., and Dower, S.K. (1988). cDNA expression cloning of the IL-1 receptor, a member of the immunoglobulin superfamily. *Science* 241, 585–589.

Smith, C.A., Davis, T., Anderson, D., Solam, L., Beckmann, M.P., Jerzy, R., Dower, S.K., Cosman, D., and Goodwin, R.G. (1990). A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. *Science* 248, 1019–1023.

Sporn, M.B., and Roberts, A.B. (1990). TGF β : problems and prospects. *Cell. Regul.* 1, 875–882.

Sprengers, E.D., and Kluft, C. (1987). Plasminogen activator inhibitors. *Blood* 69, 381–387.