

[0227] "Aryl" alone or in combination means phenyl or naphthyl optionally carbocyclic fused with a cycloalkyl of preferably 5-7, more preferably 5-6, ring members.

[0228] "Substituted aryl" refers to an aryl group as defined above independently substituted with 1 or more, e.g., 1, 2, or 3, groups or substituents such as halo, hydroxy, optionally substituted alkoxy, optionally substituted alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted amino, optionally substituted amido, amidino, urea optionally substituted with alkyl, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, alkylsulfonylamino, carboxyl, heterocycle, substituted heterocycle, nitro, cyano, thiol, sulfonylamino or the like attached at any available point to produce a stable compound.

[0229] "Alkoxy" denotes the group $-OR^{ee}$, where R^{ee} is alkyl. "Lower alkoxy" denotes the group $-OR^{ee}$, where R^{ee} is lower alkyl.

[0230] "Substituted alkoxy" denotes the group $-OR^{dd}$, where R^{dd} is substituted alkyl. "Substituted lower alkoxy" denotes the group $-OR^{ddd}$, where R^{ddd} is substituted lower alkyl.

[0231] "Alkylthio" or "thioalkoxy" refers to the group $-S-R^{ee}$, where R^{ee} is alkyl.

[0232] "Substituted alkylthio" or "substituted thioalkoxy" refers to the group $-S-R$, where R is substituted alkyl.

[0233] "Sulfinyl" denotes the group $-S(O)-$.

[0234] "Sulfonyl" denotes the group $-S(O)_2-$.

[0235] "Substituted sulfinyl" denotes the group $-S(O)-R^{ff}$, where R^{ff} is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkylalkyl, substituted cycloalkylalkyl, heterocyclyl, substituted heterocyclyl, heterocyclylalkyl, substituted heterocyclylalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heteroaralkyl, substituted heteroaralkyl, aralkyl or substituted aralkyl.

[0236] "Substituted sulfonyl" denotes the group $-S(O)_2R^{gg}$, where R^{gg} is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkylalkyl, substituted

cycloalkylalkyl, heterocyclyl, substituted heterocyclyl, heterocyclylalkyl, substituted heterocyclylalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heteroaralkyl, substituted heteroaralkyl, aralkyl or substituted aralkyl.

[0237] "Sulfonylamino" denotes the group $-S(O)_2NR^{hh}$, where R^{hh} is hydrogen or alkyl.

[0238] "Substituted sulfonylamino" denotes the group $-S(O)_2NR^{ii}-R^{jj}$, where R^{ii} is hydrogen or optionally substituted alkyl, and R^{jj} is alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heteroaralkyl, substituted heteroaralkyl, aralkyl or substituted aralkyl.

[0239] "Amino" or "amine" denotes the group $-NH_2$. A "divalent amine" denotes the group $-NH-$. A "substituted divalent amine" denotes the group $-NR^{kk}$, wherein R^{kk} is alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, sulfonyl or substituted sulfonyl.

[0240] "Substituted amino" or "substituted amine" denotes the group $-NR^{mmm}R^{nn}$, wherein R^{mmm} and R^{nn} are independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, sulfonyl, substituted sulfonyl, or cycloalkyl provided, however, that at least one of R^{mmm} and R^{nn} is not hydrogen. $R^{mmm}R^{nn}$ in combination with the nitrogen may form an optionally substituted heterocyclic or heteroaryl ring.

[0241] "Alkylsulfinyl" denotes the group $-S(O)R^{oo}$, wherein R^{oo} is optionally substituted alkyl.

[0242] "Alkylsulfonyl" denotes the group $-S(O)_2R^{pp}$, wherein R^{pp} is optionally substituted alkyl.

[0243] "Alkylsulfonylamino" denotes the group $-NR^{qq}S(O)_2R^{rr}$, wherein R^{rr} is optionally substituted alkyl, and R^{qq} is hydrogen or alkyl.

[0244] A "primary amino substituent" denotes the group $-NH_2$.

[0245] A "secondary amino substituent" denotes the group -NHR^{ss} , wherein R^{ss} is alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, sulfonyl, substituted sulfonyl, or cycloalkyl.

[0246] A "tertiary amino substituent" denotes the group $\text{-NR}^{\text{ss}}\text{R}^{\text{b}}$, wherein R^{ss} and R^{b} are independently alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, sulfonyl, substituted sulfonyl, or cycloalkyl.

[0247] "Quaternary ammonium substituent" denotes the group $\text{-N}^+\text{R}^{\text{ss}}\text{R}^{\text{b}}\text{R}^{\text{uu}}$, wherein R^{ss} , R^{b} and R^{uu} are independently alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, acyl, substituted acyl, sulfonyl, substituted sulfonyl, or cycloalkyl.

[0248] "Heteroaryl" means a monocyclic aromatic ring structure containing 5 or 6 ring atoms, or a bicyclic aromatic group having 8 to 10 atoms, containing one or more, preferably 1-4, more preferably 1-3, even more preferably 1-2, heteroatoms independently selected from the group consisting of O, S, and N. Heteroaryl is also intended to include oxidized S or N, such as sulfinyl, sulfonyl and N-oxide of a tertiary ring nitrogen. A carbon or nitrogen atom is the point of attachment of the heteroaryl ring structure such that a stable aromatic ring is retained. Examples of heteroaryl groups are pyridinyl, pyridazinyl, pyrazinyl, quinoxalyl, indolizinyl, benzo[b]thienyl, quinazolinyl, purinyl, indolyl, quinolinyl, pyrimidinyl, pyrrolyl, oxazolyl, thiazolyl, thienyl, isoxazolyl, oxathiadiazolyl, isothiazolyl, tetrazolyl, imidazolyl, triazinyl, furanyl, benzofuryl, indolyl, and the like. "Heteroarylene" means a divalent heteroaryl.

[0249] "Heterocycle" or "heterocyclyl" means a saturated or unsaturated, non-aromatic carbocyclic group having a single ring or multiple condensed rings, e.g. a cycloalkyl group having from 5 to 10 atoms in which from 1 to 3 carbon atoms in a ring are replaced by heteroatoms, such as O, S, N, and are optionally fused with benzo or heteroaryl of 5-6 ring members and/or are optionally substituted. Heterocyclyl is intended to include oxidized S or N, such as sulfinyl, sulfonyl and N-oxide of a tertiary ring nitrogen. Examples of heterocycle or heterocyclyl groups are morpholino, tetrahydrofuranyl, dihydropyridinyl, piperidinyl, pyrrolidinyl, piperazinyl, dihydrobenzofuryl, dihydroindolyl, and the like.

[0250] "Heterocyclalkyl" refers to the group -R-Het where Het is a heterocycle group and R is an alkylene group.

[0251] A "substituted heteroaryl," "substituted heterocycl," or "substituted heterocyclalkyl" is a heteroaryl, heterocycl, or heterocyclalkyl, respectively, independently substituted with 1 or more, e.g., 1, 2, or 3, groups or substituents such as halogen, hydroxy, optionally substituted alkoxy, optionally substituted alkylthio, alkylsulfinyl, alkylsulfonyl, acyloxy, optionally substituted aryl, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted amino, optionally substituted amido, amidino, urea optionally substituted with alkyl, aryl, heteroaryl or heterocycl groups, aminosulfonyl optionally N-mono- or N,N-di-substituted with alkyl, aryl or heteroaryl groups, alkylsulfonylamino, arylsulfonylamino, heteroaryl/sulfonylamino, alkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, carboxyl, heterocycle, substituted heterocycle, heteroaryl, substituted heteroaryl, nitro, cyano, thiol, sulfonylamino, optionally substituted alkyl, optionally substituted alkenyl, or optionally substituted alkynyl, attached at any available point to produce a stable compound.

[0252] "Amido" denotes the group -C(O)NH₂. "Substituted amido" denotes the group -C(O)NR^kR^l, wherein R^k and R^l are independently hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, heteroaryl, or substituted heteroaryl, provided, however, that at least one of R^k and R^l is not hydrogen. R^kR^l in combination with the nitrogen may form an optionally substituted heterocyclic or heteroaryl ring.

[0253] "Amidino" denotes the group -C(=NR^m)NRⁿR^o, wherein R^m, Rⁿ, and R^o are independently hydrogen or optionally substituted lower alkyl.

[0254] "Acyloxy" denotes the group -OC(O)R^h, where R^h is hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, heterocycl, substituted heterocycl, aryl, substituted aryl, heteroaryl, substituted heteroaryl and the like.

[0255] "Aryloxy" denotes the group -OAr, where Ar is an aryl, or substituted aryl, group. "Heteroaryloxy" denotes groups -OHet, wherein Het is an optionally substituted heteroaryl group.

[0256] "Arylsulfonylamino" denotes the group $-NR^aS(O)_2R^b$, wherein R^b is optionally substituted aryl, and R^a is hydrogen or lower alkyl. "Heteroarylsulfonylamino" denotes the group $-NR^aS(O)_2R^b$, wherein R^b is optionally substituted heteroaryl, and R^a is hydrogen or lower alkyl.

[0257] "Alkylcarbonylamino" denotes the group $-NR^aC(O)R^b$, wherein R^b is optionally substituted alkyl, and R^a is hydrogen or lower alkyl.

[0258] "Arylcarbonylamino" denotes the group $-NR^aC(O)R^b$, wherein R^b is optionally substituted aryl, and R^a is hydrogen or lower alkyl.

[0259] "Heteroarylcarbonylamino" denotes the group $-NR^aC(O)R^b$, wherein R^b is optionally substituted aryl, and R^a is hydrogen or lower alkyl.

[0260] Pharmaceutical formulations containing K_{ATP} channel openers can include the free base of a compound defined by any of Formulae I – VIII, or a salt thereof. Salts of the compounds of Formulae I – VIII as provided herein may have one or more of the following characteristics: (1) stability in solution during synthesis and formulation, (2) stability in a solid state, (3) compatibility with excipients used in the manufacture of tablet formulations, (4) quantitatively yield the K_{ATP} channel opener upon exposure to simulated or actual gastric and duodenal conditions, (5) release K_{ATP} channel opener from sufficiently small particles that are readily dissolved and absorbed, (6) provide, when incorporated into a pharmaceutical formulation, for absorption of greater than 80% of the administered dose, (7) present no elevated toxicological risk as compared to the free base of the K_{ATP} channel opener, (8) can be formulated into acceptable pharmaceutical formulations to treat obesity and other diseases of humans, (9) are acceptable to the FDA as the basis of a drug product, (10) can be recrystallized to improve purity, (11) can be used to form co-crystals of two or more salts of the K_{ATP} channel opener, (12) have limited hygroscopicity to improve stability, (13) synthetic and crystallization conditions under which the salt is formed can be varied resulting in different crystal structures (polymorphs) can be controlled in the synthesis of the salt, or (14) have improved solubility as compared to the free base in aqueous systems at physiological pH values.

[0261] The K_{ATP} channel openers provided in Formulae I – VIII are preferably formulated as pharmaceutically acceptable salts. Pharmaceutically acceptable salts are

non-toxic salts in the amounts and concentrations at which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical characteristics of a compound without preventing it from exerting its physiological effect. Useful alterations in physical properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate administering lower effective doses of the drug.

[0262] Salts of the compounds of Formulae I – IV can include metal cations, preferably alkali metal cations, such as for example, sodium or potassium. Cations can be selected from any group I alkali metal. Divalent metals cations, such as alkaline earth metals (e.g., magnesium, calcium and the like), have not been found to be useful for salt formation with the compounds of Formulae I – IV.

[0263] Salts of the compounds of Formulae I – IV which include alkali metal cations can be prepared by reacting the compounds of Formulae I – IV with an alkali metal hydroxide or alkali metal alkoxide, such as for example, NaOH, KOH or NaOCH₃, in a variety of solvents which may be selected from low molecular weight ketones (e.g., acetone, methyl ethyl ketone), tetrahydrofuran (THF), dimethylformamide (DMF), and N-methyl pyrrolidinone, and the like. Surprisingly, salt formation with an alkali metal hydroxide or alkoxide is not observed when an alcohol, particularly a lower alcohol such as for example methanol or ethanol, is used as the solvent. This result was confirmed by both X-Ray Powder Diffraction and NMR, and is contrary to the disclosure of U.S. Pat. No. 2,986,573, which purports to describe formation of diazoxide salts in alcohol.

[0264] The compounds of Formulae I – IV can also form salts with organic cations that include at least one tertiary amine or ammonium cation. Organic cation compounds can be monovalent, divalent, trivalent and tetravalent by inclusion of one, two, three or four tertiary amine or ammonium ions within the compound, respectively. When a multivalent compound is used, the tertiary amine or quaternary ammonium moieties are preferably separated by a chain of at least 4 atoms, more preferably by a chain of at least 6 atoms, such as for example, hexamethyl hexamethylene diammonium dihydroxide, wherein the quaternary ammonium moieties are separated by $-(CH_2)_6-$. Primary and secondary amines do not to effectively form salts with the compounds of Formulae I – IV.

[0265] Salts of the compounds of Formulae I – IV can be prepared by reacting the compounds of Formulae I – IV with compounds that include at least one tertiary amine or quaternary ammonium ion (e.g., choline hydroxide, hexamethylhexamethylene diammonium dihydroxide) in a solvent selected from low molecular weight ketones (e.g., acetone, methyl ethyl ketone), tetrahydrofuran, dimethylformamide, and n-methyl pyrrolidinone. As with the preparation of salts from alkali metal hydroxides, amine and ammonium containing compounds do not form salts when the solvent is an alcohol.

[0266] Pharmaceutically acceptable salts of the compounds of Formulae I – IV can also include basic addition salts such as those containing benzathine, chlorprocaine, choline, diethylamino-ethanol, hydroxyethyl pyrrolidine, ammonium, tetrapropylammonium, tetrabutylphosphonium, hexamethyl diammonium, methyldiethanamine, triethylamine, meglumine, and procaine, and can be prepared using the appropriate corresponding bases.

[0267] Preferred basic addition salts of the compounds of Formulae I – IV can include those containing hexamethyl hexamethylene diammonium, choline, sodium, potassium, methyldiethyl amine, triethylamine, diethylamino-ethanol, hydroxyethyl pyrrolidine, tetrapropylammonium and tetrabutylphosphonium ions.

[0268] Preferred basic addition salts of the compounds of Formulae I – IV can be prepared using hexamethyl hexamethylene diammonium dihydroxide, choline hydroxide, sodium hydroxide, sodium methoxide, potassium hydroxide, potassium methoxide, ammonium hydroxide, tetrapropylammonium hydroxide, and tetrabutylphosphonium hydroxide. The basic addition salts can be separated into inorganic salts (e.g., sodium, potassium and the like) and organic salts (e.g., choline, hexamethyl hexamethylene diammonium hydroxide, and the like).

[0269] The compounds of Formulae V – VIII have the unique property of being able to form both anions and cations. In basic media, the compounds of Formulae V – VIII typically form anions. Anions can be formed at either an amino or substituted amino substituent, or at the sulfonyl group. In acidic media, the compounds of Formulae V – VIII generally form cations by protonation of an amino group, thereby forming an ammonium moiety.

[0270] Salts of the anions of compounds of Formulae V – VIII can include metal cations, including monovalent metal cations of any group I alkali metal (e.g., sodium, potassium, and the like), divalent metal cations of any group II alkaline earth metal (e.g., calcium, magnesium, and the like), and aluminum cations.

[0271] Salts of the compounds of Formulae V – VIII which include metal cations can be prepared by reacting the compounds of Formulae V – VIII with a alkali or alkaline earth metal hydroxides or alkoxides, such as for example, sodium hydroxide or sodium methoxide, in an organic solvent, such as for example lower alcohols, low molecular weight ketones (e.g., acetone, methyl ethyl ketone, and the like), tetrahydrofuran, dimethylformamide, and n-methyl pyrrolidinone, and the like.

[0272] Salts of the compounds of Formulae V - VIII, may include organic or inorganic counter ions, including but not limited to, acetate, acetonide, acetyl, adipate, aspartate, besylate, biacetate, bitartrate, bromide, butoxide, butyrate, calcium, camsylate, caproate, carbonate, citrate, cypionate, decaroate, diacetate, dimegulumine, dinitrate, dipotassium, dipropionate, disodium, disulfide, edisylate, enanthate, estolate, etabonate, ethylsuccinate, fumarate, furoate, gluceptate, gluconate, hexacetonide, hippurate, hyclate, hydrobromide, hydrochloride, isethionate, lactobionate, malate, maleate, meglumine, methylbromide, methylsulfate, metrizoate, nafate, napsylate, nitrate, oleate, palmitate, pamoate, phenpropionate, phosphate, pivalate, polistirex, polygalacturonate, probutate, propionate, saccharate, sodium glycinate, sodium phosphate, sodium succinate, stearate, succinate, sulfate, sulfonate, sulfosalicylate, tartrate, tebutate, terephalate, terephthalate, tosylate, trifluate, trihydrate, trisilicate, tromethamine, valerate, or xinafolate. Preferred organic cations include compounds having tertiary amines or quaternary ammonium groups.

[0273] Other, pharmaceutically acceptable salts of the compounds of Formulae V – VIII include acid addition salts such as those containing sulfate, chloride, hydrochloride, fumarate, maleate, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluene sulfonate, cyclohexylsulfamate and quinate. Pharmaceutically acceptable salts of the compounds of Formulae V - VIII can be obtained from acids such as hydrochloric acid, maleic acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric

acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, fumaric acid, and quinic acid.

[0274] Pharmaceutically acceptable salts of the compounds of Formulae V – VIII also include basic addition salts such as those containing benzathine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine, procaine, aluminum, calcium, lithium, magnesium, potassium, sodium, ammonium, alkylamine, and zinc, when acidic functional groups, such as carboxylic acid or phenol are present. For example, see *Remington's Pharmaceutical Sciences*, 19th ed., Mack Publishing Co., Easton, PA, Vol. 2, p. 1457, 1995. Such salts of the compounds of Formulae V – VIII can be prepared using the appropriate corresponding bases.

[0275] Salts of the compounds of Formulae V – VIII can be prepared, for example, by dissolving the free-base form of a compound in a suitable solvent, such as an aqueous or aqueous-alcohol in solution containing the appropriate acid and then isolated by evaporating the solution. In another example, a salt is prepared by reacting the free base and acid in an organic solvent.

[0276] The salts of the compounds of Formulae V – VIII may be present as a complex. Examples of complexes include 8-chlorotheophylline complex (analogous to, e.g., dimenhydrinate: diphenhydramine 8-chlorotheophylline (1:1) complex; Dramamine) and various cyclodextrin inclusion complexes.

[0277] Solvents useful in the preparation of pharmaceutically acceptable salts of the compounds of Formulae V – VIII include organic solvents, such as for example, acetonitrile, acetone, alcohols (e.g., methanol, ethanol and isopropanol), tetrahydrofuran, methyl ethyl ketone (MEK), ethers (e.g., diethyl ether), benzene, toluene, xylenes, dimethylformamide (DMF), and N-methyl pyrrolidinone (NMP), and the like. Preferably, the solvents are selected from acetonitrile and MEK.

[0278] The salts of compounds of Formulae V – VIII may be present as a complex. Examples of complexes include 8-chlorotheophylline complex (analogous to, e.g., dimenhydrinate: diphenhydramine 8-chlorotheophylline (1:1) complex; Dramamine) and various cyclodextrin inclusion complexes.

[0279] Formulations of salts of the compounds of Formulae I – VIII provided herein exhibit at least one, or preferably some or even more preferably, all the following characteristics: (1) they are stable at ambient temperatures for a minimum of one year; (2) they provide for ease of oral administration; (3) they facilitate patient compliance with dosing; (4) upon administration, they consistently facilitate high levels of absorption of the pharmaceutical active; (5) upon once or twice daily oral administration they allow release of the K_{ATP} channel opener over a sustained time frame such that the circulating concentration of the K_{ATP} channel opener or its metabolically active metabolites does not fall below a therapeutically effective concentration; (6) they achieve these results independent of the pH of the gastrointestinal tract of treated subjects, and (7) they delay release until gastric transit is complete or nearly complete.

[0280] Formulations designed for oral administration of the salts of the compounds of Formulae I – VIII can be provided, for example, as capsules, tablets, or as quick dissolve tablets or films. Capsule or tablet formulations include a number of distinguishing components. One is a component to improve absorption of the K_{ATP} channel opener. Another sustains release of the drug over more than 2 hours. A third delays substantial release of the drug until gastric transit is completed.

[0281] Oral administration formulations of the salts of the compounds of Formulae I – VIII can also be provided, for example, as oral suspensions, oral solutions, encapsulated oral suspensions, and encapsulated oral solutions. Formulations can be designed for immediate release or controlled release. Preferably, such oral formulations are not produced from a liquid form of the sodium salt of diazoxide.

[0282] Formulations of the salts of the compounds of Formulae I – VIII can also be prepared for transdermal, intranasal and intravenous (I.V.) administration, provided that when the anion is diazoxide and the cation is sodium, the formulation is not for intravenous use.

[0283] In another embodiment, formulations of the salts of the compounds of Formulae I – VIII are prepared for transdermal or intranasal administration, provided that when the anion is diazoxide and the cation is sodium, the formulation is not produced using a liquid form of the salt of the compounds of Formulae I – VIII.

[0284] In another embodiment, formulations of the salts of the compounds of Formulae I – VIII are prepared for transdermal, intranasal and intravenous (I.V.) administration excluding the sodium salt of diazoxide.

[0285] Formulations of K_{ATP} channel openers prepared using salts of the compounds selected from Formulae I – VIII exhibit improved solubility and absorption compared to previous formulations of these drugs. These advantageous properties are achieved by any one or more of the following approaches: (1) reducing particle size of the formulation by comminution, spray drying, or other micronising techniques, (2) using an ion exchange resin in the formulation, (3) using inclusion complexes, for example using a cyclodextrin, (4) compaction of the salt of K_{ATP} channel opener with a solubilizing agent including low viscosity hypromellose, low viscosity methylcellulose or similarly functioning excipient and combinations thereof, (5) associating the salt of the K_{ATP} channel opener with a distinct salt prior to formulation, (6) using a solid dispersion of the salt of the K_{ATP} channel opener, (7) using a self emulsifying system, (8) adding one or more surfactants to the formulation, (9) using nanoparticles in the formulation, or (10) combinations of these approaches.

[0286] Release of K_{ATP} channel opener selected from salts of the compounds of Formulae I – VIII over a sustained period of time (e.g., 2-30 hours) can be achieved by the use of one or more approaches including, but not limited to: (1) the use of pH sensitive polymeric coatings, (2) the use of a hydrogel, (3) the use of a film coating that controls the rate of diffusion of the drug from a coated matrix, (4) the use of an erodable matrix that controls rate of drug release, (5) the use of polymer coated pellets, granules, or microparticles which can be further encapsulated or compressed into a tablet, (6) the use of an osmotic pump system, (7) the use of a compression coated tablet, or (8) combinations of these approaches.

[0287] Delay of release of K_{ATP} channel openers selected from the salts of the compounds of Formulae I – VIII from the formulation until gastric transit is complete can be achieved in the formulations provided herein by any of several mechanisms. For example, pH sensitive polymer or co-polymer can be used which when applied around the drug matrix functions as an effective barrier to release of active at pH 3.0 or lower and is unstable at pH 5.5 and above. This provides for control of release of the active compound

in the stomach but rapidly allows release once the dosage form has passed into the small intestine. An alternative to a pH sensitive polymer or co-polymer is a polymer or co-polymer that is non-aqueous-soluble. The extent of resistance to release in the gastric environment can be controlled by coating with a blend of the non-aqueous-soluble and a aqueous soluble polymer. In this approach neither of the blended polymers or co-polymers are pH sensitive. One example of a pH sensitive co-polymer is the Eudragit® methacrylic co-polymers, including Eudragit® L 100, S 100 or L 100-55 solids, L 30 D-55 or FS 30D dispersions, or the L 12,5 or S 12,5 organic solutions.

[0288] Polymers that delay release can be applied to a tablet either by spray coating (as a thin film) or by compression coating. If a capsule is used, then the polymer(s) may be applied over the surface of the capsule or applied to microparticles of the drug, which may then be encapsulated such as in a capsule or gel. If the capsule is coated, then it will resist disintegration until after gastric transit. If microparticles are coated, then the capsule may disintegrate in the stomach but little to no drug will be released until after the free microparticles complete gastric transit. Finally, an osmotic pump system that uses e.g., a swellable hydrogel can be used to delay drug release in the stomach. The swellable hydrogel takes up moisture after administration. Swelling of the gel results in displacement of the drug from the system for absorption. The timing and rate of release of the drug depend on the gel used, and the rate at which moisture reaches the gel, which can be controlled by the size of the opening in the system through which fluid enters. See Drug Delivery Technologies online article Dong et al., "L-OROS® SOFTCAP™ for Controlled Release of Non-Aqueous Liquid Formulations."

[0289] Accordingly, delay of release of formulations of K_{ATP} channel openers prepared as salts of the compounds of Formulae I - VIII until after gastric transit is complete can be achieved by any of several mechanisms, including, but not limited to: (a) a pH sensitive polymer or co-polymer applied as a compression coating on a tablet; (b) a pH sensitive polymer or co-polymer applied as a thin film on a tablet; (c) a pH sensitive polymer or co-polymer applied as a thin film to an encapsulation system; (d) a pH sensitive polymer or co-polymer applied to encapsulated microparticles, (e) a non-aqueous-soluble polymer or copolymer applied as a compression coating on a tablet; (f) a non-aqueous-soluble polymer or co-polymer applied as a thin film on a tablet; (g) a non-

aqueous soluble polymer applied as a thin film to an encapsulation system; (h) a non-aqueous soluble polymer applied to microparticles; (i) incorporation of the formulation in an osmotic pump system, or (j) use of systems controlled by ion exchange resins, or (k) combinations of these approaches, wherein the pH sensitive polymer or co-polymer is resistant to degradation under acid conditions.

[0290] Formulations are provided that are designed for administration once daily (i.e., once per 24 hours). These formulations can contain between 25 and 500 mg of K_{ATP} channel openers selected from salts of the compounds of Formulae I – VIII. Formulations intended for administration twice daily (per 24 hours) may also be provided. These can contain between 25 and 250 mg of K_{ATP} channel openers.

[0291] The formulations provided herein exhibit improved safety of the administered drug product. This improvement in safety occurs by at least two mechanisms. First, delay of release of active drug until gastric transit is complete can reduce the incidence of a range of gastrointestinal adverse side effects including nausea, vomiting, dyspepsia, abdominal pain, diarrhea and ileus. Second, by sustaining release of the active drug over 2 or more hours up to as long as 24 hours, peak drug levels are reduced relative to the peak drug levels observed for the same administered dose using any oral formulation that does not have sustained or controlled release. This reduction in peak drug levels can contribute to reductions in adverse effects that are partially or completely determined by peak drug levels. These adverse effects include: fluid retention with the associated reduced rates of excretion of sodium, chloride and uric acid, edema, hyperglycemia and the associated potential for progression to ketoacidosis, cataracts and non-ketotic hyperosmolar coma, headaches, tachycardia and palpitations.

[0292] Also provided herein are controlled release formulations of K_{ATP} channel openers prepared from salts of compounds of Formulae I – VIII, which have one feature from each of A-D as shown in Table I.

Table 1: Controlled Release Formulation Characteristics and Properties

A. Unit Form:	Tablet or Capsule
B. Dosage/unit:	10-100 mg
	100-200 mg
	200-300 mg
	300-500 mg
	500-2000mg
C. Dosing	Once daily (24 hours)
	Twice daily (24 hours)
D. Release time:	2-4 hrs
	4-8 hrs
	8-24 hours

[0293] For example, a controlled release composition can be a tablet containing 25-100 mg of a salt of a compound of Formulae I – VIII, wherein such tablet administered once daily to achieve a controlled release time of 2-4 hours. All of these formulations can further include the feature of substantially delaying pharmaceutical active release until after gastric transit is complete.

[0294] In addition, any of the above formulations from Table 1 can include at least one feature that improves the solubility or absorption of the K_{ATP} channel opener.

[0295] Exemplary controlled release formulations provided herein include the active compound (i.e., a K_{ATP} channel opener selected from a salt of a compound of any of Formulae I – VIII) and a matrix which includes a gelling agent that swells upon contact with aqueous fluid. The active compound entrapped within the gel is slowly released into the body upon dissolution of the gel. The active compound can be evenly dispersed within the matrix or can be present as pockets of drug in the matrix. For example, the drug can be formulated into small granules which are dispersed within the matrix. In addition, the granules of drug also can include a matrix, thus, providing a primary and a secondary matrix as described in U.S. Pat. No. 4,880,830 to Rhodes.

[0296] The gelling agent preferably is a polymeric material, which can include, for example, any pharmaceutically acceptable water soluble or water insoluble slow releasing polymer such as xanthan gum, gelatin, cellulose ethers, gum arabic, locust bean gum, guar gum, carboxyvinyl polymer, agar, acacia gum, tragacanth, veegum, sodium alginate or alginic acid, polyvinylpyrrolidone, polyvinyl alcohol, or film forming polymers such as methyl cellulose (MC), carboxymethyl cellulose (CMC), hydroxypropyl methylcellulose, hydroxypropyl methyl cellulose (HPMC), hydroxypropyl cellulose (HPC), hydroxyethyl cellulose (HEC), ethylcellulose (EC), acrylic resins or mixtures of the above (see e.g., U.S. Pat. No. 5,415,871).

[0297] The gelling agent of the matrix also can be a heterodisperse gum comprising a heteropolysaccharide component and a homopolysaccharide component which produces a fast-forming and rigid gel as described in U.S. Pat. No. 5,399,359. The matrix also can include a cross-linking agent such as a monovalent or multivalent metal cations to further add rigidity and decrease dissolution of the matrix, thus further slowing release of drug. The amount of crosslinking agent to add can be determined using methods routine to the ordinary skilled artisan.

[0298] The matrix of the controlled release composition also can include one or more pharmaceutically acceptable excipients recognized by those skilled in the art, i.e. formulation excipients. Such excipients include, for example, binders: polyvinylpyrrolidone, gelatin, starch paste, microcrystalline cellulose; diluents (or fillers): starch, sucrose, dextrose, lactose, fructose, xylitol, sorbitol, sodium chloride, dextrans, calcium phosphate, calcium sulphate; and lubricants: stearic acid, magnesium stearate, calcium stearate, Precirol ® and flow aids for example talc or colloidal silicon dioxide.

[0299] The matrix of the controlled release composition can further include a hydrophobic material which slows the hydration of the gelling agent without disrupting the hydrophilic nature of the matrix, as described in U.S. Pat. No. 5,399,359. The hydrophobic polymer can include, for example, alkylcellulose such as ethylcellulose, other hydrophobic cellulosic materials, polymers or copolymers derived from acrylic or methacrylic acid esters, copolymers of acrylic and methacrylic acid esters, zein, waxes, shellac, hydrogenated vegetable oils, waxes and waxy substances such as carnauba wax, spermaceti wax, candellila wax, cocoa butter, cetostearyl alcohol, beeswax, ceresin,

paraffin, myristyl alcohol, stearyl alcohol, cetylalcohol and stearic acid, and any other pharmaceutically acceptable hydrophobic material known to those skilled in the art.

[0300] The amount of hydrophobic material incorporated into the controlled release composition is that which is effective to slow the hydration of the gelling agent without disrupting the hydrophilic matrix formed upon exposure to an environmental fluid. In certain preferred embodiments, the hydrophobic material is included in the matrix in an amount from about 1 to about 20 percent by weight and replaces a corresponding amount of the formulation excipient. A solvent for the hydrophobic material may be an aqueous or organic solvent, or mixtures thereof.

[0301] Examples of commercially available alkylcelluloses are Aquacoat® (aqueous dispersion of ethylcellulose available from FMC) and Surelease® (aqueous dispersion of ethylcellulose available from Colorcon). Examples of commercially available acrylic polymers suitable for use as the hydrophobic material include Eudragit® RS and RL (copolymers of acrylic and methacrylic acid esters having a low content (e.g., 1:20 or 1:40) of quaternary ammonium compounds).

[0302] The controlled release composition also can be coated to retard access of liquids to the active compound and/or retard release of the active compound through the film-coating. The film-coating can provide characteristics of gastroresistance and enterosolubility by resisting rapid dissolution of the composition in the digestive tract. The film-coating generally represents about 5-15% by weight of the controlled release composition. Preferably, the core by weight represents about 90% of the composition with the remaining 10% provided by the coating. Such coating can be a film-coating as is well known in the art and include gels, waxes, fats, emulsifiers, combination of fats and emulsifiers, polymers, starch, and the like.

[0303] Polymers and co-polymers are useful as thin film coatings. Solution coatings and dispersion coatings can be used to coat the active compound, either alone or combined with a matrix. The coating is preferably applied to the drug or drug and matrix combination as a solid core of material as is well known in the art.

[0304] A solution for coating can include polymers in both organic solvent and aqueous solvent systems, and typically further including one or more compounds that act

as a plasticizer. Polymers useful for coating compositions include, for example, methylcellulose (Methocel® A; Dow Chemical Co.), hydroxypropylmethylcellulose with a molecular weight between 1,000 and 4,000,000 (Methocel® E; Dow Chemical Co. or Pharmacoat®; Shin Etsu), hydroxypropyl cellulose with a molecular weight between 2,000 and 2,000,000, ethyl cellulose, cellulose acetate, cellulose triacetate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose acetate trimellitate (Eastman Kodak), carboxymethylethyl cellulose (Duodcel®), hydroxypropyl methylcellulose phthalate, ethylcellulose, methylcellulose and, in general, cellulosic derivatives, olymethacrylic acid-methacrylic acid copolymer (Type A 1:1 Eudragit L100; Type B 1:2 Eudragit S100; and Type C 1:1 Eudragit L100-55, aqueous dispersion 30% solids, Eudragit L30D), poly(meth)acryl ester: poly(ethyl acrylate, methyl methacrylate 2:1), Eudragit NE30D aqueous dispersion 30% solids, polyaminomethacrylate Eudragit E100, poly(trimethylammonioethyl methacrylate chloride) ammoniomethacrylate copolymer, Eudragit RL30D and Eudragit RS30D, carboxyvinyl polymers, polyvinylalcohols, glucans scleroglucans, mannans, and xanthans.

[0305] Aqueous polymeric dispersions include Eudragit L30D and RS/RL30D, and NE30D, AQUACOAT® brand ethyl cellulose, Surelease brand ethyl cellulose, EC brand N-10F ethyl cellulose, Aquateric brand cellulose acetate phthalate, Coateric brand Poly(vinyl acetate phthalate), and Aquacoat brand hydroxypropyl methylcellulose acetate succinate. Most of these dispersions are latex, pseudolatex powder or micronized powder mediums.

[0306] A plasticizing agent may be included in the coating to improve the elasticity and the stability of the polymer film and to prevent changes in the polymer permeability over prolonged storage. Such changes may affect the drug release rate. Suitable conventional plasticizing agents include, for example, diethyl phthalate, glycerol triacetate, acetylated monoglycerides, acetyltributylcitrate, acetyltriethyl citrate, castor oil, citric acid esters, dibutyl phthalate, dibutyl sebacate, diethyloxalate, diethyl malate, diethylfumarate, diethylphthalate, diethylsuccinate, diethylmalonate, diethyltartarate, dimethylphthalate, glycerin, glycerol, glyceryl triacetate, glyceryltributyrate, mineral oil and lanolin alcohols, petrolatum and lanolin alcohols, phthalic acid esters, polyethylene glycols, propylene glycol, rape oil, sesame oil, triacetin, tributyl citrate, triethyl citrate,

and triethyl acetyl citrate, or a mixture of any two or more of the foregoing. Plasticizers which can be used for aqueous coatings include, for example, propylene glycol, polyethylene glycol (PEG 400), triacetin, polysorbate 80, triethyl citrate, and diethyl d-tartrate.

[0307] A coating solution comprising a mixture of hydroxypropylmethylcellulose and aqueous ethylcellulose (e.g. Aquacoat brand) as the polymer and dibutyl sebacate as plasticizer can be used for coating microparticles. (Aquacoat is an aqueous polymeric dispersion of ethylcellulose and contains sodium lauryl sulfate and cetyl alcohol). Preferably, the plasticizer represents about 1-2% of the composition.

[0308] In addition to the polymers, the coating layer can include an excipient to assist in formulation of the coating solution. Such excipients may include a lubricant or a wetting agent. Suitable lubricants as excipients for the film coating include, for example, talc, calcium stearate, colloidal silicon dioxide, glycerin, magnesium stearate, mineral oil, polyethylene glycol, and zinc stearate, aluminum stearate or a mixture of any two or more of the foregoing. Suitable wetting agents include, for example, sodium lauryl sulfate, acacia, benzalkonium chloride, cetomacrogol emulsifying wax, cetostearyl alcohol, cetyl alcohol, cholesterol, diethanolamine, docusate sodium, sodium stearate, emulsifying wax, glyceryl monostearate, hydroxypropyl cellulose, lanolin alcohols, lecithin, mineral oil, monoethanolamine, poloxamer, polyoxyethylene alkyl ethers, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene stearates, propylene glycol alginate, sorbitan esters, stearyl alcohol and triethanolamine, or a mixture of any two or more of the foregoing.

[0309] The specified tablet or capsule formulations of Table I may include co-formulation with an obesity treating drug (in addition to a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII). Obesity treating drugs that may be used include, but are not limited to, sibutramine hydrochloride (5-30 mg/unit), orlistat (50-360 mg/unit), phentermine hydrochloride or resin complex (15 to 40 mg/unit), zonisamide (100 to 600 mg/unit), topiramate (64 to 400 mg/unit), naltrexone hydrochloride (50 to 600 mg/unit), rimonabant (5 to 20 mg/unit), ADP356 (5 to 25 mg/unit), ATL962 (20 to 400 mg/unit), or AOD9604 (1 to 10 mg/unit). These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of

a compound of Formulae I – VIII is one half the amount included in the once daily formulation and the co-formulated obesity treating drug is half of the amount specified. Alternative obesity treating drugs may include, but are not limited to: selective serotonin 2c receptor agonists, dopamine antagonists, cannabinoid-1 receptor antagonists, leptin analogues, leptin transport and/or leptin receptor promoters, neuropeptide Y and agouti-related peptide antagonists, proopiomelanocortin and cocaine and amphetamine regulated transcript promoters, melanocyte-stimulating hormone analogues, melanocortin-4 receptor agonists, and agents that affect insulin metabolism/activity, which can include protein-tyrosine phosphatase-1B inhibitors, peroxisome proliferator activated receptor antagonists, short-acting bromocriptine (ergoset), somatostatin agonists (octreotide), and adiponectin, gastrointestinal-neural pathway agents, including those that increase cholecystikinin activity, increase glucagon-like peptide-1 activity (e.g., extendin 4, liraglutide, dipeptidyl peptidase IV inhibitors), and increase protein YY3-36 activity and those that decrease ghrelin activity, as well as amylin analogues, agents that may increase resting metabolic rate ("selective" β -3 stimulators/agonist, uncoupling protein homologues, and thyroid receptor agonists), melanin concentrating hormone antagonists, phytosterol analogues, amylase inhibitors, growth hormone fragments, synthetic analogues of dehydroepiandrosterone sulfate, antagonists of adipocyte 11 β hydroxysteroid dehydrogenase type 1 activity, corticotropin releasing hormone agonists, inhibitors of fatty acid synthesis, carboxypeptidase inhibitors, indanones/indanols, aminosterols, and other gastrointestinal lipase inhibitors.

[0310] The specified tablet or capsule formulations of Table 1 may include co-formulation with a diabetes treating drug (in addition to a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII). Diabetes treating drugs that may be used include, but are not limited to, acarbose (50 to 300 mg/unit), miglitol (25 to 300 mg/unit), metformin hydrochloride (300 to 2000 mg/unit), repaglinide (1-16 mg/unit), nateglinide (200 to 400 mg/unit), rosiglitazone (5 to 50 mg/unit), metaglidasen (100 to 400 mg/unit) or any drug that improves insulin sensitivity, or improves glucose utilization and uptake. These formulations are preferably used once daily. For a twice daily dosing, the amount of the K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is half the amount included in the once daily formulation and the co-formulated diabetes treating drug is half of the amount specified.

[0311] The specified tablet or capsule formulations of Table 1 may include co-formulation with a cholesterol lowering drug. Cholesterol lowering drugs that may be used include, but are not limited to, pravastatin, simvastatin, atorvastatin, fluvastatin, rosuvastatin, or lovastatin (all at 10 to 80 mg/unit). These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is preferably 25 to 200 mg/unit and the co-formulated cholesterol lowering drug is half of the amount specified.

[0312] The specified tablet or capsule formulations of Table 1 may include co-formulation with a depression treating drug. Depression treating drugs that may be used include, but are not limited to, citalopram hydrobromide (10 to 80 mg/unit), escitalopram hydrobromide (5 to 40 mg/unit), fluvoxamine maleate (25 to 300 mg/unit), paroxetine hydrochloride (12.5 to 75 mg/unit), fluoxetine hydrochloride (30 to 100 mg/unit), setraline hydrochloride (25 to 200 mg/unit), amitriptyline hydrochloride (10 to 200 mg/unit), desipramine hydrochloride (10 to 300 mg/unit), nortriptyline hydrochloride (10 to 150 mg/unit), duloxetine hydrochloride (20 to 210 mg/unit), venlafaxine hydrochloride (37.5 to 150 mg/unit), phenelzine sulfate (10 to 30 mg/unit), bupropion hydrochloride (200 to 400 mg/unit), or mirtazapine (7.5 to 90 mg/unit). These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is preferably half the amount included in the once daily formulation and the co-formulated depression treating drug is half of the amount specified.

[0313] The specified tablet or capsule formulations of Table 1 may include co-formulation with a hypertension treating drug. Hypertension treating drugs that may be used include, but are not limited, to enalapril maleate (2.5 to 40 mg/unit), captopril (2.5 to 150 mg/unit), lisinopril (10 to 40 mg/unit), benazepril hydrochloride (10 to 80 mg/unit), quinapril hydrochloride (10 to 80 mg/unit), perindopril erbumine (4 to 8 mg/unit), ramipril (1.25 to 20 mg/unit), trandolapril (1 to 8 mg/unit), fosinopril sodium (10 to 80 mg/unit), moexipril hydrochloride (5 to 20 mg/unit), losartan potassium (25 to 200 mg/unit), irbesartan (75 to 600 mg/unit), valsartan (40 to 600 mg/unit), candesartan cilexetil (4 to 64 mg/unit), olmesartan medoxamil (5 to 80 mg/unit), telmisartan (20 to 160 mg/unit), eprosartan mesylate (75 to 600 mg/unit), atenolol (25 to 200 mg/unit), propranolol

hydrochloride (10 to 180 mg/unit), metoprolol tartrate, succinate or fumarate (each at 25 to 400 mg/unit), nadolol (20 to 160 mg/unit), betaxolol hydrochloride (10 to 40 mg/unit), acebutolol hydrochloride (200 to 800 mg/unit), pindolol (5 to 20 mg/unit), bisoprolol fumarate (5 to 20 mg/unit), nifedipine (15 to 100 mg/unit), felodipine (2.5 to 20 mg/unit), amlodipine besylate (2.5 to 20 mg/unit), nicardipine (10 to 40 mg/unit), nisoldipine (10 to 80 mg/unit), terazosin hydrochloride (1 to 20 mg/unit), doxazosin mesylate (4 to 16 mg/unit), prazosin hydrochloride (2.5 to 10 mg/unit), or alfuzosin hydrochloride (10 to 20 mg/unit). These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener is preferably half the amount included in the once daily formulation and the co-formulated hypertension treating drug is half of the amount specified.

[0314] The specified tablet or capsule formulations of Table 1 may include co-formulation with a diuretic to treat edema. Diuretics that may be used include, but are not limited to amiloride hydrochloride (1 to 10 mg/unit), spironolactone (10 to 100 mg/unit), triamterene (25 to 200 mg/unit), bumetanide (0.5 to 4 mg/unit), furosemide (10 to 160 mg/unit), ethacrynic acid or ethacrylate sodium (each at 10 to 50 mg/unit), tosemide (5 to 100 mg/unit), chlorthalidone (10 to 200 mg/unit), indapamide (1 to 5 mg/unit), hydrochlorothiazide (10 to 100 mg/unit), chlorothiazide (50 to 500 mg/unit), bendroflumethiazide (5 to 25 mg/unit), hydroflumethiazide (10 to 50 mg/unit), mythyclothiazide (1 to 5 mg/unit), or polythiazide (1 to 10 mg/unit). These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is preferably half the amount included in the once daily formulation and the co-formulated diuretic is half of the amount specified.

[0315] The specified tablet or capsule formulations of Table 1 may include co-formulation with a drug to treat inflammation or pain. Drugs for treating inflammation or pain that may be used include, but are not limited to aspirin (100 to 1000 mg/unit), tramadol hydrochloride (25 to 150 mg/unit), gabapentin (100 to 800 mg/unit), acetaminophen (100 to 1000 mg/unit), carbamazepine (100 to 400 mg/unit), ibuprofen (100 to 1600 mg/unit), ketoprofen (12 to 200 mg/unit), fenpropfen sodium (100 to 600 mg/unit), flurbiprofen sodium or flurbiprofen (both at 50 to 200 mg/unit), or

combinations of any of these with a steroid or aspirin. These formulations are preferably used once daily. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is preferably half the amount included in the once daily formulation and the co-formulated diuretic is half of the amount specified.

[0316] The specified tablet or capsule formulations of Table 1 may include co-formulation with a drug to treat obesity associated co-morbidities include those specified above for treating diabetes, cholesterol, depression, hypertension and edema, or drugs to treat atherosclerosis, osteoarthritis, disc herniation, degeneration of knees and hips, breast, endometrial, cervical, colon, leukemia and prostate cancers, hyperlipidemia, asthma/reactive airway disease, gallstones, GERD, obstructive sleep apnea, obesity hypoventilation syndrome, recurrent ventral hernias, menstrual irregularity and infertility.

[0317] The specified tablet or capsule formulations of Table 1 may include co-formulation with an anti-psychotic drug the combination used to treat the psychotic condition and to treat or prevent weight gain, dyslipidemia or impaired glucose tolerance in the treated subject. Drugs for treating various psychotic conditions that may be used include, but are not limited to, lithium or a salt thereof (250 to 2500 mg/unit), carbamazepine or a salt thereof (50 to 1200 mg/unit), valproate, valproic acid, or divalproex (125 to 2500 mg/unit), lamotrigine (12.5 to 200 mg/unit), olanzapine (5 to 20 mg/unit), clozapine (12.5 to 450 mg/unit), or risperidone (0.25 to 4 mg/unit). These coformulations are preferably intended for once per day administration. For a twice daily dosing, the amount of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is preferably half the amount included in the once daily formulation and the co-formulated anti-psychotic is half of the amount specified.

[0318] The specified tablet or capsule formulations of Table 1 may include co-formulation with a drug to treat or prevent ischemic or reperfusion injury. Drugs for treating or preventing ischemic or reperfusion injury that may be used include, but are not limited to: low molecular weight heparins (e.g., dalteparin, enoxaparin, nadroparin, tinzaparin or danaparoid), ancrd, pentoxifylline, nimodipine, flunarizine, ebselen, tirilazad, clomethiazole, an AMPA agonist (e.g., GYKI 52466, NBQX, YM90K, zonampanel, or MPQX), SYM 2081, selfotel, Cerestat, CP-101,606, dextrophan, dextromethorphan, MK-801, NPS 1502, remacemide, ACEA 1021, GV150526, eliprodil

ifenprodil, lubeluzole, naloxone, nalfemene citicoline, acetyl-l-carnitine, nifedipine, resveratrol, a nitron derivative, clopidogrel, dabigatram, prasugrel, troxoprodil, AGY-94806, or KAI-9803.

[0319] Provided are formulations administered once or twice daily to an obese or overweight subject continuously result in a circulating concentration of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII sufficient to induce weight loss. Weight loss occurs by the preferential loss of body fat. Additional weight loss can occur when the formulation is administered in combination with a reduced caloric diet.

[0320] Provided are formulations of K_{ATP} channel openers selected from a salt of a compound of Formulae I – VIII administered as a single dose to an obese, overweight or obesity-prone subject that result in the inhibition of fasting or glucose stimulated insulin secretion for about 24 hours or for about 18 hours.

[0321] Provided are formulations of K_{ATP} channel openers selected from a salt of a compound of Formulae I – VIII administered as a single dose to an obese, overweight or obesity-prone subject that result in the elevation of energy expenditure for about 24 hours or for about 18 hours.

[0322] Provided are formulations of K_{ATP} channel openers selected from a salt of a compound of Formulae I – VIII administered as a single dose to an obese, overweight or obesity-prone subject that result in the elevation of beta oxidation of fat for about 24 hours or for about 18 hours.

[0323] Provided are formulations of K_{ATP} channel openers selected from a salt of a compound of Formulae I – VIII administered as a single dose to an obese, overweight or obesity-prone hyperphagic subject that result in the inhibition of hyperphagia for about 24 hours or for about 18 hours.

[0324] Provided are formulations suitable for continuous administration once or twice daily (per 24 hours) to a subject, resulting in a circulating concentration of K_{ATP} channel openers selected from a salt of a compound of Formulae I – VIII sufficient to induce either beta-cell rest or improved insulin sensitivity or both. Such beta-cell rest and

improvements in insulin sensitivity can contribute to effective treatment of type I diabetes, type II diabetes and prediabetes. Such beta-cell rest and improvements in insulin sensitivity can contribute to effective restoration of normal glucose tolerance in type II diabetic and prediabetic subjects.

[0325] The various pharmaceutical K_{ATP} channel opener formulations selected from a salt of a compound of Formulae I – VIII have a variety of applications, including, but not limited to: (1) treatment of obesity; (2) prevention of weight gain in subjects who are predisposed to obesity; (3) treatment of hyperinsulinemia or hyperinsulinism; (4) treatment of hypoglycemia; (5) treatment of hyperlipidemia, (6) treatment of type II diabetes, (7) preservation of pancreatic function in type I diabetics; (8) treatment of metabolic syndrome (or syndrome X); (9) prevention of the transition from prediabetes to diabetes, (10) correction of the defects in insulin secretion and insulin sensitivity contributing to prediabetes and type II diabetes, (11) treatment of polycystic ovary syndrome, (12) prevention of ischemic or reperfusion injury, (13) treat weight gain, dyslipidemia, or impairment of glucose tolerance in subjects treated with antipsychotics drugs, (14) prevent weight gain, dyslipidemia, or impairment of glucose tolerance in subjects treated with antipsychotics drugs and (15) treatment of any disease where hyperlipidemia, hyperinsulinemia, hyperinsulinism, hyperlipidemia, hyperphagia or obesity are contributing factors to the severity or progression of the disease, including but not limited to, Prader Willi Syndrome, Froelich's syndrome, Cohen syndrome, Summit Syndrome, Alstrom Syndrome, Borjesen Syndrome, Bardet-Biedl Syndrome, or hyperlipoproteinemia type I, II, III, and IV.

[0326] In one embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage once per 24 hours to induce weight loss. In further embodiments, the subject (a) is not a type I diabetic, (b) is not a type II diabetic, (c) is not experiencing chronic, recurrent or drug-induced hypoglycemia, (d) does not have metabolic syndrome, or (e) is not experiencing malignant hypertension.

[0327] In one embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage twice per 24 hours to induce weight loss. This treatment can be the sole treatment to

induce weight loss. In further embodiments, the overweight or obese subject (a) does not have an insulin secreting tumor, (b) is not suffering from Poly Cystic Ovary Syndrome, (c) is not a type I diabetic, (d) is not a type II diabetic, (e) does not have metabolic syndrome, (f) is not experiencing chronic recurrent or drug-induced hypoglycemia, (g) has not been treated for schizophrenia with haloperidol, or (h) is not experiencing malignant hypertension. In further embodiments, the overweight or obese adolescent (a) has not been diagnosed as being type I or type II diabetic, (b) is not experiencing chronic, recurrent or drug-induced hypoglycemia, or (c) has not been diagnosed as having metabolic syndrome.

[0328] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage form three times per 24 hours to induce weight loss. This treatment can be the sole treatment to induce weight loss. In further embodiments, the overweight or obese subject (a) does not have an insulin-secreting tumor, (b) is not suffering from Poly Cystic Ovary Syndrome, (c) is not a type I diabetic, (d) is not a type II diabetic, (e) does not have metabolic syndrome, or (f) is not experiencing chronic, recurrent or drug-induced hypoglycemia.

[0329] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese adolescent as an oral dosage form three times per 24 hours to induce weight loss. This treatment can be the sole treatment to induce weight loss. In further embodiments, the overweight or obese adolescent is (a) is not a type I or type II diabetic, (b) is not experiencing chronic, recurrent or drug-induced hypoglycemia or (c) does not have metabolic syndrome.

[0330] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered as an oral dosage form three times per 24 hours to induce weight loss to an overweight or obese adult who (a) is not simultaneously receiving glucagon injections, triiodothyroxin or furosemide, (b) is not being treated for schizophrenia with haloperidol, or (c) is not experiencing malignant hypertension.

[0331] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage form four times per 24 hours to induce weight loss.

[0332] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage form administered from one, two, three or four times per 24 hours to induce weight loss at a daily dose of 50 to 700 mg. In a further embodiment, the overweight or obese subject (a) is not type I diabetic, (b) is not type II diabetic, (c) is not suffering chronic, recurrent or drug-induced hypoglycemia, or (d) does not have metabolic syndrome.

[0333] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral dosage form administered from one, two, three or four times per 24 hours to induce weight loss at a daily dose of 130 to 400 mg. In a further embodiment, the overweight or obese subject (a) is not type I diabetic, (b) is not type II diabetic, (c) is not suffering chronic, recurrent or drug-induced hypoglycemia, or (d) does not have metabolic syndrome.

[0334] In other embodiments, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obesity prone subject as an oral dosage form one, two, three or four times per 24 hours to maintain a weight loss, as it is preferable to maintain weight in an obese subject once some weight loss has occurred when the alternative is to regain weight. In a further embodiment, the administered daily dose of the K_{ATP} channel opener is 50 to 275 mg.

[0335] In other embodiments, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered as an oral dosage form to an overweight, obese, or obesity prone subject to (a) elevate energy expenditure, (b) elevate beta oxidation of fat, or (c) reduce circulating triglyceride concentrations.

[0336] In other embodiments, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to a subject in need thereof to induce the loss of 25%, 50%, or 75% of initial body fat.

[0337] In another embodiment, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject in need thereof to induce (a) the preferential loss of body fat or (b) the preferential loss of visceral body fat.

[0338] In additional embodiments, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered one, two or three times per 24 hours at daily doses of 50 to 700 mg to an subject to (a) induce the loss of 25%, 50% or 75% of initial body fat, (b) induce the preferential loss of body fat, or (c) induce the preferential loss of visceral fat.

[0339] In another embodiment, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject to induce the preferential loss of body fat and to induce reduction in circulating triglycerides.

[0340] In another embodiment, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with sibutramine, orlistat, rimonabant, an appetite suppressant, an anti-depressant, an anti-epileptic, a diuretic, a drug that induces weight loss by a mechanism that is distinct from a K_{ATP} channel opener, or a drug that lowers blood pressure, to induce weight loss and/or treat obesity associated co-morbidities in an overweight, obese, or obesity prone subject. In further embodiments, the overweight, obese, or obesity prone subject (a) is a type I diabetic, (b) is not a type II diabetic, (c) is not suffering from chronic, recurrent or drug-induced hypoglycemia, or (d) does not have metabolic syndrome.

[0341] In another embodiment an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with an anti-depressant, a drug that lowers blood pressure, a drug that lowers cholesterol, a drug that raises HDL, an anti-inflammatory that is not a Cox-2 inhibitor, a drug that lowers circulating triglycerides, to an overweight, obese, or obesity prone subject to induce weight loss and/or treat obesity associated co-morbidities. In further embodiments, the overweight, obese, or obesity prone subject (a) is not a type I diabetic, (b) is not a type II diabetic, (c) is not suffering from chronic, recurrent or drug-induced hypoglycemia, or (d) does not have metabolic syndrome.

[0342] In another embodiment, an oral dosage of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that lowers blood pressure, a drug that lowers cholesterol, a drug that raises HDL, an anti-inflammatory that is not a Cox-2 inhibitor, a drug that lowers circulating triglycerides, to maintain weight and/or treat obesity associated co-morbidities in an overweight, obese, or obesity prone subject, as it is preferable to maintain weight in an obese subject once some weight loss has occurred when the alternative is to regain weight. In further embodiments, the overweight, obese, or obesity prone subject (a) is not a type I diabetic, (b) is not a type II diabetic, (c) is not suffering from chronic, recurrent or drug-induced hypoglycemia, or (d) does not have metabolic syndrome.

[0343] In additional embodiments, an oral dosage form of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is used to administer a therapeutically effective dose of a K_{ATP} channel opener to an obese, overweight or obesity prone subject in need thereof to treat obesity, to (a) provide beta cell rest, (b) treat type I or type II diabetes, or (c) prevent the occurrence of diabetes.

[0344] In additional embodiments, an oral dosage form of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with Phentermine or a derivative thereof to an obese adult or adolescent to induce weight loss and/or treat obesity and obesity-associated co-morbidities. In further embodiments, a solid oral dosage form or tablet formulation of a K_{ATP} channel opener is co-administered with Phentermine or a derivative thereof to an obese adult or adolescent to treat metabolic syndrome in a patient in need thereof.

[0345] In further embodiments, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII at doses of 50 to 700 mg/day is co-administered with Phentermine or a derivative thereof at daily doses of 15 to 37.5 mg to an overweight or obese subject to induce weight loss, to treat metabolic syndrome, or to induce weight loss and treat obesity-associated co-morbidities.

[0346] In another embodiment, a quick dissolving formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is used to provide a therapeutically effective dose to a patient in need thereof.

[0347] In further embodiments, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered once per 24 hours at doses of 50 mg to 700 mg to an overweight or obese subject.

[0348] In further embodiments, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is formulated as a tablet or capsule for oral administration. The tablet or capsule may be co-formulated with metformin. In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is formulated as an oral suspension or solution, and the oral suspension or solution may be further encapsulated in another embodiment.

[0349] In another embodiment, a pharmaceutical salt of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is formulated as a tablet or capsule for oral administration, or as an oral suspension or as an oral solution, or as an oral suspension or solution that is encapsulated.

[0350] In another embodiment a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-formulated with hydro-chlorothiazide, chlorothiazide, cyclothiazide, benzthiazide, metyclothiazide, bendro-flumethiazide, hydroflumethiazide, trichlormethiazide, or polythiazide in a pharmaceutical formulation suitable for oral administration.

[0351] Upon administration of formulations which include a salt of a compound of Formulae I – VIII provided herein to humans or animals, some or all of the following effects are observed: (1) the production of lipoprotein lipase by adipocytes is reduced; (2) enhanced lipolysis by adipocytes; (3) expression of fatty acid synthase by adipocytes is reduced; (4) glyceraldehydes phosphate dehydrogenase activity of adipocytes is reduced; (5) little or no new triglycerides are synthesized and stored by adipocytes; (6) enhanced expression of β_3 Adrenergic Receptor (β_3AR) an improvement in the adrenergic function in adipocytes; (7) reduced glucose stimulated secretion of insulin by pancreatic B-cells; (8) decreased insulinemia; (9) enhanced blood glucose levels; (10) increased expression of Uncoupling Protein 1 in adipocytes; (11) enhanced thermogenesis in white and brown adipose tissue; (12) reduction of plasma triglyceride concentration; (13) decrease in circulating leptin concentrations; (14) up-regulation of insulin receptors; (15) enhanced

glucose uptake; (16) reduced adipocyte hyperplasia; (17) reduced adipocyte hypertrophy; (18) reduced rates of conversion of preadipocytes to adipocytes; (19) reduced rates of hyperphagia; (20) increased protection of CNS, cardiac and other tissues from ischemic or reperfusion injury; (21) improved insulin sensitivity; (22) elevated CSF insulin concentrations; (23) elevated circulating adiponectin concentrations; (25) reduced circulating triglyceride concentrations; (26) enhancement of beta-cell rest.

[0352] Threshold concentrations of the current invention include those circulating concentrations of K_{ATP} channel openers resulting from the administration of the selected from salts of compounds of Formulae I – VIII as an i.v. formulation, an immediate release oral formulation, a controlled release formulation, a transdermal formulation, or an intranasal formulation to an overweight or obese subject which results in (1) measurable suppression of fasting insulin levels, (2) suppression of fasting insulin levels by at least 20% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (3) suppression of fasting insulin levels by at least 30% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (4) suppression of fasting insulin levels by at least 40% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (5) suppression of fasting insulin levels by at least 50% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (6) suppression of fasting insulin levels by at least 60% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (7) suppression of fasting insulin levels by at least 70% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (8) suppression of fasting insulin levels by at least 80% from the baseline measurement in the same subject prior to treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII, (9) loss of weight, (10) elevation of resting energy expenditure, or (11) elevation of the oxidation of fat or fatty acids. Threshold effects of the current invention include those circulating concentrations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII resulting from the administration

of an i.v. formulation of the drug, or an immediate release oral formulation of the drug, or a controlled release formulation of the drug, or a sustained release formulation, or a transdermal formulation, or an intranasal formulation of the drug to an obesity prone subject which result in (1) the loss of weight, and (2) the maintenance of weight.

Threshold effects of the current invention include those circulating concentrations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII resulting from the administration of an i.v. formulation of the drug, or an immediate release oral formulation of the drug, or a controlled release formulation of the drug, or a sustained release formulation, or a transdermal formulation, or an intranasal formulation of the drug to a prediabetic subject which result in prevention of the transition to diabetes. Threshold effects of the current invention include those circulating concentrations of K_{ATP} channel openers resulting from the administration of selected from salts of compounds of Formulae I – VIII as an i.v. formulation, or an immediate release oral formulation, or a controlled release formulation, or a sustained release formulation, or a transdermal formulation, or an intranasal formulation to a subject with type I diabetes which result in beta cell rest.

[0353] The mode of action by which weight is maintained or lost resulting from the prolonged administration of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to overweight, obese or obesity prone subjects as provided herein includes, but is not limited to, one or more of (1) enhanced energy expenditure, (2) enhanced oxidation of fat and fatty acids, (3) enhancement of lipolysis in adipose tissue, (4) enhanced glucose uptake by tissues and enhanced insulin sensitivity, and (5) improved beta adrenergic response. The mode of action by which weight is maintained or lost resulting from the prolonged administration of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to obese or obesity prone subjects as provided herein may also include the suppression of appetite.

[0354] Prolonged administration of pharmaceutical formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to overweight or obese humans or animals results in substantial and sustained weight loss including some or all of the following effects: (1) preferential loss of body fat; (2) loss of greater than 25% of initial body fat mass; (3) loss of greater than 50% of initial body fat mass; (4) loss of

greater than 75% of initial body fat mass; (5) significant increase in resting energy expenditure; (6) increase in the oxidation of fat and fatty acids; (7) reduction in blood pressure; (8) production of lipoprotein lipase by adipocytes is reduced; (9) enhanced lipolysis by adipocytes; (10) expression of fatty acid synthase by adipocytes is reduced; (11) glyceraldehyde phosphate dehydrogenase activity of adipocytes is reduced; (12) little or no new triglycerides are synthesized and stored by adipocytes; (13) enhanced expression of β_3 Adrenergic Receptor (β_3 AR) and an improvement in the adrenergic function in adipocytes; (14) reduced glucose stimulated secretion of insulin by pancreatic B-cells; (15) decreased insulinemia; (16) enhanced blood glucose levels; (17) increased expression of Uncoupling Protein 1 in adipocytes; (18) enhanced thermogenesis in white and brown adipose tissue; (19) reduction of plasma triglyceride concentration; (20) decrease in circulating leptin concentrations; (21) up-regulation of insulin receptors; (22) enhanced glucose uptake; (23) reduced adipocyte hyperplasia; (24) reduced adipocyte hypertrophy; (25) reduced rates of conversion of preadipocytes to adipocytes; (26) reduced rates of hyperphagia; (27) the sequential loss first of the metabolically most active adipose tissue (visceral), followed by the loss of less metabolically active adipose tissue; (28) elevation of circulating adiponectin concentrations; (29) elevation of cerebrospinal fluid insulin levels; (30) enhanced islet insulin mRNA and insulin content; or (31) enhanced metabolic efficiency of insulin.

[0355] Prolonged administration of formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to obesity prone humans or animals, including subjects who have undergone various types of bariatric surgery, results in sustained maintenance of weight including some or all of the following effects: (1) increased resting energy expenditure; (2) increase in the oxidation of fat and fatty acids; (3) reduction in blood pressure; (4) production of lipoprotein lipase by adipocytes is reduced; (5) enhanced lipolysis by adipocytes; (6) expression of fatty acid synthase by adipocytes is reduced; (7) glyceraldehyde phosphate dehydrogenase activity of adipocytes is reduced; (8) little or no new triglycerides are synthesized and stored by adipocytes; (9) enhanced expression of β_3 Adrenergic Receptor (β_3 AR) and improvement in the adrenergic function in adipocytes; (10) reduced glucose stimulated secretion of insulin by pancreatic B-cells; (11) decreased insulinemia; (12) enhanced blood glucose levels; (13) increased expression of Uncoupling Protein 1 in adipocytes; (14) enhanced thermogenesis

in white and brown adipose tissue; (15) reduction of plasma triglyceride concentration; (16) decreased circulating leptin concentration; (17) up-regulation of insulin receptors; (18) enhanced glucose uptake; (19) reduced adipocyte hyperplasia; (20) reduced adipocyte hypertrophy; (21) reduced rates of conversion of preadipocytes to adipocytes; (22) reduced rates of hyperphagia; (23) elevated circulating adiponectin concentration; (24) elevated cerebrospinal fluid insulin levels; (25) enhanced islet insulin mRNA and insulin content; or (26) enhanced metabolic efficiency of insulin.

[0356] Immediate or prolonged administration of formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to prediabetic or type I diabetic humans or animals results in the prevention of beta cell failure, improved glycemic control, and prevention of the transition from prediabetes to diabetes including some or all of the following effects: (1) increase in resting energy expenditure; (2) increase in the oxidation of fat and fatty acids; (3) reduction in blood pressure; (4) production of lipoprotein lipase by adipocytes is reduced; (5) enhanced lipolysis by adipocytes; (6) expression of fatty acid synthase by adipocytes is reduced; (7) glyceraldehyde phosphate dehydrogenase activity of adipocytes is reduced; (8) little or no new triglycerides are synthesized and stored by adipocytes; (9) enhanced expression of β_3 Adrenergic Receptor (β_3AR) and an improvement in the adrenergic function in adipocytes; (10) reduced glucose stimulated secretion of insulin by pancreatic B-cells; (11) decreased insulinemia; (12) enhanced blood glucose levels; (13) increased expression of Uncoupling Protein 1 in adipocytes; (14) enhanced thermogenesis in white and brown adipose tissue; (15) reduction of plasma triglyceride concentration; (16) decreased circulating leptin concentrations; (17) up-regulation of insulin receptors; (18) enhanced glucose uptake; (19) reduced adipocyte hyperplasia; (20) reduced adipocyte hypertrophy; (21) reduced rates of conversion of preadipocytes to adipocytes; (22) reduced rates of hyperphagia; (23) elevated circulating adiponectin concentrations; (24) elevated cerebrospinal fluid insulin levels; (25) enhanced islet insulin mRNA and insulin content; or (26) enhanced metabolic efficiency of insulin.

[0357] Immediate or prolonged administration of formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII to humans or animals that are at risk for myocardial infarct, or stroke, or undergoing surgical procedure that restores

blood flow to heart or brain results in improved therapeutic outcomes post-surgically, or following the occurrence of myocardial infarct or stroke by improving the survival of tissue after blood flow is restored, reduced stunning of tissue, and altering the nature of the inflammatory responses.

[0358] Pharmaceutical formulations as provided herein are designed to be used in the treatment of obesity, hyperlipidemia, hypertension, weight maintenance, type I diabetes, prediabetes, type II diabetes, metabolic syndrome or any condition where weight loss, reduction in circulating triglycerides or beta cell rest contributes to therapeutic outcomes provide for a range of critical changes in pharmacodynamic and pharmacokinetic responses to administered doses of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII which changes include one or more of the following: (1) extending the pharmacodynamic effect of an administered dose to 24 hours or longer as measured by the suppression of insulin secretion; (2) providing for substantial uptake of the active pharmaceutical ingredient in the small intestine; (3) providing for substantial uptake of the active pharmaceutical ingredient in the large intestine; (4) result in lowered C_{max} versus current oral suspension or capsule products for the same administered dose of active pharmaceutical ingredient; (5) provide for circulating concentrations of unbound active pharmaceutical ingredient above threshold concentrations for 24 or more hours from a single administered dose; and (6) provide for more consistent drug absorption by treated subjects as compared to existing capsule formulations.

[0359] Pharmaceutical co-formulations of the current invention designed to treat a range of conditions in humans and animals include the combination of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII with: (1) a diuretic, (2) a drug that lowers blood pressure, (3) a drug that suppresses appetite, (4) a cannabinoid receptor antagonist, (5) a drug that suppresses that action of gastric lipases, (6) any drug that is used to induce weight loss, (7) a drug that lowers cholesterol, (8) a drug that lowers LDL bound cholesterol, (9) a drug that improves insulin sensitivity, (10) a drug that improves glucose utilization or uptake, (11) a drug that reduces incidence of atherosclerotic plaque, (12) a drug that reduces inflammation, (13) a drug that is antidepressant, (14) a drug that is an anti-epileptic, or (15) a drug that is an anti-psychotic.

[0360] Treatment of humans or animals using pharmaceutical formulations (which include K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII) result in reduced incidence of adverse side effects including but not limited to edema, fluid retention, reduced rates of excretion of sodium, chloride, and uric acid, hyperglycemia, ketoacidosis, nausea, vomiting, dyspepsia, ileus and headaches. These reductions in frequency of adverse side effects are achieved by: (1) initiating dosing of subjects at subtherapeutic doses and in a step wise manner increasing the dose daily until the therapeutic dose is achieved where the number of days over which the step up in dose is effected is 2 to 10, (2) use of the lowest effective dose to achieve the desired therapeutic effect, (3) use of a pharmaceutical formulation that delays release of active until gastric transit is complete, (4) use of a pharmaceutical formulation that delays release of active until gastric transit is complete, (5) use of a pharmaceutical formulation that results in lower circulating peak drug levels as compared to an immediate release oral suspension or capsule formulation for the same administered dose, and (6) optimizing the timing of administration of dose within the day and relative to meals.

[0361] Treatment of patients suffering from Prader Willi Syndrome, Froelich's Syndrome, Cohen Syndrome, Summit Syndrome, Alstrom Syndrome, Borjesen Syndrome, Bardet-Biedl Syndrome, and hyperlipoproteinemia type I, II, III, and IV with the current invention using pharmaceutical formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII result in some or all of the following therapeutic outcomes: (1) weight loss, (2) reduced rates of weight gain, (3) inhibition of hyperphagia, (4) reduced incidence of impaired glucose tolerance, prediabetes or diabetes, (5) reduced incidence of congestive heart failure, (6) reduced hypertension, and (7) reduced rates of all cause mortality.

[0362] Treatment of prediabetic subjects using invention pharmaceutical formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII result in some or all of the following therapeutic outcomes: (1) weight loss, (2) restoration of normal glucose tolerance, (3) delayed rates of progression to diabetes, (4) reduced hypertension, and (5) reduced rates of all cause mortality.

[0363] Treatment of diabetic subjects using invention pharmaceutical formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII result in

some or all of the following therapeutic outcomes: (1) weight loss, (2) restoration of normal glucose tolerance, (3) delayed rates of progression of diabetes, (4) improvements in glucose tolerance, (5) reduced hypertension, and (6) reduced rates of all cause mortality.

[0364] Co-administration of drugs with formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII in the treatment of diseases of overweight, obese or obesity prone human and animal subjects involves the co-administration of a pharmaceutically acceptable formulation of K_{ATP} channel openers with an acceptable formulation of: (1) sibutramine, (2) orlistat, (3) rimonabant, (4) a drug that is an appetite suppressant, (5) any drug used to induce weight loss in an obese or overweight subject, (6) a non-thiazide diuretic, (7) a drug that lowers cholesterol, (8) a drug that raises HDL cholesterol, (9) a drug that lowers LDL cholesterol, (10) a drug that lowers blood pressure, (11) a drug that is an anti-depressant, (12) a drug that improves insulin sensitivity, (13) a drug that improves glucose utilization and uptake (14) a drug that is an anti-epileptic, (15) a drug that is an anti-inflammatory, or (16) a drug that lowers circulating triglycerides.

[0365] Co-administration of drugs with formulations of K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII in the treatment or prevention of weight gain, dyslipidemia, impaired glucose tolerance or diabetes in subjects treated with antipsychotics drugs involve the co-administration of a pharmaceutically acceptable formulation of K_{ATP} channel openers with an acceptable formulation of: lithium, carbamazepine, valproic acid and divalproex, and lamotrigine; antidepressants generally classified as monoamine oxidase inhibitors including isocarboxazid, phenelzine sulfate and tranylcypromine sulfate; tricyclic antidepressants including doxepin, clomipramine, amitriptyline, maprotiline, desipramine, nortriptyline, desipramine, doxepin, trimipramine, imipramine and protriptyline; tetracyclic antidepressants including mianserin, mirtazapine, maprotiline, oxaprotiline, delequamine, levoprotline, trifluorcarbine, setipiline, lortalaline, azipramine, aptazapine maleate and pirlindole; and major tranquilizers and atypical antipsychotics including paloproxidol, perphenazine, thioridazine, risperidone, clozapine, olanzapine and chlorpromazine.

[0366] In one embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to measurably reduce fasting insulin levels for a prolonged period. Preferably the K_{ATP} channel opener formulation reduces fasting insulin levels by at least 20%, more preferably by at least 30%, more preferably by at least 40%, more preferably by at least 50%, more preferably by at least 60%, more preferably by at least 70%, and more preferably by at least 80%. Fasting insulin levels are commonly measured using the glucose tolerance test (OGTT). After an overnight fast, a patient ingests a known amount of glucose. Initial glucose levels are determined by measuring pre-test glucose levels in blood and urine. Blood insulin levels are measured by a blood is draw every hour after the glucose is consumed for up to three hours. In a fasting glucose assay, subjects with plasma glucose values greater than 200 mg/dl at 2 hours post-glucose load indicate an impaired glucose tolerance.

[0367] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to induce weight loss for a prolonged period.

[0368] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to elevate resting energy expenditure for a prolonged period.

[0369] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to elevate fat and fatty acid oxidation for a prolonged period.

[0370] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an obesity prone subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to induce weight loss for a prolonged period.

[0371] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an obesity prone subject as an oral, transdermal or intranasal formulation to reach and maintain the threshold concentration required to maintain weight for a prolonged period.

[0372] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain a drug concentration above the threshold concentration required to induce weight loss for a prolonged period.

[0373] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation for a prolonged period of time to reduce body fat by more than 25%, more preferably by at least 50%, and more preferably by at least 75%.

[0374] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation for a prolonged period of time to preferentially reduce visceral fat deposits.

[0375] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an overweight or obese subject as an oral, transdermal or intranasal formulation for a prolonged period of time to reduce visceral fat depots and other fat deposits.

[0376] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to a normoinsulinemic overweight or obese subject as an oral, transdermal or intranasal formulation to reach and maintain a drug concentration above the threshold concentration required to induce weight loss for a prolonged period.

[0377] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to a prediabetic subject as an oral, transdermal or intranasal formulation to reach and maintain a drug concentration above

the threshold concentration required to prevent the transition to diabetes for a prolonged period.

[0378] In another embodiment, a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to a type I diabetic subject as an oral, transdermal or intranasal formulation to reach and maintain a drug concentration above the threshold concentration required to induce beta cell rest for a prolonged period.

[0379] In another embodiment, a single dose of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject in need thereof that results in circulating concentration of active drug sufficient to diminish the secretion of insulin for 24 or more hours.

[0380] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to diminish the secretion of insulin on a continuous basis.

[0381] In another embodiment, a single dose of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject in need thereof that results in circulating concentration of active drug sufficient to elevate non-esterified fatty acids in circulation for 24 or more hours.

[0382] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to elevate non-esterified fatty acids in circulation on a continuous basis.

[0383] In another embodiment, a single dose of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject in need thereof that results in circulating concentration of active drug sufficient to treat hypoglycemia in circulation for 24 or more hours.

[0384] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to treat hypoglycemia on a continuous basis.

[0385] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to induce weight loss on a continuous basis.

[0386] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to maintain weight on a continuous basis, as it is preferable to maintain weight in an obese subject once some weight loss has occurred when the alternative is to regain weight.

[0387] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that results in circulating concentration of active drug sufficient to reduce circulating triglyceride levels on a continuous basis.

[0388] In another embodiment, a single dose of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered to an subject in need thereof that results in circulating concentration of active drug sufficient to reduce or prevent ischemic or reperfusion injury in circulation for 24 or more hours.

[0389] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is administered over a prolonged basis to an subject in need thereof no more than once per 24 hours that

results in circulating concentration of active drug sufficient reduce or prevent ischemic or reperfusion injury on a continuous basis.

[0390] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation of diazoxide or its derivatives that is administered to an subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached.

[0391] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an subject in need thereof on a daily basis in which the active ingredient is not released from the formulation until gastric transit is complete.

[0392] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an subject in need thereof on a daily basis in which the maximum circulating concentration of active ingredient is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation of Proglycem®.

[0393] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached, the active ingredient is not release from the formulation until gastric transit is complete and in which the maximum circulating concentration of active ingredient is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation of Proglycem®.

[0394] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is

reduced using a pharmaceutically acceptable formulation that is administered to an overweight or obese subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached, the active ingredient is not release from the formulation until gastric transit is complete, in which the maximum circulating concentration of active ingredient is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation of Proglycem®, and in which the maximum dose is less than 5 mg/kg/day.

[0395] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an overweight or obese subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached, the active ingredient is not release from the formulation until gastric transit is complete, in which the maximum circulating concentration of active ingredient is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation, and in which the maximum dose is less than 2.5 mg/kg/day.

[0396] In another embodiment, the treatment of an overweight or obese subject is optimized for weight loss by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide continuous release for at least 6 hours.

[0397] In another embodiment, the treatment of an overweight or obese subject is optimized for weight loss by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide continuous release for at least 12 hours.

[0398] In another embodiment, the treatment of an overweight or obese subject is optimized for weight loss by administration of a pharmaceutically acceptable formulation

of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide a rising drug concentration in circulation for at least 8 hours.

[0399] In another embodiment, the treatment of an overweight or obese subject is optimized for weight loss by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide a rising drug concentration in circulation for at least 12 hours.

[0400] In another embodiment, the treatment of an overweight or obese subject is optimized for weight loss by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to match the pattern of basal insulin secretion.

[0401] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an obesity prone subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached, the active ingredient is not release from the formulation until gastric transit is complete, in which the maximum circulating concentration of active ingredient is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation, and in which the maximum dose is less than 5 mg/kg/day.

[0402] In another embodiment, the frequency of adverse effects caused by treatment with a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is reduced using a pharmaceutically acceptable formulation that is administered to an obesity prone subject in need thereof on a daily basis in which the first dose is known to be subtherapeutic and daily dose is subsequently increased stepwise until the therapeutic dose is reached, the active ingredient is not release from the formulation until gastric transit is complete, in which the maximum circulating concentration of active ingredient

is lower than what would be realized by the administration of the same dose using an oral suspension or capsule formulation, and in which the maximum dose is less than 2.5 mg/kg/day.

[0403] In another embodiment, the treatment of an obesity prone subject is optimized for weight maintenance by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide continuous release for at least 6 hours.

[0404] In another embodiment, the treatment of an obesity prone subject is optimized for weight maintenance by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide continuous release for at least 12 hours.

[0405] In another embodiment, the treatment of an obesity prone subject is optimized for weight maintenance by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide a rising drug concentration in circulation for at least 8 hours.

[0406] In another embodiment, the treatment of an obesity prone subject is optimized for weight maintenance by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to provide a rising drug concentration in circulation for at least 12 hours.

[0407] In another embodiment, the treatment of an obesity prone subject is optimized for weight maintenance by administration of a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII once per 24 hours in which the release of the active ingredient from the formulation has been modified to match the pattern of basal insulin secretion.

[0408] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with sibutramine to an overweight or obese subject to induce weight loss.

[0409] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with orlistat to an overweight or obese subject to induce weight loss.

[0410] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with rimonabant to an overweight or obese subject to induce weight loss.

[0411] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with an appetite suppressant to an overweight or obese subject to induce weight loss.

[0412] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with an anti-depressant to an overweight or obese subject to induce weight loss.

[0413] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with anti-epileptic to an overweight or obese subject to induce weight loss.

[0414] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener is selected from a salt of a compound of Formulae I – VIII is co-administered with a non-thiazide diuretic to an overweight or obese subject to induce weight loss.

[0415] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that induces weight loss by a mechanism that is distinct from diazoxide to an overweight or obese subject to induce weight loss.

[0416] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that lowers blood pressure to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0417] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that lowers cholesterol to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0418] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that raises HDL associated cholesterol to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0419] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that improves insulin sensitivity to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0420] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with an anti-inflammatory to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0421] In another embodiment, a pharmaceutically acceptable formulation of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII is co-administered with a drug that lowers circulating triglycerides to an overweight, obesity prone or obese subject to induce weight loss and treat obesity associated co-morbidities.

[0422] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with sibutramine in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0423] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with orlistat or other active that suppresses the action of gastric lipases in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0424] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a non-thiazide diuretic in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0425] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an appetite suppressant in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0426] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a cannabinoid receptor antagonist in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0427] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an anti-cholesteremic active in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0428] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an antihypertensive active in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0429] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an insulin sensitizing active in a

pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0430] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an anti-inflammatory active in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0431] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an anti-depressant active in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0432] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an anti-epileptic active in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0433] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an active that reduces the incidence of atherosclerotic plaque in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0434] In another embodiment, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with an active that lowers circulating concentrations of triglycerides in a pharmaceutically acceptable formulation that is administered to an overweight, obesity prone or obese subject to induce weight loss and treat obesity-associated co-morbidities.

[0435] The reduction of circulating triglycerides in an overweight, obese or obesity prone subject is achieved by the administration of an effective amount of an oral dosage form of a K_{ATP} channel opener, selected from a salt of a compound of Formulae I – VIII.

[0436] An oral dosage form of K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII can be used to administer a therapeutically effective dose of K_{ATP} channel opener to an overweight or obesity prone subject in need thereof to maintain weight, as it is preferable to maintain weight in an obese subject once some weight loss has occurred when the alternative is to regain weight.

[0437] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat obesity. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such obesity treatment drugs include, but are not limited to: sibutramine hydrochloride (5-30 mg), orlistat (50-360 mg), phentermine hydrochloride or resin complex (15 to 40 mg), zonisamide (100 to 600 mg), topiramate (64 to 400 mg), naltrexone hydrochloride (50 to 600 mg), or rimonabant (5 to 20 mg).

[0438] A further embodiment of the co-formulation contains K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII and a drug to treat obesity. Such co-formulations can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 12 hours. Such obesity treatment drugs include, but are not limited to: sibutramine hydrochloride (2.5 to 15 mg), orlistat (25 to 180 mg), phentermine hydrochloride or resin complex (7.5 to 20 mg), zonisamide (50 to 300 mg), topiramate (32 to 200 mg), naltrexone hydrochloride (25 to 300 mg), or rimonabant (2.5 to 10 mg).

[0439] In another embodiment of the invention K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat obesity, diabetes, metabolic syndrome or an obesity related comorbidity. Such drugs to treat these conditions include drugs that: agonizes the α_1 -noradrenergic receptor; agonizes the β_2 noradrenergic receptor; stimulates noradrenalin release; blocks noradrenalin uptake; stimulates 5-HT release; blocks 5-HT uptake; is a serotonin (5-hydroxytryptamine) 2C receptor agonist; antagonizes acetyl-CoA carboxylase 2; agonizes the D1-receptor; antagonizes the H3-receptor; is a leptin analogue; agonizes the leptin receptor, sensitizes CNS tissue to the action of leptin; agonizes the MC4 receptor; agonizes NPY-Y1; agonizes NPY-Y2; agonizes NPY-Y4; agonizes NPY-Y5; antagonizes the MCH receptor;

blocks CRH-BP; agonizes the CRH receptor; agonizes the urocortin receptor; antagonizes the galanin receptor; antagonizes the orexin receptor; agonizes the CART receptor; agonizes the Amylin receptor; agonizes the Apo(A^{IV}) receptor; antagonizes the CB-1 receptor; is an α MSH analogue; inhibits PTP-1B; antagonizes PPAR γ receptor; is a short acting bromocriptine; agonizes somatostatin; increases adiponectin; increases CCK activity; increases PYY activity; increases GLP-1 activity; decreases ghrelin activity; is a selective B3 stimulator or agonist; agonizes thyroid receptor; inhibits gastrointestinal lipases or other digestive enzymes; blocks absorption of dietary fat; or block de-novo fatty acid synthesis. Additionally, such drugs to treat obesity may include, but are not limited to those that antagonize or agonize the function or expression of 11B hydroxysteroid dehydrogenase type 1; acetyl-CoA carboxylase 1; ADAM 12, member 12 of a disintegrin and metalloprotease family or its shorter secreted form; agouti related protein; angiotensinogen; adipocyte lipid binding protein; adipocyte fatty acid binding protein; adrenergic receptors; acylation-stimulating protein; bombesin receptor subtype-3; C/EBP, CCAAT/enhancer binding protein; cocaine- and amphetamine-regulated transcript; cholecystokinin; cholecystokinin A receptor; CD36, fatty acid translocase; corticotropin-releasing hormone; diacylglycerol acyltransferases; E2F transcription factor; eukaryotic translation initiation factor 4e binding protein 1; estrogen receptor; fatty acid synthase; fibroblast growth factor; forkhead box C2; glucose-dependent insulinotropic peptide; GIP receptor; inhibitory G protein α -subunit; glucagon-like peptide-1; GLP-1 receptor; glycerol-3-phosphate acyltransferase; glycerol 3-phosphate dehydrogenase; stimulatory G protein α -subunit; high-mobility group phosphoprotein isoform I-C; hormone sensitive lipase; inducible nitric oxide synthase; Janus kinases; lipoprotein lipase; melanocortin-3 receptor; melanocortin-4 receptor; mitochondrial GPAT; metallothionein-I and -II; nescient helix-loop-helix 2; neuropeptide Y; neuropeptide Y-1 receptor; neuropeptide Y-2 receptor; neuropeptide Y-4 receptor; neuropeptide Y-5 receptor; plasminogen activator inhibitor-1; PPAR γ co-activator 1; pro-opiomelanocortin; peroxisome proliferator-activated receptor; protein tyrosine phosphatase 1B; regulatory subunit II β of protein kinase A; retinoid X receptor; steroidogenic factor 1; single-minded 1; sterol regulatory element binding protein; tyrosine hydroxylase; thyroid hormone receptor α 2; uncoupling protein; nerve growth factor induced protein; leucine zipper transcription factor; α -melanocyte-stimulating hormone.

[0440] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat diabetes. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such diabetes treatment drugs include, but are not limited to: acarbose (50 to 300 mg), miglitol (25 to 300 mg), metformin hydrochloride (300 to 2000 mg), repaglinide (1-16 mg), nateglinide (200 to 400 mg), or rosiglitazone (5 to 50 mg).

[0441] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 12 hours. Such drugs to treat diabetes include, but are not limited to: acarbose (25 to 150 mg), miglitol (12.5 to 150 mg), metformin hydrochloride (150 to 1000 mg), repaglinide (0.5 to 8 mg), nateglinide (100 to 200 mg), or rosiglitazone (2.5 to 25 mg).

[0442] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat elevated cholesterol. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such drugs to treat elevated cholesterol include, but are not limited to: pravastatin, simvastatin, atorvastatin, fluvastatin, rosuvastatin or lovastatin (10 to 80 mg).

[0443] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 12 hours. Such drugs to treat elevated cholesterol include, but are not limited to: pravastatin, simvastatin, atorvastatin, fluvastatin, rosuvastatin or lovastatin (5 to 40 mg).

[0444] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat depression. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained

release of the active over a period of 2 to 24 hours. Such drugs to treat depression include, but are not limited to: citalopram hydrobromide (10 to 80 mg), escitalopram hydrobromide (5 to 40 mg), fluvoxamine maleate (25 to 300 mg), paroxetine hydrochloride (12.5 to 75 mg), fluoxetine hydrochloride (30 to 100 mg), setraline hydrochloride (25 to 200 mg), amitriptyline hydrochloride (10 to 200 mg), desipramine hydrochloride (10 to 300 mg), nortriptyline hydrochloride (10 to 150 mg), duloxetine hydrochloride (20 to 210 mg), venlafaxine hydrochloride (37.5 to 150 mg), phenelzine sulfate (10 to 30 mg), bupropion hydrochloride (200 to 400 mg), or mirtazapine (7.5 to 90 mg).

[0445] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active for 2 to 12 hours. Such drugs to treat depression include, but are not limited to: citalopram hydrobromide (5 to 40 mg), escitalopram hydrobromide (2.5 to 20 mg), fluvoxamine maleate (12.5 to 150 mg), paroxetine hydrochloride (6.25 to 37.5 mg), fluoxetine hydrochloride (15 to 50 mg), setraline hydrochloride (12.5 to 100 mg), amitriptyline hydrochloride (5 to 100 mg), desipramine hydrochloride (5 to 150 mg), nortriptyline hydrochloride (5 to 75 mg), duloxetine hydrochloride (10 to 100 mg), venlafaxine hydrochloride (18 to 75 mg), phenelzine sulfate (5 to 15 mg), bupropion hydrochloride (100 to 200 mg), or mirtazapine (4 to 45 mg).

[0446] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat hypertension. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such drugs to treat hypertension include, but are not limited to: enalapril maleate (2.5 to 40 mg), captopril (2.5 to 150 mg), lisinopril (10 to 40 mg), benazepril hydrochloride (10 to 80 mg), quinapril hydrochloride (10 to 80 mg), perindopril erbumine (4 to 8 mg), ramipril (1.25 to 20 mg), trandolapril (1 to 8 mg), fosinopril sodium (10 to 80 mg), moexipril hydrochloride (5 to 20 mg), losartan potassium (25 to 200 mg), irbesartan (75 to 600 mg), valsartan (40 to 600 mg), candesartan cilexetil (4 to 64 mg), olmesartan medoxamil (5 to

80 mg), telmisartan (20 to 160 mg), eprosartan mesylate (75 to 600 mg), atenolol (25 to 200 mg), propranolol hydrochloride (10 to 180 mg), metoprolol tartrate, succinate or fumarate (25 to 400 mg), nadolol (20 to 160 mg), betaxolol hydrochloride (10 to 40 mg), acebutolol hydrochloride (200 to 800 mg), pindolol (5 to 20 mg), bisoprolol fumarate (5 to 20 mg), nifedipine (15 to 100 mg), felodipine (2.5 to 20 mg), amlodipine besylate (2.5 to 20 mg), nicardipine (10 to 40 mg), nisoldipine (10 to 80 mg), terazosin hydrochloride (1 to 20 mg), doxazosin mesylate (4 to 16 mg), prazosin hydrochloride (2.5 to 10 mg), or alfuzosin hydrochloride (10 to 20 mg).

[0447] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of active over a period of 2 to 12 hours. Such drugs to treat hypertension include, but are not limited to: enalapril maleate (1.25 to 20 mg), captopril (2 to 75 mg), lisinopril (5 to 20 mg), benazepril hydrochloride (5 to 40 mg), quinapril hydrochloride (5 to 40 mg), perindopril erbumine (2 to 4 mg), ramipril (1 to 10 mg), trandolapril (1 to 4 mg), fosinopril sodium (5 to 40 mg), moexipril hydrochloride (2.5 to 10 mg), losartan potassium (12.5 to 100 mg), irbesartan (37.5 to 300 mg), valsartan (20 to 300 mg), candesartan cilexetil (2 to 32 mg), olmesartan medoxamil (2.5 to 40 mg), telmisartan (10 to 80 mg), eprosartan mesylate (37.5 to 300 mg), atenolol (12.5 to 100 mg), propranolol hydrochloride (5 to 90 mg), metoprolol tartrate, succinate or fumarate (12.5 to 200 mg), nadolol (10 to 80 mg), betaxolol hydrochloride (5 to 20 mg), acebutolol hydrochloride (100 to 400 mg), pindolol (2.5 to 10 mg), bisoprolol fumarate (2.5 to 10 mg), nifedipine (7.5 to 50 mg), felodipine (1 to 10 mg), amlodipine besylate (1 to 10 mg), nicardipine (5 to 20 mg), nisoldipine (5 to 40 mg), terazosin hydrochloride (1 to 10 mg), doxazosin mesylate (2 to 8 mg), prazosin hydrochloride (1 to 5 mg), or alfuzosin hydrochloride (5 to 10 mg).

[0448] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a diuretic. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such diuretics can include, but are not limited to: amiloride hydrochloride (1 to 10 mg), spironolactone (10 to 100 mg), triamterene (25 to

200 mg), bumetanide (0.5 to 4 mg), furosemide (10 to 160 mg), ethacrynic acid or ethacrynate sodium (10 to 50 mg), tosemide (5 to 100 mg), chlorthalidone (10 to 200 mg), indapamide (1 to 5 mg), hydrochlorothiazide (10 to 100 mg), chlorothiazide (50 to 500 mg), bendroflumethiazide (5 to 25 mg), hydroflumethiazide (10 to 50 mg), mythyclothiazide (1 to 5 mg), and polythiazide (1 to 10 mg).

[0449] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 12 hours. Such diuretics include, but are not limited to: amiloride hydrochloride (0.5 to 5 mg), spironolactone (5 to 50 mg), triamterene (12 to 100 mg), bumetanide (0.2 to 2 mg), furosemide (5 to 80 mg), ethacrynic acid or ethacrynate sodium (5 to 25 mg), tosemide (2 to 50 mg), chlorthalidone (5 to 100 mg), indapamide (0.5 to 2.5 mg), hydrochlorothiazide (5 to 50 mg), chlorothiazide (25 to 250 mg), bendroflumethiazide (2 to 12.5 mg), hydroflumethiazide (5 to 25 mg), mythyclothiazide (0.5 to 2.5 mg), and polythiazide (0.5 to 5 mg).

[0450] In another embodiment of the invention, K_{ATP} channel openers selected from salts of compounds of Formulae I – VIII are co-formulated with a drug to treat inflammation or pain. Such co-formulations can be formulated for oral administration once per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 24 hours. Such drugs to treat inflammation or pain include, but are not limited to: aspirin (100 to 1000mg), tramadol hydrochloride (25 to 150 mg), gabapentin (100 to 800 mg), acetaminophen (100 to 1000 mg), carbamazepine (100 to 400 mg), ibuprofen (100 to 1600 mg), ketoprofen (12 to 200 mg), fenprofen sodium (100 to 600 mg), flurbiprofen sodium or flurbiprofen (50 to 200 mg), or combinations of these with a steroid or aspirin.

[0451] In a further embodiment, the co-formulation can be formulated for oral administration twice per 24 hours, for delayed release of the active until gastric transit is complete, and for sustained release of the active over a period of 2 to 12 hours. Such drugs to treat inflammation or pain include, but are not limited to: aspirin (100 to 650 mg), tramadol hydrochloride (12 to 75 mg), gabapentin (50 to 400 mg), acetaminophen (50 to 500 mg), carbamazepine (50 to 200 mg), ibuprofen (50 to 800 mg), ketoprofen (6

to 100 mg), fenpropfen sodium (50 to 300 mg), flurbiprofen sodium or flurbiprofen (25 to 100 mg), or combinations of these with a steroid or aspirin.

[0452] A method of inducing loss of greater than 25% of initial body fat in an overweight or obese subject can be achieved by the prolonged administration of an oral dosage form of a K_{ATP} channel opener, selected from a salt of a compound of Formulae I – VIII.

[0453] A method of inducing loss of greater than 50% of initial body fat in an overweight or obese subject can be achieved by the prolonged administration of an oral dosage form of a K_{ATP} channel opener selected from a salt of a compound of Formulae I – VIII.

[0454] A method of inducing loss of greater than 75% of initial body fat in an overweight or obese subject can be achieved by the prolonged administration of an oral dosage form of a K_{ATP} channel opener, selected from a salt of a compound of Formulae I – VIII.

[0455] A method of inducing preferential loss of visceral fat in an overweight or obese subject can be achieved by the prolonged administration of an oral dosage form of a K_{ATP} channel opener, selected from a salt of a compound of Formulae I – VIII.

[0456] A method of inducing loss of body fat and reductions in circulating triglycerides in an overweight or obese subject can be achieved by the prolonged administration of an oral dosage form of a K_{ATP} channel opener, selected from a salt of a compound of Formulae I – VIII.

[0457] In some embodiments, the invention provides a polymorph of a salt, which salt includes diazoxide and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or quaternary ammonium group. In some embodiments, the cation is choline.

[0458] In some embodiments, the polymorph of diazoxide choline salt is of Form A having characteristic peaks in the XRPD pattern at values of two-theta ($\text{Cu K}\alpha$, 40 kV, 40

mA) at approximately 9.8, 10.5, 14.9, 17.8, 17.9, 18.5, 19.5, 22.1, 22.6, 26.2, 29.6, and 31.2 degrees.

[0459] In some embodiments, the polymorph of diazoxide choline salt is of Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.9, 10.3, 12.0, 18.3, 20.6, 24.1, 24.5, 26.3, 27.1, and 28.9 degrees.

[0460] In some embodiments, the polymorph of diazoxide choline salt is of Form A having characteristic infrared absorbances at 2926, 2654, 1592, 1449, and 1248 cm⁻¹.

[0461] In some embodiments, the polymorph of diazoxide choline salt is of Form B having characteristic infrared absorbances at 3256, 2174, 2890, 1605, 1463, and 1235 cm⁻¹.

[0462] In some embodiments, the polymorph of diazoxide includes potassium as the cation.

[0463] In some embodiments, the polymorph of diazoxide potassium salt is of Form A having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 6.0, 8.1, 16.3, 17.7, 18.6, 19.1, 22.9, 23.3, 23.7, 24.7, 25.4, 26.1, 28.2, 29.6, and 30.2 degrees.

[0464] In some embodiments, the polymorph of diazoxide potassium salt is of Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 10.8, 16.9, 18.2, 21.6, 25.5, 26.1, and 28.9 degrees.

[0465] In some embodiments, the polymorph of diazoxide potassium salt is of Form C having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.7, 6.1, 17.9, 23.9, 25.1, and 37.3 degrees.

[0466] In some embodiments, the polymorph of diazoxide potassium salt is of Form D having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.7, 6.2, 8.1, 8.5, 8.8, 16.9, 18.6, 23.2, 24.5, 25.8, and 26.1 degrees.

[0467] In some embodiments, the polymorph of diazoxide potassium salt is of Form E having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 6.7, 7.1, 14.1, and 21.2 degrees.

[0468] In some embodiments, the polymorph of diazoxide potassium salt is of Form F having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 9.0, 18.7, 20.6, 23.5, 27.5, and 36.3 degrees.

[0469] In some embodiments, the polymorph of diazoxide potassium salt is of Form G having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.2, 5.5, 13.1, 16.5, 19.3, 22.8, 24.8, 26.4, 28.7, and 34.1 degrees.

[0470] In some embodiments, the polymorph of diazoxide potassium salt is of Form A having characteristic infrared absorbances at 1503, 1374, 1339, 1207, 1131, 1056, and 771 cm^{-1} .

[0471] In some embodiments, the polymorph of diazoxide potassium salt is of Form B having characteristic infrared absorbances at 1509, 1464, 1378, and 1347 cm^{-1} .

[0472] In some embodiments, the polymorph of diazoxide potassium salt is of Form C having characteristic infrared absorbances at 1706, 1208, 1146, and 746 cm^{-1} .

[0473] In some embodiments, the polymorph of diazoxide potassium salt is of Form D having characteristic infrared absorbances at 1595, 1258, 1219, and 890 cm^{-1} .

[0474] In some embodiments, the polymorph of diazoxide potassium salt is of Form E having characteristic infrared absorbances at 1550, 1508, 1268, 1101, and 1006 cm^{-1} .

[0475] In some embodiments, the polymorph of diazoxide potassium salt is of Form F having characteristic infrared absorbances at 1643, 1595, 1234, 1145, and 810 cm^{-1} .

[0476] In some embodiments, the polymorph of diazoxide potassium salt is of Form G having characteristic infrared absorbances at 1675, 1591, 1504, 1458, 1432, 1266, 999, 958, 905, and 872 cm^{-1} .

[0477] In some embodiments, the polymorph of diazoxide choline salt is of Form A having characteristic peaks in the XRPD pattern substantially as shown in FIG. 16(a), and an NMR spectrum substantially as shown in FIG. 17(a).

[0478] In some embodiments, the polymorph of diazoxide choline salt is of Form B having characteristic peaks in the XRPD pattern substantially as shown in FIG. 16(c) and an NMR spectrum substantially as shown in FIG. 17(b).

[0479] In some embodiments, the polymorph of diazoxide potassium salt includes one or more of Forms A-G, wherein each of the Forms A-G has characteristic peaks in the XRPD pattern substantially as shown in FIG. 18-19.

[0480] In some embodiments of the invention there are provided methods for producing a diazoxide choline salt, which methods include suspending diazoxide in a solvent (e.g., alcohols such as methanol, i-BuOH, i-AmOH, t-BuOH, and the like, ketones, tetrahydrofuran, dimethylformamide, n-methyl pyrrolidinone, and the like) and mixing with a choline salt (e.g., choline hydroxide), adding a co-solvent (e.g., MTBE, EtOA, IPA, c-Hexane, heptane, toluene, CH₂Cl₂, dioxane, and the like) to the suspension under conditions sufficient to cause formation and precipitation of said diazoxide choline salt, and harvesting the precipitate to provide the diazoxide choline salt.

[0481] In some embodiments, the solvent is tetrahydrofuran. In some embodiments, the solvent is 2-methyltetrahydrofuran (2-MeTHF).

[0482] In some embodiments, the diazoxide and the solvent are present at a ratio of about 1 g diazoxide per 1 mL solvent to about 1 g diazoxide per 5 mL solvent. In some embodiments, the diazoxide and the solvent are present at a ratio of about 1 g diazoxide per 3 mL solvent.

[0483] In some embodiments, the choline salt is a solution in MeOH. In some embodiments, the choline salt is choline hydroxide in about a 45% solution (e.g., 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%) in MeOH.

[0484] In some embodiments, the choline salt is added as 1 equivalent of diazoxide.

[0485] In some embodiments, the co-solvent is MTBE.

[0486] In some embodiments, the amount of co-solvent added is in a ratio to the amount of the solvent of about 3:14 (solvent:co-solvent) (e.g., 3:12, 3:13, 3:14, 3:15, 3:16).

[0487] In some embodiments, the process of making polymorphs of diazoxide choline salt includes the step of seeding with crystals of diazoxide choline salt polymorph Form B prior to the harvesting step.

[0488] In some embodiments for the method for producing a diazoxide choline salt the salt includes polymorph Form B substantially free of polymorph Form A, the polymorph Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 10.8, 16.9, 18.2, 21.6, 25.5, 26.1, and 28.9 degrees.

[0489] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the compound is a compound of Formula V.

[0490] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the compound is a compound of Formula VI.

[0491] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the compound is a compound of Formula VII.

[0492] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the compound is a compound of Formula VIII.

[0493] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the method further comprises administering a drug selected from the group consisting of Sibutramine, Orlistat, Rimonabant, an appetite suppressant, a non-thiazide diuretic, a drug that lowers cholesterol, a drug that raises HDL cholesterol, a drug that lowers LDL cholesterol, a drug that lowers blood pressure, a drug that is an anti-depressant, a drug that is an anti-epileptic, a drug that is an anti-inflammatory, a drug that is an appetite suppressant, a drug that lowers circulating triglycerides, and a drug that is used to induce weight loss in an overweight or obese individual.

[0494] In some embodiments for the method of treating obesity or obesity-related morbidity in an obese subject, the method further comprises administering a pharmaceutically active agent other than the K_{ATP} channel opener. In some embodiments, the other pharmaceutically active agent is an agent useful for the treatment of a condition selected from the group consisting of obesity, prediabetes, diabetes, hypertension, depression, elevated cholesterol, fluid retention, obesity associated co-morbidities, ischemic and reperfusion injury, epilepsy, cognitive impairment, schizophrenia, mania, and other psychotic condition.

[0495] In some embodiments for the method for treating a subject suffering from or at risk for Alzheimer's disease (AD), the method includes administration to a subject a therapeutically effective amount of a salt of diazoxide including salts provided herein. In some embodiments for the method for treating a subject suffering from or at risk for AD, the method includes administration to a subject a therapeutically effective amount of a compound according to any of Formulae I-VIII. In some embodiments, the compound is diazoxide or a salt thereof.

[0496] AD is a neurodegenerative disorder neuropathologically characterized by abnormal accumulations of intracellular neurofibrillary tangles and extracellular amyloid plaques throughout cortical and limbic brain regions and the loss of synapses and neurons. AD is further characterized by significant cognitive and memory impairment. β amyloid plaques form from the β amyloid peptide (i.e., "A β ", H-Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala-OH, SEQ ID NO: __), either 1-40 or 1-42 peptide, which is released from amyloid precursor protein following cleavage by gamma secretase. In addition to forming plaques, the β amyloid peptides are cytotoxic either as the monomer or as a short-lived oligomeric intermediate. β amyloid peptides (monomers, dimers or oligomers) can be identified both in CSF (cerebrospinal fluid) and in serum. Amyloid angiopathy is characterized by A β deposition and may contribute to the cerebrovascular abnormalities that precede the onset of AD.

[0497] The invention will now be described with reference to the following non-limiting examples.

EXAMPLES

A. Potassium ATP Channel Activator Containing Formulations

1. Compressed Tablet Formulations of Diazoxide salt or derivative

[0498] Diazoxide salt or a derivative thereof at about 15-30% by weight is mixed with hydroxypropyl methylcellulose at about 55-80% by weight, ethylcellulose at about 3-10 wt/vol % and magnesium stearate (as lubricant) and talc (as glidant) each at less than 3% by weight. The mixture is used to produce a compressed tablet as described in Reddy et al., *AAPS Pharm Sci Tech* 4(4):1-9 (2003). The tablet may be coated with a thin film as discussed below for microparticles.

[0499] A tablet containing 100 mg of diazoxide salt or a derivative thereof will also contain approximately 400 mg of hydroxypropyl cellulose and 10 mg of ethylcellulose. A tablet containing 50 mg of diazoxide salt or a derivative thereof will also contain approximately 200 mg of hydroxypropyl cellulose and 5 mg of ethylcellulose. A tablet containing 25 mg of diazoxide salt or a derivative thereof will also contain approximately 100 mg of hydroxypropyl cellulose and 2.5 mg of ethylcellulose.

2. Encapsulated Coated Microparticle Formulation of Diazoxide Salt or derivative

[0500] Diazoxide salt or a derivative thereof is encapsulated into microparticles in accordance with well known methods (see, e.g. U.S. Patent No. 6,022,562). Microparticles of between 100 and 500 microns in diameter containing diazoxide salt or a derivative thereof, alone or in combination with one or more suitable excipient, is formed with the assistance of a granulator and then sieved to separate microparticles having the appropriate size. Microparticles are coated with a thin film by spray drying using commercial instrumentation (e.g. Uniglatt Spray Coating Machine). The thin film comprises ethylcellulose, cellulose acetate, polyvinylpyrrolidone and/or polyacrylamide. The coating solution for the thin film may include a plasticizer which may be castor oil, diethyl phthalate, triethyl citrate and salicylic acid. The coating solution may also include a lubricating agent which may be magnesium stearate, sodium oleate, or polyoxyethylenated sorbitan laurate. The coating solution may further include an

excipient such as talc, colloidal silica or of a mixture of the two added at 1.5 to 3% by weight to prevent caking of the film coated particles.

3. Formulations for controlled release of diazoxide or derivative

3.1. Formulation of a tableted form of diazoxide or a derivative for controlled release

[0501] Prior to mixing, both the active ingredient and hydroxypropyl methylcellulose (Dow Methocel K4M P) are passed through an ASTM 80 mesh sieve. A mixture is formed from 1 part diazoxide salt or a derivative thereof to 4 parts hydroxypropyl methylcellulose. After thorough mixing, a sufficient volume of an ethanolic solution of ethylcellulose as a granulating agent is added slowly. The quantity of ethylcellulose per tablet in the final formulation is about 1/10th part. The mass resulting from mixing the granulating agent is sieved through 22/44 mesh. Resulting granules are dried at 40°C for 12 hours and thereafter kept in a desiccator for 12 hours at room temperature. Once dry the granules retained on 44 mesh are mixed with 15% fines (granules that passed through 44 mesh). Talc and magnesium stearate are added as glidant and lubricant at 2% of weight each. A colorant is also added. The tablets are compressed using a single punch tablet compression machine.

3.2. Formulation of a compression tableted form of diazoxide or a derivative thereof that provides for controlled release.

[0502] Diazoxide salt or a derivative thereof at 20-40% weight is mixed with 30% weight hydroxypropyl methylcellulose (Dow Methocel K100LV P) and 20-40% weight impalpable lactose. The mixture is granulated with the addition of water. The granulated mixture is wet milled and then dried 12 hours at 110°C. The dried mixture is dry milled. Following milling, 25% weight ethylcellulose resin is added (Dow Ethocel 10FP or Ethocel 100FP) followed by 0.5% weight magnesium stearate. A colorant may also be added. The tablets are compressed using a single punch tablet compression machine (Dasbach et al., Poster at AAPS Annual Meeting Nov. 10-14 (2002)).

3.3. Formulation of a compression coated tableted form of diazoxide or a derivative thereof that provides for controlled release.

[0503] The core tablet is formulated by mixing either 100 mg of diazoxide salt or a derivative thereof with 10 mg of ethylcellulose (Dow Ethocel 10FP), or by mixing 75 mg

of diazoxide or a derivative thereof with 25 mg lactose and 10 mg of ethylcellulose (Dow Ethocel 10FP), or by mixing 50 mg of diazoxide or a derivative thereof with 50 mg of lactose and 10 mg of ethylcellulose (Dow Ethocel 10FP). The core tablets are formed on an automated press with concave tooling. The compression coating consisting of 400 mg of polyethylene oxide (Union Carbide POLYOX WSR Coagulant) is applied and compressed to 3000 psi (Dashbach et al., Poster at AAPS Annual Meeting Oct. 26-30 (2003)).

3.4. Formulation of a controlled release tableted form of diazoxide choline salt

3.4.1 Controlled Release Formulations

[0504] Controlled release tableted formulations of diazoxide choline salt were developed and investigated with respect to a variety of properties known by those of skill in the pharmaceutical art relating to the manufacture of tableted formulations including, for example, ease and consistency of manufacture, appearance (e.g., sheen, compressibility, microscopic appearance), and dissolution properties (e.g., rate, order and extent of dissolution). Tablets were produced individually on a press, where the final blend of diazoxide choline salt and excipient was weighed out to the desired total tablet weight prior to compression. As shown in Table 2, Formulations A-H, J, and L contained 50.0 mg diazoxide as the choline salt (i.e., 72.5 mg total diazoxide choline salt present), and Formulations I and K contained 200.0 mg diazoxide as the choline salt (i.e., 290.0 mg total diazoxide choline salt present). The manufacture of formulation L is exemplary of the manufacturing methods available to the skilled artisan. For formulation L, diazoxide choline salt, talc, and approximately half of the colloidal silicon dioxide (Cab-o-sil) were mixed in a KG-5 mixer bowl with an impeller speed of about 300 rpm and a chopper speed of about 3000 rpm for about 4 min. The mixture was passed through a co-mil equipped with a 024R screen, square-edged paddle, and 0.175" spacer. To this milled mixture in a 8-qt V-shell blender was added Emcompress through a #20 mesh hand screen with blending for about 10 min. To this mixture was added PEO N750 and PEO 303, which had been passed through a #20 mesh hand screen, with blending for about 10 min. To this mixture was added Pruv and the remainder of the Cab-o-sil, having been passed through a #20 mesh hand screen, with blending for about 5 min. The mixture was

subjected to pressing (Manesty Beta Press) using 0.2220" x 0.5720" caplet shaped tooling (Set #21.)

Table 2. Exemplary Formulations for Diazoxide Choline salt.

Formulation	INGREDIENT	Amount per tablet (mg)
A.	Diazoxide Choline	72.50
	Kollidon SR	100.0
	PEO N303	25.00
	Emcompress	50.50
	Pruv	2.00
	TOTAL WEIGHT	250.0
B.	Diazoxide Choline	72.50
	Methocel K100M	237.3
	Emcompress	35.00
	Magnesium Stearate	3.50
	Cab-o-sil	1.75
	TOTAL WEIGHT	350.0
C.	Diazoxide Choline	72.50
	Kollidon SR	105.0
	PEO N303	35.0
	Emcompress	134.1
	Pruv	3.50
	TOTAL WEIGHT	350.1
D.	Diazoxide Choline	72.50
	Methocel K100M	175.1
	Emcompress	97.30
	Magnesium Stearate	3.50
	Cab-o-sil	1.75
	TOTAL WEIGHT	350.2
E.	Diazoxide Choline	72.50
	PEO N750 NF	105.0
	PEO N303 NF	52.50
	Emcompress	116.6
	Pruv	3.50

	TOTAL WEIGHT	350.1
F.	Diazoxide Choline	72.50
	Kollidon SR	105.0
	PEO N303	7.00
	Emcompress	162.1
	Pruv	3.50
	TOTAL WEIGHT	350.1
G.	Diazoxide Choline	72.50
	Methocel K100M	175.1
	Emcompress	105.1
	Magnesium Stearate	3.50
	Cab-o-sil	1.75
	TOTAL WEIGHT	358.0
H.	Diazoxide Choline	72.50
	PEO N750 NF	105.0
	PEO N303 NF	35.01
	Emcompress	134.1
	Pruv	3.50
	TOTAL WEIGHT	350.1
I.	Diazoxide Choline	290.0
	Methocel K100M	240.0
	Emcompress	258.0
	Magnesium Stearate	8.00
	Cab-o-sil	4.00
	TOTAL WEIGHT	800.0
J.	Diazoxide Choline	72.50
	PEO N750 NF	105.0
	PEO N303 NF	52.50
	Emcompress	116.6
	Pruv	3.50
	Talc	3.50
	Cab-o-sil	1.75

	TOTAL WEIGHT	355.4
K.	Diazoxide Choline	290.0
	PEO N750 NF	249.0
	PEO N303 NF	124.5
	Emcompress	145.3
	Pruv	8.30
	Talc	8.30
	Cab-o-sil	4.15
	TOTAL WEIGHT	829.6
L.	Diazoxide Choline	72.51
	PEO N750 NF	105.1
	PEO N303 NF	52.54
	Emcompress	111.4
	Pruv	3.50
	Talc	3.50
	Cab-o-sil	1.75
	TOTAL WEIGHT	350.3

[0505] Microscopic observation of the tablets having Formulation A revealed a grainy texture of the diazoxide choline salt, and at the 29% loading of diazoxide choline salt in Formulation A the blend had poor flow characteristics. Accordingly, the loading of diazoxide choline was reduced in Formulation B. A caplet shaped tooling, approximately 6 mm x 15 mm was found to result in acceptable tablet appearance, sheen, and ease of compressibility. However, Formulation B also exhibited poor blend flow.

[0506] Subsequent formulations (e.g., Formulations C-L) incorporated a milling step of the diazoxide choline salt prior to incorporation into the tablet blend. A milling study was conducted using a test mill equipped with different screen sizes to evaluate and determine a suitable milling process. Particle size was determined by visual comparison with 40 μ m reference beads. As shown in Table 3, use of the 024R screen resulted in the best recovery of API (i.e., "active pharmaceutical ingredient" = diazoxide as the diazoxide choline salt) providing the broadest range of particle sizes. Accordingly, material milled through a 024R screen was selected for subsequent formulation.

Table 3. Milling study for diazoxide choline salt prior to formulation

Screen/Paddle/Spacer/Speed	API loaded/API recovered (g)	Particle size (μm)
018R/square/0.200"/80%	50.0/36.5	~100-200
024R/square/0.200"/80%	50.0/42.0	~100-250
039R/square/0.200"/80%	50.0/42.0	~150-250

3.4.2. Dissolution Studies

[0507] Dissolution of tablets with formulation as set forth in Table 4 was investigated. One or more tablets (e.g., 1 or 2) of the indicated tableted formulation were placed into a volume of buffer (e.g., 900 mL) with known buffer salt concentration (e.g., 0.05 M potassium phosphate, pH 8.6; 0.05 M potassium phosphate, pH 7.5), with or without surfactant (e.g., 0.05% CTAB), at a known temperature (e.g., 37 °C). Stirring conditions employed paddles at, for example, 50 RPM. Aliquots (e.g., 10 mL) removed as a function of time were filtered (e.g., 0.45 μm GMF Filter) prior to analysis.

Table 4. Dissolution study for formulations of diazoxide choline salt (entries are % dissolved diazoxide)

Entry	Protocol	Time (hr)								
		0	1	2	3	6	9	12	18	24
C	a	0.0	2.9	4.5	5.7	9.4	13.0	16.6		
D	a	0.0	4.5	7.8	11.0	19.1	25.6	31.3		
F	b				29	36		44		54
G	b				30	39		50		67
H	b				50	75		91		101
I	b				25	37		48		57
J	b				34	53		76		104
K	b				25 24	43 41		64 62		94 92

Column 1: Entry from Table 2.

Column 2: Protocol a) 0.05 M potassium phosphate, pH 8.6, 37 °C; Protocol b) 0.05 M potassium phosphate, pH 7.5, 0.05% CTAB, 37 °C.

[0508] The dissolution profile (i.e., % dissolved with time) of Proglycem® capsules (100 mg) and controlled-release tablet formulations of diazoxide as provided herein is shown in Table 5. The 50 mg tablet entry of Table 5 refers to Formulation J (Table 2)

wherein talc and cab-o-sil was present at 2% and 1%, respectively. The 200 mg tablet entry of Table 5 refers to Formulation K (Table 2) wherein talc and cab-o-sil are present at 2% and 1%, respectively

Table 5. Dissolution profile of 100 mg Proglycem® (capsule) and diazoxide choline salt (tablet)

Time (hr)	100 mg capsule Proglycem®	200 mg tablet diazoxide choline salt	50 mg tablet diazoxide choline salt
0	0	0	0
3	79	22	25
6	85	36	39
12	92	54	70
24	98	88	99

[0509] As evidenced by Table 5, the 100-mg Proglycem® capsule provides for faster dissolution of diazoxide relative to tablets described herein. Approximately 79% diazoxide component of Proglycem® is recovered in dissolution buffer after 3-hr. In contrast, the 50 and 200-mg tablets described in Table 5 dissolved at levels of 25% and 22%, respectively, at 3-hr. At 12-hr, 100-mg Proglycem® capsule dissolved at 92%, whereas the 50 and 200-mg tablets dissolved at 70% and 54%, respectively. Approximately total dissolution is observed with 100-mg Proglycem® capsule and 50-mg tablet at 24 hrs.

3.4.3. Excipient Compatibility Studies

[0510] Studies to determine the compatibility of diazoxide choline salt for various excipients are summarized in Table 6. Each mixture of excipient (100 mg) and diazoxide choline salt (100 mg) was made in acetonitrile to 10 mL. Samples were assayed immediately (i.e., "Initial" column of Table 6) and at one-month storage under conditions a) 40 °C/75% RH (relative humidity), and b) 50 °C.

Table 6. Excipient compatibility study for diazoxide choline salt.

Excipient	Initial %(w/w)	1 Month 40 °C/75% RH %(w/w)	1 Month 50 °C %(w/w)
Hydroxypropylmethyl cellulose (HMPC)	98.8	101.4	96.6
Hydroxypropylcellulose (HPC)	103.1	97.4	100.8
Ethylcellulose (EC)	90.6 99.5*	102.0	99.7
Methylcellulose (MC)	102.3	100.3	99.1
Carboxymethyl cellulose Na (CMC Na)	96.8	101.6	99.5
Starch 1500	103.3	90.6	95.2
Kollidon SR	81.2 100.2*	99.8	100.9
Polyethyleneoxide N3 03 (PEO)	81.0 99.1*	99.2	99.0
Dibasic Calcium Phosphate	98.4	102.9	102.2
Sodium Stearyl Fumarate	98.9	101.1	103.7
Magnesium Stearate	100.1	101.4	99.3
Colloidal Silicon Dioxide (Cah-o-sil)	99.0	99.4	104.5
Microcrystalline Cellulose	96.8	98.6	107.0
Lactose Monohydrate	94.5	99.2	103.0
Mannitol	111.7	98.1	81.3
Diazoxide Control	N/A	99.0	99.4

* Re-assayed

[0511] Initial low results for ethylcellulose, Kollidon SR, and polyethyleneoxide were investigated by re-preparing samples of diazoxide choline salt and excipient. As shown in Table 6, initial sample recovery (i.e., column "Recovery") of the re-prepared samples indicates method accuracy.

[0512] No reportable impurities were detected in any sample, and no degradation was observed in the samples stored for one-month with the exception of mannitol wherein 81.3% of diazoxide was recovered after 1-month at 50 °C.

3.4.3. Stability Studies for Diazoxide Choline Controlled Release Tablets

[0513] Studies to determine the stability of formulations of diazoxide choline salt as the controlled release tablet were conducted on sample formulations as described in Table 2, results of which are shown in Table 7. In these studies, storage conditions were the following: a) 25 °C/60% RH, and b) 40 °C/75% RH. In Table 7, the term "Appearance" refers to the physical appearance of the tablets of the study. The term "Assay (%)" refers to the percentage of diazoxide choline salt assayed with respect to nominal (i.e., 50 mg or 200 mg) content of the sample. The term "Dissolution (%)" refers to the amount, expressed as a percentage, of diazoxide choline salt assayed by method described herein.

Table 7. Stability Study for Formulations of Diazoxide Choline Salt.

Test	Time point		
	Initial	1 month	2 months
<i>50 mg tablet, Formulation J, 25 °C/60% RH,</i>			
Appearance	White tablets	White tablets	White tablets
Assay (%)	95.8	102.9	99.3
Dissolution (%):			
3 hr	24	19	15
6 hr	41	30	27
12 hr	62	48	52
24 hr	92	90	94
<i>50 mg tablet, Formulation J, 40 °C/75% RH,</i>			
Appearance	White tablets	White tablets	White tablets
Assay (%)	95.8	103.2	99.4
Dissolution (%):			
3 hr	24	10	10
6 hr	41	22	21
12 hr	62	50	46
24 hr	92	90	84
<i>200 mg tablet, Formulation K, 25 °C/60% RH</i>			
Appearance	White tablets	White tablets	White tablets
Assay (%)	95.9	100.9	100.3

Dissolution (%):			
3 hr	34	15	13
6 hr	53	31	30
12 hr	76	69	68
24 hr	104	101	96
<i>200 mg tablet, Formulation K, 40 °C/75% RH</i>			
Appearance	White tablets	White tablets	White tablets
Assay (%)	95.9	99.6	100.0
Dissolution (%):			
3 hr	34	10	10
6 hr	53	26	28
12 hr	76	60	61
24 hr	104	90	90

3.5. *A controlled release dosage form of diazoxide or a derivative thereof using an osmotically controlled release system.*

[0514] Diazoxide salt or a derivative thereof is formulated as an osmotically regulated release system. In general, two components, and an expandable hydrogel that drives release of the active drug is assembled with diazoxide salt or a derivative thereof into a semipermeable bilaminar shell. Upon assembly a hole is drilled in the shell to facilitate release of active upon hydration of the hydrogel.

[0515] A dosage form adapted, designed and shaped as an osmotic delivery system is manufactured as follows: first, a diazoxide salt or a derivative thereof composition is provided by blending together into a homogeneous blend of polyethylene oxide, diazoxide salt or a derivative thereof and hydroxypropyl methylcellulose. Then, a volume of denatured anhydrous ethanol weighing 70% of the dry mass is added slowly with continuous mixing over 5 minutes. The freshly prepared wet granulation is screened through a 20 mesh screen through a 20 mesh screen, dried at room temperature for 16 hours, and again screened through a 20 mesh screen. Finally, the screened granulation is mixed with 0.5% weight of magnesium stearate for 5 minutes.

[0516] A hydrogel composition is prepared as follows: first, 69% weight of polyethylene oxide weight, 25% weight of sodium chloride and 1% weight ferric oxide

are separately screened through a 40 mesh screen. Then, all the screened ingredients are mixed with 5% weight of hydroxypropyl methylcellulose to produce a homogeneous blend. Next, a volume of denatured anhydrous alcohol equal to 50% of the dry mass is added slowly to the blend with continuous mixing for 5 minutes. The freshly prepared wet granulation is passed through a 20 mesh screen, allowed to dry at room temperature for 16 hours, and again passed through a 20 mesh screen. The screened granulation is mixed with 0.5% weight of magnesium stearate for 5 minutes (see U.S. Patent No. 6,361,795 by Kuczynski, et al.).

[0517] The diazoxide salt composition, or a derivative thereof, and the hydrogel composition, are compressed into bilaminate tablets. First the diazoxide salt or a derivative thereof composition is added and tamped. The hydrogel composition is then added and the laminae are pressed under a pressure head of 2 tons into a contacting laminated arrangement.

[0518] The bilaminate arrangements are coated with a semipermeable wall (i.e. thin film). The wall forming composition comprises 93% cellulose acetate having a 39.8% acetyl content, and 7% polyethylene glycol. The wall forming composition is sprayed onto and around the bilaminate.

[0519] Finally, an exit passageway can be drilled through the semipermeable wall to connect the diazoxide salt or a derivative thereof drug lamina with the exterior of the dosage system. Residual solvent is removed by drying at 50° C and 50% humidity. The osmotic systems are dried at 50° C to remove excess moisture (see U.S. Patent No. 6,361,795 by Kuczynski, et al.).

4. Preparation of Diazoxide Salts

4.1. Preparation of the Sodium Salt

[0520] The sodium salt of diazoxide was prepared by dissolving 300 mg of diazoxide in approximately 45 mL methyl ethyl ketone (MEK). The diazoxide/MEK solution was heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added 1.3 mL of 1M NaOH (1 molar equivalent). The combined solutions were heated at 75°C for approximately 30 minutes and allowed to cool to room temperature. The mixture was concentrated under reduced pressure, and dried *in vacuo* at 55°C and 30 in. Hg.

Elemental analysis: Calculated, 38.03% C, 2.39% H, 11.09% N and 9.1% Na; Found, 38.40% C, 2.25% H, 10.83% N and 7.4% Na.

[0521] A sodium salt of diazoxide was also prepared by dissolving 300 mg of diazoxide in approximately 45 mL acetonitrile. The diazoxide/acetonitrile solution was heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added 1.3 mL of 1M NaOH (approximately 1 molar equivalent). The combined solutions were heated at 75°C for approximately 30 minutes and allowed to cool to room temperature. The mixture was concentrated under reduced pressure, and dried *in vacuo* at 55°C and 30 in. Hg.

4.2. Preparation of the Potassium Salt

[0522] The potassium salt of diazoxide was prepared by dissolving 300 mg of diazoxide in approximately 45 mL methyl ethyl ketone (MEK). The diazoxide/MEK solution was heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added approximately 1.3 mL of 1 M KOH (1 molar equivalent) and the solution was returned to the orbital shaker, heated at 75°C for approximately 30 minutes and allowed to cool to room temperature. The solvent was removed under reduced pressure and the solid was dried *in vacuo* at 55°C and 30 in. Hg. Elemental analysis: Calculated, 33.59% C, 2.81% H, 9.77% N and 13.63% K; Found, 34.71% C, 2.62% H, 9.60% N and 10.60% K.

[0523] The potassium salt of diazoxide was also prepared by dissolving 300 mg of diazoxide in approximately 45 mL tetrahydrofuran (THF). The diazoxide/THF solution was heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added approximately 1.3 mL of 1 M KOH (1 molar equivalent) and the resulting solution was returned to the orbital shaker, heated at 75°C for approximately 30 minutes and allowed to cool to room temperature. THF was removed under reduced pressure and the solid was dried *in vacuo* at 55°C and 30 in. Hg.

4.3. Preparation of Choline Salt

4.3.1. Preparation of the Choline Salt: proof of concept

[0524] The choline salt of diazoxide was prepared by dissolving 300 mg of diazoxide in approximately 45 mL methyl ethyl ketone (MEK). The diazoxide/MEK solution was

heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added approximately 315 mg of 50 wt. % of choline hydroxide (1 molar equivalent) and the solution was returned to the orbital shaker and stirred at 75°C for approximately 30 minutes. The solvent was removed under reduced pressure, and the solid was dried *in vacuo* at 55°C and 30 in. Hg. Elemental analysis: Calculated, 46.77% C, 5.86% H, and 12.00% N; Found, 46.25% C, 6.04% H, and 12.59% N.

4.3.2. Process of preparation of the Choline Salt

[0525] An investigation of the minimum solvent volumes required to optimize the formation of the diazoxide choline salt was performed in MeCN, THF, MEK, and 2-MeTHF on small scale in the presence and absence of MTBE. Analyses by ¹H NMR and XRPD of all reactions were consistent with the assigned structure and Form B of diazoxide choline salt.

4.3.2.1. Solvent Efficiency Study with Single Solvent System.

[0526] Investigation of single solvent system synthesis of diazoxide choline salt using solvents THF, MEK, and 2-MeTHF were conducted, results of which are shown in Table 8. These results suggest that precipitation can be achieved with a single solvent system in THF or 2-MeTHF. The best results, (i.e., Table 8 entries 14-17) obtained with 2-MeTHF. However, it was observed that in 2-MeTHF at 3-volumes and above, complete dissolution was not achieved after addition of choline hydroxide.

Table 8. Solvent Efficiency of a Single Solvent System

Entry	Solvent	Solvent volume	Yield (%)	Polymorphic Form
1	THF	1.0	7	B
2	THF	2.0	n/a	n/a
3	THF	3.0	20	B
4	THF	4.0	28	B
5	THF	5.0	32	B
6	THF	6.0	42	B
7	THF	7.0	53	B
8	THF	8.0	50	B
9	MEK	4.0	n/a	n/a
10	2-MeTHF	1.0	n/a	n/a

Entry	Solvent	Solvent volume	Yield (%)	Polymorphic Form
11	2-MeTHF	2.0	48	B
12	2-MeTHF	3.0	41	B
13	2-MeTHF	4.0	48	B
14	2-MeTHF	5.0	71	B
15	2-MeTHF	6.0	71	B
16	2-MeTHF	7.0	72	B
17	2-MeTHF	8.0	67	B

n/a: Samples were either seen to form a gum, or no precipitate was observed.

4.3.2.2. Solvent Efficiency Study with Binary Solvent Systems.

[0527] The addition of a second solvent at 1-20 volumes (i.e., 1-8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20) to the solvent volumes of Table 8 was envisaged to optimize diazoxide salt production yields. Results of investigation of methods to enhance precipitation and increase yield employing binary solvent system synthesis of diazoxide choline salt using solvents THF, MeCN, MEK, and 2-MeTHF and co-solvent MTBE are shown in Table 9. Optimum conditions for the preparation of the diazoxide choline salt obtained with 1-3 volumes of THF (see Table 8, entries 1-3), with 3 volumes being the ratio of choice to eliminate excessive drag during stirring of slurry in large scale production.

Table 9. Solvent Efficiency of a Binary Solvent System

Entry	Solvent	Solvent Volume	Co-Solvent (12 Vol)	Yield (%)	Polymorphic Form
1	THF	1.0	MTBE	96	B
2	THF	2.0	MTBE	96	B
3	THF	3.0	MTBE	91	B
4	THF	4.0	MTBE	90	B
5	THF	5.0	MTBE	84	B
6	THF	6.0	MTBE	90	B
7	THF	7.0	MTBE	94	B
8	THF	8.0	MTBE	94	B
9	MeCN	1.0	MTBE	92	B

Entry	Solvent	Solvent Volume	Co-Solvent (12 Vol)	Yield (%)	Polymorphic Form
10	MeCN	2.0	MTBE	90	B
11	MeCN	3.0	MTBE	87	B
12	MeCN	4.0	MTBE	79	B
13	MeCN	5.0	MTBE	79	B
14	MeCN	6.0	MTBE	75	B
15	MeCN	7.0	MTBE	70	B
16	MeCN	8.0	MTBE	74	B
17	MEK	4.0	MTBE	83	B
18	2-MeTHF	1.0	MTBE	97	B
19	2-MeTHF	2.0	MTBE	94	B
20	2-MeTHF	3.0	MTBE	94	B
21	2-MeTHF	4.0	MTBE	95	B
22	2-MeTHF	5.0	MTBE	97	B
23	2-MeTHF	6.0	MTBE	94	B
24	2-MeTHF	7.0	MTBE	97	B
25	2-MeTHF	8.0	MTBE	>99	B

4.3.2.3. Co-Solvent Efficiency and Optimization with MTBE.

[0528] Results of optimization of co-solvent (MTBE) volume in the presence of 3 volumes of solvent are shown in Table 10.

Table 10. Co-Solvent Efficiency and Optimization with MTBE

Entry	Solvent	MTBE Volume	Yield (%)	Form
1	MeCN	8.0	76	B
2	MeCN	14.0	85	B
3	THF	6.0	93	B
4	THF	8.0	93	B
5	THF	14.0	98	B

[0529] Optimized conditions for the production of the choline salt of diazoxide in a binary solvent system obtained with 3:14 (i.e., 1:4.7) THF/MTBE (v/v).

4.3.2.4. Optimization of Cooling Profile.

[0530] Optimization for the controlled precipitation of diazoxide choline salt was conducted by varying the cooling temperatures and hold times. Reactions were carried out in MeCN, THF, and 2-MeTHF (3 vol) with MTBE (14 vol). The results are presented in Table 11. These results suggest that optimum crystal growth was achieved with cooling to 0–5 °C for eight hours in THF and 2-MeTHF (Entries 5–6) when compared to those in MeCN (Entries 1, 2, and 4). These results however did not indicate enhanced precipitation when compared to previous studies carried out in THF and 2-MeTHF when the slurries were allowed to stir for two hours. In addition no notable improvements were observed for the additional cooling to –15 °C or with the filtration at ambient temperature.

Table 11. Cooling Profile Optimization for Controlled Precipitation

Entry	Solvent	Cooling Temp. (°C)	Hold Time (hr)	Yield (%)	Form
1	MeCN	0–5	2	89	B
2	MeCN	–10 → –15	2	85	B
3	THF	RT	2	80	B
4	MeCN	RT	2	85	B
5	THF	0–5	8	94	B
6	2-MeTHF	0–5	8	95	B

4.3.2.5. Optimization of Rate of Co-solvent addition.

[0531] The effect of rate and hold time of co-solvent addition on the precipitation of diazoxide choline salt was examined. The reactions were carried out on 500 mg scale in three volumes of primary solvent and 14 volumes of MTBE (THF and 2-MeTHF) or 1:3 volume ratio (MeCN). Addition of the co-solvent was made dropwise in one portion for Entry 1, and for Entries 2 and 3 the co-solvent was added at a rate of 1 mL/20 minutes for a total hold time of 140 minutes for the addition of 7 mL. The results, presented in Table 12, indicate that optimum precipitation of diazoxide choline salt was achieved by the addition of MTBE with 20 minute hold times in between additions in THF or 2-MeTHF (entries 2-3).

Table 12. Optimization of Rate of Co-solvent addition

Entry	Solvent	Hold Time (min)	Addition Rate (mL/min)	Yield (%)	Form
-------	---------	--------------------	---------------------------	-----------	------

1	MeCN	60	n/a	87	B
2	THF	140	1.0/20	96	B
3	2-MeTHF	140	1.0/20	96	B

4.3.2.6. Thermal Stability Study.

[0532] The stability of the isolated diazoxide choline salt was investigated to determine optimal drying conditions to minimize residual solvents without degradation. Samples of the diazoxide choline salt prepared in THF and MTBE (Entries 1–4) were dried under vacuum at 40 °C for various durations on small scale (100 mg). The study was then carried out on 5 g scale (Entry 5) at 30 °C for eight hours under vacuum to show reproducibility on large scale. The results are presented in Table 13. Analyses of the samples of Table 13 by ¹H NMR, XRPD, and HPLC indicated that elevated temperatures and prolonged drying times did not degrade the compound or affect the form of the compound.

Table 13. Thermal Stability Study

Entry	Drying Temp (°C)	Time (h)	OVI (THF/MTBE) (ppm)	Polymorphic Form
1	RT	12	376/248	B
2	40	20	317/160	B
3	40	28	276/127	B
4	40	42	268/128	B
5	30	8	241/509	B

4.3.2.6. Demonstration of 50-g Scale synthesis of Diazoxide Choline Salt in THF.

[0533] The preparation of the diazoxide choline salt was carried out in a binary-solvent system of THF and MTBE with a ratio of 1:4.7 (solvent/co-solvent) volumes and cooling to 0–5 °C for two hours with stirring. A demonstration run for the large scale production utilizing the modified procedure was carried out on 50 g-scale. Diazoxide (50 g) as a hot (62 °C) suspension in THF (140 mL) was treated with choline hydroxide (45% solution in MeOH, 1.0 equiv) added (2 mL/min) over 30 minutes. The resulting solution was stirred for 30 minutes, followed by cooling to 52 °C for the addition of MTBE (14 vol) over 45 minutes. On addition of two volumes of co-solvent precipitation was

observed. The resultant slurry was then allowed to cool naturally to ambient temperature followed by further cooling to 0–5 °C with an ice/water bath and an additional 2 hr stirring. The precipitate was isolated by vacuum filtration, and the filter cake rinsed with ice cold MTBE (~50 mL) and dried under vacuum at room temperature for 12 hours. OVI analysis indicated that THF levels were above the recommended ICH guidelines (799 ppm) and the material was returned to the oven at 30 °C for eight hours to give diazoxide choline salt [70.87 g, 97% yield, 97% purity AUC] as a white crystalline solid. Analyses by ¹H NMR, XRPD, and HPLC were consistent with the assigned structure of Form B while maintaining a high purity with good yield.

4.3.2.7. Demonstration of 50-g Scale synthesis of Diazoxide Choline Salt in 2-MeTHF.

[0534] A 50-g level synthesis was duplicated in 2-MeTHF/MTBE as an alternative to THF/MTBE for the large scale production of the diazoxide choline salt described above. No issues or concerns arose during the reaction other than complete dissolution was not achieved after addition of the choline hydroxide. A white precipitate formed after addition of the co-solvent, which was isolated via vacuum filtration and dried under vacuum at 30 °C for eight hours to give diazoxide choline salt [71.51 g, 97% yield] as a white crystalline solid. Analyses by ¹H, XRPD, and HPLC were consistent with the assigned structure and Form B. OVI analysis showed 2-MeTHF and MTBE levels of 125 and 191 ppm, respectively, which were below the ICH guidelines.

4.3.2.8. 250-g Scale synthesis of Diazoxide Choline Salt in THF.

[0535] Larger scale preparation of the diazoxide choline salt was carried out in a binary-solvent system of THF and MTBE with a ratio of 1:4.7 (v/v) volumes. Diazoxide as a hot (62 °C) suspension in THF (745 mL) was treated with choline hydroxide as a 45% solution in MeOH (1.0 equiv) added over 30 minutes. The resulting solution was stirred for 30 minutes, followed by cooling to 52 °C for the addition of MTBE (14 vol) over 45 minutes. On addition of two volumes of co-solvent, precipitation was observed. The resultant slurry was then allowed to cool naturally to ambient temperature followed by cooling to 0–5 °C with an ice/water bath. The precipitate was isolated by vacuum filtration, and the filter cake rinsed with ice cold MTBE (approximately 250 mL) and dried under vacuum at 30 °C for 38 hours to give diazoxide choline salt (350.28 g, 97.7%

yield) as a white crystalline solid. Analyses by ^1H NMR, XRPD, and HPLC were consistent with the assigned structure of Form B while maintaining a high purity with good yield.

4.3.2.9. 2-kg Scale Synthesis of Diazoxide Choline Salt in THF.

[0536] A 12-L reaction flash was charged with 2.0 kg diazoxide and 5.0-L THF with stirring and heating to 55 °C. Choline hydroxide (45% solution in methanol, 2.32 L) was added dropwise to this reaction mixture over about 2.5 hr with stirring. The temperature was maintained at 60 ± 5 °C. After addition of choline hydroxide, stirring was continued for about 30 min. The reaction mixture was clarified by in-line 10 micron filtration upon transfer to a 22-L reaction flask pre-charged with 2-L pre-filtered THF, into which was added 10-L pre-filtered MTBE dropwise. This reaction mixture was transferred to another flask which was then charged with an additional 30-L pre-filtered MTBE dropwise, with adjustment of temperature to < 5 °C and stirring for about 2 hr. Diazoxide choline salt was recovered by vacuum filtration to afford 2.724 kg (94%) diazoxide choline salt (99.8%, HPLC purity), confirmed by ^1H NMR, IR, and UV/Visible analysis.

4.4. Preparation of the Hexamethyl Hexamethylene Diammonium Hydroxide Salt

[0537] The hexamethyl hexamethylene ammonium salt of diazoxide was prepared by dissolving 50 mg of diazoxide in approximately 7.5 mL methyl ethyl ketone (MEK). The diazoxide/MEK solution was heated at 75°C on an orbital shaker to ensure dissolution. To the solution was added approximately 2.17mL of 0.1M hexamethyl hexamethylene ammonium hydroxide solution (1 molar equivalent) and the solution was stirred at 75°C for an additional 10 minutes and then cooled to room temperature at the rate of 30°C/h. The solvent was removed under reduced pressure, and the solid was dried *in vacuo* at 55°C and 30 in. Hg.

4.5. Failure to Obtain Salts of Diazoxide and derivatives

4.5.1. Failure to Obtain Salts from Alkali Metal Hydroxides

[0538] U.S. Patent Number 2,986,573 ("the '573 patent") describes the synthesis of diazoxide metal salts in aqueous or non-aqueous solutions in the presence of an alkali metal alkoxide. According to the '573 Patent, diazoxide can be dissolved in an alkali metal solution, and the salt obtained upon evaporation. Also described is a method for

forming salts from non-aqueous media wherein diazoxide and sodium methoxide are dissolved in anhydrous methanol and the solvent is evaporated to obtain the sodium salt of diazoxide as a white solid.

[0539] Attempts were made to prepare a diazoxide salt from alkali metal hydroxides by the method described in the '573 patent. Salt preparation was carried out in aqueous media by dissolving diazoxide in a basic solution 1M NaOH, followed by evaporation of the solvent. A solid was obtained and analyzed by XRPD (X-Ray Powder Diffraction) and NMR. However, this analysis confirmed that the solid obtained was the diazoxide starting material and not a salt.

[0540] Salt preparation was carried out in non-aqueous media by dissolving diazoxide in anhydrous methanol in the presence of either sodium methoxide or potassium methoxide and stirring the mixture at 60°C for 15 minutes. The mixture was then cooled to room temperature while stirred. After approximately two hours, a solid was recovered, isolated by filtration, and dried in vacuo. Analysis by XRPD confirmed that the solid obtained was the diazoxide starting material and not a salt.

4.5.2. Preparation of Salts in Methanol or Ethanol According to the '573 Patent

[0541] The preparation of diazoxide salts in methanol and ethanol was attempted using 22 different counter-ions according to the methods described by the '573 Patent. For example, 20 mg of diazoxide was dissolved in 5 mL of ethanol and stirred and heated to ensure dissolution of the diazoxide. To the stirred solution was added approximately 1 molar equivalents of sodium methoxide. The solution was stirred for approximately 10-15 minutes at 60°C, and cooled to room temperature for approximately 2 hours. The resulting solid precipitate was concentrated under a nitrogen stream, and collected by filtration. The product was dried *in vacuo* and analyzed by XRPD. As shown in Figure-15, XRPD of the solids collected from the sodium methoxide experiment, (as well as the solids collected from the potassium methoxide experiment, which was run under similar conditions), revealed that the solid was the diazoxide starting material and that no sodium or potassium salt was prepared. In Figure 15, (a) is the XRPD pattern of free form diazoxide, (b) is the XRPD pattern of the product of potassium methoxide in methanol, and (c) is the XRPD pattern of the product of sodium methoxide in methanol.

[0542] Although not wishing to be bound by any theory, it is believed that a possible explanation for the failure to prepare diazoxide salts by the methods of the '573 Patent (i.e., in the presence of alcohols, e.g., methanol or ethanol), is that the alcohol may have an effect on the stability of the alkali salts of diazoxide. This was supported by UV spectroscopy analysis of alkali salts of diazoxide (sodium and potassium). Both the sodium and potassium salts of diazoxide were synthesized by reacting diazoxide with NaOH or KOH in MEK. Elemental analysis, NMR and XRPD confirmed that the salts were made.

[0543] Upon dissolution of the sodium or the potassium salt in acetonitrile, the UV spectrum shows a shift into the red region of the spectrum for the λ_{max} from approximately 268 nm for the free form of diazoxide to approximately 298 nm for both the potassium and sodium salt (see Figure 1). Similarly, these salts can be stabilized in aqueous solutions by elevating the pH to above 9.0. A similar shift in the absorption maximum at pH 9.0 is measured using UV spectroscopy. Subsequent adjustment of the pH from greater than 9.0 to less than 6.2 results in the hydrolysis of the salt as measured by the recovery of the UV absorption pattern of the diazoxide free base (see Figure 2). In contrast, when the sodium or the potassium salt is dissolved in methanol, the UV-Vis spectrum of the salt was identical to that of the diazoxide starting material, (see Figure 3).

[0544] These results demonstrate that using methanol as the solvent is incompatible with the synthesis of alkali salts of diazoxide. In addition, the results show that isolation of an alkali metal salt of diazoxide (or any other salt) in the presence of an alcohol, such as methanol, may not be possible.

4.5.3. Failure to Obtain Salts from Acidic Counter Ions

[0545] Salt formation with diazoxide was also attempted using acidic counter-ions, such as, for example, hydrochloric acid, maleic acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, fumaric acid, benzoic acid, undecylenic acid, salicylic acid and quinic acid. However, no salt formation was observed for any acid in any solvent.

[0546] For example, 100 mg of diazoxide was dissolved in 50 mL of acetone and heated to 35°C. To a stirred solution was added approximately 4.65 molar equivalents of HCl. The reaction mixture was allowed to cool to room temperature for approximately 3 hours, with no precipitation observed. Solvent was removed *in vacuo*. The resulting solid was analyzed by XRPD, and the observed XRPD pattern was consistent with the free form diazoxide starting material. In all cases, the attempted synthesis of diazoxide salts from acids was unsuccessful in all solvents attempted.

4.6. Preparation of Salts of Compounds of Formulae V-VIII

[0547] The chloride salt of 3-amino-4-methyl-1,2,4-benzothiadiazine-1,1-dioxide is prepared dissolving approximately 300 mg (1.4 mmol) in 45 mL acetonitrile. The mixture is heated to approximately 75°C and stirred for 30 min. To the stirred solution, approximately 1 molar equivalent of HCl is added dropwise, and stirred for approximately 30 min at 75°C. The mixture is cooled to room temperature and the solvent is removed under reduced pressure, affording the chloride salt as a solid.

[0548] The sodium salt of 3-amino-4-methyl-1,2,4-benzothiadiazine-1,1-dioxide is prepared dissolving approximately 300 mg (1.4 mmol) in 45 mL acetonitrile. The mixture is heated to approximately 75°C and stirred for 30 min. To the stirred solution, approximately 1 molar equivalent of NaOH is added dropwise, and stirred for approximately 30 min at 75°C. The mixture is cooled to room temperature and the solvent is removed under reduced pressure, affording the sodium salt as a solid.

5. Characterization of Prepared Diazoxide Salts

[0549] Synthesis of the desired salts was confirmed by X-Ray Powder Diffraction (XRPD), UV-Vis spectroscopy, and NMR. All spectra were compared with the spectra of the free form diazoxide (i.e., not a salt). Differential scanning calorimetry (DSC), thermal gravimetric analysis (TGA), FTIR (Fourier transform infrared spectroscopy), NMR, UV-vis spectroscopy and moisture sorption analysis were also performed.

5.1. Experimental Procedures

[0550] DSC analysis were conducted with a Mettler 822 DSC, by measuring the amount of energy released by a sample, as the sample was heated from 30°C to between 300-500°C at a rate of 10°C/min. Typical applications of DSC analysis include

determination of melting point temperature and the heat of melting; measurement of the glass transition temperature; curing and crystallization studies; and identification of phase transformations.

[0551] TGA measurements were conducted with a Mettler 851 SDTA/TGA, by measuring weight loss as a function of increasing temperature, as the samples were heated from 30°C to 230°C at a rate of 10°C/min. The TGA can be used to analyze desorption and decomposition behavior, characterize oxidation behavior, set burnout or conditioning parameters (temperature/ramp rate/time), and determine chemical composition.

[0552] XRPD samples were analyzed with a Shimadzu XRD-6000 system, using a Cu K α , 40 kV, 40 mA X-ray tube. The divergence and scatter slits were 1.00 deg, and the receiving slit was 0.30 mm. Samples were continuously scanned at a range of 3.0-45.0 deg, with a step size of 0.04 deg., at a scan rate of 2 deg/min.

[0553] Fourier Transform Infrared Spectroscopy was measured with a Thermo-Nicolet Avatar 370 with a Smart Endurance Attenuated Total Reflection (ATR) attachment. Compressed samples were analyzed, with corrections for background noise being made. Using the IR spectrum, chemical bonds and the molecular structure of organic compounds can be identified. Attenuated total reflectance (ATR) allows for the analysis of thin films, organic and inorganic, in areas as small as 10-15 microns.

[0554] Nuclear Magnetic Resonance (NMR) was performed with a 400 MHz Bruker Avance with a 4mm CP/MAS H-X probe. Acquisition of ^1H NMR spectra were performed by taking between 5-10 mg of the sample, dissolved in approximately 0.78 mL of DMSO- d_6 . Spectra were acquired with either 16 or 32 scans, using a pulse delay of 1.0 sec, with a 10 μsec (30°) pulse width.

[0555] UV spectroscopy was performed with a Perkin-Elmer Lambda 25 spectrometer. Samples were dissolved in acetonitrile, water and a buffer system having a pH between 5.6 and 10. Spectra were acquired between 340 and 190 nm, using a 1 cm path length with background correction.

[0556] Moisture Sorption Analysis was performed with a Hiden IGAsorp Moisture Sorption Instrument. Samples were first dried at 0% relative humidity at 25°C until an

equilibrium weight was reached, or for a maximum of 4 hours. Samples were then subjected to an isothermal (25°C) scan from 10 – 90% relative humidity in steps of 10%. The samples were allowed to equilibrate to an asymptotic weight at each point for a maximum of 4 hours. Following absorption, a desorption scan from 85% relative humidity (at 25°C) was run in steps of -10%, again allowing a maximum of 4 hours for the samples to equilibrate. The resulting samples after desorption were dried at 80°C for two hours and analyzed by XRPD.

3.2. Free Form Diazoxide Characterization

[0557] The free form of diazoxide was characterized by XRPD, differential scanning calorimetry (DSC), thermal gravimetric analysis (TGA), moisture sorption, ¹H NMR, FTIR and UV-vis spectroscopy to provide a baseline for comparison with the salts. Free form diazoxide is highly crystalline, as shown by the XRPD pattern. (See Figure 4(a)). The DSC shows a large endothermic event at 330°C, and TGA shows that the free form of diazoxide is anhydrous, where diazoxide shows no weight loss below 200°C, and a weight loss of only 0.2% below 230°C. Moisture absorption of the free form diazoxide shows the material to be non-hygroscopic. Absorption of water by diazoxide was tested at between 0 – 90% relative humidity (RH) at 25°C, showing absorption of approximately 0.04 wt% at 60% RH and 0.20 wt% at 90% RH. The molecule does not form a stable hydrate, as shown by the lack of hysteresis during desorption. Additionally, the XRPD pattern for the diazoxide before and after absorption of water indicate the same crystalline form.

[0558] UV-vis spectroscopy measurements taken of the free form diazoxide in neutral aqueous solution show a λ_{max} at approximately 268 nm. In acetonitrile, the λ_{max} was 264 nm, demonstrating a small solvatochromic shift. As shown in Figure 2, as pH increased the λ_{max} also increased, from approximately 265 nm to approximately 280 nm, due to a change in the electronics of the molecule.

[0559] Studies were also conducted to evaluate the likelihood for conversion and degradation under thermal stress. Samples were heated in a closed environment, protected from light, at 60°C for approximately 14 days. The diazoxide showed no conversion or degradation at 7 days or 14 days. Diazoxide samples were found to be consistent with the starting material with respect to XRPD and DSC.

[0560] Slurry studies to determine propensity of inter-conversion of the solid form were conducted on the free form diazoxide at room temperature, in the absence of light, using water, isopropyl alcohol, dichloromethane and toluene. Approximately 20 mg of the free form diazoxide was stirred for 14 days. Analysis by XRPD, DSC and HPLC were consistent with the starting material, indicating that the free form of diazoxide did not convert to alternate crystal forms.

5.3. Characterization of Sodium Diazoxide Salt

[0561] The XRPD pattern of the sodium salt of diazoxide was analyzed, showing the material to be crystalline. (See Figure 4(d)). The DSC analysis revealed a major exothermic event at 448°C. Small transitions below 400°C are likely due to sample imperfections. TGA analysis showed weight loss of 0.2% and 0.03% below 120°C, which may be the result of bound solvent. Moisture absorption performed from 0 – 90% relative humidity at 25°C showed the material to be hygroscopic as the sample deliquesced at 90% relative humidity. The sample absorbed 1.2 wt% of water at 60% RH, and 6.6 wt% water at 80% RH. Hysteresis was observed upon desorption at 65 and 55% RH, indicating possible hydrate formation. (Possible hydrate formation was noted, although the amount of water absorbed was less than 0.5 mole). ¹H NMR showed a chemical shift in the aromatic and methyl resonances of the sodium salt, as expected due to changes to the aromatic system. See Figure 5 showing NMR spectrum for the free form of diazoxide (a) and the sodium salt of diazoxide (c). FTIR showed expected changes for the sodium salt.

[0562] Elemental analysis of the salt indicated that the salt was formed in a ratio of approximately 1:1, with the percentage of sodium being slightly low (approximately 3.4%). This deficiency may be due to matrix effects, as NMR indicated the sample had a relatively high purity.

[0563] UV-vis measurements in neutral aqueous solution show a λ_{max} of approximately 271 nm. (See Figure 1). This value is slightly higher than free form diazoxide (265 nm). In acetonitrile, the λ_{max} of the sodium salt exhibits a solvatochromic shift to approximately 296 nm. (See Figure 3). An increase in the pH of the solution is expected to produce a bathochromic shift from approximately 265 nm to approximately 280 nm.

[0564] Solubility measurements performed at pH 2, 7, and 12 in 10 mM phosphate buffer at room temperature showed solubility of the sodium salt of diazoxide to be 13.0 mg/mL, 18.1 mg/mL and 48.6 mg/mL, respectively.

[0565] Form conversion and degradation under thermal stress were conducted as described for the free form diazoxide salt and showed the salt had little propensity to change form or degrade over a period of 14 days. Similarly, slurry studies were conducted as described for the free form diazoxide in n-heptane, dichloromethane and toluene showed no propensity of inter-conversion. See Figure 6, wherein (a) is the XRPD pattern for the sodium salt of diazoxide, (b) is the XRPD pattern for the sodium salt after the slurry study, and (c) the XRPD of the free form of diazoxide.

5.4. Characterization of Potassium Diazoxide Salt

[0566] The XRPD pattern for the potassium salt of diazoxide was analyzed, showing the material to be crystalline. (See Figure 4(b)). The DSC analysis revealed two major exothermic events at 128 and 354°C. (See Figure 7). Small endotherms are likely due to sample impurities or the presence of solvent. TGA analysis showed weight loss of 7.7% below 220°C, which may be the result of moisture sorption. (See Figure 8). Theoretical weight loss for a monohydrate of the diazoxide salt is 6.6%. Moisture absorption performed from 0 – 90% relative humidity at 25°C showed the material to be hygroscopic as the sample deliquesced at 90% relative humidity, showing 38.3 wt% water. The sample absorbed 5.4 wt% of water at 60% RH. Hysteresis was observed upon desorption, however the material was determined to be a hemihydrate from 0 – 30% RH and a monohydrate from 35 – 75% RH. XRPD following the desorption analysis indicated that the sample had changed to an alternate crystalline form. ¹H NMR showed a chemical shift in the aromatic and methyl resonances of the sodium salt, as expected due to changes to the aromatic system. (See Figure 5 showing spectrum of the free form diazoxide (a) and the Potassium salt of diazoxide (b)). FTIR showed expected changes for the potassium salt.

[0567] Elemental analysis of the potassium salt indicated that the salt was formed in a ratio of approximately 1:1, with the percentage of potassium being slightly low (approximately 1.6%). This deficiency may be due to matrix effects, as NMR indicated the sample had a relatively high purity.

[0568] UV-vis measurements of the potassium diazoxide salt in neutral aqueous solution show a λ_{max} of approximately 265 nm, which is equivalent to the diazoxide free form λ_{max} . (See Figure 1. In acetonitrile, the λ_{max} of the potassium salt exhibits a solvatochromic shift to approximately 296 nm. (See Figure 3). The potassium salt was used in a pH dependency study and showed that increasing the pH of the solution resulted in a bathochromic shift of the λ_{max} from approximately 265 nm to approximately 280 nm.

[0569] Solubility measurements performed at pH 2, 7, and 12 in 10 mM phosphate buffer at room temperature showed solubility of the sodium salt of diazoxide to be 9.9 mg/mL, 14.4 mg/mL and 43.0 mg/mL, respectively. The potassium salt displayed greater solubility than the free form diazoxide, and demonstrated similar solubility to the sodium diazoxide salt. The XRPD pattern of solids obtained after the solubility analysis indicated that the potassium salt had changed back to the free form diazoxide material.

[0570] Propensity for form conversion and degradation under thermal stress were conducted as described for the free form diazoxide salt. The XRPD pattern of the sample after 7 and 14 days showed unique peaks, as compared with the potassium salt starting material. Analysis by DSC after 14 days also showed unique peaks as well. Using a gradient area percent assay, HPLC did not show any significant degradation of the potassium salt.

[0571] Slurry studies were conducted as described for the free form diazoxide in n-heptane, dichloromethane and toluene showed no propensity of inter-conversion. XRPD analysis of samples after 7 and 14 days showed unique peaks similar to those observed with the thermal stress study. See Figure 9, wherein (a) is the XRPD pattern of the potassium salt of diazoxide, (b) is the XRPD pattern of the potassium salt of diazoxide after the slurry study in toluene, and (c) the XRPD pattern of the potassium salt after 14 days of the slurry study in toluene. Analysis by DSC after 14 days also showed unique peaks as compared with the starting material. HPLC using a gradient area percent assay did not show any significant degradation after the study.

5.5. Characterization of Choline Diazoxide Salt

[0572] The XRPD pattern of the choline salt of diazoxide was analyzed, showing the material to be crystalline. (See Figure 10 (b)). The DSC analysis revealed a major

exothermic events at 167°C. (See Figure 11). A smaller endothermic event was seen at 119°C and is likely due to sample impurities or the presence of residual solvent. TGA analysis showed weight loss of 0.8% between 100 and 140°C, which may be the result of residual solvents. (See Figure 12). Moisture absorption performed from 0 – 90% relative humidity at 25°C showed the material to be hygroscopic as the sample absorbed over 28% at 80% relative humidity, and deliquesced at 90% RH. Hysteresis was not observed upon desorption, indicating the choline salt does not form a stable hydrate. The XRPD pattern following the desorption analysis indicated that the sample had changed to an alternate crystalline form during the hydration and subsequent desorption. (See Figure 13 (c)) ¹H NMR was consistent with a 1:1 molar ratio of diazoxide to counter-ion, and had the expected differences in the chemical shift of the aromatic and methyl resonances, as expected due to the change in the local environment of the aromatic system due to the presence of the choline counter-ion. (See Figure 14 (b)). FTIR showed expected changes for the choline salt, similar to those seen with the sodium and potassium salts.

[0573] Elemental analysis of the choline salt indicated that the salt was formed in a ratio of approximately 1:1. This is consistent with the ¹H NMR.

[0574] UV-vis measurements of the choline diazoxide salt in neutral aqueous solution show a λ_{max} of approximately 268 nm, which is close to the λ_{max} for the diazoxide free form of 265 nm. In acetonitrile, the λ_{max} of the choline salt exhibits a solvatochromic shift to approximately 296 nm, which is consistent with the sodium and potassium diazoxide salts. The potassium salt was used in a pH dependency study and showed that increasing the pH of the solution resulted in a bathochromic shift of the λ_{max} from approximately 265 nm to approximately 280 nm.

[0575] Solubility measurements performed at pH 2, 7, and 12 in 10 mM phosphate buffer at room temperature showed solubility of the sodium salt of diazoxide to be 28.2 mg/mL, 41.5 mg/mL and greater than 293 mg/mL, respectively. The choline salt displayed greater solubility than the free form diazoxide after being allowed to equilibrate for 12 hours. The XRPD pattern of the solids obtained after the solubility analysis (at pH 2 and 7 only) indicated that the choline salt had changed back to the free form diazoxide material.

[0576] Propensity of the choline salt for form conversion and degradation under thermal stress were conducted as described for the free form diazoxide salt. XRPD analysis of the sample after 7 and 14 days showed an XRPD pattern consistent with the starting material, as well as the presence of additional unique peaks. (More unique peaks were present after 14 days than after 7 days). Analysis by DSC after 14 days did not show any significant difference, with a small endotherm at 117°C and a large endotherm at 168°C. (Initial DSC showed endotherms at 119°C and a large endotherm at 167°C). Using a gradient area percent assay, HPLC did not show any significant degradation of the choline salt.

[0577] Slurry studies were conducted as described for the free form diazoxide in n-heptane, dichloromethane and toluene showed no propensity of inter-conversion. XRPD analysis of samples after 7 and 14 days showed signal associated with the starting material, with additional unique peaks present. (See Figure 13 (c)). The XRPD pattern of slurry study samples from n-heptane showed signals associated with the starting material, as well as other additional unique signals. The XRPD pattern of slurry samples from dichloromethane and toluene were consistent with spectra obtained after the thermal studies and the moisture sorption analysis. Analysis by DSC after 14 days revealed a small endotherm at 109°C, and a major endotherm at 167°C. HPLC using a gradient area percent assay did not show any significant degradation after the study.

5.6. Characterization of Diazoxide Salt of Hexamethyl Hexamethylene Diammonium Hydroxide (HHDADH)

[0578] The XRPD pattern of the HHDADH salt of diazoxide was analyzed, showing the material to be a crystalline solid. (See Figure 10(c)). Integration of the ¹H NMR spectra was consistent with a 2:1 molar ratio of diazoxide to counter-ion (wherein the HHDADH counter-ion is divalent), and had the expected differences in the chemical shift of the aromatic and methyl resonances due to the change in the local environment of the aromatic system due to the presence of the choline counter-ion. (See Figure 14(c)). The λ_{max} of the HHDADH diazoxide salt in acetonitrile measured by UV-vis is 296 nm, which is consistent with the sodium, potassium and choline diazoxide salts.

[0579] A summary of the characterization of the free form diazoxide and the potassium, sodium, choline and hexamethyl hexamethylene diammonium hydroxide salts of diazoxide are presented in Table 14.

Table 14: Summary of Characterization and Solubility for Diazoxide and Salts

Test	Free form	Potassium diazoxide salt	Sodium diazoxide salt	Choline diazoxide salt	HHDADH
XRPD	crystalline	crystalline	crystalline	crystalline	crystalline
FTIR	consistent	consistent	consistent	consistent	N/A
DSC	single endotherm at 330°C	multiple endotherms below 300°C	Single exotherm above 400°C	two endotherms below 200°C	N/A
UV in aqueous acetonitrile	268 nm 264 nm	265 nm 296 nm	271 nm 296 nm	268 nm 296 nm	N/A 296 nm
TGA	<1%	7.7 (monohydrate 6.7%)	<1%	<1%	N/A
Moisture Sorption	non-hygroscopic – no hysteresis	deliquescent – hemi/monohydrate	deliquescent – potential hemihydrate	deliquescent – no hysteresis	N/A
Solubility					N/A
pH 2.0	0.4 mg/mL	9.9 mg/mL	13.0 mg/mL	28.2 mg/mL	
pH 7.0	0.04 mg/mL	14.4 mg/mL	18.1 mg/mL	41.5 mg/mL	
pH 12.0	4.8 mg/mL	43.0 mg/mL	48.6 mg/mL	>293 mg/mL	

[0580] Additional solubility studies were conducted for diazoxide free base form and diazoxide choline salt, as reported in Table 15. Each determination was carried out in duplicate, slurring each sample in a 100 mM phosphate buffer solution, pH 7.00. Duplicate samples were then titrated to pH 7.0 and pH 8.8 using a 0.1 N phosphoric acid solution, followed by stirring for 18h at ambient temperature. After this time all samples were centrifuged, and the supernatant was diluted with mobile phase (MeCN/ Water.). Solubility was then obtained using a calibration curve for diazoxide by HPLC analysis. In Table 15, the term "Diazoxide, free from" refers to the free base of diazoxide; the term "Diazoxide, choline salt" is the choline salt of diazoxide as described herein; the term "Diazoxide, choline salt, milled" refers to the choline salt of diazoxide which has been milled by methods described herein. Table 15 shows that in 100mM phosphate buffer, pH 7, and with titration with 0.1N phosphoric acid to a pH of about 6.8 to 8.8, solubility is notably suppressed compared to solubility at pH 10-11. Furthermore, the diazoxide

choline salts were found to have increased solubility when compared to the parent free base.

Table 15: Summary of Solubility for Diazoxide and Diazoxide Choline Salt.

Sample	Sample amount (mg)	Phosphate buffer amount (mL)	1N H_3PO_4 (mL)	pH after titration with 0.1N H_3PO_4	pH after 18h slurry	Solubility [mg/mL]
Diazoxide, free form	49.23	1	0.01	6.87	6.90	0.07
Diazoxide free form	47.64	1	---	7.30	7.31	0.07
Diazoxide, choline salt	44.34	1	0.15	7.22	7.45	0.12
Diazoxide, choline salt	48.26	1	0.11	8.81	8.64	0.18
Diazoxide, choline salt, milled	45.35	1	0.14	7.32	7.36	0.12
Diazoxide, choline salt, milled	53.15	1	0.16	8.50	8.52	0.20
Diazoxide, choline salt	50.0	1	---	---	10.57	42.11
Diazoxide, choline salt, milled	50.0	3	---	---	10.52	32.88

“---” Indicates no titration was conducted.

6. Polymorphic Forms of Diazoxide Salts

[0581] Polymorphic forms of the salts of diazoxide, characterization, and preparation thereof are described.

6.1. Polymorphic Forms of the Choline Salt of Diazoxide

6.1.1. Demonstration of Preparation of Polymorphic Form B of the Choline Salt of Diazoxide

[0582] A 1-L round bottom flask was charged with diazoxide (5 g), MEK (750 mL) and choline hydroxide (5.25 g of 50 wt % solution in water), and heated to 77°C. The mixture was allowed to cool to approximately ~30 °C, and filtered to remove insoluble brown residues. The filtrate was then concentrated under reduced pressure to afford yellow oil, which was dried *in vacuo* at 55°C and 30 in. Hg to afford approximately 7.8 g as a waxy solid. The solids were dried *in vacuo* at 55°C and 30 in. Hg to afford 7.13 g of the choline salt of diazoxide as a crystalline solid. Elemental analysis. Theoretical: C 46.77%, H 6.04% and N 12.59%. Measured: C 45.44%, H 5.98% and N 11.46%.

6.1.2. Characterization of the Polymorphic Form B of the Choline Salt of Diazoxide

[0583] The polymorphic Form B of the choline salt of diazoxide was analyzed by XRPD, DSC, and ¹H NMR. As shown in Figure 16, the two identified polymorphic forms of the choline salt of diazoxide show different XRPD patterns. See Figure 16, wherein (a) shows the polymorphic Form A of the choline salt of diazoxide and (c) shows the polymorphic Form B of the choline salt of diazoxide.

[0584] As shown in Figure 17, ¹H NMR of the polymorphic Form B of the choline salt of diazoxide shows no change from the polymorphic Form A.

[0585] Moisture absorption of the polymorphic Form B crystal structure of the choline salt of diazoxide performed from 0 – 80% relative humidity (RH) at 25°C showed the material to be hygroscopic as the sample absorbed over 14.5% at 70% relative humidity, and deliquesced at 80% RH. The XRPD pattern following the desorption analysis indicated that the sample remained in the Form B crystal structure during the hydration and subsequent desorption.

[0586] Solubility measurements performed at pH 2, 7, and 12 in 10 mM phosphate buffer at room temperature showed solubility of the Form B crystal structure of the choline salt of diazoxide to be 32.8 mg/mL, 80.1 mg/mL and 216 mg/mL, respectively.

The XRPD pattern of the solids obtained after the solubility analysis (at pH 2 and 7) indicated that Form B of the choline salt of diazoxide was still present.

[0587] Slurry experiments were performed on each form to determine the propensity for conversion and to search for possible new and/or unique forms. Upon slurrying Form B in CH₂Cl₂, n-heptane, and toluene, form conversion was not observed.

6.1.3. Demonstration of Preparation of the Polymorphic Form A of the Choline Salt of Diazoxide from the Polymorphic Form B of the Choline Salt of Diazoxide

[0588] Approximately 20 mg of the Form B polymorph of the choline salt of diazoxide was added to approximately 1 mL of acetone and heated to approximately 55°C. The mixture was filtered while hot and placed in a refrigerator (4°C) for 16 hours. No precipitate was observed. The solvent was evaporated down to dryness using a gentle stream of nitrogen. The resultant solids were also dried *in vacuo* at room temperature and 30 in. Hg. XRPD analysis showed the salt had converted from the Form B polymorph to the Form A polymorph.

6.1.4. Characterization of the Polymorphic Form A of the Choline Salt of Diazoxide

[0589] Form A is an anhydrous crystalline form of diazoxide choline, with an endothermic event at approximately 165 °C in the DSC (see Figure 20). The XRPD pattern for Form A is unique compared to that of Form B as shown in Figure 21. FTIR (ATR) spectroscopy additionally indicates differences between the two forms. ¹H NMR analysis affords a spectrum consistent with diazoxide and a 1:1 ratio of compound/counterion. NMR data also indicate that the magnetic environment of the diazoxide structure changes between free and polymorph forms, as evidenced by a movement in chemical shift of the aromatic and methyl proton resonances. In addition, the resonance due to the amine proton is not observed which suggests deprotonation in solution. Weight loss by TGA is less than 1% and may be due to residual solvent. The temperature of weight loss is above 100 °C which suggests that solvent may have been bound (i.e., solvate material). Moisture-sorption analysis conducted at 25 °C from 0 to 80% RH (adsorption) and 75 to 0% RH (desorption) shows Form A to be a hygroscopic solid, showing 2.4 wt % water at 60% RH. The sample was found to have deliquesced

above 75% RH. In comparison, Form B is also hygroscopic and showed 7.4 wt % water at 60% RH and deliquesced at 80% RH. XRPD analysis following the moisture-sorption experiment affords a pattern consistent with Form A. Solubility studies conducted on both forms at pH 2, 7, and 12 in phosphate buffer showed differences, with Form A showing 28, 41, and >293 mg/mL respectively. Solubility concentrations were determined using area-percent calculations with HPLC calibration curves. Slurry experiments were performed on each form to determine propensity for conversion and to see if a unique form could be generated. Upon slurrying Form A in CH₂Cl₂, *n*-heptane, and toluene, conversion to Form B was observed after seven days. These results suggest that Form A is less thermodynamically stable under these conditions than Form B according to Ostwald's Rule of Stages.

6.1.5. Screening for Polymorphic Forms of Diazoxide Choline Salt.

[0590] A polymorph screening study of diazoxide choline salt was conducted with a series of crystallization and slurry conditions. As described herein, interconversion of diazoxide choline salt forms A and B was observed during this investigation. Each polymorphic form of diazoxide choline salt resulting from this study was characterized using the techniques and procedures described herein. A summary of characterization tests is listed in Table 16.

Table 16. Characterization of Forms A and B of Diazoxide Choline Salt In Screening Study

Experiment	Form A	Form B
XRPD*	Crystalline	Crystalline
DSC	Endotherm $\approx$ 65 °C	Endotherm $\approx$ 60 °C
TGA	<1%	$\approx$ 4%
FTIR (ATR)**	Consistent w/ structure	Consistent w/ structure
¹ H NMR	Consistent w/ 1:1 ratio	Consistent w/ 1:1 ratio
Moisture Sorption	Hygroscopic – deliquesced at 90% RH	Hygroscopic – deliquesced at 80% RH
Solubility	pH 2: 28 mg/mL pH 7: 41 mg/mL pH 12: >293 mg/mL	pH 2: 33 mg/mL pH 7: 80 mg/mL pH 12: 216 mg/mL

Thermal	Started to convert to Form B after 14 days at 60 °C	Stable after 14 days at 60 °C
Slurries	Converted to Form B after 7 days in <i>n</i> -heptane, toluene, and CH ₂ Cl ₂	Stable after 14 days in CH ₂ Cl ₂ , THF, and <i>t</i> -AmOH

*Major peaks (2-θ):

Form A (9.8, 10.5, 14.9, 17.8, 17.9, 18.5, 19.5, 22.1, 22.6, 26.2, 29.6, 31.2);

Form B (8.9, 10.3, 12.0, 18.3, 20.6, 24.1, 24.5, 26.3, 27.1, 28.9).

** Unique FTIR (ATR) absorbances (cm⁻¹):

Form A (2926, 2654, 1592, 1449, 1248);

Form B (3256, 2174, 2890, 1605, 1463, 1235).

6.1.5.1. Solubility Screen in organic solvents.

[0591] Diazoxide choline, prepared in MEK using choline hydroxide as 50 wt % solution in water (see above) displayed some solubility in the following solvents: acetonitrile, acetone, ethanol, IPA, MEK, DMF, and methanol. These solvents were chosen due to differences in functionality, polarity, and boiling points and their ability to dissolve diazoxide. Other solvents which showed poor ability to dissolve salts were used as antisolvents and in slurry experiments where some solubility was observed: dioxane, MTBE, EtOAc, IPAc, THF, water, cyclohexane, heptane, CH₂Cl₂, and toluene.

[0592] Solvents for crystallizations during screening were chosen based on the solubility screen summarized in Table 17. Crystallizations of diazoxide choline from all conditions afforded a total of two forms, A and B. Forms A and B were found to be anhydrous polymorphs of diazoxide choline. Form B was observed to be generated from most solvents used. It was difficult to isolate pure Form A on large scales (>50 mg) as conditions observed to produce Form A on a smaller scale (approximately 50 mg or less) were found to result in Form B or mixtures of both forms on larger scales. Based on room-temperature slurry experiments, anhydrous Form B was found to be the most thermodynamically stable form in this study. Form A readily converted to Form B in all slurry solvents utilized.

Table 17. Solubility Screen for Diazoxide Choline Salt

Solvent	Compd (mg)	Solvent (mL)	Conc. (mg/mL)	Temp. (°C)	Soluble
MeCN	1.7	0.25	>6.80	rt	Yes
Dioxane	1.4	5.00	0.28	55	No
Acetone	1.9	0.25	7.60	55	Yes
MTBE	2.4	5.00	0.48	55	No
EtOH	1.5	0.25	>6.00	rt	Yes
EtOAc	1.2	5.00	0.24	55	No
IPAc	1.4	5.00	0.28	55	No
IPA	1.8	0.25	7.20	55	Yes
THF	1.1	5.00	0.22	55	No
MEK	1.8	1.00	1.80	55	Yes
DMF	1.2	0.25	>4.80	rt	Yes
Water	2.0	5.00	0.40	55	No
MeOH	1.9	0.25	7.60	55	Yes
n-Hexane	2.0	5.00	0.40	55	No
Heptane	1.9	5.00	0.38	55	No
CH ₂ Cl ₂	1.3	5.00	0.26	55	Partially
Toluene	1.4	5.00	0.28	55	No

6.1.5.2. Single-Solvent Crystalizations

[0593] *Fast cooling procedure:* Diazoxide (approximately 20 mg) was weighed out into vials and enough solvent (starting with 0.25 mL) was added until the material completely dissolved at elevated temperature. After hot filtration the vials were placed in a refrigerator (4 °C) for 16 hours. After the cooling-process the samples were observed for precipitates which were isolated by filtration. Vials not demonstrating precipitates were evaporated down to dryness using a gentle stream of nitrogen. All solids were dried *in vacuo* at ambient temperature and 30 in. Hg.

[0594] *Slow cooling procedure:* Diazoxide (approximately 30 mg of choline salt) was weighed out into vials and enough solvent was added until the material went into solution at elevated temperature. After hot filtration the vials were then slowly cooled to

room temperature at the rate of 20 °C/h and stirred at room temperature for 1–2 hours. All solids were dried *in vacuo* at ambient temperature and 30 in. Hg.

[0595] Based on the initial solubility study, seven solvents were selected for the fast-cooling crystallization: acetonitrile, acetone, ethanol, IPA, MEK, DMF, and methanol. Table 18 shows a list of the solvents that were used and the amount of solvent needed to dissolve the material. After the cooling-process precipitates were noticed in samples # 2, 3, 5, and 6, the solids were isolated by filtration. The other samples (# 1, 4, and 7) were evaporated down to dryness using a gentle stream of nitrogen. The diazoxide choline salts were found to be consistent with Form A by XRPD analysis for all solids with the exception of sample #2 (consistent with the freeform) and sample #5 (consistent with Form B with preferred orientation observed).

Table 18. Single-Solvent Crystallization of Diazoxide Choline Salt Using Fast-Cooling Procedure

Entry	Solvent	BP (°C)	Cmpd (mg)	Solvent Amt (mL)	Conc (mg/mL)	Temp. (°C)	Precipitate	Form
1	Acetone	56	21.0	1.00	21.00	55	No/Evap	A
2	MeOH	64	20.3	0.25	81.20	55	Yes	FF*
3	EtOH	78	21.3	0.25	85.20	62	Yes	A
4	MEK	80	19.6	1.25	15.68	75	No/Evap	A
5	MeCN	81	20.6	0.25	82.40	55	Yes	Unique
6	IPA	82	22.8	0.25	91.20	62	Yes	A
7	DMF	153	26.0	0.25	104.00	55	No/Evap	A

[0596] In accordance with the data obtained from fast-cooling experiments, four solvents which showed precipitation of solids were chosen for the slow-cooling experiments: MeOH, EtOH, MeCN, and IPA (Table 19). All obtained analyzable solids of the choline salt were found to be consistent with Form B by XRPD with the exception of Entry #1 which was consistent with diazoxide freeform and Entry #2 which was not analyzable. Mother liquor of Entry #2 was concentrated to dryness and the residual solids were analyzed by XRPD and found to be Form B material. As a result of obtaining freeform material from the single-solvent crystallizations in methanol, three more alcohols were tested for the single-solvent crystallizations using fast- and slow-cooling

procedures. Tables 20 and 21 provide a list of the solvents that were used and the amount of solvent needed to dissolve the material. XRPD patterns of the fast-cooling procedure showed freeform of diazoxide from isobutanol, Form B from isoamyl alcohol, and Form A from *tert*-amyl alcohol compared to the slow-cooling procedure, which afforded Form B material from all three solvents.

Table 19. Single-Solvent Crystallization of Diazoxide Choline Salt Using Slow-Cooling Procedure

Solvent	Boiling Point (°C)	Material Amount (mg)	Solvent Amount (mL)	Conc. (mg/mL)	Temp. (°C)	Precipitate	Form
MeOH	64	32.1	0.3	107.00	62	Yes	FF*
EtOH	78	33.3	0.3	111.00	75	Yes	NA**
MeCN	81	30.9	0.3	103.00	62	Yes	B
IPA	82	33.7	0.3	112.33	80	Yes	B

Table 20. Single-Solvent Crystallization of Diazoxide Choline Salt Using Fast-Cooling Procedure

Solvent	Boiling Point (°C)	Material Amount (mg)	Solvent Amount (mL)	Conc. (mg/mL)	Temp. (°C)	Precipitate	Form
<i>i</i> -BuOH	108	29.7	0.3	99.00	78	Yes	(sm)*
<i>i</i> -AmOH	130	29.6	0.3	98.67	82	Yes	B
<i>t</i> -AmOH	102	29.5	0.3	98.33	95	No/Evap	A

Table 21. Single-Solvent Crystallization of Diazoxide Choline Salt Using Slow-Cooling Procedure

Solvent	Boiling Point (°C)	Material Amount (mg)	Solvent Amount (mL)	Conc. (mg/mL)	Temp. (°C)	Precipitate	Form
<i>i</i> -BuOH	108	33.0	0.3	110.00	92	Yes	B
<i>i</i> -AmOH	130	28.2	0.3	94.00	92	Yes	B
<i>t</i> -AmOH	102	29.0	0.4	72.50	92	Yes	B

[0597] The results of the choline salt single-solvent fast- and slow-cooling crystallizations (see Tables 19 to 21) indicated that Form A was more likely to be isolated with fast-cooling profiles and Form B with slow-cooling profiles.

6.1.5.3. Binary Solvent Crystallizations

[0598] Binary-solvent crystallizations of the choline salt were performed using four primary solvents (MeOH, EtOH, IPA, and MeCN) and nine cosolvents (MTBE, EtOAc, IPAc, THF, *n*-hexane, heptane, toluene, CH₂Cl₂, and dioxane) with a fast-cooling profile (*supra*). XRPD patterns showed that Form B was obtained from mixtures of MeOH with MTBE, EtOAc, IPAc, toluene, and dioxane. As shown in Table 22, Form A was obtained from mixtures of MeOH with THF and with CH₂Cl₂ after evaporating the solvent to dryness. The mixtures of MeOH with cyclohexane and heptane provided the freeform of diazoxide. All solids obtained from fast-cooling procedures with EtOH, IPA, and MeCN as primary solvents provided Form B material.

Table 22. Binary-Solvent Crystallizations of Choline Salt of Diazoxide Using Fast-Cooling Procedure and MeOH as a Primary Solvent

Diazoxide Amt (mg)	MeOH* (mL)	Antisolvent	Amount (mL)	Precipitate	Form
27.8	0.3	MTBE	1.4	Yes	B
30.7	0.3	EtOAc	6.0	Yes	B
32.0	0.3	IPAc	6.0	Yes	B
31.9	0.3	THF	6.0	No/Evap to Dry	A
29.5	0.3	<i>n</i> -Hexane	2.0	Yes (small)	FF**
30.2	0.3	Heptane	2.0	Yes	FF**
29.3	0.3	Toluene	6.0	Yes	B
32.0	0.3	CH ₂ Cl ₂	6.0	No/Evap to Dry	A
28.8	0.3	Dioxane	6.0	Yes	B

* Solids were dissolved at 62 °C.

** Freeform of diazoxide.

[0599] Binary-solvent recrystallizations of the choline salt with the slow-cooling procedure were performed using two primary solvents (IPA and MeCN) and nine cosolvents (MTBE, EtOAc, IPAc, THF, *n*-hexane, heptane, toluene, CH₂Cl₂, and

dioxane). All solids obtained from a slow-cooling procedure with IPA and MeCN as primary solvents provided Form B material based on XRPD analysis. The results of binary-solvent crystallizations indicated that Form B was the most thermodynamically stable form of diazoxide choline.

6.1.5.4. Binary Solvent Crystallizations Using Water as a Cosolvent

[0600] In an attempt to investigate the formation of hydrates of the choline salt, experiments were performed using fast- and slow-cooling procedures and water as a cosolvent.

[0601] The fast cooling procedure (*supra*) was used with the exception of using different primary solvents which were miscible with water: acetone, acetonitrile, DMF, IPA, *i*-BuOH, *j*-AmOH, and *t*-AmOH. Water was utilized in these crystallizations as a cosolvent. All solids obtained from the fast-cooling procedure with water as the cosolvent provided diazoxide freeform material by XRPD analysis.

[0602] To compare the results obtained from the fast-cooling procedure a set of experiments was performed using a slow-cooling procedure and water as a cosolvent. All obtained solids were analyzed by XRPD and afforded patterns consistent with diazoxide freeform. Without wishing to be bound by theory, these results suggest that the conditions used for crystallization caused dissociation of the choline salt. A small amount of a second crop was obtained in each sample, but only two samples were analyzable by XRPD and indicated that the samples were freeform material. All mother liquors were evaporated to dryness and the residual solids were also analyzed by XRPD to afford patterns consistent with Form B of the choline salt.

6.1.5.5. Metastable Zone Width Estimation

[0603] *Form B:* To produce a robust process, an understanding of the solubility profiles of the various solid forms under consideration is required. From a practical standpoint, this involves the measurement of the metastable zone width (MSZW) of pure forms, whereby the saturation and supersaturation curves of the different forms are generated over a well defined concentration and temperature range. This knowledge can then be used to design a crystallization protocol that should ideally favor a selective crystal growth of the desired form.

[0604] Form B of diazoxide choline salt showed moderate solubility in a solvent mixture made of MeCN/MeOH/MtBE (10:1:12, volume ratios). The wide width of the metastable zone as shown in Table 23 gives many seeding options. During the MSZW measurement, aliquots from the crystallizing material were withdrawn and analyzed by XRPD to ensure that no form conversion occurred during the experiment. Indeed, the material remained unchanged during the test.

Table 23. Meta-Stable Zone Width For Form B Diazoxide Choline Salt in MeCN/MeOH/MtBE (10:1:12) (v/v).

Conc.] (mg/mL)	Temp. In (°C)	Temp. Out (°C)	Temp. Range (°C)
30.8	53.2	35.0	18.2
28.5	49.0	33.6	15.4
26.5	47.0	32.0	15.0
24.7	43.8	29.1	14.7
23.2	40.5	28.5	12.0
21.9	38.0	26.0	12.0

[0605] *Form A*: The metastable zone width for Form could not be estimated because this polymorphic form converted during the experiment to Form B.

6.1.5.6. Crystallization of Form A of Diazoxide Choline Salt

[0606] The choline salt of diazoxide (160.3 mg) was dissolved in 1 mL of IPA at 55 °C which was then passed through a Millipore 0.45 µm filter into a clean vial. This vial was placed in freezer at -20 °C overnight. Solids were not noticed and the flask was scratched with a micro-spatula. The vial was placed back in the freezer and nucleation was noticed after ten minutes. The solids were collected by vacuum filtration and washed with 1 mL of MtBE. The solids were dried *in vacuo* at 40 °C and 30 in. Hg to afford 70 mg (43.6% recovery) of Form A as determined by XRPD.

6.1.5.7. 500-mg Scale Crystallization of Form B of Diazoxide Choline Salt

[0607] The choline salt of diazoxide (524.3 mg) was dissolved in 3 mL of IPA at 78 °C and this solution was then cooled to 55 °C for the addition of MtBE. The MtBE (4 mL) was added until nucleation was observed. After nucleation the batch was allowed to cool to room temperature at a rate of 20 °C/h. The solids were collected by vacuum filtration and washed with 1 mL of MtBE. The solids were dried *in vacuo* at 40 °C and 30 in. of Hg to afford 426.7 mg (81.3% recovery) of Form B as determined by XRPD.

6.1.5.8. 3-g Scale Crystallization of Form B of Diazoxide Choline Salt

[0608] The choline salt of diazoxide (2.0015 g) was dissolved in 5.5 mL of IPA at 78 °C to afford a clear solution. This solution was passed through a Millipore Millex FH 0.45 µm filter. This solution was then cooled to 55 °C. MtBE was added in 1 mL portions, with a two minute interval between portions. Nucleation was noted after the second addition of MtBE. This suspension was allowed to cool to room temperature at a rate of 20 °C/h and stirred at this temperature for 16 hours. The solids were collected by vacuum filtration and washed with 1 mL of MtBE. The solids were dried *in vacuo* at 40 °C and 30 in. of Hg to afford 1.6091 g (80.4% recovery) of Form B as determined by XRPD.

6.1.5.9. Detection of Form Impurities

[0609] Mixtures of diazoxide choline Forms A and B were prepared by adding a minor amount of Form A to Form B. Samples were lightly ground by hands with a mortar and pestle for approximately one minute. Samples were then analyzed by XRPD analysis. XRPD analysis was found to be suitable for detecting 5% of Form A in Form B.

6.2. Polymorphic Forms of the Potassium Salt of Diazoxide

[0610] A summary of characterization tests for three common crystalline forms of diazoxide potassium salt are listed in Table 24. Solvents for crystallizations were chosen based on the solubility screen summarized below. Crystallizations of diazoxide potassium salt from all conditions provided herein afforded a total of seven unique crystalline forms, A through G. Forms C, D, and F were found to be the most common during the crystallization screen, and were therefore scaled up for further characterization.

Table 24. Results Summary of Characterization Tests for Salt of Diazoxide Potassium

Experiment	Form C	Form D	Form F
XRPD*	Crystalline	Crystalline	Crystalline
DSC	187, 360 °C	130, 191, 352 °C	191, 363 °C
TGA	8.4%**	4.5%**	13.1%**
FTIR (ATR)*	Consistent w/ structure	Consistent w/ structure	Consistent w/ structure
¹ H NMR	Consistent w/ structure**	Consistent w/ structure	Consistent w/ structure**
Moisture Sorption	Hygroscopic – deliquesced at 90% RH Hygroscopic – deliquesced at 90% RH	Hygroscopic – deliquesced at 90% RH	
Solubility	N/A	pH 2: 29 mg/mL pH 7: 33 mg/mL pH 12: 59 mg/mL	N/A
Thermal	Stable (7 days)	Stable (7 days)	Stable (7 days)
Slurries	Converted to Form D	Converted to Form D	Converted to Form D

*XRPD Major peaks (2-θ):

Form A (6.0, 8.1, 16.3, 17.7, 18.6, 19.1, 22.9, 23.3, 23.7, 24.7, 25.4, 26.1, 28.2, 29.6, 30.2);

Form B (8.5, 10.8, 16.9, 18.2, 21.6, 25.5, 26.1, 28.9);

Form C (5.7, 6.1, 17.9, 23.9, 25.1, 37.3);

Form D (5.7, 6.2, 8.1, 8.5, 8.8, 16.9, 18.6, 23.2, 24.5, 25.8, 26.1);

Form E (6.7, 7.1, 14.1, 21.2);

Form F (8.5, 9.0, 18.7, 20.6, 23.5, 27.5, 36.3);

Form G (5.2, 5.5, 13.1, 16.5, 19.3, 22.8, 24.8, 26.4, 28.7, 34.1);

*Unique FTIR (ATR) absorbances (cm⁻¹):

Form A (1503, 1374, 1339, 1207, 1131, 1056, 771);

Form B (1509, 1464, 1378, 1347);

Form C (1706, 1208, 1146, 746);

Form D (1595, 1258, 1219, 890);

Form E (1550, 1508, 1268, 1101, 1006).

Form F (1643, 1595, 1234, 1145, 810).

Form G (1675, 1591, 1504, 1458, 1432, 1266, 999, 958, 905, 872).

** Data indicates a half-molar equivalent of acetone, water, or dioxane for Forms C, D, and F respectively.

[0611] Diazoxide potassium Forms C, D, and F were observed to be an acetone solvate, a hemihydrate, and a dioxane solvate of diazoxide potassium, respectively. Form C is an acetone solvate that was generated predominantly when acetone was used in the crystallization. Form D, a hemihydrate, was observed to be generated from most solvents used. Form F is a dioxane solvate generated when dioxane was used as an antisolvent. Forms A, B, E, and G were not commonly observed during the crystallization. Elemental analysis data indicated that the unique forms observed may be mixtures and/or have residual solvent(s) present.

[0612] Based on room-temperature slurry experiments presented, Form D was found to be the most thermodynamically stable form of those discovered in this study. Forms C and F readily converted to Form D in all slurry solvents utilized. Without wishing to be bound by theory, since nonaqueous solvents were used, the material may have converted to the hemihydrate, Form D, upon removal from the solvent.

6.2.1. Demonstration of Preparation of Polymorphic Form A of the Potassium Salt of Diazoxide

[0613] The polymorphic Form A of the potassium salt of diazoxide was prepared as described above.

6.2.2. Demonstration of Preparation of the Polymorphic Form B of the Potassium Salt of Diazoxide

[0614] Diazoxide (2.95 g) was combined with 450 mL of methyl ethyl ketone and heated to approximately 77°C to dissolve the diazoxide. To the solution was added approximately 13 mL of 1M potassium hydroxide at a rate of approximately 20 mL/min, stirred and allowed to cool to room temperature. The solution was stirred at room temperature for ~16 h. The solvent was removed under reduced pressure, and the residual solids were dried *in vacuo* at 57 °C and 30 in. Hg to afford 3.7 g of the potassium salt.

6.2.3. Characterization of the Polymorphic Form B of the Potassium Salt of Diazoxide

[0615] The Form B polymorph of the potassium salt of diazoxide was analyzed by XRPD, and ^1H NMR. Figure 18 shows the XRPD pattern of (a) the Form A polymorph of the potassium salt of diazoxide and (b) the Form B polymorph of the potassium salt of diazoxide.

[0616] The ^1H NMR of the Form B polymorph of the potassium salt of diazoxide shows no change from the Form A polymorph.

6.2.4. Demonstration of Preparation of the Polymorphic Form A of the Potassium Salt of Diazoxide from Form B

[0617] Approximately 20 mg of the Form B polymorph of the diazoxide potassium salt was added to 2 mL of acetone and heated until the material completely dissolved at 55°C. The solution was hot filtered and placed in a refrigerator (4°C) for 16 hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 18(a).

6.2.5. Preparation of the Polymorphic Form C of the Potassium Salt of Diazoxide from Form B

[0618] Approximately 20 mg of the Form B polymorph of the diazoxide potassium salt was added to 6 mL of ethyl acetate and heated until the material completely dissolved at 75°C. The solution was hot filtered and placed in a refrigerator (4°C) for 16 hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 18(c).

6.2.6. Preparation of the Polymorphic Form D of the Potassium Salt of Diazoxide from Form B

[0619] Approximately 20 mg of the Form B polymorph of the diazoxide potassium salt was added to 0.3 mL of isopropyl alcohol and heated until the material completely dissolved at 52°C. The solution was hot filtered and placed in a refrigerator (4°C) for 16

hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 19(a).

6.2.7. Preparation of the Polymorphic Form E of the Potassium Salt of Diazoxide from Form B

[0620] Approximately 20 mg of the Form B polymorph of the diazoxide potassium salt was added to 2.5 mL of tert-amyl alcohol and heated until the material completely dissolved at 95°C. The solution was hot filtered and placed in a refrigerator (4°C) for 16 hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 19(b).

6.2.8. Preparation of the Polymorphic Form F of the Potassium Salt of Diazoxide from Form B

[0621] Approximately 20 mg of the Form B polymorph of the diazoxide potassium salt was added to 0.6 mL of acetonitrile and heated until the material completely dissolved at 80°C. The solution was hot filtered, 6 mL of dioxane was added, and the solution was placed in a refrigerator (4°C) for 16 hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 19(c).

6.2.9. Preparation of the Polymorphic Form G of the Potassium Salt of Diazoxide from Form B

[0622] Approximately 22.3 mg of the Form B polymorph of the diazoxide potassium salt was added to 0.5 mL of isoamyl alcohol and heated until the material completely dissolved at 73°C. The solution was hot filtered, 6 mL of isopropyl acetate was added, and the solution was placed in a refrigerator (4°C) for 16 hours. No precipitate was formed. The solvent was evaporated down to dryness using a gentle stream of nitrogen and the resultant solids were dried *in vacuo* at room temperature and 30 in. Hg. The solid was analyzed by XRPD to determine the physical form. See Figure 19(d).

[0623] As shown in Tables 25 and 26, both the solvent used for recrystallization and the rate of cooling (i.e., fast cooling vs. slow cooling during the recrystallization) effect the crystal structure obtained once the product is isolated.

Table 25. Fast cooling of Potassium Salt of Diazoxide in Various Solvents

Solvent	Cmpd (mg)	Solvent (mL)	Recovery (mg)	Form
THF	20.8	6.5	14.2	B
EtOAc	21.1	6.0	10.6	C
MeCN	21.2	0.5	n/a	C + D
IPA	22.7	0.3	n/a	D
Water	20.6	1.5	n/a	Free form diazoxide
tAA	21.9	2.5	16.7	E
IAA	19.7	0.3	n/a	D
DMF	21.6	0.3	15.9	G

Table 26. Slow cooling of Potassium Salt of Diazoxide in Various Solvents

Solvent	Cmpd (mg)	Solvent (mL)	Recovery (mg)	Form
EtOAc	20.4	6.0	7.1	D
MeCN	22.8	0.5	13.8	C
IPA	21.5	0.3	13.2	C
Water	20.9	1.5	1.2	Free form diazoxide
tAA	21.5	2.5	15.4	E
IAA	20.6	0.3	6.6	D

6.2.10. Polymorphs Obtained by Recrystallization of the Potassium Salt of Diazoxide in Binary Solvents

[0624] As shown in Tables 27 and 28, recrystallization of the potassium salt of diazoxide from a variety of binary solvent systems also demonstrated conversion of the potassium salt to an alternate form. Use of acetonitrile as the primary solvent is shown in

Table 27 and use of acetone as the primary solvent is shown in Table 28. As shown in Table 27, recrystallization of the Form B polymorph of the potassium salt of diazoxide from acetonitrile using methyl tert-butyl ether, ethyl acetate, isopropyl acetate, tetrahydrofuran, c-hexane, heptane, toluene and dichloromethane as the secondary solvent all yielded the D Form polymorph of the potassium salt. Recrystallization from acetonitrile using dioxane as the secondary solvent yielded the F Form polymorph of the diazoxide salt of potassium.

Table 27. Recrystallization in Acetonitrile

Acetonitrile (mL)	Secondary solvent	Cmpd (mg)	Recovery (mg)	Form
0.6	MTBE	20.7	13.7	D
0.6	EtOAc	23.4	9.5	D
0.6	IPAc	20.0	13.3	D
0.6	THF	20.3	6.4	D
0.6	c-Hexane	20.4	9.6	D
0.6	Heptane	20.3	10.8	D
0.6	Toluene	23.6	16.1	D
0.6	Dichloromethane	21.3	12.7	D
0.6	Dioxane	20.7	12.6	F

[0625] As shown in Table 28, recrystallization of the Form B polymorph of the potassium salt of diazoxide from acetone using methyl tert-butyl ether, tetrahydrofuran, and c-hexane as the secondary solvent all yielded the Form A polymorph of the potassium salt of diazoxide. Recrystallization from acetone using ethyl acetate, heptane, toluene and dichloromethane as the secondary solvent all yielded the C Form polymorph of the potassium salt. Recrystallization from acetone using isopropyl acetate as the secondary solvent yielded the D Form polymorph of the diazoxide salt of potassium. Recrystallization from acetone using dioxane as the secondary solvent yielded the F Form polymorph of the diazoxide salt of potassium.

Table 28. Recrystallization in Acetone

Acetone (mL)	Other Solvent	Amt (mg)	Recovery (mg)	Form
2.0	MTBE	20.2	13.9	A
2.0	EtOAc	21.6	4.8	C
2.0	IPAc	20.6	11.6	D
2.0	THF	20.9	12.0	A
2.0	n-Hexane	21.3	12.3	A
2.0	Heptane	20.6	12.7	C
2.0	Toluene	20.4	13.3	C
2.0	Dichloromethane	21.1	13.0	C
2.0	Dioxane	20.4	12.5	F

6.2.11. Screening for Polymorphic Forms of Diazoxide Potassium Salt.

[0626] A polymorphic screening study of diazoxide potassium salt was conducted with a series of crystallization conditions described below.

6.2.11.1. Solubility Screening for Polymorphic Forms of Diazoxide Potassium Salt.

[0627] Diazoxide potassium, prepared in MEK using 1 M potassium hydroxide solution in water, displayed some solubility in the following ten solvents: acetone, THF, EtOAc, MEK, MeCN, IPA, water, *t*-AmOH, *i*-AmOH, and DMF. These solvents were chosen due to differences in functionality, polarity, and boiling points and their ability to dissolve diazoxide. Solvents affording poor to fair solubility were used as antisolvents in binary/ternary crystallizations as well as slurry studies. Table 29 summarizes the results of the solubility screen.

Table 29. Solubility of Diazoxide Potassium Salt in Various Solvent

Solvent	Cmpd Amt (mg)	Solvent (mL)	Cone. (mg/mL)	Temp. (°C)	Soluble
MeCN	1.7	2.00	0.85	55	Yes
Dioxane	1.4	5.00	0.28	55	No
Acetone	1.6	4.00	0.40	55	Yes

Solvent	Cmpd Amt (mg)	Solvent (mL)	Conc. (mg/mL)	Temp. (°C)	Soluble
MTBE	1.8	5.00	0.36	55	No
EtOH	2.2	0.75	2.93	55	Yes
EtOAc	1.8	5.00	0.36	55	No
IPAc	1.7	5.00	0.34	55	No
IPA	2.1	1.00	2.10	55	Yes
THF	1.8	5.00	0.36	55	Partially
MEK	1.5	5.00	0.30	55	Partially
DMF	1.6	0.25	>6.40	rt	Yes
Water	1.5	5.00	0.30	55	No
MeOH	1.5	0.25	6.00	55	Yes
<i>n</i> -Hexane	1.5	5.00	0.30	55	No
Heptane	1.2	5.00	0.24	55	No
CH ₂ Cl ₂	1.3	5.00	0.26	55	No
Toluene	1.4	5.00	0.28	55	No

*6.2.11.2. Single-Solvent Screening for Polymorphic Forms of
Diazoxide Potassium Salt.*

[0628] Single-solvent crystallizations of potassium salt were performed using ten solvents: acetone, THF, EtOAc, MEK, MeCN, IPA, water, *t*-AmOH, *l*-AmOH, and DMF for the fast-cooling procedure and six solvents (EtOAc, MeCN, IPA, water, *t*-AmOH, and *l*-AmOH) for the slow-cooling procedures. The "fast" and "slow" cooling procedures were as described above. Four of the solvents were excluded from the slow-cooling experiments because they did not provide solids during fast-cooling experiments and needed to be evaporated to dryness. Tables 30 and 31 provide a list of the solvents that were used and the amount of solvent needed to dissolve the material. All solids were analyzed by XRPD to determine the physical form and six unique patterns (Forms A–E, G) were observed.

Table 30. Single-Solvent Crystallizations of Potassium Salt of Diazoxide Using Fast-Cooling

Solvent	BP (°C)	Cmpd Amt (mg)	Solvent Amt (mL)	Conc (mg/mL)	Temp (°C)	Precipitate	Form
Acetone	56	20.7	2.0	10.35	55	No/Evap	A
THF*	65	20.8	6.5	3.20	63	No/Evap	B
EtOAc	76	21.1	6.0	3.52	75	Yes	C
MEK	80	20.2	4.0	5.05	75	No/Evap	A
MeCN	81	21.2	0.5	42.40	80	Yes	C+D
IPA	82	22.7	0.3	75.67	62	Yes	D
Water	100	20.6	1.5	13.73	95	Yes	FF
<i>i</i> -AmOH	103	21.9	2.5	8.76	95	Yes	E
<i>i</i> -AmOH	130	19.7	0.3	65.67	73	Yes	D
DMF	153	21.6	0.3	72.00	RT	No/Evap	G

*Solids were not completely dissolved.

Table 31. Single-Solvent Crystallizations of Potassium Salt of Diazoxide Using Slow-

Solvent	BP (°C)	Cmpd Amt (mg)	Solvent Amt (mL)	Conc (mg/mL)	Temp (°C)	Precipitate	Form
EtOAc	76	20.4	6.0	3.40	75	Yes	D
MeCN	81	22.8	0.5	45.60	80	Yes	C
IPA	82	21.5	0.3	71.67	80	No/Evap	C
Water	100	20.9	1.5	13.93	95	Yes	FF
<i>i</i> -AmOH	103	21.5	2.5	8.60	95	Yes	E
<i>i</i> -AmOH	130	20.6	0.3	68.67	80	Yes	D

6.2.11.3. Binary-Solvent Screening for Polymorphic Forms of Diazoxide Potassium Salt.

[0629] Binary-solvent crystallizations of the potassium salt utilizing fast-cooling procedure were performed using MeCN, acetone, and isoamyl alcohol as primary solvents and the following nine cosolvents: MTBE, EtOAc, IPAc, THF, *n*-hexane, heptane, toluene, CH₂Cl₂, and dioxane. Table 32 is representative, employing acetonitrile as primary solvent. XRPD patterns from crystallizations using acetonitrile as a primary

solvent were consistent with Form D with only one exception being the solids obtained from the mixture of MeCN/dioxane afforded a unique pattern (Form F) material.

Table 32. Binary-Solvent Crystallizations of Potassium Salt of Diazoxide Using Fast-cooling Procedure and MeCN as a Primary Solvent

Cmpd Amt (mg)	MeCN* (mL)	Anti-solvent	Amt (mL)	Precipitate	Form
20.7	0.6	MTBE	1.0	Yes	D
23.4	0.6	EtOAc	6.0	Yes	D
20.0	0.6	IPAc	3.0	Yes	D
20.3	0.6	THF	6.0	No/Evap. To ppt	D
20.4	0.6	<i>n</i> -Hexane	2.0	Yes	D
20.3	0.6	Heptane	2.0	Yes	D
23.6	0.6	Toluene	1.0	Yes	D
21.3	0.6	CH ₂ Cl ₂	1.0	Yes	D
20.7	0.6	Dioxane	6.0	Yes	F

* Solids were dissolved at 80 °C.

[0630] XRPD patterns from binary crystallizations with acetone as primary solvent employing a fast cooling procedure afforded four forms A, C, D, and F. Mixtures of acetone with MTBE, THF, and cyclohexane provided Form A material; Form C was obtained from the mixtures of acetone with EtOAc, heptane, toluene, and CH₂Cl₂. The mixture of acetone with IPAc afforded Form D, and the mixture of acetone with dioxane afforded Form F solids.

[0631] XRPD patterns from binary crystallizations with isoamyl alcohol as the primary solvent employing a fast cooling procedure afforded five forms: C, D, E, F, and G. Form C was obtained from crystallizations using MTBE and EtOAc as cosolvents; Form D was obtained from mixtures of isoamyl alcohol with heptane, toluene, and CH₂Cl₂; Form E was crystallized out of *i*-AmOH/THF and *i*-AmOH/cyclohexane; Form G was obtained from *i*-AmOH/IPAc and Form F was obtained from *i*-AmOH/dioxane. Form D was the most common form observed from the crystallizations, and Form F was observed only when dioxane was used as an antisolvent.

[0632] Binary solvent recrystallizations of the potassium salt with the slow-cooling procedure were performed using three primary solvents (MeCN, acetone, and *i*-AmOH) and eight cosolvents (MTBE, EtOAc, IPAc, *c*-hexane, heptane, toluene, CH₂Cl₂, and dioxane). All solids were analyzed by XRPD to determine the physical form. Two patterns were observed to be Forms C and D respectively with additional peaks present. Other crystallizations provided Forms D, C, or F. Form D was obtained from the following solvent mixtures: MeCN/MTBE, MeCN/IPAc, MeCN/toluene, MeCN/CH₂Cl₂ and also from the mixtures of *i*-AmOH with MTBE, IPAc, cyclohexane, heptane, and toluene. Form C was obtained from MeCN/heptane, *i*-AmOH/EtOAc, and mixtures of acetone with MTBE, EtOAc, IPAc, cyclohexane, heptane, toluene, and CH₂Cl₂. Form F was crystallized from the mixtures of MeCN/dioxane and *i*-AmOH/dioxane. The solvent mixture of MeCN/EtOH provided amorphous material. Elemental analysis results indicate that the forms observed may not be pure and/or have bound or residual solvents present. Forms C, D, and F were found to be the most common forms of the potassium salt isolated and based on the results, these forms were chosen for scale-up and further characterization. Differences were found in the XRPD patterns and FTIR spectra of the scale-up lots which were attributed to differences in impurity profiles, crystallinity, and form purity.

6.2.11.4. Characterization of Polymorphic Form C of Diazoxide Potassium Salt.

[0633] Form C of diazoxide potassium salt is an acetone solvate with a 2:1 ratio of diazoxide/solvent. It is a crystalline form of diazoxide potassium, with endothermic events at 187 and 360 °C in the DSC. The XRPD pattern for Form C is unique compared to all other forms observed. FTIR (ATR) spectroscopy showed differences between forms. ¹H NMR spectra were found to be consistent with the structure of diazoxide with a half-molar equivalent of acetone present. NMR data also indicated that the magnetic environment of the diazoxide structure had changed evidenced by a movement in chemical shift of the aromatic and methyl proton resonances. In addition, the resonance due to the amine proton was not observed which suggested deprotonation in solution. Weight loss by TGA was 8.9%, consistent with a half-molar equivalent of acetone, and occurred near 180 °C consistent with the endotherm observed in the DSC experiment. Moisture-sorption analysis conducted at 25 °C from 0 to 90% RH (adsorption) and 85 to

0% RH (desorption) showed Form C to be a hygroscopic solid, showing approximately 47 wt % water at 90% RH which indicated the sample deliquesced. In comparison, Forms D and F (hemihydrate and dioxane solvate) showed approximately 26 and 22 wt % water at 90% RH respectively. XRPD analysis following the moisture-sorption experiment afforded a pattern consistent with Form D. Slurry experiments were performed on 50/50 mixtures of Forms C, D, and F to determine propensity for conversion and to see if a unique form could be generated. Upon slurrying Form C mixtures in ethyl acetate, acetonitrile, and isopropanol conversion to Form D was observed in all solvents. Without wishing to be bound by theory, these results as well as conversion of Form F to Form D in these conditions suggest that Form D is more thermodynamically stable than Form C and F according to Ostwald's Rule of Stages. A thermal-stability experiment on Form C at 60 °C found the form to be stable. Conversion to another form was not observed.

6.2.11.5. Characterization of Polymorphic Form D of Diazoxide Potassium Salt.

[0634] Form D of diazoxide potassium salt is a hemihydrate. It is a crystalline form of diazoxide potassium, with endothermic events at 130, 191, and 352 °C in the DSC. FTIR (ATR) spectroscopy showed differences between forms. ¹H NMR spectra were found to be consistent with the structure of diazoxide. NMR data also indicated that the magnetic environment of the diazoxide structure had changed evidenced by a movement in chemical shift of the aromatic and methyl proton resonances. In addition, the resonance due to the amine proton was not observed which suggested deprotonation in solution. Weight loss by TGA was 4.5%, consistent with a half-molar equivalent of water, and occurred near 110 °C consistent with the endotherm observed in the DSC experiment. Moisture-sorption analysis conducted at 25 °C from 0 to 90% RH (adsorption) and 85 to 0% RH (desorption) showed Form D to be a hygroscopic, showing approximately 26 wt % water at 90% RH. In comparison, Forms C and F (acetone and dioxane solvates) showed approximately 47 and 22 wt % water at 90% RH, respectively. XRPD analysis following the moisture-sorption experiment afforded a pattern consistent with Form D. Solubility studies were conducted at pH 2, 7, and 12 for Form D and showed 29, 33, and 59 mg/mL respectively. Solubility concentrations were determined using area-percent calculations with an HPLC calibration curve. Slurry experiments were performed on

50/50 mixtures of Forms C, D, and F to determine their propensity for conversion and to see if a unique form could be generated. Upon slurrying mixtures of Form D with Form C or Form F in ethyl acetate, acetonitrile, and isopropanol conversion to Form D was observed in all solvents.

6.2.11.6. Characterization of Polymorphic Form F of Diazoxide Potassium Salt.

[0635] Form F of diazoxide potassium salt is a dioxane solvate with a 2:1 ratio of diazoxide/solvent. It is a crystalline form of diazoxide potassium, with endothermic events at 191 and 363 °C in the DSC. The XRPD pattern for Form F is unique compared to all other forms observed. FTIR (ATR) spectroscopy showed differences between forms. ¹H NMR spectra were found to be consistent with the structure of diazoxide with a half-molar equivalent of dioxane present. NMR data also indicated that the magnetic environment of the diazoxide structure had changed as evidenced by a movement in chemical shift of the aromatic and methyl proton resonances. In addition, the resonance due to the amine proton was not observed which suggested deprotonation in solution. Weight loss by TGA was 13.1%, consistent with a half-molar equivalent of dioxane, and occurs near 180 °C consistent with the endotherm observed in the DSC experiment. Moisture-sorption analysis conducted at 25 °C from 0 to 90% RH (adsorption) and 85 to 0% RH (desorption) showed Form F to be a hygroscopic solid, showing approximately 22 wt % water at 90% RH. In comparison, Forms C and D (acetone solvate and hemihydrate) showed approximately 47 and 26 wt % water at 90% RH, respectively. XRPD analysis following the moisture-sorption experiment afforded a pattern consistent with Form D. Slurry experiments were performed on 50/50 mixtures of Forms C, D, and F to determine their propensity for conversion and to see if a unique form could be generated. Upon slurrying mixtures of Form F with Form C and Form D in ethyl acetate, acetonitrile, and isopropanol conversion to Form D was observed in all solvents.

B. *In vivo* Obesity Testing

1. Obesity animal model

[0636] Formulations of salts of any of the compounds of Formulae I–IV prepared as described herein can be tested for efficacy in an animal model of obesity as described by Surwit et al. (*Endocrinology* 141:3630-3637 (2000)). Briefly, 4-week-old B6 male mice

are housed 5/cage in a temperature-controlled (22°C) room with a 12-h light, 12-h dark cycle. The high fat (HF) and low fat (LF) experimental diets contain 58% and 11% of calories from fat, respectively. A group of mice are fed the HF diet for the first 4 weeks of the study; the remaining 15 mice are fed the LF diet. The mice assigned to the LF diet are maintained on this diet throughout the study as a reference group of lean control mice. At week 4, all HF-fed mice are reassigned to 2 groups of mice. The first group remains on the HF diet throughout the study as the obese control group. The remaining 3 groups of mice are fed the HF diet and administered the controlled release formulation of salts of any of the compounds of Formulae I – IV at about 150 mg of active per kg per day as a single dose administered by oral gavage. Animals are weighed weekly, and food consumption is measured per cage twice weekly until the diets are changed at week 4, whereupon body weight and food intake are determined daily. The feed efficiency (grams of body weight gained per Cal consumed) is calculated on a per cage basis. Samples for analysis of insulin, glucose, and leptin are collected on day 24 (4 days before the diets are changed), on day 32 (4 days after the change), and biweekly thereafter. In all cases food is removed 8 h before samples are collected. Glucose is analyzed by the glucose oxidase method. Insulin and leptin concentrations are determined by double antibody RIA. The insulin assay is based on a rat standard, and the leptin assay uses a mouse standard. At the termination of the study, a postprandial plasma sample is collected and analyzed for triglyceride and nonesterified fatty acid concentrations. After 4 weeks of drug treatment, a subset of 10 animals from each group is killed. The epididymal white adipose tissue (EWAT), retroperitoneal (RP) fat, interscapular brown adipose tissue (IBAT) fat pads, and gastrocnemius muscle are removed, trimmed, and weighed. The percent body fat is estimated from the weight of the epididymal fat pad. A subset of five animals from each group is injected i.p. with 0.5 g/kg glucose. At 30 min post injection, a plasma sample is collected and analyzed for glucose content by the glucose oxidase method.

2. Treatment of obesity in humans

[0637] Formulations of salts of any of the compounds of Formulae I – IV prepared as described herein can be tested for efficacy in obese humans, as described by Alemzadeh (Alemzadeh et al., *J Clin Endocr Metab* 83:1911-1915 (1998)). Subjects consist of moderate-to-morbidly obese adults with a body mass index (BMI) greater than or equal to

30 kg/m². Each subject undergoes a complete physical examination at the initial evaluation, body weight being measured on a standard electronic scale and body composition being measured by DEXA.

[0638] Before initiation of the study, all subjects are placed on a hypocaloric diet for a lead-in period of 1 week. This is designed to exclude subjects who are unlikely to be compliant and to ensure stable body weight before treatment. Up to 50 subjects are tested at each dosage of drug. Daily dosage is set at 100, 200, and 300 mg/day. The daily dose is divided into 2 doses for administration. The dose is administered as either one, two or three 50 mg capsules or tablets at each time of administration. Subjects are dosed daily for up to 12 months. Subjects are reviewed weekly, weighed, and asked about any side effects or concurrent illnesses.

[0639] Twenty-four-hour dietary recall is obtained from each subject. The dietary recalls are analyzed using a standard computer software program. All subjects are placed on a hypocaloric diet and encouraged to participate in regular exercise.

[0640] Before commencing, and after completion of the study, the following laboratory tests are obtained: blood pressure, fasting plasma glucose, insulin, cholesterol, triglycerides, free fatty acids (FFA), and glycohemoglobin, and measures of rate of appearance and oxidation of plasma derived fatty acids. Additionally, routine chemistry profiles and fasting plasma glucose are obtained weekly to identify those subjects with evidence of glucose intolerance and/or electrolyte abnormalities. Glucose is analyzed in plasma, by the glucose oxidase method.

[0641] Insulin concentration is determined by RIA using a double-antibody kit. Cholesterol and triglycerides concentrations are measured by an enzymatic method. Plasma FFA is determined by an enzymatic colorimetric method. SI was assessed by an iv glucose tolerance test (IVGTT) using the modified minimal model. After an overnight fast, a glucose bolus (300 mg/kg) was administered iv, followed (20 min later) by a bolus of insulin. Blood for determination of glucose and insulin is obtained from a contra lateral vein at -30, -15, 0, 2, 3, 4, 5, 6, 8, 10, 19, 22, 25, 30, 40, 50, 70, 100, 140, and 180 min. SI and glucose effectiveness (SG) are calculated using Bergman's modified minimal-model computer program before and after the completion of the study. Acute

insulin response to glucose is determined over the first 19 min of the IVGTT, and the glucose disappearance rate (Kg) is determined from 8–19 min of the IVGTT. Body composition is measured by bioelectrical impedance before and at the completion of the study. Resting energy expenditure (REE) is measured by indirect calorimetry after an overnight 12-h fast, with subjects lying supine for a period of 30 min. Urine is collected over the corresponding 24 h, for measurement of total nitrogen and determination of substrate use, before and after the study.

3. *Treatment of obesity in humans by coadministering diazoxide and phentermine*

[0642] Evaluation of a prolonged co-administration of solid oral dosage form of salts of any of the compounds of Formulae I – IV thereof in combination with phentermine can be conducted in humans with moderate-to-morbid obesity and a body mass index (BMI) greater than or equal to 30 kg/m². Each subject undergoes a complete physical examination at the initial evaluation, body weight being measured on a standard electronic scale and body composition by DEXA.

[0643] Before initiation of the study, all subjects are placed on a hypocaloric diet for a lead-in period of 1 week. This is designed to exclude subjects who are unlikely to be compliant and to ensure stable body weight before treatment. Up to 100 subjects are tested. Daily dosage of salts of any of the compounds of Formulae I – IV is set at 200 mg. The daily dose is divided into 2 doses for administration. The dose is administered as either a 100 mg capsule or a 100 mg tablet at each time of administration. Subjects are dosed daily for up to 12 months. Phentermine is administered as a single daily dose of 15 mg. Subjects are reviewed every two weeks, weighed, and asked about any side effects or concurrent illnesses.

[0644] All subjects are continued on a hypocaloric diet and encouraged to participate in regular exercise. Before commencing, and after completion of the study, laboratory tests as described in the example above are obtained.

4. *Prevention of diabetes in prediabetic humans*

[0645] The example describes use of salts of any of the compounds of Formulae I – IV in a prediabetic subject to prevent the occurrence of diabetes. Subjects included in the study all have elevated risk of developing diabetes as measured by one of two methods.

In a fasting glucose assay they have plasma glucose values between 100 and 125 mg/dl indicating impaired fasting glucose, or in an oral glucose tolerance test they have plasma glucose values between 140 and 199 mg/dl at 2 hours post-glucose load indicating they have impaired glucose tolerance. Treatment is initiated in any subject meeting either criteria. Treated subjects receive either 200 mg diazoxide per day as a 100 mg capsule or tablet twice per day or as two 100 mg capsules or tablets once per day. Placebo treated subjects receive either one placebo capsule or tablet twice per day or two placebo capsules or tablets once per day.

[0646] Treatment is continued for once year with OGTT or fasting glucose measured monthly.

5. A sustained release coformulation of diazoxide HCl and metformin HCl use to treat diabetic patients

[0647] A sustained release co-formulation of diazoxide HCl and metformin HCl is produced by forming a compressed tablet matrix that includes 750 mg of metformin HCl and 100 mg of diazoxide HCl. These active ingredients are blended with sodium carboxymethyl cellulose (about 5% (w/w)), hypromellose (about 25% (w/w)), and magnesium stearate (<2% (w/w)). The compressed tablet is further coated with a combination of ethylcellulose (80% (w/w)) and methyl cellulose (20% (w/w)) as a thin film to control rate of hydration and drug release.

[0648] Type II diabetic patients are treated with the oral dosage form by administration of two tablets once per day or one tablet every 12 hours. Treatment of the patient with the drug is continued until one of two therapeutic endpoints is reached, or for so long as the patient derives therapeutic benefit from administration. The two therapeutic endpoints that would serve as the basis for the decision to cease treatment include the patient reaching a Body Mass Index (BMI (kg/m²)) between 18 and 25 or the re-establishment of normal glucose tolerance in the absence of treatment. The patient is monitored periodically for (a) glucose tolerance using an oral glucose tolerance test, (b) glycemic control using a standard blood glucose assay, (c) weight gain or loss, (d) progression of diabetic complications, and (e) adverse effects associated with the use of these active ingredients.

6. *Prevention or treatment of weight gain in a patient treated with olanzapine*

[0649] Pharmacotherapy for schizophrenia is initiated for a patient meeting DSM III-R criteria for schizophrenia. The patient is administered 10 mg of olanzapine (Zyprexa, Lilly) once per day. Adjunctive therapy to the patient for schizophrenia includes 250 mg equivalent of valproic acid as divalproex sodium (Depakote, Abbott Labs). Weight gain, dyslipidemia and impaired glucose tolerance, and metabolic syndrome are high frequency adverse events in patients treated with this combination of anti-psychotics. Weight gain, dyslipidemia, impaired glucose tolerance or metabolic syndrome are treated by the co-administration of a therapeutically effective dose of a K_{ATP} channel opener. The patient is treated with administration of 200 mg/day of salts of any of the compounds of Formulae I – IV as a once daily tablet formulation. Administration of salts of any of the compounds of Formulae I – IV continues until the weight gain, dyslipidemia, impaired glucose tolerance or metabolic syndrome is corrected or until treatment of the patient with olanzapine is discontinued. Dyslipidemia is detected by measuring circulating concentrations of total, HDL, and LDL cholesterol, triglycerides and non-esterified fatty acids. Impaired glucose tolerance is detected through the use of oral or IV glucose tolerance tests. Metabolic syndrome is detected by measuring its key risk factors including central obesity, dyslipidemia, impaired glucose tolerance, and circulating concentrations of key proinflammatory cytokines.

7. *Comparison of Single Doses of Diazoxide Administered as Proglycem® Oral Suspension or as Diazoxide Choline Controlled-Release Tablets in Obese Subjects.*

7.1 *Experimental Design*

7.1.1 *Objective of Study.*

[0650] Clinical studies employing a randomized, open-label, parallel protocol comparing the safety, tolerability and bioavailability (i.e., pharmacokinetics) in obese subjects of Proglycem® (oral suspension) and diazoxide choline salt (controlled-release tablet) were conducted. The study evaluated the safety and tolerability of a single 200 mg does (approximately 2 mg/kg) of diazoxide. The study further compared the single dose pharmacokinetics of an oral suspension of the free base of diazoxide (Proglycem®) with a controlled-release tableted formulation of diazoxide choline salt under fasting conditions in obese subjects.

7.1.2. Rationale for Study.

[0651] Diazoxide choline was selected as an alternative molecule for the oral administration of diazoxide because of significantly greater aqueous solubility over diazoxide free base, rapid conversion to diazoxide base on exposure to an aqueous environment, and incompatibility with incorporation into sustained release formulations. The fate of diazoxide choline in an aqueous medium in vitro has been extensively characterized. Without wishing to be bound by theory, once solubilized from the controlled-release tablet formulation, prior to absorption, the salt hydrolyzes to the free base of diazoxide and choline hydroxide. This was extensively characterized using UV/Vis absorbance, and occurred at all physiological relevant pH values from pH 2.0 to pH 8.5. This hydrolysis occurred in deionized water and buffered aqueous solutions, and in polar solvents that contained trace amounts of water. Thus, diazoxide, as the free base, is the molecular form absorbed following oral administration of the choline salt of diazoxide to animals or humans. Serum and plasma assays used in TK and PK analysis measure diazoxide free base concentrations. Dissolution assays measure diazoxide free base. Differences in plasma concentration-time profiles of diazoxide will be dependent on the time course of release of diazoxide choline from the tablet matrix in the intestinal tract. Historically, both oral suspension and immediate release capsule formulations of diazoxide have been commercially available, marketed as Proglycem®. The oral suspension has been shown to be highly bioavailable and rapidly absorbed upon administration. Because of the difficulty in characterizing dissolution from the oral suspension, the in-vitro dissolution of Proglycem® capsules was compared under standardized conditions to that of diazoxide choline controlled-release tablets (50 mg and 200 mg).

7.1.3. Inclusion Criteria.

[0652] Inclusion criteria for the present study included age 18 to 65 years old inclusive and BMI between 30 and 45 kg/m², inclusive, and signed informed consent. Female participants were required to be either postmenopausal for at least 1 year, surgically sterile [bilateral tubal ligation, bilateral oophorectomy, or hysterectomy], or practicing a medically acceptable method of birth control. Inclusion criteria further included freedom from serious medical disorders involving the kidneys, digestive system, heart and blood vessels, lungs, liver, eyes, nerves, brain, skin, endocrine system, bones or

blood. Subjects, other than their obese condition, were generally healthy as documented by medical history, physical examination, vital sign, 12-lead ECG, and clinical laboratory assessments. Characteristics of the subjects randomized in the study are provided in Table 33.

Table 33. Characteristics of subjects randomized in the study

Parameter	Proglycem Oral Suspension (n=15)	Diazoxide Choline Controlled-Release Tablet (n=15)
	Mean±SD (range)	Mean±SD (range)
Age (yr)	32.5±12.1 (18-56)	27.9±12.6 (19-54)
BMI (kg/m ²)	32.9±4.3 (30.2-44.8)	33.8±3.2 (31.4-42.1)
Gender (male/female)	9/6	7/8

7.1.4. Exclusion Criteria.

[0653] Exclusion criteria included treatment with an investigational drug within 28 days prior to dosing, presence or history of a clinically significant disorder, clinical laboratory test values outside of the accepted reference range, reactive screen for HBV, HCV, or HIV, use of any medication affecting body weight, lipid or glucose metabolism within 2 months, use of any systemic prescription medication from 14 days prior to screening to dosing except hormonal contraceptives, use of any drug known to induce or inhibit hepatic drug metabolism from 28 days prior to screening until dosing, positive test for drug of abuse, current tobacco use, positive pregnancy screen, or pregnant or breastfeeding.

7.1.5. Randomization.

[0654] A total of 30 subjects were randomized in the present study. Fifteen subjects were randomized to each arm. For convenience, the 30 subjects were broken up into two cohorts. Cohort 1 checked into the clinic on October 12, 2006, was dosed on October 14, 2006, remained in the clinic for 72 hours following dose administration, returning at 96 and 120 hours after dose administration. Cohort 1 included 10 subjects randomized to receive Proglycem Oral Suspension and 5 subjects randomized to receive a Diazoxide Choline Controlled-Release Tablet. Cohort 2 checked into the clinic on October 14,

2006, were dosed on October 16, 2006, remained in the clinic for 72 hours following dose administration, returning at 96 and 120 hours after dose administration. Cohort 2 included 5 subjects randomized to receive Proglycem® Oral Suspension and 10 subjects randomized to receive a diazoxide choline controlled-release tablet. All subjects completed the study.

7.1.6. Dosing.

[0655] Subjects randomized to the Proglycem® Oral Suspension arm received a single 200 mg dose (4 mL) taken with 240 mL of room temperature water. Subjects randomized to the diazoxide choline controlled-release tablet arm received a single tablet containing 290 mg of diazoxide choline, which is equivalent to 200 mg of diazoxide. Doses were administered after an overnight fast, and fasting continued until approximately 4.25 hours after dose administration at which time a standardized meal was served.

7.1.7. Safety Monitoring.

[0656] Clinical laboratory tests including hematology, clinical serum chemistry and urinalysis were conducted at screening and at end of study. Hematology included measurements of hemoglobin, hematocrit, white blood cell count with differential, red blood cell count and platelet count. Clinical serum chemistry included evaluation of sodium, potassium, BUN, creatinine, total bilirubin, total protein, albumin, alkaline phosphatase, AST, ALT, glucose, total cholesterol, HDL-cholesterol, and triglycerides. Urinalysis included pH, specific gravity, protein, glucose, ketones, bilirubin, blood, nitrites, urobilinogen, leukocytes, and microscopic urine analysis if sample was dipstick positive. All adverse events were recorded. Vital signs were collected at screening, at one hour prior to dosing, and after dose administration at 1, 3, 6, 9, 12, 24, 48, 72 and 120 hours. Continuous two-lead cardiac monitoring (telemetry) was performed on subjects from 24 hours prior to dose administration (to establish baseline data) until 24 hours after dose administration.

7.1.8. Exploratory Endpoints

[0657] Three exploratory endpoints were evaluated in the study. Blood glucose, insulin and non-esterified fatty acids (NEFA) were measured prior to dosing and at 1, 3, 6, 9, 12, 24, and 48 hours post dosing. Blood glucose and insulin levels were also

assessed at end of study. Data collected at times 1, 3, 24, and 48 hours after dose administration were obtained under fasting conditions. The remaining data points were post-prandial measures. Samples for pharmacokinetic analysis to evaluate bioavailability were collected prior to dosing and after dose administration at 1, 2, 4, 6, 8, 12, 16, 20, 24, 32, 40, 48, 72, 96, and 120 hours.

7.2. Results

7.2.1. Adverse Events.

[0658] The adverse events observed with each of the two formulations are summarized in Tables 34 and 35. With the single exception, a moderate headache requiring treatment with acetaminophen in a subject treated with Proglycem® Oral Suspension, all adverse events were mild. Both formulations appeared to be well tolerated in the study. One subject administered with diazoxide choline controlled-release tablet experienced mild loss of appetite. Because most obese and obese diabetic animal model studies of diazoxide show some reduction in feed consumption, loss of appetite was, therefore, considered part of the pharmacodynamic response to the drug in obese patients, rather than an adverse event.

Table 34. Adverse events (AE) in subjects administered Proglycem® Oral Suspension (n=15)

Adverse Event	Hours Post Dosing		Corrective Treatment	Outcome
Gastrointestinal				
Nausea ^a	2.75	Mild	None	Resolved
Neurological				
Headache	20.0	Moderate	Therapy required acetaminophen	Resolved
Headache ^b	6.25	Mild	None	Resolved
Dizziness ^c	1.0	Mild	Assisted to supine position	Resolved
Dizziness ^c	3.25	Mild	Assisted to supine position	Resolved
Dizziness ^d	1.0	Mild	Assisted to supine position	Resolved

Dizziness ^d	4.25	Mild	Assisted to supine position	Resolved
Miscellaneous				
Chills ^b	6.25	Mild	None	Resolved
Dermatitis from ECG patches ^a	4.75	Mild	None	Resolved

Number of subjects with at least one AE = 5, AEs followed by the same superscript letter are the same subject

Table 35. Adverse events (AE) in subjects administered Diazoxide Choline Controlled-Release Tablet (n=15)

Adverse Event	Hours Post Dosing		Corrective Treatment	Outcome
Gastrointestinal				
Nausea ^b	1.5	Mild	None	Resolved
Dry Heaves ^c	1.6	Mild	None	Resolved
Neurological				
Lightheaded	12.0	Mild	Assisted to supine position	Resolved
Dizziness ^b	20.0	Mild	None	Resolved
Miscellaneous				
Chills ^f	2.0	Mild	None	Resolved
Back Pain ^f	2.0	Mild	None	Resolved
Warm ^g	20.0	Mild	None	Resolved

Number of subjects with at least one AE = 4, AEs followed by the same superscript letter are the same subject

7.2.2. Clinical Chemistry.

[0659] A summary of fasting glucose, fasting insulin, NEFA, and serum sodium (Na), potassium (K) and creatinine level is provided in Table 36. Neither treatment resulted in

significant changes in serum Na, K or creatinine from baseline to end of study. Treatment with diazoxide primarily impacts glucose-stimulated insulin secretion rather than basal insulin secretion. Fasting glucose levels increased slightly in the first 3 hours following dose administration. This increase was more pronounced in the Proglycem® Oral Suspension arm as compared to the diazoxide choline controlled-release tablet arm. These results are consistent with the differences in measured rates of dissolution from these formulations (see Table 5 herein) and the PK levels (see Figures 24 and 25 herein). In this study, there is no evidence for a reduction in fasting insulin after the administration of the single dose of diazoxide. There is also no evidence for a significant increase in fasting glucose as measured at 24 or 48 hours after dose administration. The most sensitive pharmacodynamic response measure is NEFA. Diazoxide treatment normally results in a short-term increase in NEFA, which returns to levels at or below baseline in about 6 hours. Both formulations showed this transient increase in NEFA. Treatment with Proglycem® Oral Suspension resulted in a statistically significant increase ($p<0.001$) in NEFA at 3 hours, which by 6 hours after dose administration had returned to levels well below baseline. Similarly the increase in NEFA at 3 hours after administration of diazoxide choline controlled-release tablets was statistically significant ($p<0.0012$). NEFA levels in the diazoxide choline controlled-release tablet treated subjects also returned to levels well below baseline by 6 hours after dose administration.

Table 36. Fasting glucose, fasting insulin, non-esterified fatty acid (NEFA), serum sodium (Na), potassium (K) and creatinine of subjects

Parameter	Proglycem Oral Suspension (n=15)	Diazoxide Choline Controlled-Release Tablet (n=15)
Fasting Glucose (mg/dL)	Mean±SD (range)	Mean±SD (range)
Baseline	93±6 (83-104)	93±6 (83-108)
1 h post-dose	99.4±7 (89-117)	97.9±7 (92-117)
3 h post-dose	102.8±6 (93-113)	98.0±7 (90-112)
24 h post-dose	97±5 (87-106)	98±8 (88-118)
48 h post-dose	92±5 (84-103)	93±5 (83-104)
Fasting Insulin (μIU/ml)		
Baseline	8.6±5.2 (2.9-24.1)	9.9±3.3 (4.4-16.3)

	24 h post-dose	12.0±8.9 (6.1-41.9)	11.1±3.3 (5.5-17.0)
	48 h post-dose	9.3±5.7 (4.0-28.9)	11.1±2.9 (5.3-17.8)
NEFA (μmol/L)			
	1 h post-dose	0.36±0.11 (0.21-0.65)	0.34±0.14 (0.09-0.67)
	3 h post-dose	0.53±0.13 (0.31-0.78)	0.55±0.19 (0.28-0.88)
	6 h post-dose	0.13±0.07 (0.07-0.32)	0.11±0.04 (0.07-0.21)
	24 h post-dose	0.36±0.10 (0.23-0.62)	0.40±0.18 (0.25-0.97)
	48 h post-dose	0.38±0.13 (0.2-0.66)	0.39±0.15 (0.17-0.65)
Na (mEq/L)			
	Baseline	138.0±2.0 (135-142)	138.1±2.6 (134-143)
	End of Study	138.7±1.1 (137-141)	139.3±2.3 (135-144)
K (mmol/L)			
	Baseline	4.4±0.4 (3.9-5.5)	4.2±0.3 (3.8-5.1)
	End of Study	4.4±0.3(3.9-5.2)	4.2±0.3(3.8-5.2)
Creatinine (mg/dL)			
	Baseline	0.9±0.1 (0.7-1.1)	0.9±0.2 (0.7-1.4)
	End of Study	1.0±0.1 (0.8-1.3)	0.9±0.2 (0.6-1.3)

[0660] All other clinical laboratory tests, including hematology, clinical serum chemistry, and urinalysis were within normal ranges for obese subjects.

7.2.3. Vital Signs.

[0661] A graph depicting sitting blood pressure at baseline and at various times following dose administration is provided in **Figure 22**. Neither treatment was associated with a sustained reduction or increase in blood pressure, nor was there any clear trend from baseline over 6, 9, 12, or 24 hours after dose administration. Pulse rates dropped incrementally from baseline levels in the period from dose administration to 3 hours post dosing (**Figure 23**). Pulse rates rose to levels equivalent to baseline or slightly above baseline in the period from 6 to 12 hours after dose administration, and remained at baseline levels from 24 hours after dose administration to the end of the study (**Figure 23**). Review of the baseline cardiac telemetry data for all subjects was conducted the morning of dosing. Based on the absence of clinical significant abnormalities, all subjects were allowed to proceed with dosing. The cardiac telemetry data from study

time 0 to 24 hours after dose administration was evaluated. No clinically significant abnormalities (e.g. arrhythmias) were observed.

7.2.4. Preliminary Evaluation of Diazoxide Plasma Pharmacokinetics.

[0662] Compared to the Proglycem® Oral Suspension, the diazoxide choline controlled-release tablet had a 30% lower peak exposure (C_{max}) and a 15% lower total exposure (AUC) when the formulations were administered orally as 200 mg equivalents of diazoxide (Table 37). The timing of the peak plasma concentration (T_{max}) occurred at a median time of 4 hr after administration of the oral suspension and 20 hr after administration of the diazoxide choline controlled-release tablet. The terminal half-life for diazoxide was similar with both formulations (29 hr for the oral suspension and 32 hr for the tablet). The two formulations had similar between-patient variability for C_{max} and AUC. The most notable difference between the two formulations following a single dose was the lower, broader peak of the concentration-time profile of the tablet formulation (Figures 24 and 25). The peak concentration for the tablet averaged about 30% less and occurred about 16 hours later than the suspension. Figures 24 and 25 illustrate that the mean peak concentration was virtually unchanged between 12-hr and 24-hr post-dose following administration of the diazoxide choline controlled-release tablet formulation. Because of a more sustained release pattern, the tablet formulation was predicted to have a greater accumulation factor with chronic dosing than was the oral suspensions (3.96 vs. 2.84). Simulations of repeated once-daily dosing with the two formulations (assuming linear pharmacokinetics) predicted that the two formulations would have similar trough concentrations at steady-state (22-23 $\mu\text{g/mL}$) (see Figure 26).

Table 37. Summary Statistics for Plasma Diazoxide Pharmacokinetic Parameters after a Single Dose of Two Formulations

	Diazoxide Choline Controlled-Release Tablets							
	C_{max}	T_{max}	AUC(0-24)	AUC(0-120)	AUC(0- ∞)	λ_z	$T_{1/2}$	Accum
	$\mu\text{g/mL}$	hr	$\mu\text{g}\cdot\text{hr/mL}$	$\mu\text{g}\cdot\text{hr/mL}$	$\mu\text{g}\cdot\text{hr/mL}$	1/hr	hr	Factor
N	15	15	15	15	15	15	15	15

Mean	9.25	22.1	166	553	618	0.0242	31.9	3.96
Geomean	9.07	19.6	157	531	588	0.0229	30.2	3.74
St Dev	1.89	11.5	54	160	198	0.0085	10.6	1.48
SEM	0.49	3.0	14	41	51	0.0022	2.7	0.38
CV%	20.5%	52.0%	32.5%	28.9%	32.0%	35.2%	33.3%	37.5%
Median	9.08	20.0	164	518	586	0.0237	29.3	3.80
Min	5.53	8.00	74.6	261	274	0.0144	15.9	2.26
Max	13.2	48.0	266	842	1024	0.0437	48.3	7.84

Diazoxide Oral Suspension (Proglycem)								
	C _{max}	T _{max}	AUC(0-24)	AUC(0-120)	AUC(0-∞)	λ _z	T _{1/2}	Accum
	μg/mL	hr	μg·hr/mL	μg·hr/mL	μg·hr/mL	1/hr	hr	Factor
N	15	15	15	15	15	15	15	15
Mean	13.5	6.67	245	643	696	0.0256	28.6	2.84
Geomean	13.3	5.36	243	629	677	0.0249	27.9	2.79
St Dev	2.3	5.49	35	142	173	0.0068	6.3	0.57
SEM	0.6	1.42	9	37	45	0.0018	1.6	0.15
CV%	17.1%	82.3%	14.4%	22.1%	24.9%	26.7%	22.0%	20.0%
Median	13.3	4.00	245	620	671	0.0234	29.6	2.86
Min	10.1	2.00	183	464	488	0.0180	16.1	1.84
Max	18.0	24.0	314	926	1054	0.0430	38.5	4.25

7.3 Conclusions of Comparison of Single Doses of Diazoxide Administered as Proglycem® Oral Suspension or as Diazoxide Choline Controlled-Release Tablets in Obese Subjects.

[0663] The present clinical study on the comparison of a single dose of diazoxide choline controlled-release tablet with an equivalent dose of diazoxide administered as Proglycem® Oral Suspension in obese patients indicates that both formulations were well tolerated. With a single exception, moderate headache requiring treatment with acetaminophen in one subject treated with Proglycem®, all adverse events were mild. Diazoxide choline controlled-release tablets appear to have a better CNS safety profile than Proglycem® Oral Suspension as evidenced by the absence of headaches and reduced rate of dizziness. Exploratory endpoints showed no detrimental impact. Preliminary pharmacokinetic analysis showed that compared to Proglycem® Oral Suspension, the diazoxide choline controlled-release tablet had a 30% lower peak exposure (C_{max}) and a 15% lower total exposure (AUC) for the same 200 mg diazoxide equivalent dose. The timing of the peak plasma concentration (T_{max}) occurred at a median time of 4 hr after administration of the oral suspension and 20 hr after administration of the diazoxide choline tablet. The terminal half-life for diazoxide is similar with both formulations (29 hr for the oral suspension and 32 hr for the tablet). The two formulations had similar between-patient variability for C_{max} and AUC.

[0664] All patents and other references cited in the specification are indicative of the level of skill of those skilled in the art to which the invention pertains, and are incorporated by reference in their entireties, including any tables and figures, to the same extent as if each reference had been incorporated by reference in its entirety individually.

[0665] One skilled in the art would readily appreciate that the present invention is well adapted to obtain the ends and advantages mentioned, as well as those inherent therein. The methods, variances, and compositions described herein as presently representative of preferred embodiments are exemplary and are not intended as limitations on the scope. Changes therein and other uses will occur to those skilled in the art, which are encompassed within the spirit of the invention, are defined by the scope of the claims.

[0666] Definitions provided herein are not intended to be limiting from the meaning commonly understood by one of skill in the art unless indicated otherwise.

[0667] The inventions illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising", "including," "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the inventions embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

[0668] The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein. Other embodiments are within the following claims. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

THAT WHICH IS CLAIMED IS:

1. A salt comprising:
an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV; and
a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group,
wherein when said salt comprises diazoxide as the anion and sodium as the cation, said salt is not in a form suitable for intravenous use.
2. The salt according to claim 1 wherein when said salt comprises diazoxide as the anion and sodium as the cation, said salt is not in a liquid form.
3. The salt according to claim 1 wherein said anion is an anion of Formula III.
4. The salt according to claim 1 wherein said anion is an anion of Formula IV.
5. The salt according to claim 1 wherein said anion is an anion of diazoxide.
6. The salt according to claim 1 wherein said compound comprising a tertiary amine or ammonium group is selected from choline or hexamethyl hexamethylene diammonium.
7. The salt according to claim 1 wherein said alkali metal cation is selected from a sodium cation or a potassium cation.
8. The salt according to claim 1 wherein said salt is selected from a sodium, potassium, choline, or hexamethyl hexamethylene diammonium salt of Formula III or Formula IV.
9. A pharmaceutical formulation comprising the salt according to claim 1.
10. A salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula V, Formula VI, Formula VII and Formula VIII.

11. The salt according to claim 10 wherein at least one substituent comprises an amino group.
12. The salt according to claim 10 wherein R^1 comprises an amino group.
13. The salt according to claim 10 wherein said anion is an anion of Formula VII.
14. The salt according to claim 10 wherein said anion is an anion of Formula VIII.
15. The salt according to claim 10 wherein said salt comprises a cation of an alkali metal.
16. A pharmaceutical formulation comprising the salt according to claim 10.
17. A salt comprising a cation of a K_{ATP} channel opener selected from the group consisting of Formula V, Formula VI, Formula VII and Formula VIII, wherein at least one substituent comprises an amino group.
18. The salt according to claim 17 wherein R^1 comprises an amino group.
19. The salt according to claim 17 wherein said cation is an cation of Formula VII.
20. The salt according to claim 17 wherein said cation is an cation of Formula VIII.
21. The salt according to claim 17 wherein said salt comprises an anion selected from the group consisting of sulfate, chloride, hydrochloride, fumarate, malate, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluene sulfonate, cyclohexylsulfamate and quinate ions.
22. The salt according to claim 17 wherein said salt comprises an anion selected from a halogen.
23. A pharmaceutical formulation comprising the salt according to claim 17.

24. The formulation according to any one of claims 9, 16 and 23 for a single administration that contains between 10 and 100 mg of the salt.
25. The formulation according to any one of claims 9, 16 and 23 for a single administration that contains between 100 and 200 mg of the salt.
26. The formulation according to any one of claims 9, 16 and 23 for a single administration that contains between 200 and 300 mg of the salt.
27. The formulation according to any one of claims 9, 16 and 23 for a single administration that contains between 300 and 500 mg of the salt.
28. The formulation according to any one of claims 9, 16 and 23 that contains between 500 mg and 2000 mg of the salt.
29. The formulation according to any one of claims 9, 16 and 23 that is suitable for oral administration.
30. The formulation according to claim 29 further comprising at least one component that substantially inhibits release of the K_{ATP} channel opener from the formulation until after gastric transit.
31. The formulation according to claim 30 wherein the component is selected from the group consisting of: (a) a pH sensitive polymer or co-polymer applied as a compression coating on a tablet, (b) a pH sensitive polymer or co-polymer applied as a thin film on a tablet, (c) a pH sensitive polymer or co-polymer applied as a thin film to an encapsulation system, (d) a pH sensitive polymer or co-polymer applied to encapsulated microparticles, (e) a non-aqueous-soluble polymer or copolymer applied as a compression coating on a tablet, (f) a non-aqueous-soluble polymer or co-polymer applied as a thin film on a tablet, (g) a non-aqueous soluble polymer applied as a thin film to an encapsulation system, and (h) combinations of these approaches, wherein the pH sensitive polymer or co-polymer is resistant to degradation under acid conditions.
32. The formulation according to either of claims 29 or 30 further comprising a component that contributes to sustained release of the salt over a period of 2-30 hours following administration.

33. The formulation according to claim 32 wherein the time period is 2-4 hours following administration.

34. The formulation according to claim 32 wherein the time period is 4-8 hours following administration.

35. The formulation according to claim 32 wherein the time period is 8-24 hours following administration.

36. The formulation according to claim 32 wherein the component is selected from the group consisting of: (a) a pH sensitive polymeric coating, (b) a hydrogel, (c) a film coating that controls the rate of diffusion of the drug from a coated matrix, (d) an erodable matrix that controls rate of drug release, (e) polymer coated pellets, granules or microparticles of drug which can be further encapsulated or compressed into a tablet, (f) an osmotic pump system containing the drug, (g) a compression coated tablet form of the drug, and (h) combinations thereof.

37. A method of (a) inhibiting fasting insulin secretion (b) inhibiting glucose stimulated insulin secretion, (c) elevating energy expenditure, (d) elevating beta oxidation of fat, or (e) inhibiting hyperphagia, in an obese, overweight or obesity prone subject, comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

38. The method according to claim 37 wherein the duration of said inhibition or elevation is at least about 18 hours.

39. The method according to claim 37 wherein the duration of said inhibition or elevation is at least about 24 hours.

40. A method of maintaining weight loss in an obese, overweight, or obesity prone subject, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

41. The method according to claim 40 wherein said formulation is administered once per 24 hours.

42. The method according to claim 40 wherein said formulation is administered two times per 24 hours.

43. A method of inducing weight loss in an obese or overweight subject, comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

44. The method according to claim 43 wherein the formulation is administered once per 24 hours.

45. The method according to claim 43 wherein the formulation is administered two times per 24 hours.

46. A method of treating obesity associated co-morbidities in an obese, overweight or obesity prone subject, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III

and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

47. The method according to claim 46 wherein the formulation is administered once per 24 hours.

48. The method according to claim 46 wherein the formulation is administered two times per 24 hours.

49. A method of elevating energy expenditure in an overweight, obese or obesity prone subject, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

50. The method according to claim 49 wherein the formulation is administered once per 24 hours.

51. The method according to claim 49 wherein the formulation is administered two times per 24 hours.

52. A method of elevating beta oxidation of fat in an overweight, obese or obesity prone subject, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

53. The method according to claim 52 wherein the formulation is administered once per 24 hours.

54. The method according to claim 52 wherein the formulation is administered two times per 24 hours.

55. A method of preventing the transition to diabetes of a prediabetic subject comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

56. The method according to claim 55 wherein the formulation is administered once per 24 hours.

57. The method according to claim 55 wherein the formulation is administered two times per 24 hours.

58. A method of restoring normal glucose tolerance to a diabetic subject comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

59. The method according to claim 58 wherein the formulation is administered once per 24 hours.

60. The method according to claim 58 wherein the formulation is administered two times per 24 hours.

61. A method of restoring normal glucose tolerance to a prediabetic subject comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

62. The method according to claim 61 wherein the formulation is administered once per 24 hours.

63. The method according to claim 61 wherein the formulation is administered two times per 24 hours.

64. A method of delaying or preventing the progression of diabetes in a subject comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

65. The method according to claim 64 wherein the formulation is administered once per 24 hours.

66. The method according to claim 64 wherein the formulation is administered two times per 24 hours.

67. A method of treating obesity or obesity associated co-morbidities or other diseases or conditions involving K_{ATP} channels by co-administration to a subject in need thereof a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III

and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23, and a drug selected from the group consisting of an amphetamine or amphetamine mixture, Sibutramine, Orlistat, Rimonabant, a CB-1 agonist, a 5HT_{2c} receptor agonist, a drug used to treat addiction, a beta adrenergic receptor agonist, an ACC inhibitor, leptin, a leptin analogue, a leptin agonist, a somatostatin agonist, an adiponectin agonist or secretagogue, Amylin, PYY or a PYY analogue, a ghrelin antagonist, a drug that inhibits gastrointestinal lipases or other digestive enzymes, a de-novo lipogenesis inhibitor, a drug that blocks absorption of dietary fat, growth hormone or a growth hormone analogue, a growth hormone secretagogue, a CCK agonist, an oleoylethanolamine receptor agonist, a fatty acid synthase inhibitor, a thyroid receptor agonist, a selective androgen receptor modulator, a PPAR agonist, oxymodulin, oleoylestrone, a NPY2 receptor antagonist, a NPY5 receptor antagonist, a NPY agonist, a monoamine uptake inhibitor, a MTP inhibitor, a MC4 receptor agonist, a MCH1 receptor antagonist, a 5HT-6 antagonist, a histamine-3 antagonist, a glycine analog, a fgfl inhibitor, a DGAT-1 inhibitor, a carboxypeptidase inhibitor, an appetite suppressant, a non-thiazide diuretic, a drug that lowers cholesterol, a drug that raises HDL cholesterol, a drug that lowers LDL cholesterol, a drug that lowers blood pressure, a drug that is an anti-depressant, a drug that improves insulin sensitivity, a drug that improves glucose utilization or uptake, a drug that is an anti-epileptic, a drug that is an anti-inflammatory, a drug that is an appetite suppressant, a drug that lowers circulating triglycerides, a drug that is used to induce weight loss in an overweight or obese individual, and pharmaceutically acceptable salts thereof.

68. A method of preventing weight gain, dyslipidemia, diabetes or impaired glucose tolerance in a subject treated with an anti-psychotic drug, the method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

69. A method of treating weight gain, dyslipidemia, diabetes or impaired glucose tolerance in a subject treated with an anti-psychotic drug, the method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

70. A method of treating a subject having one or more diseases characterized by obesity, hyperphagia, dyslipidemia, or decreased energy expenditure including (a) Prader Willi Syndrome, (b) Froelich's syndrome, (c) Cohen syndrome, (d) Summit Syndrome, (e) Alstrom, Syndrome, (f) Borjesen Syndrome, (g) Bardet-Biedl Syndrome, or (h) hyperlipoproteinemia type I, II, III, and IV, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

71. The formulation according to any one of claims 9, 16 and 23 further comprising another pharmaceutically active agent.

72. The formulation according to claims 71 wherein the other pharmaceutically active agent is an agent useful for the treatment of a condition selected from the group consisting of obesity, prediabetes, diabetes, hypertension, depression, elevated cholesterol, fluid retention, obesity associated co-morbidities, ischemic and reperfusion injury, epilepsy, cognitive impairment, schizophrenia, mania, and other psychotic condition.

73. The pharmaceutical formulation according to any one of claims 9, 16 and 23 further comprising a pharmaceutically active agent other than the K_{ATP} channel opener.

74. The formulation according to claim 73 wherein the other pharmaceutically active agent is an agent useful for the treatment of a condition selected from the group consisting of obesity, prediabetes, diabetes, hypertension, depression, elevated cholesterol, fluid retention, obesity associated co-morbidities, ischemic and reperfusion injury, epilepsy, cognitive impairment, schizophrenia, mania, and other psychotic condition.

75. The formulation according to claim 9 wherein said cation of the salt is selected from the sodium, potassium, choline and hexamethyl hexamethylene diammonium hydroxide of diazoxide.

76. The formulation according to claim 75 for a single administration that contains between 10 and 2000 mg of the salt.

77. The formulation according to claim 75 wherein said formulation is a controlled release formulation further comprising at least one component that substantially inhibits release of the K_{ATP} channel opener from the formulation until after gastric transit.

78. The formulation according to claim 75 further comprising a component that contributes to the sustained release of the K_{ATP} channel opener over a period of 2-24 hours.

79. A method for the preparation of a salt of diazoxide comprising:
reacting one equivalent of a cation source selected from the group consisting of sodium hydroxide, potassium hydroxide, choline hydroxide, and hexamethyl hexamethylene diammonium dihydroxide with diazoxide, wherein said diazoxide is dissolved in a solvent selected from the group consisting of acetonitrile, low molecular weight ketones, tetrahydrofuran, dimethylformamide and n-methyl pyrrolidinone, and removing the solvent.

80. A polymorph of a salt, said salt comprising:
diazoxide, and

a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or quaternary ammonium group.

81. The polymorph of Claim 80, wherein said cation is choline.

82. The polymorph of Claim 81 comprising Form A having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 9.8, 10.5, 14.9, 17.8, 17.9, 18.5, 19.5, 22.1, 22.6, 26.2, 29.6, and 31.2 degrees.

83. The polymorph of Claim 81 comprising Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.9, 10.3, 12.0, 18.3, 20.6, 24.1, 24.5, 26.3, 27.1, and 28.9 degrees.

84. The polymorph of Claim 81 comprising Form A having characteristic infrared absorbances at 2926, 2654, 1592, 1449, and 1248 cm⁻¹.

85. The polymorph of Claim 81 comprising Form B having characteristic infrared absorbances at 3256, 2174, 2890, 1605, 1463, and 1235 cm⁻¹.

86. The polymorph of Claim 80, wherein said cation is potassium.

87. The polymorph of Claim 86 comprising Form A having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 6.0, 8.1, 16.3, 17.7, 18.6, 19.1, 22.9, 23.3, 23.7, 24.7, 25.4, 26.1, 28.2, 29.6, and 30.2 degrees.

88. The polymorph of Claim 86 comprising Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 10.8, 16.9, 18.2, 21.6, 25.5, 26.1, and 28.9 degrees.

89. The polymorph of Claim 86 comprising Form C having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.7, 6.1, 17.9, 23.9, 25.1, and 37.3 degrees.

90. The polymorph of Claim 86 comprising Form D having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.7, 6.2, 8.1, 8.5, 8.8, 16.9, 18.6, 23.2, 24.5, 25.8, and 26.1 degrees.

91. The polymorph of Claim 86 comprising Form E having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 6.7, 7.1, 14.1, and 21.2 degrees.

92. The polymorph of Claim 86 comprising Form F having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 9.0, 18.7, 20.6, 23.5, 27.5, and 36.3 degrees.

93. The polymorph of Claim 86 comprising Form G having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 5.2, 5.5, 13.1, 16.5, 19.3, 22.8, 24.8, 26.4, 28.7, and 34.1 degrees.

94. The polymorph of Claim 86 comprising Form A having characteristic infrared absorbances at 1503, 1374, 1339, 1207, 1131, 1056, and 771 cm⁻¹.

95. The polymorph of Claim 86 comprising Form B having characteristic infrared absorbances at 1509, 1464, 1378, and 1347 cm⁻¹.

96. The polymorph of Claim 86 comprising Form C having characteristic infrared absorbances at 1706, 1208, 1146, and 746 cm⁻¹.

97. The polymorph of Claim 86 comprising Form D having characteristic infrared absorbances at 1595, 1258, 1219, and 890 cm⁻¹.

98. The polymorph of Claim 86 comprising Form E having characteristic infrared absorbances at 1550, 1508, 1268, 1101, and 1006 cm⁻¹.

99. The polymorph of Claim 86 comprising Form F having characteristic infrared absorbances at 1643, 1595, 1234, 1145, and 810 cm⁻¹.

100. The polymorph of Claim 86 comprising Form G having characteristic infrared absorbances at 1673, 1591, 1504, 1458, 1432, 1266, 999, 958, 905, and 872 cm⁻¹.

101. The polymorph of Claim 81 comprising Form A having characteristic peaks in the XRPD pattern substantially as shown in FIG. 16(a), and an NMR spectrum substantially as shown in FIG. 17(a).

102. The polymorph of Claim 81 comprising Form B having characteristic peaks in the XRPD pattern substantially as shown in FIG. 16(c) and an NMR spectrum substantially as shown in FIG. 17(b).

103. The polymorph of Claim 86 comprising one or more of Forms A-G, wherein each said Form A-G has characteristic peaks in the XRPD pattern substantially as shown in FIG. 18-19.

104. A process for producing a diazoxide choline salt, said process comprising:
suspending diazoxide in a solvent and mixing with a choline salt;
adding a co-solvent to the suspension under conditions sufficient to cause
formation and precipitation of said diazoxide choline salt;
harvesting the precipitate to provide said diazoxide choline salt.

105. The process of Claim 104 wherein said solvent is THF.

106. The process of Claim 104, wherein said solvent is 2-MeTHF.

107. The process of Claim 104, wherein said diazoxide and said solvent are present at a ratio of about 1 g diazoxide per 1 mL solvent to about 1 g diazoxide per 5 mL solvent.

108. The process of Claim 107, wherein said diazoxide and said solvent are present at a ratio of about 1 g diazoxide per 3 mL solvent.

109. The process of Claim 104, wherein said choline salt is choline hydroxide.

110. The process of Claim 104, wherein said choline salt is a solution in MeOH.

111. The process of Claim 110, wherein said choline salt is choline hydroxide in a 45% solution in MeOH.

112. The process of Claim 104, wherein said choline salt is added as 1 equivalent of said diazoxide.
113. The process of Claim 104, wherein said co-solvent is MTBE.
114. The process of Claim 104, wherein the amount of said co-solvent added is in a ratio to the amount of said solvent of about 3:14 (solvent:co-solvent).
115. The process of Claim 104, further comprising seeding with crystals of diazoxide choline salt polymorph Form B prior to said harvesting step.
116. The process of Claim 104, wherein said choline salt of diazoxide comprises polymorph Form B substantially free of polymorph Form A, said polymorph Form B having characteristic peaks in the XRPD pattern at values of two-theta (Cu K α , 40 kV, 40 mA) at approximately 8.5, 10.8, 16.9, 18.2, 21.6, 25.5, 26.1, and 28.9 degrees.
117. A method of treating obesity or obesity-related morbidity in an obese subject, said method comprising administering to a subject in need thereof an effective amount of a compound of Formula V-VIII.
118. The method of Claim 117, wherein said compound is a compound of Formula V.
119. The method of Claim 117, wherein said compound is a compound of Formula VI.
120. The method of Claim 117, wherein said compound is a compound of Formula VII.
121. The method of Claim 117, wherein said compound is a compound of Formula VIII.
122. The method of Claim 117, further comprising administering a drug selected from the group consisting of an amphetamine or amphetamine mixture, Sibutramine, Orlistat, Rimonabant, a CB-1 agonist, a 5HT_{2c} receptor agonist, a drug used to treat addiction, a beta adrenergic receptor agonist, an ACC inhibitor, leptin, a leptin analogue, a leptin agonist, a somatostatin agonist, an adiponectin agonist or secretagogue,

Amylin, PYY or a PYY analogue, a ghrelin antagonist, a drug that inhibits gastrointestinal lipases or other digestive enzymes, a de-novo lipogenesis inhibitor, a drug that blocks absorption of dietary fat, growth hormone or a growth hormone analogue, a growth hormone secretagogue, a CCK agonist, an oleoylethanolamine receptor agonist, a fatty acid synthase inhibitor, a thyroid receptor agonist, a selective androgen receptor modulator, a PPAR agonist, oxyntomodulin, oleoylestrone, a NPY2 receptor antagonist, a NPY5 receptor antagonist, a NPY agonist, a monoamine uptake inhibitor, a MTP inhibitor, a MC4 receptor agonist, a MCH1 receptor antagonist, a 5HT-6 antagonist, a histamine-3 antagonist, a glycine analog, a fgfl inhibitor, a DGAT-1 inhibitor, a carboxypeptidase inhibitor, an appetite suppressant, a non-thiazide diuretic, a drug that lowers cholesterol, a drug that raises HDL cholesterol, a drug that lowers LDL cholesterol, a drug that lowers blood pressure, a drug that is an anti-depressant, a drug that is an anti-epileptic, a drug that is an anti-inflammatory, a drug that lowers circulating triglycerides, a drug that is used to induce weight loss in an overweight or obese individual, and pharmaceutically acceptable salts thereof.

123. The method of Claim 117, further comprising administering a pharmaceutically active agent other than a K_{ATP} channel opener.

124. The method of Claim 123, wherein said other pharmaceutically active agent is an agent useful for the treatment of a condition selected from the group consisting of obesity, prediabetes, diabetes, hypertension, depression, elevated cholesterol, fluid retention, obesity associated co-morbidities, ischemic and reperfusion injury, epilepsy, cognitive impairment, schizophrenia, mania, and other psychotic condition.

125. A method of treatment of a subject suffering from or at risk for Alzheimer's disease, said method comprising administering to said subject a therapeutically effective amount of a compound according to any of Formulas I-VIII.

126. The method of claim 125 wherein said compound is diazoxide.

127. A method of treatment of a subject suffering from or at risk for Alzheimer's disease, said method comprising administering to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a)

a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

128. A method for treatment of a subject suffering from hypoglycemia, said method comprising administration to said subject a therapeutically effective amount of a formulation selected from the group consisting of: a) a formulation comprising a salt, said salt comprising an anion of a K_{ATP} channel opener selected from the group consisting of Formula I, Formula II, Formula III and Formula IV, and a cation selected from the group consisting of an alkali metal and a compound comprising a tertiary amine or ammonium group; b) a formulation according to claim 16; and c) a formulation according to claim 23.

129. The method of claim 128, wherein said hypoglycemia is selected from the group consisting of a) nighttime hypoglycemia; b) hypoglycemia attributable to a defect in insulin secretion; c) hypoglycemia attributable to an insulin secreting tumor; and d) drug-induced hypoglycemia.

1/51




FIG. 1