

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		
The application data sheet is part of the provisional or nonprovisional application for which it is being submitted. The following form contains the bibliographic data arranged in a format specified by the United States Patent and Trademark Office as outlined in 37 CFR 1.76. This document may be completed electronically and submitted to the Office in electronic format using the Electronic Filing System (EFS) or the document may be printed and included in a paper filed application.			

Secrecy Order 37 CFR 5.2:

☐	Portions or all of the application associated with this Application Data Sheet may fall under a Secrecy Order pursuant to 37 CFR 5.2 (Paper filers only. Applications that fall under Secrecy Order may not be filed electronically.)
--------------------------	---

Inventor Information:

Inventor 1					Remove		
Legal Name							
Prefix	Given Name		Middle Name		Family Name		Suffix
	Alexis				HOWERTON		
Residence Information (Select One)							☒ US Residency ☐ Non US Residency ☐ Active US Military Service
City	San Francisco		State/Province	CA	Country of Residence ⁱ	US	
Mailing Address of Inventor:							
Address 1		611 Gateway Boulevard, Suite #740					
Address 2							
City	South San Francisco			State/Province	CA		
Postal Code	94080			Country ⁱ	US		
Inventor 2					Remove		
Legal Name							
Prefix	Given Name		Middle Name		Family Name		Suffix
	Hal				GERBER		
Residence Information (Select One)							☒ US Residency ☐ Non US Residency ☐ Active US Military Service
City	San Francisco		State/Province	CA	Country of Residence ⁱ	US	
Mailing Address of Inventor:							
Address 1		611 Gateway Boulevard, Suite #740					
Address 2							
City	South San Francisco			State/Province	CA		
Postal Code	94080			Country ⁱ	US		
All Inventors Must Be Listed - Additional Inventor Information blocks may be generated within this form by selecting the Add button.							Add

Correspondence Information:

Enter either Customer Number or complete the Correspondence Information section below. For further information see 37 CFR 1.33(a).

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

☐ An Address is being provided for the correspondence information of this application.			
Customer Number	21971		
Email Address	patentdocket@wsgr.com	<input type="button" value="Add Email"/>	<input type="button" value="Remove Email"/>

Application Information:

Title of the Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		
Attorney Docket Number	50535-710.306	Small Entity Status Claimed	☐
Application Type	Nonprovisional		
Subject Matter	Utility		
Total Number of Drawing Sheets (if any)	14	Suggested Figure for Publication (if any)	

Filing By Reference:

Only complete this section when filing an application by reference under 35 U.S.C. 111(c) and 37 CFR 1.57(a). Do not complete this section if application papers including a specification and any drawings are being filed. Any domestic benefit or foreign priority information must be provided in the appropriate section(s) below (i.e., "Domestic Benefit/National Stage Information" and "Foreign Priority Information").

For the purposes of a filing date under 37 CFR 1.53(b), the description and any drawings of the present application are replaced by this reference to the previously filed application, subject to conditions and requirements of 37 CFR 1.57(a).

Application number of the previously filed application	Filing date (YYYY-MM-DD)	Intellectual Property Authority or Country

Publication Information:

☐ Request Early Publication (Fee required at time of Request 37 CFR 1.219)
☐ Request Not to Publish. I hereby request that the attached application not be published under 35 U.S.C. 122(b) and certify that the invention disclosed in the attached application has not and will not be the subject of an application filed in another country, or under a multilateral international agreement, that requires publication at eighteen months after filing.

Representative Information:

Representative information should be provided for all practitioners having a power of attorney in the application. Providing this information in the Application Data Sheet does not constitute a power of attorney in the application (see 37 CFR 1.32). Either enter Customer Number or complete the Representative Name section below. If both sections are completed the customer Number will be used for the Representative Information during processing.			
Please Select One:	☒ Customer Number	☐ US Patent Practitioner	☐ Limited Recognition (37 CFR 11.9)
Customer Number	21971		

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

Domestic Benefit/National Stage Information:

This section allows for the applicant to either claim benefit under 35 U.S.C. 119(e), 120, 121, 365(c), or 386(c) or indicate National Stage entry from a PCT application. Providing benefit claim information in the Application Data Sheet constitutes the specific reference required by 35 U.S.C. 119(e) or 120, and 37 CFR 1.78.

When referring to the current application, please leave the "Application Number" field blank.

Prior Application Status	Pending	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
	Continuation of	18941832	2024-11-08		
Prior Application Status	Abandoned	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
18941832	Continuation of	17667285	2022-02-08		
Prior Application Status	Patented	Remove			
Application Number	Continuity Type	Prior Application Number	Filing Date (YYYY-MM-DD)	Patent Number	Issue Date (YYYY-MM-DD)
17/667285	Continuation of	17/081694	2020-10-27	11304950	2022-04-19
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
17/081694	Continuation of	PCT/US2019/029486	2019-04-26		
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
PCT/US2019/029486	Claims benefit of provisional	62/822815	2019-03-23		
Prior Application Status	Expired	Remove			
Application Number	Continuity Type	Prior Application Number	Filing or 371(c) Date (YYYY-MM-DD)		
PCT/US2019/029486	Claims benefit of provisional	62/663951	2018-04-27		
Additional Domestic Benefit/National Stage Data may be generated within this form by selecting the Add button.					Add

Foreign Priority Information:

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

This section allows for the applicant to claim priority to a foreign application. Providing this information in the application data sheet constitutes the claim for priority as required by 35 U.S.C. 119(b) and 37 CFR 1.55. When priority is claimed to a foreign application that is eligible for retrieval under the priority document exchange program (PDX)ⁱ the information will be used by the Office to automatically attempt retrieval pursuant to 37 CFR 1.55(i)(1) and (2). Under the PDX program, applicant bears the ultimate responsibility for ensuring that a copy of the foreign application is received by the Office from the participating foreign intellectual property office, or a certified copy of the foreign priority application is filed, within the time period specified in 37 CFR 1.55(g)(1).

			Remove
Application Number	Country ⁱ	Filing Date (YYYY-MM-DD)	Access Code ⁱ (if applicable)

Additional Foreign Priority Data may be generated within this form by selecting the **Add** button.

Add

Statement under 37 CFR 1.55 or 1.78 for AIA (First Inventor to File) Transition Applications

<p>This application (1) claims priority to or the benefit of an application filed before March 16, 2013 and (2) also contains, or contained at any time, a claim to a claimed invention that has an effective filing date on or after March 16, 2013.</p> <p>☐ NOTE: By providing this statement under 37 CFR 1.55 or 1.78, this application, with a filing date on or after March 16, 2013, will be examined under the first inventor to file provisions of the AIA.</p>
--

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

Authorization or Opt-Out of Authorization to Permit Access:

When this Application Data Sheet is properly signed and filed with the application, applicant has provided written authority to permit a participating foreign intellectual property (IP) office access to the instant application-as-filed (see paragraph A in subsection 1 below) and the European Patent Office (EPO) access to any search results from the instant application (see paragraph B in subsection 1 below).

Should applicant choose not to provide an authorization identified in subsection 1 below, applicant **must opt-out** of the authorization by checking the corresponding box A or B or both in subsection 2 below.

NOTE: This section of the Application Data Sheet is **ONLY** reviewed and processed with the **INITIAL** filing of an application. After the initial filing of an application, an Application Data Sheet cannot be used to provide or rescind authorization for access by a foreign IP office(s). Instead, Form PTO/SB/39 or PTO/SB/69 must be used as appropriate.

1. Authorization to Permit Access by a Foreign Intellectual Property Office(s)

A. Priority Document Exchange (PDX) - Unless box A in subsection 2 (opt-out of authorization) is checked, the undersigned hereby **grants the USPTO authority** to provide the European Patent Office (EPO), the Japan Patent Office (JPO), the Korean Intellectual Property Office (KIPO), the State Intellectual Property Office of the People's Republic of China (SIPO), the World Intellectual Property Organization (WIPO), and any other foreign intellectual property office participating with the USPTO in a bilateral or multilateral priority document exchange agreement in which a foreign application claiming priority to the instant patent application is filed, access to: (1) the instant patent application-as-filed and its related bibliographic data, (2) any foreign or domestic application to which priority or benefit is claimed by the instant application and its related bibliographic data, and (3) the date of filing of this Authorization. See 37 CFR 1.14(h)(1).

B. Search Results from U.S. Application to EPO - Unless box B in subsection 2 (opt-out of authorization) is checked, the undersigned hereby **grants the USPTO authority** to provide the EPO access to the bibliographic data and search results from the instant patent application when a European patent application claiming priority to the instant patent application is filed. See 37 CFR 1.14(h)(2).

The applicant is reminded that the EPO's Rule 141(1) EPC (European Patent Convention) requires applicants to submit a copy of search results from the instant application without delay in a European patent application that claims priority to the instant application.

2. Opt-Out of Authorizations to Permit Access by a Foreign Intellectual Property Office(s)

☐ A. Applicant **DOES NOT** authorize the USPTO to permit a participating foreign IP office access to the instant application-as-filed. If this box is checked, the USPTO will not be providing a participating foreign IP office with any documents and information identified in subsection 1A above.

☐ B. Applicant **DOES NOT** authorize the USPTO to transmit to the EPO any search results from the instant patent application. If this box is checked, the USPTO will not be providing the EPO with search results from the instant application.

NOTE: Once the application has published or is otherwise publicly available, the USPTO may provide access to the application in accordance with 37 CFR 1.14.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

Applicant Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.			
Applicant 1			Remove
<p>If the applicant is the inventor (or the remaining joint inventor or inventors under 37 CFR 1.45), this section should not be completed. The information to be provided in this section is the name and address of the legal representative who is the applicant under 37 CFR 1.43; or the name and address of the assignee, person to whom the inventor is under an obligation to assign the invention, or person who otherwise shows sufficient proprietary interest in the matter who is the applicant under 37 CFR 1.46. If the applicant is an applicant under 37 CFR 1.46 (assignee, person to whom the inventor is obligated to assign, or person who otherwise shows sufficient proprietary interest) together with one or more joint inventors, then the joint inventor or inventors who are also the applicant should be identified in this section.</p>			
Clear			
☒ Assignee	☐ Legal Representative under 35 U.S.C. 117	☐ Joint Inventor	
☐ Person to whom the inventor is obligated to assign.		☐ Person who shows sufficient proprietary interest	
If applicant is the legal representative, indicate the authority to file the patent application, the inventor is:			
Name of the Deceased or Legally Incapacitated Inventor: <input type="text"/>			
If the Applicant is an Organization check here. ☒			
Organization Name	Spruce Biosciences, Inc.		
Mailing Address Information For Applicant:			
Address 1	611 Gateway Boulevard, Suite #740		
Address 2			
City	South San Francisco	State/Province	CA
Country	US	Postal Code	94080
Phone Number		Fax Number	
Email Address			
Additional Applicant Data may be generated within this form by selecting the Add button. Add			

Assignee Information including Non-Applicant Assignee Information:

Providing assignment information in this section does not substitute for compliance with any requirement of part 3 of Title 37 of CFR to have an assignment recorded by the Office.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

Assignee 1

Complete this section if assignee information, including non-applicant assignee information, is desired to be included on the patent application publication. An assignee-applicant identified in the "Applicant Information" section will appear on the patent application publication as an applicant. For an assignee-applicant, complete this section only if identification as an assignee is also desired on the patent application publication.

[Remove](#)If the Assignee or Non-Applicant Assignee is an Organization check here. ☐

Prefix	Given Name	Middle Name	Family Name	Suffix

Mailing Address Information For Assignee including Non-Applicant Assignee:

Address 1			
Address 2			
City		State/Province	
Country ⁱ		Postal Code	
Phone Number		Fax Number	
Email Address			

Additional Assignee or Non-Applicant Assignee Data may be generated within this form by selecting the Add button.

[Add](#)**Signature:**[Remove](#)

NOTE: This Application Data Sheet must be signed in accordance with 37 CFR 1.33(b). **However, if this Application Data Sheet is submitted with the INITIAL filing of the application and either box A or B is not checked in subsection 2 of the "Authorization or Opt-Out of Authorization to Permit Access" section, then this form must also be signed in accordance with 37 CFR 1.14(c).**

This Application Data Sheet **must** be signed by a patent practitioner if one or more of the applicants is a **juristic entity** (e.g., corporation or association). If the applicant is two or more joint inventors, this form must be signed by a patent practitioner, **all** joint inventors who are the applicant, or one or more joint inventor-applicants who have been given power of attorney (e.g., see USPTO Form PTO/AIA/81) on behalf of **all** joint inventor-applicants.

See 37 CFR 1.4(d) for the manner of making signatures and certifications.

Signature	/Victoria Cheng/		Date (YYYY-MM-DD)	2025-01-31	
First Name	Victoria	Last Name	Cheng	Registration Number	79641

Additional Signature may be generated within this form by selecting the Add button.

[Add](#)

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Application Data Sheet 37 CFR 1.76		Attorney Docket Number	50535-710.306
		Application Number	
Title of Invention	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

This collection of information is required by 37 CFR 1.76. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 23 minutes to complete, including gathering, preparing, and submitting the completed application data sheet form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. Accordingly, pursuant to the requirements of the Act, please be advised that: (1) the general authority for the collection of this information is 35 U.S.C. 2(b)(2); (2) furnishing of the information solicited is voluntary; and (3) the principal purpose for which the information is used by the U.S. Patent and Trademark Office is to process and/or examine your submission related to a patent application or patent. If you do not furnish the requested information, the U.S. Patent and Trademark Office may not be able to process and/or examine your submission, which may result in termination of proceedings or abandonment of the application or expiration of the patent.

The information provided by you in this form will be subject to the following routine uses:

1. The information on this form will be treated confidentially to the extent allowed under the Freedom of Information Act (5 U.S.C. 552) and the Privacy Act (5 U.S.C. 552a). Records from this system of records may be disclosed to the Department of Justice to determine whether the Freedom of Information Act requires disclosure of these records.
2. A record from this system of records may be disclosed, as a routine use, in the course of presenting evidence to a court, magistrate, or administrative tribunal, including disclosures to opposing counsel in the course of settlement negotiations.
3. A record in this system of records may be disclosed, as a routine use, to a Member of Congress submitting a request involving an individual, to whom the record pertains, when the individual has requested assistance from the Member with respect to the subject matter of the record.
4. A record in this system of records may be disclosed, as a routine use, to a contractor of the Agency having need for the information in order to perform a contract. Recipients of information shall be required to comply with the requirements of the Privacy Act of 1974, as amended, pursuant to 5 U.S.C. 552a(m).
5. A record related to an International Application filed under the Patent Cooperation Treaty in this system of records may be disclosed, as a routine use, to the International Bureau of the World Intellectual Property Organization, pursuant to the Patent Cooperation Treaty.
6. A record in this system of records may be disclosed, as a routine use, to another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)).
7. A record from this system of records may be disclosed, as a routine use, to the Administrator, General Services, or his/her designee, during an inspection of records conducted by GSA as part of that agency's responsibility to recommend improvements in records management practices and programs, under authority of 44 U.S.C. 2904 and 2906. Such disclosure shall be made in accordance with the GSA regulations governing inspection of records for this purpose, and any other relevant (i.e., GSA or Commerce) directive. Such disclosure shall not be used to make determinations about individuals.
8. A record from this system of records may be disclosed, as a routine use, to the public after either publication of the application pursuant to 35 U.S.C. 122(b) or issuance of a patent pursuant to 35 U.S.C. 151. Further, a record may be disclosed, subject to the limitations of 37 CFR 1.14, as a routine use, to the public if the record was filed in an application which became abandoned or in which the proceedings were terminated and which application is referenced by either a published application, an application open to public inspections or an issued patent.
9. A record from this system of records may be disclosed, as a routine use, to a Federal, State, or local law enforcement agency, if the USPTO becomes aware of a violation or potential violation of law or regulation.

**CERTIFICATION AND REQUEST FOR PRIORITIZED EXAMINATION
UNDER 37 CFR 1.102(e)** (Page 1 of 1)

First Named Inventor:	Alexis HOWERTON	Nonprovisional Application Number (if known):	
Title of Invention:	METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS		

APPLICANT HEREBY CERTIFIES THE FOLLOWING AND REQUESTS PRIORITIZED EXAMINATION FOR THE ABOVE-IDENTIFIED APPLICATION.

1. The processing fee set forth in 37 CFR 1.17(i)(1) and the prioritized examination fee set forth in 37 CFR 1.17(c) have been filed with the request. The publication fee requirement is met because that fee, set forth in 37 CFR 1.18(d), is currently \$0. The basic filing fee, search fee, and examination fee are filed with the request or have been already been paid. I understand that any required excess claims fees or application size fee must be paid for the application.
2. I understand that the application may not contain, or be amended to contain, more than four independent claims, more than thirty total claims, or any multiple dependent claims, and that any request for an extension of time will cause an outstanding Track I request to be dismissed.
3. The applicable box is checked below:

I. ☒ Original Application (Track One) - Prioritized Examination under § 1.102(e)(1)

i. (a) The application is an original nonprovisional utility application filed under 35 U.S.C. 111(a). This certification and request is being filed with the utility application via USPTO patent electronic filing system.

---OR---

(b) The application is an original nonprovisional plant application filed under 35 U.S.C. 111(a). This certification and request is being filed with the plant application in paper.

ii. An executed inventor's oath or declaration under 37 CFR 1.63 or 37 CFR 1.64 for each inventor, or the application data sheet meeting the conditions specified in 37 CFR 1.53(f)(3)(i) is filed with the application.

II. ☐ Request for Continued Examination - Prioritized Examination under § 1.102(e)(2)

- i. A request for continued examination has been filed with, or prior to, this form.
- ii. If the application is a utility application, this certification and request is being filed via USPTO patent electronic filing system.
- iii. The application is an original nonprovisional utility application filed under 35 U.S.C. 111(a), or is a national stage entry under 35 U.S.C. 371.
- iv. This certification and request is being filed prior to the mailing of a first Office action responsive to the request for continued examination.
- v. No prior request for continued examination has been granted prioritized examination status under 37 CFR 1.102(e)(2).

Signature <u>/Victoria Cheng/</u>	Date <u>1/31/2025</u>
Name (Print/Typed) <u>Victoria Cheng</u>	Practitioner Registration Number <u>79641</u>
Note: This form must be signed in accordance with 37 CFR 1.33. See 37 CFR 1.4(d) for signature requirements and certifications. Submit multiple forms if more than one signature is required.*	
☒ *Total of <u>1</u> forms are submitted.	

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. The United States Patent and Trademark Office (USPTO) collects the information in this record under authority of 35 U.S.C. 2. The USPTO's system of records is used to manage all applicant and owner information including name, citizenship, residence, post office address, and other information with respect to inventors and their legal representatives pertaining to the applicant's/owner's activities in connection with the invention for which a patent is sought or has been granted. The applicable Privacy Act System of Records Notice for the information collected in this form is COMMERCE/PAT-TM-7 Patent Application Files, available in the Federal Register at 78 FR 19243 (March 29, 2013). <https://www.govinfo.gov/content/pkg/FR-2013-03-29/pdf/2013-07341.pdf>

Routine uses of the information in this record may include disclosure to:

- 1) law enforcement, in the event that the system of records indicates a violation or potential violation of law;
- 2) a federal, state, local, or international agency, in response to its request;
- 3) a contractor of the USPTO having need for the information in order to perform a contract;
- 4) the Department of Justice for determination of whether the Freedom of Information Act (FOIA) requires disclosure of the record;
- 5) a Member of Congress submitting a request involving an individual to whom the record pertains, when the individual has requested the Member's assistance with respect to the subject matter of the record;
- 6) a court, magistrate, or administrative tribunal, in the course of presenting evidence, including disclosures to opposing counsel in the course of settlement negotiations;
- 7) the Administrator, General Services Administration (GSA), or their designee, during an inspection of records conducted by GSA under authority of 44 U.S.C. 2904 and 2906, in accordance with the GSA regulations and any other relevant (i.e., GSA or Commerce) directive, where such disclosure shall not be used to make determinations about individuals;
- 8) another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c));
- 9) the Office of Personnel Management (OPM) for personnel research purposes; and
- 10) the Office of Management and Budget (OMB) for legislative coordination and clearance.

If you do not furnish the information requested on this form, the USPTO may not be able to process and/or examine your submission, which may result in termination of proceedings, abandonment of the application, and/or expiration of the patent.



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC ACKNOWLEDGEMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 01/31/2025 07:18:35 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Colleen Mensching
PATENT CENTER #	69010935	FILING DATE	-
CUSTOMER #	21971	FIRST NAMED INVENTOR	Alexis HOWERTON
CORRESPONDENCE ADDRESS	-	AUTHORIZED BY	Victoria Cheng

Documents

TOTAL DOCUMENTS: 6

DOCUMENT	PAGES	DESCRIPTION	SIZE (KB)
Spruce Biosciences Inc. 50535-710.306 - ADS.pdf	9	Application Data Sheet	2180 KB
Spruce Biosciences Inc. 50535-710.306 - Track 1 Request.pdf	2	Track One Request	174 KB
Spruce Biosciences Inc. 50535-710.306 - Specification.pdf	116	-	1638 KB
Spruce Biosciences Inc. 50535-710.306 - Specification-SPEC.pdf	(1-112) 112	Specification	1663 KB
Spruce Biosciences Inc. 50535-710.306 - Specification-CLM.pdf	(113-115) 3	Claims	140 KB

Spruce Biosciences Inc. 50535-710.306 - Specification-ABST.pdf	(116- 116)	1	Abstract	87 KB
Spruce Biosciences Inc. 50535-710.306 - Figures.pdf		14	Drawings-other than black and white line drawings	1089 KB

Digest

DOCUMENT	MESSAGE DIGEST(SHA-512)
Spruce Biosciences Inc. 50535-710.306 - ADS.pdf	FD697774AF7606E8E8CDF02FEB56CA5F898D56A8356E15745922F7406A6718BA86736EC293C087EEAEAE6C873F60F6BE11233F7A269CACE66B572FD4BBA1D44E
Spruce Biosciences Inc. 50535-710.306 - Track 1 Request.pdf	F590884EC8F63664AAF363574AA2CC95573007AEBDC3F891673325DA36303E6FC55764F2457868A5AE0DEAC68FE2BD7E8A732E6ED252414D60078C78D4F2D9A9
Spruce Biosciences Inc. 50535-710.306 - Specification.pdf	B2AADBCB431DB71E15B34A216EA2D7D608EAC813B0182F7CE5932D4419DB4301CE7F6AF02D3C9FF1A1BCC442445D8BBE62C6E90CF4A539F3A42F02418510B5F9
Spruce Biosciences Inc. 50535-710.306 - Specification-SPEC.pdf	B0CDF5F980B0B5BFF4605288488A027ECDDE1D854EDC94D2CE8E1948B67F8734184DDCBFD6B4FB9C7B9C05A3137ED44CB221101A6BC1A4FBFDAAA76402EAC5F0
Spruce Biosciences Inc. 50535-710.306 - Specification-CLM.pdf	40B1E72637B76D62086EC159AA237D12B0BEB28D9B2A0C4EA311D7BC714E0F13F50A40DD7A32AF0CC9E860FD2387BE9AA049F427ADFDC76DF12E6F0D4D053D59
Spruce Biosciences Inc. 50535-710.306 - Specification-ABST.pdf	4870AE468D7B983C7CFD5C4D7E10C01E1BFE15206984B6D1D9E8C86DF9C80F73E32EAA7084797F96CA5919D2AB84D13955F2F9F9FAE9EC70DE17F4BF0D3E832D
Spruce Biosciences Inc. 50535-710.306 - Figures.pdf	7037A1FF19D8EB20A63B100B88165C9505B47B42AAAD9BFEBEF3F5C07BD33F054652182B37ED1DF5BFB495E110C70289B6B01374FEB0059FC480C4439501C77

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

CROSS-REFERENCE

[0001] This application is a continuation of U.S. Application No. 18/941,832, filed November 8, 2024, which is a continuation of U.S. Application No. 17/667,285, filed on February 8, 2022, which is a continuation of U.S. Application No. 17/081,694 filed on October 27, 2020, which issued as U.S. Pat. No. 11,304,950 on April 19, 2022, and which is a continuation of International Application No. PCT/US2019/029486 filed on April 26, 2019, which claims the benefit of U.S. Provisional Patent Application No. 62/663,951, filed April 27, 2018, and U.S. Provisional Patent Application No. 62/822,815, filed March 23, 2019, each of which is entirely incorporated herein by reference.

BACKGROUND

[0002] Corticotropes are basophilic cells in the anterior pituitary that produce melanocyte-stimulating hormone, adrenocorticotrophic hormone (ACTH) and lipotropin. The cells produce pro-opiomelanocortin (POMC) which undergoes cleavage to ACTH and β -lipotropin (β -LPH). These cells respond to corticotropin-releasing factor (CRF) and make up about 20% of the cells in the anterior pituitary. CRF is a releasing hormone that belongs to the corticotropin-releasing factor family and is the primary physiological regulator of proopiomelanocortin (POMC).

[0003] CRF is produced by parvocellular neuroendocrine cells within the paraventricular nucleus of the hypothalamus and is released at the median eminence from neurosecretory terminals of these neurons into the primary capillary plexus of the hypothalamo-hypophyseal portal system. The portal system carries the CRF to the anterior lobe of the pituitary, where it stimulates corticotropes to secrete ACTH and other biologically-active substances. ACTH, in turn, stimulates the synthesis of cortisol, glucocorticoids, mineralocorticoids and DHEA.

[0004] Testicular adrenal rest tumors (TART) and ovarian adrenal rest tumors (OART) are ACTH-responsive lesions of the testes and ovaries derived from adrenal tissue interchelated within these organs during embryogenesis. In response to high ACTH, hyperplasia of this tissue occurs, resulting in single or multiple lesions that may cause pain and infertility. The identification of compounds that modulate CRF function and downstream processes is an ongoing challenge. Currently, there are no non-steroidal, non-surgical solutions to control, shrink, or reduce TART. Disclosed herein, *inter alia*, are solutions to these and other problems in the art.

SUMMARY


[0005] Congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency is a serious genetic disorder characterized by impaired adrenal synthesis of cortisol and consequent overproduction of adrenocorticotrophic hormone (ACTH), 17-hydroxyprogesterone (17OHP), and adrenal androgens such as androstenedione (A4). CAH is an orphan condition with no FDA-approved therapy. To suppress androgen production, patients are often prescribed long-term supraphysiologic doses of glucocorticoid (GC), which has adverse effects and fail to suppress 17OHP and A4 in the majority of patients.

[0006] Compound 1 may be a potent, selective, nonsteroidal, oral corticotropin-releasing factor type-1 (CRF1) receptor antagonist that reduces excess ACTH, 17OHP, and A4 to improve clinical and biochemical sequelae of CAH. This may obviate the need for high dose steroids.

[0007] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0008] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0009] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):




wherein:

R¹ and R² are independently ethyl or n-propyl;

R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R⁴ is hydrogen, Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,




R^a and R^b are independently hydrogen, C₁-C₃alkyl, H₂NCH₂CH₂-, (CH₃)₃COC(O)NHCH₂CH₂-, or CH₃CH₂CH₂NHCH₂CH₂-.


[0010] In some embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl. In some embodiments, R^3 is Cl, Br, or methyl. In some embodiments, R^4 is Br, R^aR^bN- , pyridin-4-yl, morpholin-4-yl,




[0011] In some embodiments, R^4 is hydrogen, Br, R^aR^bN- and R^a and R^b are independently C_1 - C_3 alkyl. In some embodiments, the compound is



[0012] In some embodiments, the compound is



[0013] In some embodiments, the compound is



[0014] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0015] In some embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically


acceptable salt thereof, is administered to the subject. In some embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

{0016} In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 5 mg/day to about 25 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 10 mg/day to about 400 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day is administered to the subject.

{0017} In some embodiments, the compound is administered orally. In some embodiments, the compound is formulated as a capsule or tablet. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state.

{0018} In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

{0019} One aspect described herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I) and a pharmaceutically acceptable excipient:




(I), or a pharmaceutically acceptable salt thereof,

wherein:

R^1 and R^2 are independently ethyl or n-propyl;


R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R^4 is hydrogen, Br, $R^a R^b N-$, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,



R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$.

[0020] In some embodiments, the compound of Formula (I) has the structure:



, or a pharmaceutically acceptable salt thereof.

[0021] In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0022] In some embodiments, the pharmaceutical composition comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265

mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.


[0023] In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0024] In some embodiments, the pharmaceutical composition is administered orally. In some embodiments, the pharmaceutical composition is formulated as capsule or a tablet. In some embodiments, the pharmaceutical composition is administered in the fed state. In some embodiments, the pharmaceutical composition is administered in the fasted state. In some embodiments, the pharmaceutical composition is administered at least once per day.

[0025] In some embodiments, the pharmaceutical composition is administered at bedtime. In some embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 2 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep. In some embodiments, the pharmaceutical composition is administered in the evening. In some embodiments, the pharmaceutical composition is administered at about 10 pm at night.


[0026] In some embodiments, the method further comprises administering to the subject an additional chemotherapeutic agent. In some embodiments, the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In some embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In some embodiments, the mineralocorticoid is fludrocortisone.

[0027] In some embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist. In some embodiments, the CRF₁ antagonist is



pharmaceutically acceptable salt thereof.

[0028] In some embodiments, the CRF₁ antagonist is



[0029] In some embodiments, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof.

[0030] In some embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof.

[0031] In some embodiments, the surgical resection is prior to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is concurrent with

administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0032] In some embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In some embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0033] In some embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the radiation therapy is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0034] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure.

Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0035] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0037] FIG. 1A-1B. A schematic representation of intra-patient dose escalation study to evaluate safety, pharmacokinetics, and pharmacodynamics **(A)**. A schematic representation of inhibition of CRF₁ with a CRF₁ antagonist (Compound 1) **(B)**.

[0038] FIG. 2. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0039] FIG. 3. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0040] FIG. 4. Graphic representation of patient-level response (measuring ACTH levels) to a CRF₁ antagonist (Compound 1).

[0041] FIG. 5A-5B. Graphic representation of patient-level response of sample responders (measuring ACTH levels) to a CRF₁ antagonist [Subject 003001 **(A)**; Subject 007002 **(B)**].

[0042] FIG. 6. Graphic representation of patient-level response (measuring 17-OHP levels) to a CRF₁ antagonist (Compound 1).

[0043] FIG. 7A-7B. Graphic representation of patient-level response of sample responders (measuring 17-OHP levels) to a CRF₁ antagonist [Subject 003001 **(A)**; Subject 007002 **(B)**].

[0044] FIG. 8. Graphic representation of patient-level response (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1).

[0045] FIG. 9A-9B. Graphic representation of patient-level response of sample responders (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1) [Subject 003001 **(A)**; Subject 007002 **(B)**].

[0046] FIG. 10A-10B. Images of a patient with TART pre-treatment with a CRF₁ antagonist (Compound 1) **(A)** and 6-weeks post-treatment with a CRF₁ antagonist (Compound 1) **(B)**.

[0047] FIG. 11A-11B. Graphic representation of an *in vitro* binding assay of a CRF₁ antagonist (Compound 1); ¹²⁵I-sauvagine binding to rat cerebellar membranes one hour after oral administration of increasing Compound 1 doses **(A)** and *in vivo* inhibition of CRF-induced activity after treatment with a CRF₁ antagonist; Compound 1 (oral, 10 mg/kg) **(B)**.

[0048] FIG. 12A-12B. Graphic representation of ACTH levels **(A)** and 17-OHP levels **(B)** after treatment with a CRF₁ antagonist (Compound 1).

[0049] FIG. 13. Graphic representation of hormone responses after treatment with a CRF₁ antagonist (Compound 1).

[0050] FIG. 14 depicts the proof of concept and escalating QD dosing at 10 pm.

[0051] FIG. 15A-15D are graphic representations of changes in androstenedione seen in participants in Cohort A and Cohort B.

[0052] FIG. 16A-16D are graphic representations of changes in 17-hydroxyprogesterone seen in participants in Cohort A and Cohort B.

[0053] FIG. 17A-17D are graphic representations of changes in adrenocorticotrophic hormone seen in participants in Cohort A and Cohort B.

DETAILED DESCRIPTION

[0054] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

I. Definitions

[0055] The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are constructed according to the standard rules of chemical valency known in the chemical arts.

[0056] Where substituent groups are specified by their conventional chemical formulae, written from left to right, they equally encompass the chemically identical substituents that would result from writing the structure from right to left, e.g., $-\text{CH}_2\text{O}-$ is equivalent to $-\text{OCH}_2-$.

[0057] The term “alkyl,” by itself or as part of another substituent, means, unless otherwise stated, a straight (i.e., unbranched) or branched carbon chain (or carbon), or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include mono-, di- and multivalent radicals, having the number of carbon atoms designated (i.e., $\text{C}_1\text{-C}_{10}$ means one to ten carbons). Alkyl is an uncyclized chain. Examples of saturated hydrocarbon radicals include, but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, (cyclohexyl)methyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. An alkoxy is an alkyl attached to the remainder of the molecule via an oxygen linker ($-\text{O}-$).

[0058] The term “alkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyl, as exemplified, but not limited by, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred herein. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene group, generally having eight or fewer

carbon atoms. The term “alkenylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkene.

[0059] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or combinations thereof, including at least one carbon atom and at least one heteroatom (e.g., O, N, P, Si, and S), and wherein the nitrogen and sulfur atoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) (e.g., N, S, Si, or P) may be placed at any interior position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Heteroalkyl is an uncyclized chain. Examples include, but are not limited to: -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃, -O-CH₃, -O-CH₂-CH₃, and -CN. Up to two or three heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. A heteroalkyl moiety may include one heteroatom (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include two optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include three optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include four optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include five optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include up to 8 optionally different heteroatoms (e.g., O, N, S, Si, or P).

[0060] Similarly, the term “heteroalkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from heteroalkyl, as exemplified, but not limited by, -CH₂-CH₂-S-CH₂-CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (e.g., alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied by the direction in which the formula of the linking group is written. For example, the formula -C(O)₂R'- represents both -C(O)₂R'- and -R'C(O)₂-. As described above, heteroalkyl groups, as used herein, include those groups that are attached to the remainder of the molecule through a heteroatom, such as -C(O)R', -C(O)NR', -NR'R'', -OR', -SR', and/or -SO₂R'. Where “heteroalkyl” is recited, followed by recitations of specific heteroalkyl groups, such as -NR'R'' or the like, it will be understood that the terms heteroalkyl and -NR'R'' are not redundant or mutually exclusive. Rather, the specific heteroalkyl groups are recited to add clarity. Thus, the term “heteroalkyl” should not be interpreted herein as excluding specific heteroalkyl groups, such as -NR'R'' or the like.

[0061] The terms “cycloalkyl” and “heterocycloalkyl,” by themselves or in combination with other terms, mean, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl,”

respectively. Cycloalkyl and heterocycloalkyl are not aromatic. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like. A “cycloalkylene” and a “heterocycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a cycloalkyl and heterocycloalkyl, respectively. “Cycloalkyl” is also meant to refer to bicyclic and polycyclic hydrocarbon rings such as, for example, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, etc.

[0062] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl” are meant to include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” includes, but is not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0063] The term “acyl” means, unless otherwise stated, -C(O)R where R is a substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

[0064] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent, which can be a single ring or multiple rings (preferably from 1 to 3 rings) that are fused together (i.e., a fused ring aryl) or linked covalently. A fused ring aryl refers to multiple rings fused together wherein at least one of the fused rings is an aryl ring. The term “heteroaryl” refers to aryl groups (or rings) that contain at least one heteroatom such as N, O, or S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. Thus, the term “heteroaryl” includes fused ring heteroaryl groups (i.e., multiple rings fused together wherein at least one of the fused rings is a heteroaromatic ring). A 5,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 5 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. Likewise, a 6,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. And a 6,5-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 5 members, and wherein at least one ring is a heteroaryl ring. A heteroaryl group can be attached to the remainder of the molecule through a carbon or heteroatom. Non-


limiting examples of aryl and heteroaryl groups include phenyl, naphthyl, pyrrolyl, pyrazolyl, pyridazinyl, triazinyl, pyrimidinyl, imidazolyl, pyrazinyl, purinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, benzofuran, isobenzofuranyl, indolyl, isoindolyl, benzothiophenyl, isoquinolyl, quinoxalyl, quinolyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below. An “arylene” and a “heteroarylene,” alone or as part of another substituent, mean a divalent radical derived from an aryl and heteroaryl, respectively. A heteroaryl group substituent may be -O- bonded to a ring heteroatom nitrogen.

[0065] Spirocyclic rings are two or more rings wherein adjacent rings are attached through a single atom. The individual rings within spirocyclic rings may be identical or different. Individual rings in spirocyclic rings may be substituted or unsubstituted and may have different substituents from other individual rings within a set of spirocyclic rings. Possible substituents for individual rings within spirocyclic rings are the possible substituents for the same ring when not part of spirocyclic rings (e.g. substituents for cycloalkyl or heterocycloalkyl rings). Spirocyclic rings may be substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkylene, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heterocycloalkylene and individual rings within a spirocyclic ring group may be any of the immediately previous list, including having all rings of one type (e.g. all rings being substituted heterocycloalkylene wherein each ring may be the same or different substituted heterocycloalkylene). When referring to a spirocyclic ring system, heterocyclic spirocyclic rings means a spirocyclic rings wherein at least one ring is a heterocyclic ring and wherein each ring may be a different ring. When referring to a spirocyclic ring system, substituted spirocyclic rings means that at least one ring is substituted and each substituent may optionally be different.

[0066] The symbol “” denotes the point of attachment of a chemical moiety to the remainder of a molecule or chemical formula.

[0067] The term “oxo,” as used herein, means an oxygen that is double bonded to a carbon atom.

[0068] The term “alkylarylene” as an arylene moiety covalently bonded to an alkylene moiety (also referred to herein as an alkylene linker). In embodiments, the alkylarylene group has the formula:



[0069] An alkylarylene moiety may be substituted (e.g. with a substituent group) on the alkylene moiety or the arylene linker (e.g. at carbons 2, 3, 4, or 6) with halogen, oxo, $-N_3$, $-CF_3$, $-CCl_3$, $-CBr_3$, $-Cl_3$, $-CN$, $-CHO$, $-OH$, $-NH_2$, $-COOH$, $-CONH_2$, $-NO_2$, $-SH$, $-SO_2CH_3$, $-SO_3H$, $-OSO_3H$, $-SO_2NH_2$, $-NHNH_2$, $-ONH_2$, $-NHC(O)NHNH_2$, substituted or unsubstituted C_1 - C_5 alkyl or substituted or unsubstituted 2 to 5 membered heteroalkyl). In embodiments, the alkylarylene is unsubstituted.

[0070] Each of the above terms (e.g., “alkyl,” “heteroalkyl,” “cycloalkyl,” “heterocycloalkyl,” “aryl,” and “heteroaryl”) includes both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

[0071] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be one or more of a variety of groups selected from, but not limited to, $-OR'$, $=O$, $=NR'$, $=N-OR'$, $-NR'R''$, $-SR'$, $-halogen$, $-SiR'R''R'''$, $-OC(O)R'$, $-C(O)R'$, $-CO_2R'$, $-CONR'R''$, $-OC(O)NR'R''$, $-NR''C(O)R'$, $-NR'-C(O)NR''R'''$, $-NR''C(O)_2R'$, $-NR-C(NR'R''R''')=NR'''$, $-NR-C(NR'R'')=NR'''$, $-S(O)R'$, $-S(O)_2R'$, $-S(O)_2NR'R''$, $-NRSO_2R'$, $-NR'NR''R'''$, $-ONR'R''$, $-NR'C(O)NR''NR'''R''''$, $-CN$, $-NO_2$, $-NR'SO_2R''$, $-NR'C(O)R''$, $-NR'C(O)-OR''$, $-NR'OR''$, in a number ranging from zero to $(2m+1)$, where m is the total number of carbon atoms in such radical. R , R' , R'' , R''' , and R'''' each preferably independently refer to hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl (e.g., aryl substituted with 1-3 halogens), substituted or unsubstituted heteroaryl, substituted or unsubstituted alkyl, alkoxy, or thioalkoxy groups, or arylalkyl groups. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R' , R'' , R''' , and R'''' group when more than one of these groups is present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 4-, 5-, 6-, or 7-membered ring. For example, $-NR'R''$ includes, but is not limited to, 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term “alkyl” is meant to include groups including carbon atoms bound to groups other than hydrogen groups, such as haloalkyl (e.g., $-CF_3$ and $-CH_2CF_3$) and acyl (e.g., $-C(O)CH_3$, $-C(O)CF_3$, $-C(O)CH_2OCH_3$, and the like).

[0072] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups are varied and are selected from, for example: $-OR'$, $-NR'R''$, $-SR'$, $-halogen$, $-$

SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R''R''')=NR''', -NR-C(NR'R'')=NR'', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -NR'NR''R''', -ONR'R'', -NR'C(O)NR''NR''R''', -CN, -NO₂, -R', -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, -NR'SO₂R'', -NR'C(O)R'', -NR'C(O)-OR'', -NR'OR'', in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'', R''', and R'''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', and R'''' groups when more than one of these groups is present.

[0073] Substituents for rings (e.g. cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylene, heterocycloalkylene, arylene, or heteroarylene) may be depicted as substituents on the ring rather than on a specific atom of a ring (commonly referred to as a floating substituent). In such a case, the substituent may be attached to any of the ring atoms (obeying the rules of chemical valency) and in the case of fused rings or spirocyclic rings, a substituent depicted as associated with one member of the fused rings or spirocyclic rings (a floating substituent on a single ring), may be a substituent on any of the fused rings or spirocyclic rings (a floating substituent on multiple rings). When a substituent is attached to a ring, but not a specific atom (a floating substituent), and a subscript for the substituent is an integer greater than one, the multiple substituents may be on the same atom, same ring, different atoms, different fused rings, different spirocyclic rings, and each substituent may optionally be different. Where a point of attachment of a ring to the remainder of a molecule is not limited to a single atom (a floating substituent), the attachment point may be any atom of the ring and in the case of a fused ring or spirocyclic ring, any atom of any of the fused rings or spirocyclic rings while obeying the rules of chemical valency. Where a ring, fused rings, or spirocyclic rings contain one or more ring heteroatoms and the ring, fused rings, or spirocyclic rings are shown with one more floating substituents (including, but not limited to, points of attachment to the remainder of the molecule), the floating substituents may be bonded to the heteroatoms. Where the ring heteroatoms are shown bound to one or more hydrogens (e.g. a ring nitrogen with two bonds to ring atoms and a third bond to a hydrogen) in the structure or formula with the floating substituent, when the heteroatom is bonded to the floating substituent, the substituent will be understood to replace the hydrogen, while obeying the rules of chemical valency.

[0074] Two or more substituents may optionally be joined to form aryl, heteroaryl, cycloalkyl, or heterocycloalkyl groups. Such so-called ring-forming substituents are typically, though not

necessarily, found attached to a cyclic base structure. In one embodiment, the ring-forming substituents are attached to adjacent members of the base structure. For example, two ring-forming substituents attached to adjacent members of a cyclic base structure create a fused ring structure. In another embodiment, the ring-forming substituents are attached to a single member of the base structure. For example, two ring-forming substituents attached to a single member of a cyclic base structure create a spirocyclic structure. In yet another embodiment, the ring-forming substituents are attached to non-adjacent members of the base structure.

[0075] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally form a ring of the formula $-T-C(O)-(CRR')_q-U-$, wherein T and U are independently $-NR'$ -, $-O-$ -, $-CRR'$ -, or a single bond, and q is an integer of from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-A-(CH_2)_r-B-$, wherein A and B are independently $-CRR'$ -, $-O-$ -, $-NR'$ -, $-S-$ -, $-S(O)-$ -, $-S(O)_2-$ -, $-S(O)_2NR'$ -, or a single bond, and r is an integer of from 1 to 4. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-(CRR')_s-X'-(C''R''R''')_d-$, where s and d are independently integers of from 0 to 3, and X' is $-O-$ -, $-NR'$ -, $-S-$ -, $-S(O)-$ -, $-S(O)_2-$ -, or $-S(O)_2NR'$ -. The substituents R, R', R'', and R''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

[0076] As used herein, the terms “heteroatom” or “ring heteroatom” are meant to include oxygen (O), nitrogen (N), sulfur (S), phosphorus (P), and silicon (Si).

[0077] A “substituent group,” as used herein, means a group selected from the following moieties:

(A) oxo, halogen, $-CF_3$ -, $-CN$ -, $-OH$ -, $-NH_2$ -, $-COOH$ -, $-CONH_2$ -, $-NO_2$ -, $-SH$ -, $-SO_3H$ -, $-SO_4H$ -, $-SO_2NH_2$ -, $-NHNH_2$ -, $-ONH_2$ -, $-NHC(=O)NHNH_2$ -, $-NHC(=O)NH_2$ -, $-NHSO_2H$ -, $-NHC(=O)H$ -, $-NHC(O)-OH$ -, $-NHOH$ -, $-OCF_3$ -, $-OCHF_2$ -, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(B) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(i) oxo, halogen, $-CF_3$ -, $-CN$ -, $-OH$ -, $-NH_2$ -, $-COOH$ -, $-CONH_2$ -, $-NO_2$ -, $-SH$ -, $-SO_3H$ -, $-SO_4H$ -, $-SO_2NH_2$ -, $-NHNH_2$ -, $-ONH_2$ -, $-NHC(=O)NHNH_2$ -, $-NHC(=O)NH_2$ -, $-NHSO_2H$ -, $-NHC(=O)H$ -, $-NHC(O)-OH$ -, $-NHOH$ -, $-OCF_3$ -, $-OCHF_2$ -, unsubstituted alkyl, unsubstituted heteroalkyl,

unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(ii) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(a) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC=(O)NHNH₂, -NHC=(O)NH₂, -NHSO₂H, -NHC=(O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(b) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from: oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC=(O)NHNH₂, -NHC=(O)NH₂, -NHSO₂H, -NHC=(O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl.

[0078] A “size-limited substituent” or “size-limited substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl.

[0079] A “lower substituent” or “lower substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl.

[0080] In some embodiments, each substituted group described in the compounds herein is substituted with at least one substituent group. More specifically, in some embodiments, each substituted alkyl, substituted heteroalkyl, substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl, substituted heteroaryl, substituted alkylene, substituted heteroalkylene, substituted cycloalkylene, substituted heterocycloalkylene, substituted arylene, and/or substituted heteroarylene described in the compounds herein are substituted with at least one substituent group. In other embodiments, at least one or all of these groups are substituted with at least one size-limited substituent group. In other embodiments, at least one or all of these groups are substituted with at least one lower substituent group.

[0081] In other embodiments of the compounds herein, each substituted or unsubstituted alkyl may be a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl. In some embodiments of the compounds herein, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₂₀ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 20 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₈ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a substituted or unsubstituted 3 to 8 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 10 membered heteroarylene.

[0082] In some embodiments, each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl. In some embodiments, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₈ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 8 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₇ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a

substituted or unsubstituted 3 to 7 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 9 membered heteroarylene. In some embodiments, the compound is a chemical species set forth in the Examples section, figures, or tables below.

[0083] Certain compounds of the present invention possess asymmetric carbon atoms (optical or chiral centers) or double bonds; the enantiomers, racemates, diastereomers, tautomers, geometric isomers, stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)- or, as (D)- or (L)- for amino acids, and individual isomers are encompassed within the scope of the present invention. The compounds of the present invention do not include those that are known in art to be too unstable to synthesize and/or isolate. The present invention is meant to include compounds in racemic and optically pure forms. Optically active (R)- and (S)-, or (D)- and (L)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers.

[0084] As used herein, the term “isomers” refers to compounds having the same number and kind of atoms, and hence the same molecular weight, but differing in respect to the structural arrangement or configuration of the atoms.

[0085] The term “tautomer,” as used herein, refers to one of two or more structural isomers which exist in equilibrium and which are readily converted from one isomeric form to another.

[0086] It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention.

[0087] Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention.

[0088] Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention.

[0089] The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the

compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I), or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are encompassed within the scope of the present invention.

[0090] It should be noted that throughout the application that alternatives are written in Markush groups, for example, each amino acid position that contains more than one possible amino acid. It is specifically contemplated that each member of the Markush group should be considered separately, thereby comprising another embodiment, and the Markush group is not to be read as a single unit.

[0091] “Analog” or “analogue” is used in accordance with its plain ordinary meaning within Chemistry and Biology and refers to a chemical compound that is structurally similar to another compound (i.e., a so-called “reference” compound) but differs in composition, e.g., in the replacement of one atom by an atom of a different element, or in the presence of a particular functional group, or the replacement of one functional group by another functional group, or the absolute stereochemistry of one or more chiral centers of the reference compound. Accordingly, an analog is a compound that is similar or comparable in function and appearance but not in structure or origin to a reference compound.

[0092] The terms “a” or “an,” as used in herein means one or more. In addition, the phrase “substituted with a[n],” as used herein, means the specified group may be substituted with one or more of any or all of the named substituents. For example, where a group, such as an alkyl or heteroaryl group, is “substituted with an unsubstituted $\text{C}_1\text{-C}_{20}$ alkyl, or unsubstituted 2 to 20 membered heteroalkyl,” the group may contain one or more unsubstituted $\text{C}_1\text{-C}_{20}$ alkyls, and/or one or more unsubstituted 2 to 20 membered heteroalkyls.

[0093] Moreover, where a moiety is substituted with an R substituent, the group may be referred to as “R-substituted.” Where a moiety is R-substituted, the moiety is substituted with at least one R substituent and each R substituent is optionally different. Where a particular R group is present in the description of a chemical genus (such as Formula (I)), a Roman alphabetic symbol may be used to distinguish each appearance of that particular R group. For example, where multiple R^{13} substituents are present, each R^{13} substituent may be distinguished as $\text{R}^{13\text{A}}$, $\text{R}^{13\text{B}}$, $\text{R}^{13\text{C}}$, $\text{R}^{13\text{D}}$, etc., wherein each of $\text{R}^{13\text{A}}$, $\text{R}^{13\text{B}}$, $\text{R}^{13\text{C}}$, $\text{R}^{13\text{D}}$, etc. is defined within the scope of the definition of R^{13} and optionally differently.

[0094] A “detectable moiety” as used herein refers to a moiety that can be covalently or noncovalently attached to a compound or biomolecule that can be detected for instance, using techniques known in the art. In embodiments, the detectable moiety is covalently attached. The detectable moiety may provide for imaging of the attached compound or biomolecule. The detectable moiety may indicate the contacting between two compounds. Exemplary detectable

moieties are fluorophores, antibodies, reactive dies, radio-labeled moieties, magnetic contrast agents, and quantum dots. Exemplary fluorophores include fluorescein, rhodamine, GFP, coumarin, FITC, Alexa fluor, Cy3, Cy5, BODIPY, and cyanine dyes. Exemplary radionuclides include Fluorine-18, Gallium-68, and Copper-64. Exemplary magnetic contrast agents include gadolinium, iron oxide and iron platinum, and manganese.

[0095] Descriptions of compounds of the present invention are limited by principles of chemical bonding known to those skilled in the art. Accordingly, where a group may be substituted by one or more of a number of substituents, such substitutions are selected so as to comply with principles of chemical bonding and to give compounds which are not inherently unstable and/or would be known to one of ordinary skill in the art as likely to be unstable under ambient conditions, such as aqueous, neutral, and several known physiological conditions. For example, a heterocycloalkyl or heteroaryl is attached to the remainder of the molecule via a ring heteroatom in compliance with principles of chemical bonding known to those skilled in the art thereby avoiding inherently unstable compounds.

[0096] The term “pharmaceutically acceptable salts” is meant to include salts of the active compounds that are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, oxalic, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see*, for example, Berge *et al.*, “Pharmaceutical Salts”, *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0097] Thus, the compounds of the present invention may exist as salts, such as with pharmaceutically acceptable acids. The present invention includes such salts. Non-limiting examples of such salts include hydrochlorides, hydrobromides, phosphates, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, propionates, tartrates (e.g., (+)-tartrates, (-)-tartrates, or mixtures thereof including racemic mixtures), succinates, benzoates, and salts with amino acids such as glutamic acid, and quaternary ammonium salts (e.g. methyl iodide, ethyl iodide, and the like). These salts may be prepared by methods known to those skilled in the art.

[0098] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound may differ from the various salt forms in certain physical properties, such as solubility in polar solvents. In embodiments, compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts. The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in a conventional manner. The parent form of the compounds differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but, unless specifically indicated, the salts disclosed herein are equivalent to the parent form of the compound for the purposes of the present invention.

[0099] In addition to salt forms, the present invention provides compounds, which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Prodrugs of the compounds described herein may be converted *in vivo* after administration. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment, such as, for example, when contacted with a suitable enzyme or chemical reagent.

[0100] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0101] “Pharmaceutically acceptable excipient” and “pharmaceutically acceptable carrier” refer to a substance that aids the administration of a compound to and absorption by a subject and can be included in the compositions of the present invention without causing a significant adverse toxicological effect on the patient. Non-limiting examples of pharmaceutically acceptable

excipients include water, NaCl, normal saline solutions, lactated Ringer's, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer's solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with the compounds of the invention. One of skill in the art will recognize that other pharmaceutical excipients are useful in the present invention.

[0102] The term “preparation” is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

[0103] A “CRF1 inhibitor” refers to a compound (e.g., compounds described herein) that reduces the activity of CRF1 when compared to a control, such as absence of the compound or a compound with known inactivity.

[0104] The terms “polypeptide,” “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues, wherein the polymer may optionally be conjugated to a moiety that does not consist of amino acids. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. In embodiments, the terms “polypeptide,” “peptide,” and “protein”, used interchangeably herein, refer to a polymeric form of amino acids of any length, which can include genetically coded and non-genetically coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified polypeptide backbones. The terms include fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence; fusion proteins with heterologous and homologous leader sequences, with or without N-terminus methionine residues; immunologically tagged proteins; and the like.

[0105] A polypeptide, or a cell is “recombinant” when it is artificial or engineered, or derived from or contains an artificial or engineered protein or nucleic acid (*e.g.* non-natural or not wild type). For example, a polynucleotide that is inserted into a vector or any other heterologous location, *e.g.*, in a genome of a recombinant organism, such that it is not associated with nucleotide sequences that normally flank the polynucleotide as it is found in nature is a

recombinant polynucleotide. A protein expressed *in vitro* or *in vivo* from a recombinant polynucleotide is an example of a recombinant polypeptide. Likewise, a polynucleotide sequence that does not appear in nature, for example a variant of a naturally occurring gene, is recombinant.

[0106] “Contacting” is used in accordance with its plain ordinary meaning and refers to the process of allowing at least two distinct species (e.g. chemical compounds including biomolecules or cells) to become sufficiently proximal to react, interact or physically touch. It should be appreciated; however, the resulting reaction product can be produced directly from a reaction between the added reagents or from an intermediate from one or more of the added reagents that can be produced in the reaction mixture.

[0107] The term “contacting” may include allowing two species to react, interact, or physically touch, wherein the two species may be a compound as described herein and a protein or enzyme. In some embodiments contacting includes allowing a compound described herein to interact with a protein or enzyme that is involved in a signaling pathway (e.g., MAP kinase pathway).

[0108] As defined herein, the term “activation”, “activate”, “activating” and the like in reference to a protein refers to conversion of a protein into a biologically active derivative from an initial inactive or deactivated state. The terms reference activation, or activating, sensitizing, or up-regulating signal transduction or enzymatic activity or the amount of a protein decreased in a disease.

[0109] The terms “agonist,” “activator,” “upregulator,” etc. refer to a substance capable of detectably increasing the expression or activity of a given gene or protein. The agonist can increase expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the agonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or higher than the expression or activity in the absence of the agonist. In embodiments, an agonist is a molecule that interacts with a target to cause or promote an increase in the activation of the target. In embodiments, activators are molecules that increase, activate, facilitate, enhance activation, sensitize, or up-regulate, e.g., a gene, protein, ligand, receptor, or cell.

[0110] As defined herein, the term “inhibition”, “inhibit”, “inhibiting” and the like in reference to a protein-inhibitor interaction means negatively affecting (e.g. decreasing) the activity or function of the protein relative to the activity or function of the protein in the absence of the inhibitor. In embodiments inhibition means negatively affecting (e.g. decreasing) the concentration or levels of the protein relative to the concentration or level of the protein in the absence of the inhibitor. In embodiments inhibition refers to reduction of a disease or symptoms of disease. In embodiments, inhibition refers to a reduction in the activity of a particular protein

target. Thus, inhibition includes, at least in part, partially or totally blocking stimulation, decreasing, preventing, or delaying activation, or inactivating, desensitizing, or down-regulating signal transduction or enzymatic activity or the amount of a protein. In embodiments, inhibition refers to a reduction of activity of a target protein resulting from a direct interaction (e.g. an inhibitor binds to the target protein). In embodiments, inhibition refers to a reduction of activity of a target protein from an indirect interaction (e.g. an inhibitor binds to a protein that activates the target protein, thereby preventing target protein activation).

[0111] The terms “inhibitor,” “repressor” or “antagonist” or “downregulator” interchangeably refer to a substance capable of detectably decreasing the expression or activity of a given gene or protein. The antagonist can decrease expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the antagonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or lower than the expression or activity in the absence of the antagonist. An antagonist prevents, reduces, inhibits, or neutralizes the activity of an agonist, and an antagonist can also prevent, inhibit, or reduce constitutive activity of a target, e.g., a target receptor, even where there is no identified agonist. In embodiments, inhibitors are molecules that decrease, block, prevent, delay activation, inactivate, desensitize, or down-regulate, e.g., a gene, protein, ligand, receptor, or cell. An inhibitor may also be defined as a molecule that reduces, blocks, or inactivates a constitutive activity. An “antagonist” is a molecule that opposes the action(s) of an agonist.

[0112] The terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” refer to a protein (including homologs, isoforms, and functional fragments thereof) and is a receptor for corticotropin releasing-hormone. The terms include any recombinant or naturally-occurring form of CRF₁ variants thereof that maintain CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). The terms include any mutant form of CRF₁ variants (e.g., frameshift mutations) thereof that maintain CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). In embodiments, the CRF₁ protein encoded by the *CRHR1* gene has the amino acid sequence set forth in or corresponding to Entrez 1233, UniProt P34998.

[0113] The term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion. Expression can be detected using conventional techniques for detecting protein (e.g., ELISA, Western blotting, flow cytometry, immunofluorescence, immunohistochemistry, *etc.*).

[0114] The terms “disease” or “condition” refer to a state of being or health status of a patient or subject capable of being treated with the compounds or methods provided herein. The disease may be a malignant or benign tumor (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)). The disease may be cancer. In some further instances, “cancer” refers to human cancers and carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, etc., including solid and lymphoid cancers, kidney, breast, lung, bladder, colon, ovarian, prostate, pancreas, adrenal, stomach, brain, head and neck, skin, uterine, testicular, glioma, esophagus, and liver cancer, including hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin’s lymphomas (*e.g.*, Burkitt’s, Small Cell, and Large Cell lymphomas), Hodgkin’s lymphoma, leukemia (including MDS, AML, ALL, ATLL and CML), or multiple myeloma.

[0115] As used herein, the term “cancer” refers to all types of cancer, neoplasm or malignant tumors found in mammals (*e.g.* humans), including leukemia, carcinomas and sarcomas. Exemplary cancers that may be treated with a compound or method provided herein include adrenal cancer and tumors (*e.g.*, testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) brain cancer, glioma, glioblastoma, neuroblastoma, prostate cancer, colorectal cancer, pancreatic cancer, cervical cancer, gastric cancer, ovarian cancer, lung cancer, and cancer of the head. Exemplary cancers that may be treated with a compound or method provided herein include cancer of the thyroid, endocrine system, brain, breast, cervix, colon, head & neck, liver, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus, Medulloblastoma, colorectal cancer, pancreatic cancer. Additional examples include, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung adenocarcinoma, lung squamous cell carcinoma, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, glioma, glioblastoma multiforme, ovarian cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, primary brain tumors, cancer, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, endometrial cancer, adrenal cortical cancer, neoplasms of the endocrine or exocrine pancreas, medullary thyroid cancer, medullary thyroid carcinoma, melanoma, colorectal cancer, papillary thyroid cancer, hepatocellular carcinoma, or prostate cancer.

[0116] The term “congenital adrenal hyperplasia” or “CAH” are any of several autosomal recessive diseases resulting from mutations of genes for enzymes mediating the biochemical

steps of production of mineralocorticoids, glucocorticoids or sex steroids from cholesterol by the adrenal glands (steroidogenesis). These conditions involve excessive or deficient production of sex steroids and can alter development of primary or secondary sex characteristics in some affected infants, children, or adults.

[0117] “Adrenal rest tumor” refers to a neoplasm of the testis (or ovaries) consisting of adrenal tissue. It may be secondary to hyperplasia of an adrenal gland.

[0118] “Testicular adrenal rest tumor(s)” or “TART” and/or ovarian adrenal rest tumor(s) are a complication of congenital adrenal hyperplasia (CAH), which may develop from ectopic remnants of intra-testicular adrenal tissue stimulated by adrenocorticotrophic hormone (ACTH) hypersecretion. TART lesions are typically located within the rete testis and are bilateral, synchronous, nodular and multiple. TART can lead to testicular structural damage, spermatogenesis disorders, infertility and mass-forming lesions.

[0119] The term “leukemia” refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease-acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number abnormal cells in the blood-leukemic or aleukemic (subleukemic). Exemplary leukemias that may be treated with a compound or method provided herein include, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemetic leukemia, basophilic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, multiple myeloma, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, or undifferentiated cell leukemia.

[0120] The term “sarcoma” generally refers to a tumor which is made up of a substance like the embryonic connective tissue and is generally composed of closely packed cells embedded in a fibrillar or homogeneous substance. Sarcomas that may be treated with a compound or method

provided herein include a chondrosarcoma, fibrosarcoma, lymphosarcoma, melanosarcoma, myxosarcoma, osteosarcoma, Abemethy's sarcoma, adipose sarcoma, liposarcoma, alveolar soft part sarcoma, ameloblastic sarcoma, botryoid sarcoma, chloroma sarcoma, chorio carcinoma, embryonal sarcoma, Wilms' tumor sarcoma, endometrial sarcoma, stromal sarcoma, Ewing's sarcoma, fascial sarcoma, fibroblastic sarcoma, giant cell sarcoma, granulocytic sarcoma, Hodgkin's sarcoma, idiopathic multiple pigmented hemorrhagic sarcoma, immunoblastic sarcoma of B cells, lymphoma, immunoblastic sarcoma of T-cells, Jensen's sarcoma, Kaposi's sarcoma, Kupffer cell sarcoma, angiosarcoma, leukosarcoma, malignant mesenchymoma sarcoma, parosteal sarcoma, reticulocytic sarcoma, Rous sarcoma, serocystic sarcoma, synovial sarcoma, or telangiectatic sarcoma.

[0121] The term “melanoma” is taken to mean a tumor arising from the melanocytic system of the skin and other organs. Melanomas that may be treated with a compound or method provided herein include, for example, acral-lentiginous melanoma, amelanotic melanoma, benign juvenile melanoma, Cloudman's melanoma, S91 melanoma, Harding-Passey melanoma, juvenile melanoma, lentigo maligna melanoma, malignant melanoma, nodular melanoma, subungal melanoma, or superficial spreading melanoma.

[0122] The term “carcinoma” refers to a malignant new growth made up of epithelial cells tending to infiltrate the surrounding tissues and give rise to metastases. Exemplary carcinomas that may be treated with a compound or method provided herein include, for example, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung adenocarcinoma, lung squamous cell carcinoma, medullary thyroid carcinoma, familial medullary thyroid carcinoma, acinar carcinoma, acinous carcinoma, adenocystic carcinoma, adenoid cystic carcinoma, carcinoma adenomatosum, carcinoma of adrenal cortex, alveolar carcinoma, alveolar cell carcinoma, basal cell carcinoma, carcinoma basocellulare, basaloid carcinoma, basosquamous cell carcinoma, bronchioalveolar carcinoma, bronchiolar carcinoma, bronchogenic carcinoma, cerebriiform carcinoma, cholangiocellular carcinoma, chorionic carcinoma, colloid carcinoma, comedo carcinoma, corpus carcinoma, cribriform carcinoma, carcinoma en cuirasse, carcinoma cutaneum, cylindrical carcinoma, cylindrical cell carcinoma, duct carcinoma, carcinoma durum, embryonal carcinoma, encephaloid carcinoma, epiermoid carcinoma, carcinoma epitheliale adenoides, exophytic carcinoma, carcinoma ex ulcere, carcinoma fibrosum, gelatiniform carcinoma, gelatinous carcinoma, giant cell carcinoma, carcinoma gigantocellulare, glandular carcinoma, granulosa cell carcinoma, hair-matrix carcinoma, hematoid carcinoma, hepatocellular carcinoma, Hurthle cell carcinoma, hyaline carcinoma, hypernephroid carcinoma,

infantile embryonal carcinoma, carcinoma in situ, intraepidermal carcinoma, intraepithelial carcinoma, Krompecher's carcinoma, Kulchitzky-cell carcinoma, large-cell carcinoma, lenticular carcinoma, carcinoma lenticulare, lipomatous carcinoma, lymphoepithelial carcinoma, carcinoma medullare, medullary carcinoma, melanotic carcinoma, carcinoma molle, mucinous carcinoma, carcinoma muciparum, carcinoma mucocellulare, mucoepidermoid carcinoma, carcinoma mucosum, mucous carcinoma, carcinoma myxomatodes, nasopharyngeal carcinoma, oat cell carcinoma, carcinoma ossificans, osteoid carcinoma, papillary carcinoma, periportal carcinoma, preinvasive carcinoma, prickle cell carcinoma, pultaceous carcinoma, renal cell carcinoma of kidney, reserve cell carcinoma, carcinoma sarcomatodes, schneiderian carcinoma, scirrhus carcinoma, carcinoma scroti, signet-ring cell carcinoma, carcinoma simplex, small-cell carcinoma, solanoid carcinoma, spheroidal cell carcinoma, spindle cell carcinoma, carcinoma spongiosum, squamous carcinoma, squamous cell carcinoma, string carcinoma, carcinoma telangiectaticum, carcinoma telangiectodes, transitional cell carcinoma, carcinoma tuberosum, tuberos carcinoma, verrucous carcinoma, or carcinoma villosum.

[0123] The terms “treating” or “treatment” refer to any indicia of success in the therapy or amelioration of an injury, disease, pathology or condition, including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the injury, pathology or condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; improving a patient’s physical or mental well-being. The treatment or amelioration of symptoms can be based on objective or subjective parameters; including the results of a physical examination, neuropsychiatric exams, and/or a psychiatric evaluation. The term “treating” and conjugations thereof, may include prevention of an injury, pathology, condition, or disease. In embodiments, treating is preventing. In embodiments, treating does not include preventing.

[0124] “Treating” or “treatment” as used herein (and as well-understood in the art) also broadly includes any approach for obtaining beneficial or desired results in a subject’s condition, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of the extent of a disease, stabilizing (*i.e.*, not worsening) the state of disease, prevention of a disease’s transmission or spread, delay or slowing of disease progression, amelioration or palliation of the disease state, diminishment of the reoccurrence of disease, and remission, whether partial or total and whether detectable or undetectable. In other words, “treatment” as used herein includes any cure, amelioration, or prevention of a disease. Treatment may prevent the disease from occurring; inhibit the disease’s spread; relieve the disease’s symptoms (*e.g.*, ocular pain, seeing

halos around lights, red eye, very high intraocular pressure), fully or partially remove the disease's underlying cause, shorten a disease's duration, or do a combination of these things.

[0125] “Treating” and “treatment” as used herein include prophylactic treatment. Treatment methods include administering to a subject a therapeutically effective amount of a compound described herein. The administering step may consist of a single administration or may include a series of administrations. The length of the treatment period depends on a variety of factors, such as the severity of the condition, the age of the patient, the concentration of the compound, the activity of the compositions used in the treatment, or a combination thereof. It will also be appreciated that the effective dosage of an agent used for the treatment or prophylaxis may increase or decrease over the course of a particular treatment or prophylaxis regime. Changes in dosage may result and become apparent by standard diagnostic assays known in the art. In some instances, chronic administration may be required. For example, the compositions are administered to the subject in an amount and for a duration sufficient to treat the patient.

[0126] The term “prevent” refers to a decrease in the occurrence of disease symptoms in a patient. As indicated above, the prevention may be complete (no detectable symptoms) or partial, such that fewer symptoms are observed than would likely occur absent treatment. In embodiments, prevent refers to slowing the progression of the disease, disorder or condition or inhibiting progression thereof to a harmful or otherwise undesired state.

[0127] “Patient” or “subject in need thereof” refers to a living organism suffering from or prone to a disease or condition that can be treated by administration of a pharmaceutical composition as provided herein. Non-limiting examples include humans, other mammals, bovines, rats, mice, dogs, monkeys, goat, sheep, cows, deer, and other non-mammalian animals. In some embodiments, a patient is human.

[0128] An “effective amount” is an amount sufficient for a compound to accomplish a stated purpose relative to the absence of the compound (e.g., achieve the effect for which it is administered, treat a disease, reduce enzyme activity, increase enzyme activity, reduce a signaling pathway, or reduce one or more symptoms of a disease or condition). An example of an “effective amount” is an amount sufficient to contribute to the treatment, prevention, or reduction of a symptom or symptoms of a disease, which could also be referred to as a “therapeutically effective amount.” A “reduction” of a symptom or symptoms (and grammatical equivalents of this phrase) means decreasing of the severity or frequency of the symptom(s), or elimination of the symptom(s). A “prophylactically effective amount” of a drug is an amount of a drug that, when administered to a subject, will have the intended prophylactic effect, e.g., preventing or delaying the onset (or reoccurrence) of an injury, disease, pathology or condition, or reducing the likelihood of the onset (or reoccurrence) of an injury, disease, pathology, or

condition, or their symptoms. The full prophylactic effect does not necessarily occur by administration of one dose, and may occur only after administration of a series of doses. Thus, a prophylactically effective amount may be administered in one or more administrations. An “activity decreasing amount,” as used herein, refers to an amount of antagonist required to decrease the activity of an enzyme relative to the absence of the antagonist. A “function disrupting amount,” as used herein, refers to the amount of antagonist required to disrupt the function of an enzyme or protein relative to the absence of the antagonist. The exact amounts will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (*see, e.g.,* Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); and *Remington: The Science and Practice of Pharmacy*, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins). The therapeutically effective amount can be ascertained by measuring relevant physiological effects, and it can be adjusted in connection with the dosing regimen and diagnostic analysis of the subject’s condition, and the like. By way of example, measurement of the serum level of a CRF1 inhibitor (or, e.g., a metabolite thereof) at a particular time post-administration may be indicative of whether a therapeutically effective amount has been administered.

[0129] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0130] Therapeutically effective amounts for use in humans can also be determined from animal models. For example, a dose for humans can be formulated to achieve a concentration that has been found to be effective in animals. The dosage in humans can be adjusted by monitoring compounds effectiveness and adjusting the dosage upwards or downwards, as described above. Adjusting the dose to achieve maximal efficacy in humans based on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan. Adjusting the dose to achieve maximal therapeutic window efficacy or toxicity in humans based on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan.

[0131] The term “therapeutically effective amount,” as used herein, refers to that amount of the therapeutic agent sufficient to ameliorate the disorder, as described above. For example, for the given parameter, a therapeutically effective amount will show an increase or decrease of at least 5%, 10%, 15%, 20%, 25%, 40%, 50%, 60%, 75%, 80%, 90%, or at least 100%. Therapeutic efficacy can also be expressed as “-fold” increase or decrease. For example, a therapeutically

effective amount can have at least a 1.2-fold, 1.5-fold, 2-fold, 5-fold, or more effect over a control.

[0132] The terms “QD,” “qd,” and “q.d.” refer to once a day, i.e., a drug or therapeutic agent is administered once a day for the duration of the therapeutic regimen.

[0133] The terms “BD,” “bd,” and “b.i.d.” refer to twice a day, i.e., a drug or therapeutic agent is administered two times a day for the duration of the therapeutic regimen.

[0134] The terms “QID,” “qid,” and “q.i.d.” refer to three times a day, i.e., a drug or therapeutic agent is administered three times a day for the duration of the therapeutic regimen.

[0135] Dosages may be varied depending upon the requirements of the patient and the compound being employed. The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial therapeutic response in the patient over time. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects. Determination of the proper dosage for a particular situation is within the skill of the practitioner. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. Dosage amounts and intervals can be adjusted individually to provide levels of the administered compound effective for the particular clinical indication being treated. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease state.

[0136] As used herein, the term “administering” means oral administration, administration as a suppository, topical contact, intravenous, parenteral, intraperitoneal, intramuscular, intralesional, intrathecal, intracranial, intranasal or subcutaneous administration, or the implantation of a slow-release device, *e.g.*, a mini-osmotic pump, to a subject. Administration is by any route, including parenteral and transmucosal (*e.g.*, buccal, sublingual, palatal, gingival, nasal, vaginal, rectal, or transdermal). Parenteral administration includes, *e.g.*, intravenous, intramuscular, intra-arteriole, intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial. Other modes of delivery include, but are not limited to, the use of liposomal formulations, intravenous infusion, transdermal patches, *etc.* By “co-administer” it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies (*e.g.* anti-cancer agent, chemotherapeutic, or treatment for a neurodegenerative disease). The compound of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compound individually or in combination (more than one compound or agent). Thus, the preparations can also be combined, when desired, with other active substances (*e.g.* to reduce metabolic degradation). The compositions of the present invention can be

delivered by transdermally, by a topical route, formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols. Oral preparations include tablets, pills, powder, dragees, capsules, liquids, lozenges, cachets, gels, syrups, slurries, suspensions, etc., suitable for ingestion by the patient. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. The compositions of the present invention may additionally include components to provide sustained release and/or comfort. Such components include high molecular weight, anionic mucomimetic polymers, gelling polysaccharides and finely-divided drug carrier substrates. These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841; 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by reference in their entirety for all purposes. The compositions of the present invention can also be delivered as microspheres for slow release in the body. For example, microspheres can be administered via intradermal injection of drug-containing microspheres, which slowly release subcutaneously (see Rao, *J. Biomater Sci. Polym. Ed.* 7:623-645, 1995; as biodegradable and injectable gel formulations (see, e.g., Gao *Pharm. Res.* 12:857-863, 1995); or, as microspheres for oral administration (see, e.g., Eyles, *J. Pharm. Pharmacol.* 49:669-674, 1997). In another embodiment, the formulations of the compositions of the present invention can be delivered by the use of liposomes which fuse with the cellular membrane or are endocytosed, *i.e.*, by employing receptor ligands attached to the liposome, that bind to surface membrane protein receptors of the cell resulting in endocytosis. By using liposomes, particularly where the liposome surface carries receptor ligands specific for target cells, or are otherwise preferentially directed to a specific organ, one can focus the delivery of the compositions of the present invention into the target cells *in vivo*. (See, e.g., Al-Muhammed, *J. Microencapsul.* 13:293-306, 1996; Chonn, *Curr. Opin. Biotechnol.* 6:698-708, 1995; Ostro, *Am. J. Hosp. Pharm.* 46:1576-1587, 1989). The compositions of the present invention can also be delivered as nanoparticles.

[0137] By “co-administer” it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies. The compounds of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compounds individually or in combination (more than one compound). The compositions of the present invention can be delivered transdermally, by a topical route, or formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols.

[0138] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0139] Utilizing the teachings provided herein, an effective prophylactic or therapeutic treatment regimen can be planned that does not cause substantial toxicity and yet is effective to treat the clinical symptoms demonstrated by the particular patient. This planning should involve the careful choice of active compound by considering factors such as compound potency, relative bioavailability, patient body weight, presence and severity of adverse side effects, preferred mode of administration and the toxicity profile of the selected agent.

[0140] The compounds described herein can be used in combination with one another, with other active agents known to be useful in treating cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0141] In some embodiments, co-administration includes administering one active agent within 0.5, 1, 2, 4, 6, 8, 10, 12, 16, 20, 24 hours, 2 days, 4 days, 1 week or 1 month of a second active agent. Co-administration includes administering two active agents simultaneously, approximately simultaneously (e.g., within about 1, 5, 10, 15, 20, or 30 minutes of each other), or sequentially in any order. In some embodiments, co-administration can be accomplished by co-formulation, i.e., preparing a single pharmaceutical composition including both active agents. In other embodiments, the active agents can be formulated separately. In another embodiment, the active and/or adjunctive agents may be linked or conjugated to one another. In some embodiments, the compounds described herein may be combined with treatments for cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0142] The compounds described herein can be administered to treat a fertility disease or disorder and/or cancer. In this regard, the compounds disclosed herein may be administered either alone to treat such diseases or disorders or may be co-administered with another therapeutic agent to treat such diseases or disorders.

[0143] The compounds disclosed herein may be co-administered with an agonist or antagonist of CRF₁ function. The compounds described herein can be co-administered with conventional anti-cancer/chemotherapeutic.

[0144] “Anti-cancer agent” is used in accordance with its plain ordinary meaning and refers to a composition (e.g. compound, drug, antagonist, inhibitor, modulator) having antineoplastic properties or the ability to inhibit the growth or proliferation of cells. In some embodiments, an anti-cancer agent is a chemotherapeutic. In some embodiments, an anti-cancer agent is an agent

identified herein having utility in methods of treating cancer. In some embodiments, an anti-cancer agent is an agent approved by the FDA or similar regulatory agency of a country other than the USA, for treating cancer. Examples of anti-cancer agents include, but are not limited to, MEK (e.g. MEK1, MEK2, or MEK1 and MEK2) inhibitors (e.g. XL518, CI-1040, PD035901, selumetinib/ AZD6244, GSK1120212/ trametinib, GDC-0973, ARRY-162, ARRY-300, AZD8330, PD0325901, U0126, PD98059, TAK-733, PD318088, AS703026, BAY 869766), alkylating agents (e.g., cyclophosphamide, ifosfamide, chlorambucil, busulfan, melphalan, mechlorethamine, uramustine, thiotepa, nitrosoureas, nitrogen mustards (e.g., mechlorethamine, cyclophosphamide, chlorambucil, melphalan), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomustine, semustine, streptozocin), triazenes (decarbazine)), anti-metabolites (e.g., 5-azathioprine, leucovorin, capecitabine, fludarabine, gemcitabine, pemetrexed, raltitrexed, folic acid analog (e.g., methotrexate), or pyrimidine analogs (e.g., fluorouracil, floxouridine, Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin), *etc.*), plant alkaloids (e.g., vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, paclitaxel, docetaxel, *etc.*), topoisomerase inhibitors (e.g., irinotecan, topotecan, amsacrine, etoposide (VP16), etoposide phosphate, teniposide, *etc.*), antitumor antibiotics (e.g., doxorubicin, adriamycin, daunorubicin, epirubicin, actinomycin, bleomycin, mitomycin, mitoxantrone, plitacimycin, *etc.*), platinum-based compounds (e.g. cisplatin, oxaloplatin, carboplatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant (e.g., mitotane, aminoglutethimide), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), inhibitors of mitogen-activated protein kinase signaling (e.g. U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-9006, wortmannin, or LY294002, Syk inhibitors, mTOR inhibitors, antibodies (e.g., rituxan), gossypol, genasense, polyphenol E, Chlorofusin, all trans-retinoic acid (ATRA), bryostatin, tumor necrosis factor-related apoptosis-inducing ligand (TRAIL), 5-aza-2'-deoxycytidine, all trans retinoic acid, doxorubicin, vincristine, etoposide, gemcitabine, imatinib (Gleevec.RTM.), geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), flavopiridol, LY294002, bortezomib, trastuzumab, BAY 11-7082, PKC412, PD184352, 20-epi-1, 25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense

oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators;
 apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine;
 axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III
 derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine;
 beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor;
 bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate;
 bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin
 derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole;
 CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS);
 castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-
 porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B;
 combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol;
 cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam;
 cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine;
 dehydrididemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil;
 diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; 9-dioxamycin;
 diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol;
 duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur;
 epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists;
 etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim;
 finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride;
 forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate;
 galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam;
 heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene;
 idramantone; ilmofofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides;
 insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins;
 iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole;
 isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide;
 leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor;
 leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole;
 linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds;
 lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
 loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin
 A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors;
 menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone;

miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol;
 mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone;
 mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin;
 monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene
 inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B;
 mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides;
 nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin;
 nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide
 modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone;
 oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer;
 ormaplatin; osaterone; oxaliplatin; oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic
 acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan
 polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol;
 phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride;
 pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum
 complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin;
 prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based
 immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein
 tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins;
 pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists;
 raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP
 inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII
 retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol;
 saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence
 derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction
 modulators; single chain antigen-binding protein; sizofuran; sobuzoxane; sodium borocaptate;
 sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid;
 spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-
 cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive
 intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans;
 tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur;
 tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide;
 tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin
 mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone;
 tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem

cell factor; translation inhibitors; tretinoin; triacetyluridine; tricyriline; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; zinostatin; stimalamer, Adriamycin, Dactinomycin, Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqune; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; iimofosine; interleukin II (including recombinant interleukin II, or rIL.sub.2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1; interferon alfa-n3; interferon beta-1a; interferon gamma-1b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedpa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazoie; nogalamycin; ormaplatin; oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprime; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spirolatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone

hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricitabine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinylicinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride, agents that arrest cells in the G2-M phases and/or modulate the formation or stability of microtubules, (e.g. Taxol.TM (i.e. paclitaxel), Taxotere.TM, compounds comprising the taxane skeleton, Erbulozole (i.e. R-55104), Dolastatin 10 (i.e. DLS-10 and NSC-376128), Mivobulin isethionate (i.e. as CI-980), Vincristine, NSC-639829, Discodermolide (i.e. as NVP-XX-A-296), ABT-751 (Abbott, i.e. E-7010), Altorhyrtins (e.g. Altorhyrtin A and Altorhyrtin C), Spongistatins (e.g. Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5, Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride (i.e. LU-103793 and NSC-D-669356), Epothilones (e.g. Epothilone A, Epothilone B, Epothilone C (i.e. desoxyepothilone A or dEpoA), Epothilone D (i.e. KOS-862, dEpoB, and desoxyepothilone B), Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide, 16-aza-epothilone B, 21-aminoepothilone B (i.e. BMS-310705), 21-hydroxyepothilone D (i.e. Desoxyepothilone F and dEpoF), 26-fluoroepothilone, Auristatin PE (i.e. NSC-654663), Soblidotin (i.e. TZT-1027), LS-4559-P (Pharmacia, i.e. LS-4577), LS-4578 (Pharmacia, i.e. LS-477-P), LS-4477 (Pharmacia), LS-4559 (Pharmacia), RPR-112378 (Aventis), Vincristine sulfate, DZ-3358 (Daiichi), FR-182877 (Fujisawa, i.e. WS-9885B), GS-164 (Takeda), GS-198 (Takeda), KAR-2 (Hungarian Academy of Sciences), BSF-223651 (BASF, i.e. ILX-651 and LU-223651), SAH-49960 (Lilly/Novartis), SDZ-268970 (Lilly/Novartis), AM-97 (Armad/Kyowa Hakko), AM-132 (Armad), AM-138 (Armad/Kyowa Hakko), IDN-5005 (Indena), Cryptophycin 52 (i.e. LY-355703), AC-7739 (Ajinomoto, i.e. AVE-8063A and CS-39.HCl), AC-7700 (Ajinomoto, i.e. AVE-8062, AVE-8062A, CS-39-L-Ser.HCl, and RPR-258062A), Vitilevuamide, Tubulysin A, Canadensol, Centaureidin (i.e. NSC-106969), T-138067 (Tularik, i.e. T-67, TL-138067 and TI-138067), COBRA-1 (Parker Hughes Institute, i.e. DDE-261 and WHI-261), H10 (Kansas State University), H16 (Kansas State University), Oncocidin A1 (i.e. BTO-956 and DIME), DDE-313 (Parker Hughes Institute), Fijianolide B, Laulimalide, SPA-2 (Parker Hughes Institute), SPA-1 (Parker Hughes Institute, i.e. SPIKET-P), 3-IAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-569), Narcosine (also known as NSC-5366), Nascapine, D-24851 (Asta Medica), A-105972 (Abbott), Hemiasterlin, 3-BAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-191), TMPN (Arizona State University), Vanadocene acetylacetonate, T-138026 (Tularik), Monsatrol, Inanocine (i.e. NSC-698666), 3-IAABE (Cytoskeleton/Mt. Sinai

School of Medicine), A-204197 (Abbott), T-607 (Tuiarik, i.e. T-900607), RPR-115781 (Aventis), Eleutherobins (such as Desmethyleleutherobin, Desaetyeleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, D-64131 (Asta Medica), D-68144 (Asta Medica), Diazonamide A, A-293620 (Abbott), NPI-2350 (Nereus), Taccalonolide A, TUB-245 (Aventis), A-259754 (Abbott), Diozostatin, (-)-Phenylahistin (i.e. NSCL-96F037), D-68838 (Asta Medica), D-68836 (Asta Medica), Myoseverin B, D-43411 (Zentaris, i.e. D-81862), A-289099 (Abbott), A-318315 (Abbott), HTI-286 (i.e. SPA-110, trifluoroacetate salt) (Wyeth), D-82317 (Zentaris), D-82318 (Zentaris), SC-12983 (NCI), Resverastatin phosphate sodium, BPR-OY-007 (National Health Research Institutes), and SSR-250411 (Sanofi), steroids (*e.g.*, dexamethasone), finasteride, aromatase inhibitors, gonadotropin-releasing hormone agonists (GnRH) such as goserelin or leuprolide, adrenocorticosteroids (*e.g.*, prednisone), progestins (*e.g.*, hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (*e.g.*, diethylstilbestrol, ethinyl estradiol), antiestrogen (*e.g.*, tamoxifen), androgens (*e.g.*, testosterone propionate, fluoxymesterone), antiandrogen (*e.g.*, flutamide), immunostimulants (*e.g.*, Bacillus Calmette-Guérin (BCG), levamisole, interleukin-2, alpha-interferon, *etc.*), monoclonal antibodies (*e.g.*, anti-CD20, anti-HER2, anti-CD52, anti-HLA-DR, and anti-VEGF monoclonal antibodies), immunotoxins (*e.g.*, anti-CD33 monoclonal antibody-calicheamicin conjugate, anti-CD22 monoclonal antibody-pseudomonas exotoxin conjugate, *etc.*), radioimmunotherapy (*e.g.*, anti-CD20 monoclonal antibody conjugated to ^{111}In , ^{90}Y , or ^{131}I , *etc.*), triptolide, homoharringtonine, dactinomycin, doxorubicin, epirubicin, topotecan, itraconazole, vindesine, cerivastatin, vincristine, deoxyadenosine, sertraline, pitavastatin, irinotecan, clofazimine, 5-nonyloxytryptamine, vemurafenib, dabrafenib, erlotinib, gefitinib, EGFR inhibitors, epidermal growth factor receptor (EGFR)-targeted therapy or therapeutic (*e.g.* gefitinib (IressaTM), erlotinib (TarcevaTM), cetuximab (ErbixTM), lapatinib (TykerbTM), panitumumab (VectibixTM), vandetanib (CaprelsaTM), afatinib/BIBW2992, CI-1033/canertinib, neratinib/HKI-272, CP-724714, TAK-285, AST-1306, ARRY334543, ARRY-380, AG-1478, dacomitinib/PF299804, OSI-420/desmethyl erlotinib, AZD8931, AEE788, pelitinib/EKB-569, CUDC-101, WZ8040, WZ4002, WZ3146, AG-490, XL647, PD153035, BMS-599626), sorafenib, imatinib, sunitinib, dasatinib, or the like.

[0145] “Chemotherapeutic” or “chemotherapeutic agent” is used in accordance with its plain ordinary meaning and refers to a chemical composition or compound having antineoplastic properties or the ability to inhibit the growth or proliferation of cells.

[0146] The compounds disclosed herein may be co-administered with an antiproliferative/antineoplastic drug or a combination thereof, as used in medical oncology, such as an alkylating agent (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard,

melphalan, chlorambucil, busulphan or a nitrosourea); an antimetabolite (for example an antifolate such as a fluoropyrimidine like 5-fluorouracil or tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, gemcitabine or paclitaxel); an antitumour antibiotic (for example an anthracycline such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin or mithramycin); an antimitotic agent (for example a vinca alkaloid such as vincristine, vinblastine, vindesine or vinorelbine, or a taxoid such as taxol or taxotere); or a topoisomerase inhibitor (for example an epipodophyllotoxin such as etoposide, teniposide, amsacrine, topotecan or a camptothecin); (ii) a cytostatic agent such as an antioestrogen (for example tamoxifen, toremifene, raloxifene, droloxifene or idoxifene), an oestrogen receptor down regulator (for example fulvestrant), an antiandrogen (for example bicalutamide, flutamide, nilutamide or cyproterone acetate), a LHRH antagonist or LHRH agonist (for example goserelin, leuporelin or buserelin), a progestogen (for example megestrol acetate), an aromatase inhibitor (for example as anastrozole, letrozole, vorazole or exemestane) or an inhibitor of 5.alpha.-reductase such as finasteride; (iii) an agent which inhibits cancer cell invasion (for example a metalloproteinase inhibitor like marimastat or an inhibitor of urokinase plasminogen activator receptor function); (iv) an inhibitor of growth factor function, for example: a growth factor antibody (for example the anti-erbB2 antibody trastuzumab, or the anti-erbB1 antibody cetuximab [C225]), a farnesyl transferase inhibitor, a tyrosine kinase inhibitor or a serine/threonine kinase inhibitor, an inhibitor of the epidermal growth factor family (for example an EGFR family tyrosine kinase inhibitor such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD 1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) or 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), an inhibitor of the platelet-derived growth factor family, or an inhibitor of the hepatocyte growth factor family; (v) an antiangiogenic agent such as one which inhibits the effects of vascular endothelial growth factor (for example the anti-vascular endothelial cell growth factor antibody bevacizumab, a compound disclosed in WO 97/22596, WO 97/30035, WO 97/32856 or WO 98/13354), or a compound that works by another mechanism (for example linomide, an inhibitor of integrin .alpha.v.beta.3 function or an angiostatin); (vi) a vascular damaging agent such as combretastatin A4, or a compound disclosed in WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 or WO 02/08213; (vii) an agent used in antisense therapy, for example one directed to one of the targets listed above, such as ISIS 2503, an anti-ras antisense; (viii) an agent used in a gene therapy approach, for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a

bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; or (ix) an agent used in an immunotherapeutic approach, for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

[0147] In embodiments, the compounds disclosed herein can be co-administered with an antibody, such as a monoclonal antibody targeting B-Lymphocytes (such as CD20 (rituximab), MRA-aIL16R and T-Lymphocytes, CTLA4-Ig, HuMax II-15) or antibody modulating Ig function such as anti-IgE (for example omalizumab).

[0148] In embodiments, treatment of cancer includes administration of an effective amount of at least two of the following: a CRF1 inhibitor, an inhibitor of the PD-L1/PD-1 pathway, an inhibitor of CTLA-4, an agonistic antibody of CD137 (4-1BB). In some embodiments, the method may comprise the use of two or more combinations.

[0149] In a further embodiment, the compounds described herein can be co-administered with conventional radiotherapeutic agents including, but not limited to, radionuclides such as ⁴⁷Sc, ⁶⁴Cu, ⁶⁷Cu, ⁸⁹Sr, ⁸⁶Y, ⁸⁷Y, ⁹⁰Y, ¹⁰⁵Rh, ¹¹¹Ag, ¹¹¹In, ^{117m}Sn, ¹⁴⁹Pm, ¹⁵³Sm, ¹⁶⁶Ho, ¹⁷⁷Lu, ¹⁸⁶Re, ¹⁸⁸Re, ²¹¹At, and ²¹²Bi, optionally conjugated to antibodies directed against tumor antigens.

[0150] In addition, a CRF₁ inhibitor may be combined with the therapeutic administration of immune cells, sometimes referred to as adoptive cell transfer. These cells may be cells from the patient, a genetically related or unrelated donor, they may be genetically modified (e.g. CAR-T cells, NK cells, etc), cell lines, genetically modified cell lines and live or dead versions of the above. CRF₁ inhibitors may also be combined with vaccines of any kind (e.g. protein/peptide, viral, bacterial, cellular) to stimulate immune responses to cancer.

[0151] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramuscular, intranasal, intraocular, intrapericardial, intraperitoneal, intrapleural, intraprostatic, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenous, intravesicular, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0152] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramuscular, intranasal, intraocular, intrapericardial, intraperitoneal, intrapleural, intraprostatic, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenous, intravesicular, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0153] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered once daily until study reached endpoint. The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered at least three times but in some studies four or more times depending on the length of the study and/or the design of the study.

[0154] The methods disclosed herein may be used in combination with additional cancer therapy. In some embodiments, the distinct cancer therapy comprises surgery, radiotherapy, chemotherapy, toxin therapy, immunotherapy, cryotherapy or gene therapy. In some embodiments, the cancer is a chemotherapy-resistant or radio-resistant cancer.

[0155] A “cell” as used herein, refers to a cell carrying out metabolic or other function sufficient to preserve or replicate its genomic DNA. A cell can be identified by well-known methods in the art including, for example, presence of an intact membrane, staining by a particular dye, ability to produce progeny or, in the case of a gamete, ability to combine with a second gamete to produce a viable offspring. Cells may include prokaryotic and eukaryotic cells. Prokaryotic cells include but are not limited to bacteria. Eukaryotic cells include but are not limited to yeast cells and cells derived from plants and animals, for example mammalian, insect (*e.g.*, spodoptera) and human cells. Cells may be useful when they are naturally nonadherent or have been treated not to adhere to surfaces, for example by trypsinization.

[0156] “Control” or “control experiment” is used in accordance with its plain ordinary meaning and refers to an experiment in which the subjects or reagents of the experiment are treated as in a parallel experiment except for omission of a procedure, reagent, or variable of the experiment. In some instances, the control is used as a standard of comparison in evaluating experimental effects. In some embodiments, a control is the measurement of the activity of a protein in the absence of a compound as described herein (including embodiments and examples).

[0157] The term “modulator” refers to a composition that increases or decreases the level of a target molecule or the function of a target molecule or the physical state of the target of the molecule. In some embodiments, a CRF₁ associated disease modulator is a compound that

reduces the severity of one or more symptoms of a disease associated with CRF₁ (e.g. an endocrine disorder, a cancer including but not limited to TART or OART). A CRF₁ modulator is a compound that increases or decreases the activity or function or level of activity or level of function of CRF₁. A modulator may act alone, or it may use a cofactor, e.g., a protein, metal ion, or small molecule. Examples of modulators include small molecule compounds and other bioorganic molecules. Numerous libraries of small molecule compounds (e.g., combinatorial libraries) are commercially available and can serve as a starting point for identifying a modulator. The skilled artisan is able to develop one or more assays (e.g., biochemical or cell-based assays) in which such compound libraries can be screened in order to identify one or more compounds having the desired properties; thereafter, the skilled medicinal chemist is able to optimize such one or more compounds by, for example, synthesizing and evaluating analogs and derivatives thereof. Synthetic and/or molecular modeling studies can also be utilized in the identification of an activator.

[0158] The term “modulate” is used in accordance with its plain ordinary meaning and refers to the act of changing or varying one or more properties. “Modulation” refers to the process of changing or varying one or more properties. For example, as applied to the effects of a modulator on a target protein, to modulate means to change by increasing or decreasing a property or function of the target molecule or the amount of the target molecule. In embodiments, the terms “modulate”, “modulation” and the like refer to the ability of a molecule (e.g., an activator or an inhibitor) to increase or decrease the function or activity of CRF₁, either directly or indirectly, relative to the absence of the molecule.

[0159] The term “associated” or “associated with” in the context of a substance or substance activity or function associated with a disease (e.g. a protein associated disease, an endocrine disorder, a disease associated with CRF₁ activity, a CRF₁ associated disease (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) means that the disease (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) is caused by (in whole or in part), or a symptom of the disease is caused by (in whole or in part) the substance or substance activity or function. For example, an endocrine disorder associated with CRF₁ activity or function may be an endocrine disorder that results (entirely or partially) from aberrant CRF₁ function (e.g. enzyme activity, protein-protein interaction, signaling pathway) or an endocrine disorder wherein a particular symptom of the disease is caused (entirely or partially) by aberrant CRF₁ activity or function. As used herein, what is described as being associated with a disease, if a causative agent, could be a target for treatment of the disease. For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated disease (e.g., an endocrine disorder including but not limited to TART or OART), may be treated with a

compound described herein (e.g., CRF₁ modulator, CRF₁ inhibitor, or CRF₁ antagonist), in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease (e.g., an endocrine disorder such as TART or OART). For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated endocrine disorder, may be treated with an CRF₁ modulator or CRF₁ inhibitor, in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease.

[0160] The term “aberrant” as used herein refers to different from normal. When used to describe enzymatic activity or protein function, aberrant refers to activity or function that is greater or less than a normal control or the average of normal non-diseased control samples. Aberrant activity may refer to an amount of activity that results in a disease, wherein returning the aberrant activity to a normal or non-disease-associated amount (e.g. by administering a compound or using a method as described herein), results in reduction of the disease or one or more disease symptoms.

[0161] The term “signaling pathway” as used herein refers to a series of interactions between cellular and optionally extra-cellular components (e.g. proteins, nucleic acids, small molecules, ions, lipids) that conveys a change in one component to one or more other components, which in turn may convey a change to additional components, which is optionally propagated to other signaling pathway components. For example, binding of a CRF₁ with a compound as described herein may reduce the level of a product of the CRF₁ catalyzed reaction or the level of a downstream derivative of the product or binding may reduce the interactions between the CRF₁ or a reaction product and downstream effectors or signaling pathway components (e.g., MAP kinase pathway), resulting in changes in cell growth, proliferation, or survival.

[0162] As used herein, the terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” and all other related art-accepted terms, many of which are set forth below, refer to a compound capable of modulating, either directly or indirectly, the CRF₁ receptor in an *in vitro* assay, an *in vivo* model, and/or other means indicative of therapeutic efficacy. The terms also refer to a compound that exhibits at least some therapeutic benefit in a human subject.

[0163] The phrase “in a sufficient amount to effect a change” means that there is a detectable difference between a level of an indicator measured before (e.g., a baseline level) and after administration of a particular therapy. Indicators include any objective parameter (e.g., serum concentration) or subjective parameter (e.g., a subject’s feeling of well-being).

[0164] The “activity” of a molecule may describe or refer to the binding of the molecule to a ligand or to a receptor; to catalytic activity; to the ability to stimulate gene expression or cell signaling, differentiation, or maturation; to antigenic activity; to the modulation of activities of

other molecules; and the like. The term “proliferative activity” encompasses an activity that promotes, that is necessary for, or that is specifically associated with, for example, normal cell division, as well as cancer, tumors, dysplasia, cell transformation, metastasis, and angiogenesis. **[0165]** “Substantially pure” indicates that a component makes up greater than about 50% of the total content of the composition, and typically greater than about 60% of the total polypeptide content. More typically, “substantially pure” refers to compositions in which at least 75%, at least 85%, at least 90% or more of the total composition is the component of interest. In some cases, the polypeptide will make up greater than about 90%, or greater than about 95% of the total content of the composition (percentage in a weight per weight basis).

[0166] The terms “specifically binds” and “selectively binds,” when referring to a ligand/receptor, antibody/antigen, or other binding pair, indicates a binding reaction which is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated conditions, a specified ligand binds to a particular receptor and does not bind in a significant amount to other proteins present in the sample. The antibody, or binding composition derived from the antigen-binding site of an antibody, of the contemplated method binds to its antigen, or a variant or mutein thereof, with an affinity that is at least two-fold greater, at least 10-times greater, at least 20-times greater, or at least 100-times greater than the affinity with any other antibody, or binding composition derived therefrom. In embodiments, the antibody will have an affinity that is greater than about 10^9 liters/mol, as determined by, e.g., Scatchard analysis (Munsen, et al. (1980) *Analyt. Biochem.* 107:220-239).

[0167] The terms “DNA,” “nucleic acid,” “nucleic acid molecule,” “polynucleotide” and the like are used interchangeably herein to refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Non-limiting examples of polynucleotides include linear and circular nucleic acids, messenger RNA (mRNA), complementary DNA (cDNA), recombinant polynucleotides, vectors, probes, primers and the like.


[0168] As used herein, the terms “variants” and “homologs” are used interchangeably to refer to amino acid or nucleic acid sequences that are similar to reference amino acid or nucleic acid sequences, respectively. The term encompasses naturally-occurring variants and non-naturally-occurring variants. Naturally-occurring variants include homologs (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one species to another), and allelic variants (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one individual to another within a species). Thus, variants and homologs encompass naturally occurring amino acid and nucleic acid sequences encoded thereby and their isoforms, as well as splice variants of a protein or gene. The terms also

encompass nucleic acid sequences that vary in one or more bases from a naturally-occurring nucleic acid sequence but still translate into an amino acid sequence that corresponds to the naturally-occurring protein due to degeneracy of the genetic code. Non-naturally-occurring variants and homologs include polypeptides and nucleic acids that comprise a change in amino acid or nucleotide sequence, respectively, where the change in sequence is artificially introduced (e.g., muteins); for example, the change is generated in the laboratory by human intervention (“hand of man”). Therefore, non-naturally occurring variants and homologs may also refer to those that differ from the naturally-occurring sequences by one or more conservative substitutions and/or tags and/or conjugates.

[0169] The term “muteins” as used herein refers broadly to mutated recombinant proteins. These proteins usually carry single or multiple amino acid substitutions and are frequently derived from cloned genes that have been subjected to site-directed or random mutagenesis, or from completely synthetic genes.



II. Compounds

[0170] In an aspect provided herein, is a compound of Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 and R^2 are independently ethyl or n-propyl. R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R^4 is hydrogen,

Br, R^aR^bN- , methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl,




, R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$. In embodiments, the inhibitor forms an irreversible bond with the amino acid residue. In embodiments, the bond is a covalent bond.


[0171] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0172] In certain embodiments, R^3 is Cl, Br, or methyl.


[0173] In certain embodiments, R^4 is Br, R^aR^bN- , pyridin-4-yl, morpholin-4-yl, or



[0174] In certain embodiments, R⁴ is morpholin-4-yl or




[0175] In certain embodiments, R⁴ is hydrogen, Br, R^aR^bN- and R^a and R^b are independently C₁-C₃alkyl.




[0176] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0177] In certain embodiments, the compound is



[0178] In certain embodiments, the compound is



[0179] In one aspect described herein, there is a method of treating a subject in need thereof comprising administering an astressin or a pharmaceutically acceptable salt or solvate thereof. An astressin generally refers to a nonselective corticotropin releasing hormone antagonist that reduces the synthesis of ACTH and cortisol.

[0180] In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, the method comprising: (i) measuring a hormone

level in the subject in need thereof; (ii) administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof; and repeating steps (i) and (ii) until the hormone level reaches a pre-determined range followed by a maintenance therapy of a daily dosing of said astressin.

[0181] In one aspect described herein, there is a method of improving hyperandrogenic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating menstrual irregularity, ovulatory dysfunction or infertility, in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of improving metabolic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto; or a pharmaceutically acceptable salt or solvate thereof.

[0182] In one aspect described herein, there is a method of improving quality of life in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

III. Pharmaceutical Compositions

[0183] In an aspect, there is provided a pharmaceutical composition, including a compound as described herein, including embodiments, or the structural Formula (I), and a pharmaceutically acceptable excipient.

[0184] The compounds (e.g., CRF₁ inhibitors/antagonists) of the present disclosure may be in the form of compositions suitable for administration to a subject. In general, such compositions are “pharmaceutical compositions” comprising a compound (e.g., CRF₁ inhibitor(s)/antagonist(s)) and one or more pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients. In certain embodiments, the compound (e.g., CRF₁ inhibitor/antagonist) are present in a therapeutically acceptable amount. The pharmaceutical compositions may be used in the methods of the present disclosure; thus, for example, the pharmaceutical compositions can be administered *ex vivo* or *in vivo* to a subject in order to practice the therapeutic and prophylactic methods and uses described herein.

[0185] In one aspect described herein, there is a pharmaceutical composition in the form of a capsule comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a pharmaceutical composition in the form of a tablet comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

[0186] The pharmaceutical compositions of the present disclosure can be formulated to be compatible with the intended method or route of administration; exemplary routes of administration are set forth herein.

[0187] The pharmaceutical compositions containing the active ingredient (e.g., an inhibitor of CRF₁ function) may be in a form suitable for oral use, for example, as tablets, capsules, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups, solutions, microbeads or elixirs. Pharmaceutical compositions intended for oral use may be prepared for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents such as, for example, sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets, capsules and the like contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture thereof. These excipients may be, for example, diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc.

[0188] The tablets, capsules and the like suitable for oral administration may be uncoated or coated to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action. For example, a time-delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated to form osmotic therapeutic tablets for controlled release. Additional agents include biodegradable or biocompatible particles or a polymeric substance such as polyesters, polyamine acids, hydrogel, polyvinyl pyrrolidone, polyanhydrides, polyglycolic acid, ethylene-vinylacetate, methylcellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymers, polylactide/glycolide copolymers, or ethylenevinylacetate copolymers in order to control delivery of an administered composition. For example, the oral agent can be entrapped in microcapsules prepared by coacervation techniques or by interfacial polymerization, by the use of hydroxymethylcellulose or gelatin-microcapsules or poly(methylmethacrolate) microcapsules, respectively, or in a colloid drug delivery system. Colloidal dispersion systems include macromolecule complexes, nano-capsules, microspheres, microbeads, and lipid-based systems, including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Methods for the preparation of the above-mentioned formulations will be apparent to those skilled in the art.

[0189] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium

phosphate, kaolin or microcrystalline cellulose, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

[0190] Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture thereof. Such excipients can be suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents, for example a naturally-occurring phosphatide (e.g., lecithin), or condensation products of an alkylene oxide with fatty acids (e.g., polyoxy-ethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols (e.g., for heptadecaethyleneoxycetanol), or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol (e.g., polyoxyethylene sorbitol monooleate), or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides (e.g., polyethylene sorbitan monooleate). The aqueous suspensions may also contain one or more preservatives.

[0191] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

[0192] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, and optionally one or more suspending agents and/or preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified herein.

[0193] The pharmaceutical compositions of the present disclosure may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example, liquid paraffin, or mixtures of these. Suitable emulsifying agents may be naturally occurring gums, for example, gum acacia or gum tragacanth; naturally occurring phosphatides, for example, soy bean, lecithin, and esters or partial esters derived from fatty acids; hexitol anhydrides, for example, sorbitan monooleate; and condensation products of partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate.

[0194] The pharmaceutical compositions typically comprise a therapeutically effective amount of a CRF₁ inhibitor/antagonist contemplated by the present disclosure and one or more pharmaceutically and physiologically acceptable formulation agents. Suitable pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients include, but are not limited to, antioxidants (e.g., ascorbic acid and sodium bisulfate), preservatives (e.g., benzyl

alcohol, methyl parabens, ethyl or n-propyl, p-hydroxybenzoate), emulsifying agents, suspending agents, dispersing agents, solvents, fillers, bulking agents, detergents, buffers, vehicles, diluents, and/or adjuvants. For example, a suitable vehicle may be physiological saline solution or citrate-buffered saline, possibly supplemented with other materials common in pharmaceutical compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Those skilled in the art will readily recognize a variety of buffers that can be used in the pharmaceutical compositions and dosage forms contemplated herein. Typical buffers include, but are not limited to, pharmaceutically acceptable weak acids, weak bases, or mixtures thereof. As an example, the buffer components can be water soluble materials such as phosphoric acid, tartaric acids, lactic acid, succinic acid, citric acid, acetic acid, ascorbic acid, aspartic acid, glutamic acid, and salts thereof. Acceptable buffering agents include, for example, a Tris buffer; N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) (HEPES); 2-(N-Morpholino)ethanesulfonic acid (MES); 2-(N-Morpholino)ethanesulfonic acid sodium salt (MES); 3-(N-Morpholino)propanesulfonic acid (MOPS); and N-tris[Hydroxymethyl]methyl-3-aminopropanesulfonic acid (TAPS).

[0195] After a pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form, a lyophilized form requiring reconstitution prior to use, a liquid form requiring dilution prior to use, or other acceptable form. In some embodiments, the pharmaceutical composition is provided in a single-use container (e.g., a single-use vial, ampule, syringe, or autoinjector (similar to, e.g., an EpiPen®)), whereas a multi-use container (e.g., a multi-use vial) is provided in other embodiments.

[0196] Formulations can also include carriers to protect the composition against rapid degradation or elimination from the body, such as a controlled release formulation, including liposomes, hydrogels, prodrugs and microencapsulated delivery systems. For example, a time-delay material such as glyceryl monostearate or glyceryl stearate alone, or in combination with a wax, may be employed. Any drug delivery apparatus may be used to deliver a CRF₁ inhibitor/antagonist, including implants (e.g., implantable pumps) and catheter systems, slow injection pumps and devices, all of which are well known to the skilled artisan.

[0197] Depot injections, which are generally administered subcutaneously or intramuscularly, may also be utilized to release the compound (e.g., CRF₁ inhibitor/antagonist) disclosed herein over a defined period of time. Depot injections are usually either solid- or oil-based and generally comprise at least one of the formulation components set forth herein. One of ordinary skill in the art is familiar with possible formulations and uses of depot injections.

[0198] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents mentioned herein. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Acceptable diluents, solvents and dispersion media that may be employed include water, Ringer's solution, isotonic sodium chloride solution, Cremophor® EL (BASF, Parsippany, NJ) or phosphate buffered saline (PBS), ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. In addition, sterile fixed oils are conventionally employed as a solvent or suspending medium; for this purpose, any bland fixed oil may be employed, including synthetic mono- or diglycerides. Moreover, fatty acids, such as oleic acid, find use in the preparation of injectables. Prolonged absorption of particular injectable formulations can be achieved by including an agent that delays absorption (e.g., aluminum monostearate or gelatin).

[0199] The present disclosure contemplates the administration of the compound (e.g., CRF₁ inhibitor/antagonist) in the form of suppositories for rectal administration. The suppositories can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include, but are not limited to, cocoa butter and polyethylene glycols.

[0200] The compound (e.g., CRF₁ inhibitor/antagonist) contemplated by the present disclosure may be in the form of any other suitable pharmaceutical composition (e.g., sprays for nasal or inhalation use) currently known or developed in the future.

IV. Methods

[0201] In certain embodiments, provided herein are methods for treating or preventing a disease comprising administering an effective amount of a compound of Formula (I), or an isotopic variant thereof or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0202] In one embodiment, provided herein are methods for treating or preventing a tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a benign tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a malignant tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal

tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal rest tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

{0203} In one embodiment, provided herein are methods for treating or preventing infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing male infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm count, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm motility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm morphology, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing female infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

{0204} In one embodiment, provided herein are methods for treating or preventing a proliferative or hyperproliferative disease or disorder, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, the proliferative disease is cancer. In certain embodiments, the proliferative disease is a cancer of the breast, skin, prostate, cervix, uterus, ovary, testes, bladder, lung, liver, larynx, oral cavity, colon and gastrointestinal tract (e.g., esophagus, stomach, pancreas), brain, thyroid, blood, and lymphatic system.

{0205} In certain embodiments, the cancer treatable with the methods provided herein includes, but is not limited to, (1) leukemias, including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome or a symptom thereof (such as anemia, thrombocytopenia, neutropenia, bicytopenia or pancytopenia), refractory anemia (RA), RA with ringed sideroblasts (RARS), RA with excess blasts (RAEB),

RAEB in transformation (RAEB-T), preleukemia, and chronic myelomonocytic leukemia (CMML); (2) chronic leukemias, including, but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, and hairy cell leukemia; (3) polycythemia vera; (4) lymphomas, including, but not limited to, Hodgkin's disease and non-Hodgkin's disease; (5) multiple myelomas, including, but not limited to, smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma, and extramedullary plasmacytoma; (6) Waldenstrom's macroglobulinemia; (7) monoclonal gammopathy of undetermined significance; (8) benign monoclonal gammopathy; (9) heavy chain disease; (10) bone and connective tissue sarcomas, including, but not limited to, bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, metastatic cancers, neurilemmoma, rhabdomyosarcoma, and synovial sarcoma; (11) brain tumors, including, but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, and primary brain lymphoma; (12) breast cancer, including, but not limited to, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, primary cancers, Paget's disease, and inflammatory breast cancer; (13) adrenal cancer, including, but not limited to, pheochromocytoma and adrenocortical carcinoma; (14) thyroid cancer, including, but not limited to, papillary or follicular thyroid cancer, medullary thyroid cancer, and anaplastic thyroid cancer; (15) pancreatic cancer, including, but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; (16) pituitary cancer, including, but limited to, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; (17) eye cancer, including, but not limited to, ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; (18) vaginal cancer, including, but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; (19) vulvar cancer, including, but not limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; (20) cervical cancers, including, but not limited to, squamous cell carcinoma, and adenocarcinoma; (21) uterine cancer, including, but not limited to, endometrial carcinoma and uterine sarcoma; (22) ovarian cancer, including, but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; (23) esophageal cancer, including, but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous

carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; (24) stomach cancer, including, but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; (25) colon cancer; (26) rectal cancer; (27) liver cancer, including, but not limited to, hepatocellular carcinoma and hepatoblastoma; (28) gallbladder cancer, including, but not limited to, adenocarcinoma; (29) cholangiocarcinomas, including, but not limited to, papillary, nodular, and diffuse; (30) lung cancer, including, but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma, and small-cell lung cancer; (31) testicular cancer, including, but not limited to, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, and choriocarcinoma (yolk-sac tumor); (32) prostate cancer, including, but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; (33) penile cancer; (34) oral cancer, including, but not limited to, squamous cell carcinoma; (35) basal cancer; (36) salivary gland cancer, including, but not limited to, adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; (37) pharynx cancer, including, but not limited to, squamous cell cancer and verrucous; (38) skin cancer, including, but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, and acral lentiginous melanoma; (39) kidney cancer, including, but not limited to, renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, and transitional cell cancer (renal pelvis and/or uterine); (40) Wilms' tumor; (41) bladder cancer, including, but not limited to, transitional cell carcinoma, squamous cell cancer, adenocarcinoma, and carcinosarcoma; and other cancer, including, not limited to, myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangio-endotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, and papillary adenocarcinomas (*See Fishman et al.*, 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy *et al.*, 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

[0206] Depending on the disorder, disease, or condition to be treated, and the subject's condition, the compounds or pharmaceutical compositions provided herein can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), inhalation, nasal, vaginal, rectal, sublingual, or topical (e.g., transdermal or local) routes of administration and can be formulated, alone or

together, in suitable dosage unit with pharmaceutically acceptable excipients, carriers, adjuvants, and vehicles appropriate for each route of administration as described elsewhere herein.

[0207] The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. Recommended routes of administration for the second active agents are known to those of ordinary skill in the art. See, e.g., Physicians' Desk Reference, 1755-1760 (56th ed., 2002).


[0208] In one embodiment, provided herein are methods for treating or preventing testicular adrenal rest tumors (TART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).

[0209] In one embodiment, provided herein are methods for treating or preventing ovarian adrenal rest tumors (OART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).

[0210] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.


[0211] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.


[0212] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R¹ and R² are independently ethyl or n-propyl. R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R⁴ is hydrogen,


Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl,






, R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $\text{H}_2\text{NCH}_2\text{CH}_2$ -, $(\text{CH}_3)_3\text{COC}(\text{O})\text{NHCH}_2\text{CH}_2$ -, or $\text{CH}_3\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2$ -.

[0213] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments.

[0214] In an aspect provided herein, is a method of decreasing or reducing adrenocorticotrophic hormone (ACTH) levels in a subject in need thereof, comprising administering to the subject contacting the ACTH with a compound of structural Formula (I):




, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 60% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 70% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 80% of the subjects demonstrate reduction in ACTH. In embodiments, at least about 10% of subjects were in the standard or normal range for ACTH after treatment.

[0215] In embodiments, at least about 20% of subjects were in the standard or normal range for ACTH after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for ACTH after treatment. In certain embodiments, ACTH levels are decreased or reduced within about 30 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 1 hour post dosing. In certain embodiments, ACTH levels are


decreased or reduced within about 90 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 2 hours post dosing.

[0216] In an aspect provided herein, is a method of decreasing or reducing 17 α -hydroxyprogesterone (17-OHP) levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R¹, R², R³ and R⁴ are as described herein, including embodiments. R¹, R², R³ and R⁴ are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 60% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 70% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 80% of the subjects demonstrate reduction in 17-OHP. In embodiments, at least about 30% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 40% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 60% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 60% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment.

[0217] In an aspect provided herein, is a method of decreasing or reducing androstenedione levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):




, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 60% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 70% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 80% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 90% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 100% of the subjects demonstrate reduction in androstenedione. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 60% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 40% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range for androstenedione after treatment.


[0218] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0219] In certain embodiments, R^3 is Cl, Br, or methyl.


[0220] In certain embodiments, R^4 is Br, $R^a R^b N-$, pyridin-4-yl, morpholin-4-yl, or



[0221] In certain embodiments, R^4 is morpholin-4-yl or




[0222] In certain embodiments, R^4 is hydrogen, Br, $R^a R^b N-$ and R^a and R^b are independently C_1 - C_3 alkyl.




[0223] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0224] In certain embodiments, the compound is



[0225] In certain embodiments, the compound is



[0226] In certain embodiments, a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is cyclically administered to a patient. Cycling therapy involves the administration of an active agent or a combination of active agents for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

[0227] In certain embodiments, in the treatment, prevention, or amelioration of one or more symptoms of the disorders, diseases, or conditions described herein (e.g., treatment or prevention of testicular adrenal rest tumors (TART) or treatment or prevention of ovarian adrenal rest tumors (OART)), an appropriate dosage level of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof generally is ranging from about 1 mg to 3000 mg, from about 1 mg to 2000 mg, from about 1 mg to 1000 mg, from about 1 mg to about 500 mg, from about 5 mg to about 500 mg,

from about 5 mg to about 400 mg, from about 5 mg to about 300 mg, from about 5 mg to about 250 mg, from about 5 mg to about 200 mg, from about 50 mg to about 200 mg, from about 10 mg to about 400 mg, from about 5 mg to about 25 mg, from about 10 mg to about 25 mg, or from about 25 to about 200 mg, which can be administered in single or multiple doses. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg/day.

[0228] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

[illegible]

administered in an amount of about 600 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg.

[0229] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, or about 1000 mg.

[0230] For oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing from about 1.0 mg to about 1,000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about

350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 980 mg, about 990 mg, about 990 mg, or about 1000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof for the symptomatic adjustment of the dosage to the patient to be treated. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 50 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 200 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. The pharmaceutical compositions can be administered on a regimen of one (1) to four (4) times per day, including once, twice, three times, and four times per day. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about

310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 980 mg, about 990 mg, about 995 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg,

about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered three times per day.

[0231] In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of one day. In some embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of two days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of three days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of four days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of five days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of six days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of seven days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of eight days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of nine days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of ten days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 11 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 12 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 13 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 14 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of 15 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 16 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 17 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

[illegible]

hydrate, or prodrug thereof is administered over a period of 34 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 35 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 36 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 37 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 38 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 39 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 40 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 41 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 42 days or more.

[0232] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days.

[0233] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least

about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 14 days.

{0234} In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily

for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily

for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 14 days.

[0235] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330

mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 28 days.

{0236} In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 220 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula

(I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 28 days.

{0237} In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 255 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230

mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 42 days.

[0238] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 5 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula

(I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 370 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 42 days.

[0239] In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, or about 400 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200

mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, or about 800 mg of the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered at least once per day.

[0240] In certain embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0241] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered orally.

[0242] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is formulated as a capsule or tablet.

[0243] In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.


[0244] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0245] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day.

[0246] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 6 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 5 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 3 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 1 hour before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep.

[0247] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

[0248] In another aspect is provided a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I): and a pharmaceutically acceptable excipient comprising with a compound as described herein, including embodiments, or the structural Formulae (I) and a pharmaceutically acceptable excipient:



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 , and R^4 are as described herein, including embodiments.

[0249] In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments,

the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

{0250} In certain embodiments, the pharmaceutical composition comprises about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

{0251} In certain embodiments, the pharmaceutical composition is administered orally.

{0252} In certain embodiments, the pharmaceutical composition is formulated as a capsule or tablet. In certain embodiments, the capsule or tablet comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, or about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 50 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 200 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

{0253} In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically

acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0254] In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0255] In certain embodiments, the pharmaceutical composition is administered in the fed state. In certain embodiments, the pharmaceutical composition is administered in the fasted state. In certain embodiments, the pharmaceutical composition is administered at least once per day.

[0256] In certain embodiments, the pharmaceutical composition is administered at bedtime. In certain embodiments, the pharmaceutical composition is administered less than about 6 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 5 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 3 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 2 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 1 hour before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep.


[0257] In certain embodiments, the pharmaceutical composition is administered in the evening. In certain embodiments, the pharmaceutical composition is administered at about 10 pm at night.

[0258] In certain embodiments, the subject has congenital adrenal hyperplasia (CAH).


[0259] In certain embodiments, the size and/or number of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced.

[0260] In certain embodiments of the methods disclosed herein, the method further comprises administering to the subject an additional chemotherapeutic agent.

[0261] In certain embodiments, the the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In certain embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In certain embodiments, the mineralocorticoid is fludrocortisone. In certain embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist. In certain embodiments, the CRF₁ antagonist is:



pharmaceutically acceptable salt thereof. In certain embodiments, the CRF₁ antagonist is



[0262] In certain embodiments of the methods disclosed herein, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof. In certain embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the surgical resection is prior

to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0263] In certain embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0264] In one embodiment, the pharmaceutical compositions are provided in a dosage form for oral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. The pharmaceutical compositions provided herein that are formulated for oral administration may be in tablet, capsule, powder, or liquid form. In some embodiments, a tablet comprises a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil, or synthetic oil. Physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol, or polyethylene glycol may be included. In some embodiments, a capsule comprises a solid carrier such as gelatin.

[0265] In another embodiment, the pharmaceutical compositions are provided in a dosage form for parenteral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. Where pharmaceutical compositions may be formulated for intravenous, cutaneous or subcutaneous injection, the active ingredient will be in the form of a parenterally acceptable aqueous solution, which is pyrogen-free and has a suitable pH, isotonicity, and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles, such as Sodium Chloride injection, Ringer's injection, or Lactated Ringer's injection. In some embodiments, preservatives, stabilisers, buffers, antioxidants, and/or other additives are included as required.

[0266] In yet another embodiment, the pharmaceutical compositions are provided in a dosage form for topical administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers.

[0267] The pharmaceutical compositions can also be formulated as modified release dosage forms, including delayed-, extended-, prolonged-, sustained-, pulsatile-, controlled-, accelerated-, fast-, targeted-, and programmed-release, and gastric retention dosage forms. These dosage forms can be prepared according to conventional methods and techniques known to those skilled in the art (*see, Remington: The Science and Practice of Pharmacy, supra; Modified-Release Drug Delivery Technology*, 2nd Edition, Rathbone *et al.*, Eds., Marcel Dekker, Inc.: New York, NY, 2008).

[0268] The pharmaceutical compositions provided herein can be provided in a unit-dosage form or multiple-dosage form. A unit-dosage form, as used herein, refers to physically discrete a unit suitable for administration to a human and animal subject, and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of an active ingredient(s) sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carriers or excipients. Examples of a unit-dosage form include an ampoule, syringe, and individually packaged tablet and capsule. A unit-dosage form may be administered in fractions or multiples thereof. A multiple-dosage form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dosage form. Examples of a multiple-dosage form include a vial, bottle of tablets or capsules, or bottle of pints or gallons.

[0269] The pharmaceutical compositions provided herein can be administered at once, or multiple times at intervals of time. It is understood that the precise dosage and duration of treatment may vary with the age, weight, and condition of the patient being treated, and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test or diagnostic data. It is further understood that for any particular individual, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations.

[0270] In certain embodiments, the pharmaceutical compositions provided herein further comprise one or more chemotherapeutic agents as defined herein.

A. Oral Administration

[0271] The pharmaceutical compositions provided herein for oral administration can be provided in solid, semisolid, or liquid dosage forms for oral administration. As used herein, oral administration also includes buccal, lingual, and sublingual administration. Suitable oral dosage

forms include, but are not limited to, tablets, fastmelts, chewable tablets, capsules, pills, strips, troches, lozenges, pastilles, cachets, pellets, medicated chewing gum, bulk powders, effervescent or non-effervescent powders or granules, oral mists, solutions, emulsions, suspensions, wafers, sprinkles, elixirs, and syrups. In addition to the active ingredient(s), the pharmaceutical compositions can contain one or more pharmaceutically acceptable carriers or excipients, including, but not limited to, binders, fillers, diluents, disintegrants, wetting agents, lubricants, glidants, coloring agents, dye-migration inhibitors, sweetening agents, flavoring agents, emulsifying agents, suspending and dispersing agents, preservatives, solvents, non-aqueous liquids, organic acids, and sources of carbon dioxide.

[0272] Binders or granulators impart cohesiveness to a tablet to ensure the tablet remaining intact after compression. Suitable binders or granulators include, but are not limited to, starches, such as corn starch, potato starch, and pre-gelatinized starch (e.g., STARCH 1500); gelatin; sugars, such as sucrose, glucose, dextrose, molasses, and lactose; natural and synthetic gums, such as acacia, alginic acid, alginates, extract of Irish moss, panwar gum, ghatti gum, mucilage of isabgol husks, carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone (PVP), Veegum, larch arabogalactan, powdered tragacanth, and guar gum; celluloses, such as ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose, methyl cellulose, hydroxyethylcellulose (HEC), hydroxypropylcellulose (HPC), hydroxypropyl methyl cellulose (HPMC); microcrystalline celluloses, such as AVICEL-PH-101, AVICEL-PH-103, AVICEL RC-581, AVICEL-PH-105 (FMC Corp., Marcus Hook, PA); and mixtures thereof. Suitable fillers include, but are not limited to, talc, calcium carbonate, microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The amount of a binder or filler in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The binder or filler may be present from about 50 to about 99% by weight in the pharmaceutical compositions provided herein.

[0273] Suitable diluents include, but are not limited to, dicalcium phosphate, calcium sulfate, lactose, sorbitol, sucrose, inositol, cellulose, kaolin, mannitol, sodium chloride, dry starch, and powdered sugar. Certain diluents, such as mannitol, lactose, sorbitol, sucrose, and inositol, when present in sufficient quantity, can impart properties to some compressed tablets that permit disintegration in the mouth by chewing. Such compressed tablets can be used as chewable tablets. The amount of a diluent in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art.

[0274] Suitable disintegrants include, but are not limited to, agar; bentonite; celluloses, such as methylcellulose and carboxymethylcellulose; wood products; natural sponge; cation-exchange

resins; alginic acid; gums, such as guar gum and Veegum HV; citrus pulp; cross-linked celluloses, such as croscarmellose; cross-linked polymers, such as crospovidone; cross-linked starches; calcium carbonate; microcrystalline cellulose, such as sodium starch glycolate; polacrilin potassium; starches, such as corn starch, potato starch, tapioca starch, and pre-gelatinized starch; clays; aligns; and mixtures thereof. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The pharmaceutical compositions provided herein may contain from about 0.5 to about 15% or from about 1 to about 5% by weight of a disintegrant.

[0275] Suitable lubricants include, but are not limited to, calcium stearate; magnesium stearate; mineral oil; light mineral oil; glycerin; sorbitol; mannitol; glycols, such as glycerol behenate and polyethylene glycol (PEG); stearic acid; sodium lauryl sulfate; talc; hydrogenated vegetable oil, including peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil; zinc stearate; ethyl oleate; ethyl laureate; agar; starch; lycopodium; silica or silica gels, such as AEROSIL® 200 (W.R. Grace Co., Baltimore, MD) and CAB-O-SIL® (Cabot Co. of Boston, MA); and mixtures thereof. The pharmaceutical compositions provided herein may contain about 0.1 to about 5% by weight of a lubricant.

[0276] Suitable glidants include, but are not limited to, colloidal silicon dioxide, CAB-O-SIL® (Cabot Co. of Boston, MA), and asbestos-free talc. Suitable coloring agents include, but are not limited to, any of the approved, certified, water soluble FD&C dyes, and water insoluble FD&C dyes suspended on alumina hydrate, and color lakes and mixtures thereof. A color lake is the combination by adsorption of a water-soluble dye to a hydrous oxide of a heavy metal, resulting in an insoluble form of the dye. Suitable flavoring agents include, but are not limited to, natural flavors extracted from plants, such as fruits, and synthetic blends of compounds which produce a pleasant taste sensation, such as peppermint and methyl salicylate. Suitable sweetening agents include, but are not limited to, sucrose, lactose, mannitol, syrups, glycerin, and artificial sweeteners, such as saccharin and aspartame. Suitable emulsifying agents include, but are not limited to, gelatin, acacia, tragacanth, bentonite, and surfactants, such as polyoxyethylene sorbitan monooleate (TWEEN® 20), polyoxyethylene sorbitan monooleate 80 (TWEEN® 80), and triethanolamine oleate. Suitable suspending and dispersing agents include, but are not limited to, sodium carboxymethylcellulose, pectin, tragacanth, Veegum, acacia, sodium carbomethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable preservatives include, but are not limited to, glycerin, methyl and propylparaben, benzoic add,

sodium benzoate and alcohol. Suitable wetting agents include, but are not limited to, propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate, and polyoxyethylene lauryl ether. Suitable solvents include, but are not limited to, glycerin, sorbitol, ethyl alcohol, and syrup. Suitable non-aqueous liquids utilized in emulsions include, but are not limited to, mineral oil and cottonseed oil. Suitable organic acids include, but are not limited to, citric and tartaric acid. Suitable sources of carbon dioxide include, but are not limited to, sodium bicarbonate and sodium carbonate.

[0277] It should be understood that many carriers and excipients may serve several functions, even within the same formulation.

[0278] The pharmaceutical compositions provided herein for oral administration can be provided as compressed tablets, tablet triturates, chewable lozenges, rapidly dissolving tablets, multiple compressed tablets, or enteric-coating tablets, sugar-coated, or film-coated tablets. Enteric-coated tablets are compressed tablets coated with substances that resist the action of stomach acid but dissolve or disintegrate in the intestine, thus protecting the active ingredients from the acidic environment of the stomach. Enteric-coatings include, but are not limited to, fatty acids, fats, phenyl salicylate, waxes, shellac, ammoniated shellac, and cellulose acetate phthalates. Sugar-coated tablets are compressed tablets surrounded by a sugar coating, which may be beneficial in covering up objectionable tastes or odors and in protecting the tablets from oxidation. Film-coated tablets are compressed tablets that are covered with a thin layer or film of a water-soluble material. Film coatings include, but are not limited to, hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000, and cellulose acetate phthalate. Film coating imparts the same general characteristics as sugar coating. Multiple compressed tablets are compressed tablets made by more than one compression cycle, including layered tablets, and press-coated or dry-coated tablets.

[0279] The tablet dosage forms can be prepared from the active ingredient in powdered, crystalline, or granular forms, alone or in combination with one or more carriers or excipients described herein, including binders, disintegrants, controlled-release polymers, lubricants, diluents, and/or colorants. Flavoring and sweetening agents are especially useful in the formation of chewable tablets and lozenges.

[0280] The pharmaceutical compositions provided herein for oral administration can be provided as soft or hard capsules, which can be made from gelatin, methylcellulose, starch, or calcium alginate. The hard gelatin capsule, also known as the dry-filled capsule (DFC), consists of two sections, one slipping over the other, thus completely enclosing the active ingredient. The soft elastic capsule (SEC) is a soft, globular shell, such as a gelatin shell, which is plasticized by the addition of glycerin, sorbitol, or a similar polyol. The soft gelatin shells may contain a

preservative to prevent the growth of microorganisms. Suitable preservatives are those as described herein, including methyl- and propyl-parabens, and sorbic acid. The liquid, semisolid, and solid dosage forms provided herein may be encapsulated in a capsule. Suitable liquid and semisolid dosage forms include solutions and suspensions in propylene carbonate, vegetable oils, or triglycerides. Capsules containing such solutions can be prepared as described in U.S. Pat. Nos. 4,328,245; 4,409,239; and 4,410,545. The capsules may also be coated as known by those of skill in the art in order to modify or sustain dissolution of the active ingredient.

[0281] Coloring and flavoring agents can be used in all of the above dosage forms.

[0282] The pharmaceutical compositions provided herein for oral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

B. Parenteral Administration

[0283] The pharmaceutical compositions provided herein can be administered parenterally by injection, infusion, or implantation, for local or systemic administration. Parenteral administration, as used herein, include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular, intrasynovial, intravesical, and subcutaneous administration.

[0284] The pharmaceutical compositions provided herein for parenteral administration can be formulated in any dosage forms that are suitable for parenteral administration, including solutions, suspensions, emulsions, micelles, liposomes, microspheres, nanosystems, and solid forms suitable for solutions or suspensions in liquid prior to injection. Such dosage forms can be prepared according to conventional methods known to those skilled in the art of pharmaceutical science (*see, Remington: The Science and Practice of Pharmacy, supra*).

[0285] The pharmaceutical compositions intended for parenteral administration can include one or more pharmaceutically acceptable carriers and excipients, including, but not limited to, aqueous vehicles, water-miscible vehicles, non-aqueous vehicles, antimicrobial agents or preservatives against the growth of microorganisms, stabilizers, solubility enhancers, isotonic agents, buffering agents, antioxidants, local anesthetics, suspending and dispersing agents, wetting or emulsifying agents, complexing agents, sequestering or chelating agents, cryoprotectants, lyoprotectants, thickening agents, pH adjusting agents, and inert gases.

[0286] Suitable aqueous vehicles include, but are not limited to, water, saline, physiological saline or phosphate buffered saline (PBS), sodium chloride injection, Ringers injection, isotonic dextrose injection, sterile water injection, dextrose and lactated Ringers injection. Suitable non-aqueous vehicles include, but are not limited to, fixed oils of vegetable origin, castor oil, corn oil, cottonseed oil, olive oil, peanut oil, peppermint oil, safflower oil, sesame oil, soybean oil,

hydrogenated vegetable oils, hydrogenated soybean oil, and medium-chain triglycerides of coconut oil, and palm seed oil. Suitable water-miscible vehicles include, but are not limited to, ethanol, 1,3-butanediol, liquid polyethylene glycol (e.g., polyethylene glycol 300 and polyethylene glycol 400), propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, *N,N*-dimethylacetamide, and dimethyl sulfoxide.

{0287} Suitable antimicrobial agents or preservatives include, but are not limited to, phenols, cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl *p*-hydroxybenzoates, thimerosal, benzalkonium chloride (e.g., benzethonium chloride), methyl-and propyl-parabens, and sorbic acid. Suitable isotonic agents include, but are not limited to, sodium chloride, glycerin, and dextrose. Suitable buffering agents include, but are not limited to, phosphate and citrate. Suitable antioxidants are those as described herein, including bisulfite and sodium metabisulfite. Suitable local anesthetics include, but are not limited to, procaine hydrochloride. Suitable suspending and dispersing agents are those as described herein, including sodium carboxymethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable emulsifying agents are those described herein, including polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate 80, and triethanolamine oleate. Suitable sequestering or chelating agents include, but are not limited to EDTA. Suitable pH adjusting agents include, but are not limited to, sodium hydroxide, hydrochloric acid, citric acid, and lactic acid. Suitable complexing agents include, but are not limited to, cyclodextrins, including α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, sulfobutylether- β -cyclodextrin, and sulfobutylether 7- β -cyclodextrin (CAPTISOL®, CyDex, Lenexa, KS).

{0288} When the pharmaceutical compositions provided herein are formulated for multiple dosage administration, the multiple dosage parenteral formulations must contain an antimicrobial agent at bacteriostatic or fungi static concentrations. All parenteral formulations must be sterile, as known and practiced in the art.

{0289} In one embodiment, the pharmaceutical compositions for parenteral administration are provided as ready-to-use sterile solutions. In another embodiment, the pharmaceutical compositions are provided as sterile dry soluble products, including lyophilized powders and hypodermic tablets, to be reconstituted with a vehicle prior to use. In yet another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile suspensions. In yet another embodiment, the pharmaceutical compositions are provided as sterile dry insoluble products to be reconstituted with a vehicle prior to use. In still another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile emulsions.

[0290] The pharmaceutical compositions provided herein for parenteral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

[0291] The pharmaceutical compositions provided herein for parenteral administration can be formulated as a suspension, solid, semi-solid, or thixotropic liquid, for administration as an implanted depot. In one embodiment, the pharmaceutical compositions provided herein are dispersed in a solid inner matrix, which is surrounded by an outer polymeric membrane that is insoluble in body fluids but allows the active ingredient in the pharmaceutical compositions diffuse through.

[0292] Suitable inner matrixes include, but are not limited to, polymethylmethacrylate, polybutyl-methacrylate, plasticized or unplasticized polyvinylchloride, plasticized nylon, plasticized polyethylene terephthalate, natural rubber, polyisoprene, polyisobutylene, polybutadiene, polyethylene, ethylene-vinyl acetate copolymers, silicone rubbers, polydimethylsiloxanes, silicone carbonate copolymers, hydrophilic polymers, such as hydrogels of esters of acrylic and methacrylic acid, collagen, cross-linked polyvinyl alcohol, and cross-linked partially hydrolyzed polyvinyl acetate.

[0293] Suitable outer polymeric membranes include but are not limited to, polyethylene, polypropylene, ethylene/propylene copolymers, ethylene/ethyl acrylate copolymers, ethylene/vinyl acetate copolymers, silicone rubbers, polydimethyl siloxanes, neoprene rubber, chlorinated polyethylene, polyvinylchloride, vinyl chloride copolymers with vinyl acetate, vinylidene chloride, ethylene and propylene, ionomer polyethylene terephthalate, butyl rubber epichlorohydrin rubbers, ethylene/vinyl alcohol copolymer, ethylene/vinyl acetate/vinyl alcohol terpolymer, and ethylene/vinylalcohol copolymer.

V. Examples

[0294] While preferred embodiments of the present disclosure have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.


Example 1. A Phase 2, Open-Label Study to Evaluate the Efficacy and Safety of Compound 1 in Subjects with Testicular Adrenal Rest Tumors

[0295] Testicular adrenal rest tumors (TART) is a common complication in congenital adrenal hyperplasia (CAH) that is caused by adrenocorticotropin hormone (ACTH)–driven

overstimulation of aberrant adrenal cells within the testes and results in impaired spermatogenesis and infertility. TARTS can be found anywhere on the testes, though a typical (very common) location is within the mediastinum. This is a particularly problematic location. Because of the central location of TARTs being in the mediastinum, compression of the nearby seminiferous tubules is common and can lead to obstructive azoospermia, irreversible damage to surrounding tissue, and permanent infertility in males.

[0296] Current treatment for TART consists of chronic high-dose glucocorticoids, a problematic therapy with significant side effects and a narrow therapeutic window. Given the serious nature of TART and the limitations and risks of chronic steroid therapy, new treatment modalities are needed for patients with TART. Compound 1, a high-affinity and selective small-molecule antagonist of the corticotropin-releasing factor type-1 (CRF₁) receptor, may be effective in blocking ACTH synthesis, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of glucocorticoids. This mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-hydroxyprogesterone (17-OHP) in subjects with CAH. Compound 1 has been shown to have an acceptable safety and tolerability profile in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. These promising results provide a rationale for investigating the use of Compound 1 in the treatment of TART, in combination with replacement glucocorticoids and mineralocorticoids.

[0297] The structure of Compound 1 is:



[0298] Building upon the 2 completed Phase 1 studies of Compound 1 in healthy volunteers and the ongoing Phase 2 proof-of-concept study of Compound 1 in subjects with CAH, this Phase 2 will study Compound 1 specifically in male CAH subjects with TART for the first time. A sample size of approximately 12 subjects was selected to provide sufficient data for an initial evaluation of efficacy and safety in subjects with TART while maintaining feasible subject recruitment goals for this rare condition. Given the rarity of the disease and thus the small sample size that can be recruited, this study will be open label, and all enrolled subjects will

receive active study drug. Subjects will serve as their own controls in that their outcome measures after 12 weeks of study drug dosing will be compared to their baseline measures.

[0299] The schedule of study visits (every 2 weeks through Week 4, then every 4 weeks for the remainder of the study) provides an appropriate balance between minimizing subject burden and closely monitoring subject safety and response to therapy. The 4-week post-treatment washout period will allow for adequate safety follow-up and analysis of any HPA axis rebound after the last dose of study drug.

[0300] The specification of study drug dosing at 10pm and after consumption of a snack is based on information from the Phase 1 studies in healthy volunteers. The plasma C_{max} (maximum concentration) for Compound 1 in the Phase 1 studies occurred approximately 5 hours after dosing. Therefore, Compound 1 will be dosed daily at 10pm in this study in order to achieve peak plasma study drug concentration at the time of day when most adrenal androgens are produced. Subjects will consume a standardized snack before taking study drug because the Phase 1 studies showed minimal study drug absorption when subjects were dosed in a fasted state. Specific dose reduction criteria, strict individual treatment stopping criteria, and appropriate laboratory and medical assessments have been established to ensure subject safety. (FIGS. 1A-1B).

Objectives	Endpoints
Primary	
<ul style="list-style-type: none"> To assess the efficacy of Compound 1 in subjects with TART 	<ul style="list-style-type: none"> Reduction in the size/volume of TART on testicular ultrasound from screening to Week 12
Secondary	
<ul style="list-style-type: none"> To evaluate the safety of Compound 1 in subjects with TART To assess the pharmacodynamic (PD) biomarker effects of Compound 1 	<ul style="list-style-type: none"> Adverse events (AEs), serious adverse events (SAEs), AEs leading to withdrawal, AEs of special interest (AESIs) Physical examination findings Vital signs Changes in electrocardiogram (ECG) Changes in clinical laboratory values Changes in plasma concentrations of key PD biomarkers, including ACTH, 17-OHP, testosterone, androstenedione, luteinizing hormone (LH),

- To assess changes on semen analysis
 - To assess changes in metabolism
 - To evaluate changes in quality of life (QoL) and mood
- follicle-stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, and aldosterone from baseline to Week 12
 - Changes in sperm count, morphology, and motility from baseline to Week 12
 - Changes in metabolic parameters, including fasting glucose, HbA1c, insulin sensitivity, and lipid profile (including LDL cholesterol, HDL cholesterol, and triglycerides), from baseline to Week 12
 - Changes in body weight, body mass index (BMI), waist circumference, and blood pressure
 - Changes in QoL measured using the Short Form 36 (SF-36)
 - Changes in mood measured using the Hospital Anxiety and Depression Scale (HADS)

Study Population

[0301] Approximately 12 male subjects with TART will be enrolled at approximately 6 investigative sites in the United States, United Kingdom, and possibly other countries. Up to 12 additional subjects may be added to further evaluate specific safety, PD, or efficacy findings.

Eligibility Criteria

Inclusion Criteria

- Adult male subjects aged 18 to 30, inclusive
- Documented historical diagnosis of CAH based on documented genetic mutation or historical documentation of elevated 17-OHP
- Diagnosis of TART via testicular ultrasound and/or magnetic resonance imaging (MRI) at screening with ≥ 1 lesion(s) measuring ≥ 2 cm in diameter
- On a stable regimen of GC replacement for ≥ 30 days before baseline
- Agree to follow contraception guidelines and refrain from donating sperm throughout the treatment period and for 90 days after the last dose of study drug
- Subject is able to understand all study procedures and risks involved and provides written informed consent indicating willingness to comply with all aspects of the protocol

Exclusion Criteria

- Any clinically significant unstable medical condition, medically significant illness, or chronic disease within 30 days of screening, including but not limited to:

- A malignancy or <3 years of remission history from any malignancy, other than successfully treated localized skin cancer
- Presence of clinically significant renal disease, as evidenced by an estimated glomerular filtration rate (eGFR) of <60 mL/min/1.73m²
- Current or chronic history of liver disease or known hepatic or biliary abnormalities (with the exception of Gilbert's syndrome or asymptomatic gallstones)
- Confirmed positive test at screening for active hepatitis B, hepatitis C, or human immunodeficiency virus (HIV)
- Any clinically significant psychiatric disorders, including any current major depressive episode, bipolar disorder, schizophrenia, schizoaffective disorder, major depressive disorder with psychotic features, or any other psychotic disorder within the preceding 6 months
- HADS score >12 for either depression or anxiety
- At increased risk of suicide on the basis of the Investigator's judgment or the results of the Columbia-Suicide Severity Rating Scale (C-SSRS) conducted at screening and baseline (eg, C-SSRS Type 3, 4, or 5 ideation during the preceding 6 months or any suicidal behavior within the past 12 months)
- Clinically significant and abnormal clinical or laboratory assessments must be discussed with the Medical Monitor to determine eligibility for this study. Abnormal assessments that must be reviewed to determine eligibility include but are not limited to:
 - Clinically meaningful abnormal ECG results, in the opinion of the Investigator
 - Fridericia-corrected QT interval (QTcF) >450 msec
 - Alanine aminotransferase (ALT) >2x upper limit of normal (ULN)
 - Bilirubin >1.5x ULN (isolated bilirubin >1.5x ULN is acceptable if bilirubin is fractionated and direct bilirubin is <35%)
- A history that includes bilateral adrenalectomy or hypopituitarism
- Use of any other investigational drug within 30 days or 5 half-lives, whichever is longer, before initial screening
- Require use of prohibited concomitant medications, including strong inhibitors and/or inducers of CYP3A4 (with the exception of glucocorticoids, hydrocortisone, and dexamethasone) within 30 days or 5 half-lives (whichever is longer) of first dose of study drug. Medications that are sensitive substrates or substrates with narrow therapeutic ranges (metabolized by CYP3A4, 2C8, 2C9, or 2C19) should be discussed on a case-by-case basis with the Medical Monitor to determine whether the medication should be discontinued or may be continued with caution. If washout is feasible, then the medication should be

withdrawn at least 30 days or 5 half-lives (whichever is longer) before the first dose of study drug.

- Donation of blood within 60 days before first dose of study drug, or donation of platelets, white blood cells, or plasma within 15 days before first dose of study drug.

Study Duration

[0302] The expected duration of study participation for each subject is up to approximately 20 weeks. This includes a screening period of ≤ 4 weeks, a treatment period of 12 weeks, and a safety follow-up period of 4 weeks.

Study Design

[0303] This is a Phase 2, open-label efficacy and safety study of Compound 1 for the treatment of TART. All subjects will receive a daily oral dose of Compound 1 for 12 weeks. In-clinic study visits will occur for screening and at Day 1 (baseline) and Weeks 2, 4, 8, 12 (end of treatment), and 16 (follow-up).

Dose, Route, Regimen

[0304] All subjects will receive a daily oral dose of Compound 1 for 12 weeks. Subjects will receive 600 mg/day during the first 2 weeks and 1000 mg/day during the remaining 10 weeks of the treatment period. Study drug will be taken daily at 10pm (or bedtime, if earlier), 5 to 15 minutes after consumption of a standardized snack.

[0305] On mornings of study visits with 8am blood draws for laboratory assessments, subjects should take their morning glucocorticoid replacement after the blood draws. For all other days during the study, subjects may take their morning glucocorticoid replacement at their usual time (generally between 6am and 8am).

[0306] If a subject experiences a Grade 3 or higher TEAE considered at least possibly related to study drug, the Investigator may reduce the subject's daily dose of Compound 1 down in increments of 200 mg. Study drug will be discontinued in subjects who experience clinically significant liver chemistry, QTcF, or other individual treatment stopping criteria.

Investigational Drug

[0307] Compound 1 is a small-molecule CRF₁ receptor antagonist and will be supplied as 200-mg, size 1, white, hard gelatin capsules.

Example 2.

[0308] Since elevated ACTH is the primary driver for TART growth, blocking ACTH synthesis using Compound 1, a small-molecule antagonist of the CRF₁ receptor, may be effective in reducing ACTH levels, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of GCs. In CAH, this mechanism of action has been

validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH. The safety profile of Compound 1 has been established in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. A Phase 2 study in subjects with CAH is currently ongoing.

Nonclinical Toxicology of Compound 1

[0309] The nonclinical safety profile of Compound 1 has been evaluated in repeat-dose range-finding and definitive toxicology studies of up to 91 days in duration in rats and dogs using the clinically relevant route (oral) and schedule of administration (once daily). In rats and beagle dogs administered oral Compound 1 once daily for 91 consecutive days at doses up to 2000 mg/kg/day, toxicologic findings were all mild and limited to the male reproductive tract, and all Compound 1-related findings were reversible following a 6-week treatment-free recovery period.

[0310] Additionally, a fertility toxicology study was conducted in rats. In this study, male rats were administered oral Compound 1 once daily at 0, 5, 20, or 1000 mg/kg for 85 to 87 days (70 days before mating, during mating, and ≤ 4 days between mating and necropsy). Male reproductive performance (mean day of mating, mating and fertility indices, and conception rates) was unaffected by Compound 1 treatment. Untreated females mated with the Compound 1-treated males had no ovarian and uterine effects.

[0311] In addition, a complete battery of standard genotoxicity tests consisting of in vitro bacterial reverse mutation and chromosome aberration assays and an in vivo (rat) bone marrow micronucleus assay has been conducted. Exploratory and definitive embryo-fetal development studies in rabbits and a bovine corneal opacity and permeability assay have also been conducted.

Clinical Experience with Compound 1

[0312] To date, a total of 53 subjects, all healthy adults aged 21 to 65 years, have been treated with at least 1 dose of Compound 1 in 2 Phase 1 studies. The first-in-human study of Compound 1 (Study I3C-FW-BLAA) evaluated single ascending doses of 2 mg to 800 mg. In this study, a small number of subjects experienced diarrhea and headache of mild or moderate severity. A subsequent study (Study I3C-FW-BLAB) evaluated repeat doses of 50 mg, 100 mg, or 200 mg Compound 1 once daily for 14 consecutive days. A small number of subjects in the repeat-dose study experienced dyspnea, rhinorrhea, palpitations, and headache. The Phase 1 studies showed no clinically significant alterations in safety laboratory values, vital signs, or electrocardiograms (ECGs). Compound 1 has been well tolerated in healthy human subjects, with no SAEs, AEs leading to discontinuation, or dose-limiting toxicities (DLTs). A maximum tolerated dose has not been reached in humans.

[0313] An ongoing Phase 2 open-label study (Study Compound 2) is currently evaluating the safety and efficacy of Compound 1 dosed once daily for 6 weeks in adults with classic CAH. In the initial Cohort A of this study, all subjects are dosed starting at 200 mg/day for 14 days, escalating for each subject to 600 mg/day for 14 days, then 1000 mg/day for 14 days if the subject experiences no DLTs at the previous dose level. (**FIGS. 2-3**).

Known Potential Risks

[0314] Because the reversible testicular effects (degeneration/atrophy) observed in rats and dogs treated with high-dose Compound 1 (2,000 mg/kg/day) are likely associated with on-target (mechanism-driven) effects, these effects are likely to be relevant to human risk. However, male reproductive studies in rodents showed no impairment in fertility at the highest dose tested (1,000 mg/kg/day). Additionally, there were no effects on clinical laboratory markers of testicular function (LH and FSH) when Compound 1 was administered to healthy male volunteers at doses up to 200 mg per day for 14 consecutive days. Clinical laboratory measures of testicular function (LH, FSH, inhibin B, and SHBG) will be assessed throughout this study to monitor the risk of testicular injury. Testicular ultrasounds and semen analysis will also be performed at screening/baseline and the end of the treatment period.

[0315] Laboratory abnormalities in liver function tests (LFTs) have been observed at significantly higher doses in nonclinical studies. However, these events were not observed in the Phase 1 clinical studies. Clinical laboratory measures of liver chemistry will be assessed throughout this study, and strict individual treatment stopping criteria for liver chemistry findings are in place.

[0316] Compound 1-induced effects on the thyroid gland were noted in dogs after 91 days of high-dose (2,000 mg/kg/day) treatment but not in any other nonclinical studies. Clinical pathology parameters of thyroid function (eg, T4, T3, thyroid-stimulating hormone [TSH]) were not evaluated in the Phase 1 studies but will be monitored throughout this study.

Known Potential Benefits

[0317] Early clinical studies suggests that Compound 1 is effective as a CRF₁ receptor antagonist, a mechanism of action that supports the clinical investigation of Compound 1 for treatment of TART. In CAH, this mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH.

Assessment of Potential Risks and Benefits

[0318] The mild and reversible toxicologic effects of Compound 1 on the male reproductive tract in nonclinical studies occurred in the context of animals with normal, rather than elevated, baseline hormone function. As a CRF₁ receptor antagonist, Compound 1 is intended to reduce

abnormally elevated ACTH levels, which in TART drives overstimulation of aberrant adrenal cells within the testes. In nonclinical models where baseline hormone levels are within normal ranges, clinical benefits are not expected, and indeed, toxicities may be expected as a result of androgen reduction outside of the normal range. In TART, where baseline ACTH and androgen levels are pathologically elevated, androgen reduction would be considered a positive, therapeutic effect.

[0319] Given the serious nature of TART and the limitations (and risks) of chronic steroid therapy, new treatment modalities are needed for patients with TART. Given the acceptable safety and tolerability profile of Compound 1 in healthy volunteers from the Phase 1 program, the low probability of demonstrating meaningful PD marker changes in healthy subjects, and the lack of any clear benefit to exposing more healthy subjects to study drug, the Sponsor believes the benefit-to-risk profile favors the entry of Compound 1 into Phase 2 in subjects with TART.

Example 3. Congenital Adrenal Hyperplasia (CAH)

[0320] CAH results from a genetic defect in the 21-hydroxylase enzyme. CAH patients are thus unable to produce cortisol. Without cortisol to provide negative feedback, patients over-produce ACTH. High ACTH promotes the overproduction of potent androgens (male sex hormones). Without wishing to be bound by theory, Compound 1 blocks CRF₁ receptor to prevent ACTH release by the pituitary and associated drive on the hypothalamic-pituitary-adrenal axis. Replacement of cortisol to physiological levels will correct cortisol deficiency, however, will not lower ACTH, resulting in hyperplasia of adrenal gland and associated overproduction of adrenal androgens.

[0321] As shown in **FIGS. 4** and **5A-5B**, the attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism. 80% of subjects demonstrated reduction in ACTH. 70% of subjects demonstrated > 25% reduction in ACTH. 30% of subjects were in the normal range (*) after treatment with a CRF₁ receptor antagonist.

[0322] **FIGS. 6** and **7A-7B** demonstrate “control” of disease based on standard guidelines, which would then allow steroid taper. 80% of subjects demonstrated reduction in 17-OHP. 50% of subjects demonstrated > 25% reduction in ACTH. 50% of subjects were in the guideline range (*; 1200 ng/dL) after treatment with a CRF₁ receptor antagonist.

[0323] **FIGS. 8** and **9A-9B** demonstrate attenuation of morning rise in A4 upon treatment with a CRF₁ antagonist, indicative of its ability to control excess androgen production and associated symptoms. 100% of subjects demonstrated reduction in androstenedione (at various doses). 60% of subjects demonstrated > 25% reduction in androstenedione. 50% of subjects were within the normal reference range after treatment with a CRF₁ receptor antagonist.

[0324] FIGS. 10A-10B show approximately 25% shrinkage in TART in a 25 year old male with classic congenital adrenal hyperplasia following 6 weeks treatment with a CRF₁ receptor antagonist.

[0325] FIGS. 11A-11B demonstrate Compound 1 is a potent, selective, and functional CRF₁ receptor antagonist. In in vitro and ex vivo studies, Compound 1 was determined to be a potent and selective CRF₁ receptor antagonist with high affinity and selective for the hCRF₁ receptor. Compound 1 was found to have functional potency to inhibit adenylate cyclase activity (CRF₁ receptor-mediated downstream cascade), demonstrating in vitro efficacy as a CRF₁ receptor antagonist. Compound 1 was found to significantly inhibit CRF-induced locomotor behavior in response to i.c.v. administration of CRF (**FIG. 11B**).

[0326] FIGS. 12A-12B demonstrate single pill of CRF₁ receptor antagonist improves ACTH and 17-OHP in CAH patients.

[0327] FIG. 13 shows the subject level hormone changes (ACTH, 17-OHP, and androstenedione) by dose of a CRF₁ receptor antagonist.

Example 4. Assess reduction in excess ACTH and adrenal steroid production

[0328] The objectives of the study are: to assess the ability of Compound 1 to reduce excess ACTH and adrenal steroid production in participants with classic CAH on a stable glucocorticoid regimen; to evaluate the safety and tolerability of tildacerfont in participants with classic CAH; and to evaluate the pharmacokinetics (PK) of tildacerfont.

ELIGIBILITY CRITERIA

[0329] Inclusion Criteria:

- Adult males and females 18 years or older
- Diagnosis of classic CAH due to 21-hydroxylase deficiency
- Morning 17OHP $\geq$ 800 ng/dL, before taking any morning GC dose
- On a stable regimen of glucocorticoid replacement for $\geq$ 30 days before enrollment

[0330] Exclusion Criteria:

- Clinically significant unstable medical condition, illness, or chronic disease within 30 days of screening
- Clinically significant psychiatric disorder
- History of bilateral adrenalectomy or hypopituitarism

STUDY DESIGN

[0331] FIG. 14 depicts the proof of concept and escalating QD dosing at 10 PM for Cohort A and the single dose level and BID dosing at 10 AM and 10 pm for Cohort B.

BASELINE CHARACTERISTICS

[0332] Table 1 provides a description of the baseline characteristics displayed by Cohort A and Cohort B. In Cohort A, 10 participants were enrolled and received at least 1 dose of Compound 1. One participant withdrew early because of an occupation-related scheduling conflict. In Cohort B, 9 participants were enrolled and received at least one dose of Compound 1. One participant was withdrawn upon discovery of a major protocol violation at Baseline and did not have postbaseline efficacy/PK data. Thus, baseline characteristics are calculated for n=8 in Cohort B. One participant in Cohort B was taking both HC and prednisone.

Table 1 – Baseline characteristics of each cohort

Parameter	Cohort A (n=10)	Cohort B (n=8)
Demographics		
Age (years), mean (range)	39.3 (24-66)	36.8 (19-66)
Sex	5M, 5F	1M, 7F
Bmi (kg/m ²), mean $\pm$ SD	31.5 $\pm$ 12.0	31.1 $\pm$ 6.7
Daily GC dose (mg HC equivalents), mean range	28 (13-40)	35 (20-50)
Types of GCs	4 JC, 3 pred, 3 dex	2HC, 4 pred, 3 dex
17OHP at screening ^a (ng/dL), mean $\pm$ SD	5,985 $\pm$ 6,347	4,592 $\pm$ 4,624

Abbreviations: 17OHP: 17-hydroxyprogesterone; BMI: body mass index; dex: dexamethasone; F: female; GC: glucocorticoid; HC: hydrocortisone; M: male; pred: prednisone; SD: standard deviation.

^a 17OHP reference ranges: 27-199 ng/dL males, 15-290 ng/dL females.

SAFETY

[0333] Across both cohorts, Compound 1 was generally well tolerated. The most common type of adverse event (AE) was gastrointestinal disorders. There were no serious AEs and no AEs leading to withdrawal.

PHARMACOKINETICS

[0334] Over a wide range of doses (200 to 1000 mg QD), Compound 1 exhibited a predictable, dose-dependent PK profile.

EFFICACY

[0335] **FIG. 15A-15D** show the changes in levels of Androstenedione (A4) across both cohorts. Levels of A4 improved at ≥ 1 dose level for 10 out of 10 participants in Cohort A and 6 out of 8 participants in Cohort B. Excess A4 is associated with clinical sequelae in CAH. Attenuation of A4 demonstrates control of adrenal androgen.

[0336] **FIG. 16A-16D** show the changes in levels of 17-hydroxyprogesterone (17-OHP) across both cohorts. Levels of 17OHP improved at ≥ 1 dose level for 8 out of 10 participants in Cohort A and 6 out of 7 participants in Cohort B. 17OHP is a critical hormone in CAH diagnosis and management. Attenuation of 17OHP demonstrates control of adrenal steroid production.

[0337] **FIG. 17A-17D** show the changes in levels of adrenocorticotrophic hormone (ACTH) across both cohorts. Levels of ACTH improved at ≥ 1 dose level for 8 out of 10 participants in

Cohort A and 5 out of 7 participants in Cohort B. Attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism.

Table 2 – Percentage change from baseline in ACTH, 17OHP and A4.

	Cohort A			Cohort B
	200 mg QD (Week 2)	600 mg QD (Week 4)	1000 mg QD (Week 6)	200 mg BID (Week 2)
ACTH	-25.7 (±51.6)	-60.1 (±37.5)	-32.7 (±47.0)	-36.5 (±58.0)
17OHP	+22.2 (±172.3)	-23.7 (±64.2)	-13.8 (±74.7)	-47.8 (±63.3)
A4	-12.1 (±47.4)	-33.2 (±36.5)	-28.6 (±44.6)	-16.0 (±81.4)

CONCLUSIONS


[0338] Compound 1, a potent and selective CRF₁ receptor antagonist, offers a novel nonsteroidal oral treatment approach for CAH. Compound 1 was well tolerated and efficacious in reducing excess ACTH, 17OHP, and androstenedione in patients with classic CAH. Compound 1 exhibited a predictable, dose-dependent PK profile suitable for the target population.

[0339] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS


WHAT IS CLAIMED IS:

1. A method of treating menstrual irregularity in a subject, comprising administering to the subject a small molecule corticotropin-releasing factor type-1 (CRF₁) antagonist, or a pharmaceutically acceptable salt thereof, wherein the small molecule CRF₁ antagonist is:




or a pharmaceutically acceptable salt thereof.

2. The method of claim 1, wherein the small molecule CRF₁ antagonist is:



3. The method of claim 1, wherein the small molecule CRF₁ antagonist is:





, or a pharmaceutically acceptable salt thereof.


4. The method of claim 1, wherein the metabolic parameter is insulin sensitivity.
5. The method of claim 1, wherein the subject has congenital adrenal hyperplasia (CAH).
6. The method of claim 1, wherein the subject is an adolescent.
7. The method of claim 1, wherein the subject is at least 18 years or older.
8. The method of claim 1, wherein the small molecule CRF₁ antagonist is administered at a dose of about 10 mg/day to about 400 mg/day to the subject.
9. The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 50 mg.
10. The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 100 mg.
11. The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 200 mg.
12. The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 400 mg.
13. The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject once per day.
14. The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject twice per day.
15. The method of claim 13, wherein the subject is in a fed state.
16. The method of claim 14, wherein the subject is in a fed state.
17. The method of claim 1, wherein the small molecule CRF₁ antagonist is administered orally.
18. The method of claim 1, wherein the small molecule CRF₁ antagonist is formulated as a capsule or tablet.
19. The method of claim 1, wherein the administering reduces a plasma concentration of a biomarker comprising 17 α -hydroxyprogesterone (17-OHP), adrenocorticotrophic hormone (ACTH), testosterone, androstenedione (A4), luteinizing hormone (LH), follicle-stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, or aldosterone, or a combination thereof.

20. The method of claim 19, wherein the biomarker is selected from the group consisting of 17-OHP, ACTH, and A4.
21. The method of claim 19, wherein the plasma concentration of the biomarker is reduced by at least 10% from baseline.

ABSTRACT OF THE DISCLOSURE

Provided herein are compounds and pharmaceutical compositions for the prevention and treatment of testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART).

**FIG. 1A****FIG. 1B**

**FIG. 2**




FIG. 3







FIG. 4

**FIG. 5A****FIG. 5B****FIG. 6**




FIG. 7A




FIG. 7B








FIG. 8

**FIG. 9A****FIG. 9B****FIG. 10A****FIG. 10B**


In Vitro Binding Assay

FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B

SPR001-201 Hormone responses




FIG. 13

ACTH: 7.2-63.3 pg/mL 17-OHP: <1200 ng/dL A4: 41-262 ng/dL (M) 27-152 ng/dL (F)

SPR001-201 Hormone responses




FIG. 13 (Cont.)

ACTH: 7.2-63.3 pg/mL 17-OHP: <1200 ng/dL A4: 41-262 ng/dL (M) 27-152 ng/dL (F)
















FIG. 14

**FIG. 15A****FIG. 15B****FIG. 15C****FIG. 15D**

**FIG. 16A****FIG. 16B****FIG. 16C****FIG. 16D**

**FIG. 17A****FIG. 17B****FIG. 17C****FIG. 17D**



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC PAYMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 01/31/2025 07:18:35 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Colleen Mensching
PATENT CENTER #	69010935	AUTHORIZED BY	Victoria Cheng
CUSTOMER #	21971	FILING DATE	-
CORRESPONDENCE ADDRESS	-	FIRST NAMED INVENTOR	Alexis HOWERTON

Payment Information

PAYMENT METHOD DA / 232415	PAYMENT TRANSACTION ID E20251UJ24457420	PAYMENT AUTHORIZED BY Colleen Mensching
-------------------------------	--	--

FEE CODE	DESCRIPTION	ITEM PRICE(\$)	QUANTITY	ITEM TOTAL(\$)
1011	BASIC FILING FEE - UTILITY (PAPER FILING ALSO REQUIRES NON-ELECTRONIC FILING FEE UNDER 1.16(T))	350.00	1	350.00
1111	UTILITY PATENT APPL. SEARCH FEE	770.00	1	770.00
1311	PATENT APPL. EXAMINATION FEE	880.00	1	880.00
1817	REQUEST FOR PRIORITIZED EXAMINATION	4515.00	1	4515.00
1051	SURCHARGE- LATE FILING FEE, SEARCH FEE, EXAMINATION FEE, INVENTOR'S OATH OR DECLARATION, OR APPLICATION FILED WITHOUT AT LEAST ONE	170.00	1	170.00

CLAIM OR BY REFERENCE

1054	Non-DOC-X Filing Surcharge	430.00	1	430.00
1830	PROCESSING FEE, EXCEPT IN PROVISIONAL APPLICATIONS	150.00	1	150.00
1202	EACH CLAIM IN EXCESS OF 20	200.00	1	200.00
TOTAL AMOUNT:				\$7,465.00

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961
21971 7590 03/25/2025 WILSON SONSINI GOODRICH & ROSATI 650 PAGE MILL ROAD PALO ALTO, CA 94304-1050			EXAMINER PIHONAK, SARAH	
			ART UNIT 1627	PAPER NUMBER
			NOTIFICATION DATE 03/25/2025	DELIVERY MODE ELECTRONIC

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Notice of the Office communication was sent electronically on above-indicated "Notification Date" to the following e-mail address(es):

patentdocket@wsgr.com

<i>Decision Granting Request for Prioritized Examination (Track I)</i>	Application No. 19/043,264	Applicant(s) HOWERTON et al.	
	Examiner IRVIN DINGLE	Art Unit OPET	AIA (FITF) Status Yes
<p>1. THE REQUEST FILED <u>31 January 2025</u> IS GRANTED .</p> <p>The above-identified application has met the requirements for prioritized examination</p> <p>A. ☒ for an original nonprovisional application (Track I).</p> <p>B. ☐ for an application undergoing continued examination (RCE).</p> <p>2. The above-identified application will undergo prioritized examination. The application will be accorded special status throughout its entire course of prosecution until one of the following occurs:</p> <p>A. filing a <u>petition for extension of time</u> to extend the time period for filing a reply;</p> <p>B. filing an <u>amendment to amend the application to contain more than four independent claims, more than thirty total claims</u>, or a multiple dependent claim;</p> <p>C. filing a <u>request for continued examination</u> ;</p> <p>D. filing a notice of appeal;</p> <p>E. filing a request for suspension of action;</p> <p>F. mailing of a notice of allowance;</p> <p>G. mailing of a final Office action;</p> <p>H. completion of examination as defined in 37 CFR 41.102; or</p> <p>I. abandonment of the application.</p> <p>Telephone inquiries with regard to this decision should be directed to IRVIN DINGLE at (571)272-3210. In his/her absence, calls may be directed to Petition Help Desk at (571) 272-3282.</p>			
/IRVIN DINGLE/ Lead Paralegal Specialist, OPET			



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961
21971 7590 04/25/2025 WILSON SONSINI GOODRICH & ROSATI 650 PAGE MILL ROAD PALO ALTO, CA 94304-1050			EXAMINER PIHONAK, SARAH	
			ART UNIT 1627	PAPER NUMBER
			NOTIFICATION DATE 04/25/2025	DELIVERY MODE ELECTRONIC

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Notice of the Office communication was sent electronically on above-indicated "Notification Date" to the following e-mail address(es):

patentdocket@wsgr.com

Notice of Pre-AIA or AIA Status

The present application, filed on or after March 16, 2013, is being examined under the first inventor to file provisions of the AIA.

Priority

This application, filed 01/31/2025 is a Continuation of 18941832, filed 11/08/2024.
18941832 is a Continuation of 17667285, filed 02/08/2022.
17667285 is a Continuation of 17081694, filed 10/27/2020.
17081694 is a Continuation of PCT/US2019/029486, filed 04/26/2019.
PCT/US2019/029486 Claims Priority from Provisional Application 62822815, filed 03/23/2019;
and Provisional Application 62663951, filed 04/27/2018. The provisional applications don't
provide support to treating menstrual irregularities; therefore, the effective filing and priority
date of the claims is 4/26/2019.

Status of Claims

1. Claims 1-21 are currently pending. A track one status has been granted.
2. Claims 1-21 were examined and are rejected.

Claim Rejections-35 USC § 112

3. The following is a quotation of 35 U.S.C. 112(b):
(b) CONCLUSION.—The specification shall conclude with one or more claims particularly pointing out and distinctly claiming the subject matter which the inventor or a joint inventor regards as the invention.

The following is a quotation of 35 U.S.C. 112 (pre-AIA), second paragraph:

The specification shall conclude with one or more claims particularly pointing out and distinctly claiming the subject matter which the applicant regards as his invention.

Claim 4 is rejected under 35 U.S.C. 112(b) or 35 U.S.C. 112 (pre-AIA), second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject matter which the inventor or a joint inventor (or for applications subject to pre-AIA 35 U.S.C. 112, the applicant), regards as the invention.

Claim 4 depends directly from claim 1 and recites “wherein the metabolic parameter is insulin sensitivity”. However, claim 1 doesn’t mention or refer to a metabolic parameter, therefore, there is insufficient antecedent basis for the limitation of claim 4, and the claim is indefinite.

Claim Rejections-35 USC § 102

4. The following is a quotation of the appropriate paragraphs of 35 U.S.C. 102 that form the basis for the rejections under this section made in this Office action:

A person shall be entitled to a patent unless –

(a)(1) the claimed invention was patented, described in a printed publication, or in public use, on sale, or otherwise available to the public before the effective filing date of the claimed invention.


(a)(2) the claimed invention was described in a patent issued under section 151, or in an application for patent published or deemed published under section 122(b), in which the patent or application, as the case may be, names another inventor and was effectively filed before the effective filing date of the claimed invention.

Claim(s) 1-2 and 5-21 is/are rejected under 35 U.S.C. 102(a)(1) as being anticipated by Howerton et. al., WO 2019036503 A1, publ. 2/21/2019, international filing date 8/14/2018, cited in an IDS.

Claim(s) 1-2 and 5-21 is/are rejected under 35 U.S.C. 102(a)(2) as being anticipated by Howerton et. al., WO 2019036503 A1, publ. 2/21/2019, international filing date 8/14/2018.

The applied reference has a common inventor with the instant application. Based upon the earlier effectively filed date of the reference, it constitutes prior art under 35 U.S.C. 102(a)(2). This rejection under 35 U.S.C. 102(a)(2) might be overcome by: (1) a showing under 37 CFR 1.130(a) that the subject matter disclosed in the reference was obtained directly or indirectly from the inventor or a joint inventor of this application and is thus not prior art in accordance with 35 U.S.C. 102(b)(2)(A); (2) a showing under 37 CFR 1.130(b) of a prior public disclosure under 35 U.S.C. 102(b)(2)(B) if the same invention is not being claimed; or (3) a statement pursuant to 35 U.S.C. 102(b)(2)(C) establishing that, not later than the effective filing date of the claimed invention, the subject matter disclosed in the reference and the claimed invention were either owned by the same person or subject to an obligation of assignment to the same person or subject to a joint research agreement.

Howerton discloses pharmaceutical compositions comprising 3-(4-chloro-2-(morpholin-4-yl)thiazol-5-yl)-7-(1-ethylpropyl)-2,5-dimethylpyrazolo(1,5-a)pyrimidine, and methods of treating congenital adrenal hyperplasia (CAH) and menstrual irregularity comprising administering the compound in a pharmaceutical composition (para [0003-0004], [0041]):



Howerton discloses treatment of an adolescent from about 12 years of age (para [0039], [0170]), as well as treatment of an adult of about 20 years of age (para [0039]). Howerton discloses a dose of the above shown compound between about 5-500 mg. (para [0027]), which includes the range recited in instant claim 8. Howerton discloses administration of the compound in a pharmaceutical composition once or twice per day (para [0030]), and administration of the composition to a subject in the fed state (para [0012]). Howerton discloses oral administration in dosage forms inclusive of tablets and capsules (para [0076]). Howerton discloses administration of the compound reduces 17-OHP (para [0047-0048]) and ACTH, with 50% of subjects experiencing a greater than 25% reduction in 17-OHP (para [0165-0166]). Howerton as such anticipates the claims.

Claim Rejections-35 USC § 103

5. The following is a quotation of 35 U.S.C. 103 which forms the basis for all obviousness rejections set forth in this Office action:

A patent for a claimed invention may not be obtained, notwithstanding that the claimed invention is not identically disclosed as set forth in section 102, if the differences between the claimed invention and the prior art are such that the claimed invention as a whole would have been obvious before the effective filing date of the claimed invention to a person having ordinary skill in the art to which the claimed invention pertains. Patentability shall not be negated by the manner in which the invention was made.

The factual inquiries for establishing a background for determining obviousness under 35 U.S.C. 103 are summarized as follows:

1. Determining the scope and contents of the prior art.
2. Ascertaining the differences between the prior art and the claims at issue.
3. Resolving the level of ordinary skill in the pertinent art.


4. Considering objective evidence present in the application indicating obviousness or nonobviousness.

This application currently names joint inventors. In considering patentability of the claims the examiner presumes that the subject matter of the various claims was commonly owned as of the effective filing date of the claimed invention(s) absent any evidence to the contrary. Applicant is advised of the obligation under 37 CFR 1.56 to point out the inventor and effective filing dates of each claim that was not commonly owned as of the effective filing date of the later invention in order for the examiner to consider the applicability of 35 U.S.C. 102(b)(2)(C) for any potential 35 U.S.C. 102(a)(2) prior art against the later invention.

Claim(s) 1, 3-14, and 17-21 is/are rejected under 35 U.S.C. 103 as being unpatentable over Grigoriadis, WO 2015112642 A1, publ. 7/30/2015, cited in an IDS.

Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) and inhibition of ACTH release in these patients (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often exhibit 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance,

and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis exemplifies the following compound as a CRF1 receptor antagonist (p. 14, see SSR-125543):




. Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20).

It would have been prima facie obvious to one of ordinary skill in the art, before the effective filing date of the claims to have treated menstrual irregularity in a subject with CAH comprising administering the CRF1 receptor antagonist shown in instant claims 1 and 3, in consideration of Grigoriadis. Grigoriadis teaches CRF1 receptor antagonists for the treatment of CAH, including in females, and exemplifies the compound recited in instant claim 3. Grigoriadis further teaches subjects with CAH have elevated ACTH levels and suppression of ACTH elevation can prevent menstrual disturbances. As Grigoriadis further teaches administering a CRF1 receptor antagonist can also suppress elevated ACTH levels, it would have been prima facie obvious that this treatment would have similarly treated or prevented menstrual disturbances, i.e., menstrual irregularities in a female with CAH and elevated ACTH levels. One of ordinary skill in the art would have arrived at the claimed method of treatment and have had a reasonable expectation of success. Since menstruating females include adolescents and subjects aged 18 years and older, it would have been prima facie obvious to have treated menstrual irregularities in a female with CAH who is an adolescent, as recited by instant claim 6, or who is at least 18 years of age, as recited by instant claim 7.

As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14.

6. Claim(s) 1-2, 5-14, and 17-21 is/are rejected under 35 U.S.C. 103 as being unpatentable over Chen et. al., WO 2008036579 A1, publ. 3/27/2008, cited in an IDS in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015.

Chen teaches thiazole pyrazolopyrimidine derivatives as CRF1 antagonists (title & abstract; p. 1, lines 3-7). Chen further teaches CRF1 antagonists to have activity for treating numerous conditions, including congenital adrenal hyperplasia (p. 3, lines 11-19; p. 3, line 28-p. 4, line 6). Chen exemplifies the following compound as a CRF1 antagonist (Ex. 16, p. 37, line 19-p. 38, line 18):



Oral administration of the CRF1 antagonist in a pharmaceutical composition is taught (p. 7, lines 24-28).

Chen doesn't explicitly teach or suggest treating menstrual irregularity.

Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often exhibit 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual

disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20).

It would have been prima facie obvious to one of ordinary skill in the art, before the effective filing date of the claims to have arrived at the instantly claimed method of treating menstrual irregularity in a subject with CAH comprising administering a CRF1 receptor antagonist of instant claims 1-2, in consideration of Chen and Grigoriadis. Chen teaches the compound of instant claim 2 as a CRF1 receptor antagonist for treating CAH, while Grigoriadis also teaches treatment of CAH by administering a CRF1 receptor antagonist. Grigoriadis further teaches subjects with CAH have elevated ACTH levels and suppression of ACTH elevation can prevent menstrual disturbances. As Grigoriadis further teaches administering a CRF1 receptor

antagonist can also suppress elevated ACTH levels, it would have been prima facie obvious that this treatment would have similarly treated or prevented menstrual disturbances, i.e., menstrual irregularities in a female with CAH and elevated ACTH levels. One of ordinary skill in the art would have arrived at the claimed method of treatment and have had a reasonable expectation of success. Since menstruating females include female adolescents and female subjects aged 18 years and older, it would have been prima facie obvious to have treated menstrual irregularities in a female with CAH who is an adolescent, as recited by instant claim 6, or who is at least 18 years or older, as recited by instant claim 7.

The dose of CRF1 receptor antagonist from about 50-1000 mg. as taught by Grigoriadis overlaps with the amounts recited by instant claims 8-12. Therefore, it would have been prima facie obvious to one of ordinary skill in the art to have administered the CRF1 receptor antagonist taught by Chen within the same dosage range, for treating CAH in a subject having menstrual irregularities. As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14.

Claim Rejections-Nonstatutory Double Patenting

7. The nonstatutory double patenting rejection is based on a judicially created doctrine grounded in public policy (a policy reflected in the statute) so as to prevent the unjustified or improper timewise extension of the “right to exclude” granted by a patent and to prevent possible harassment by multiple assignees. A nonstatutory double patenting rejection is

appropriate where the conflicting claims are not identical, but at least one examined application claim is not patentably distinct from the reference claim(s) because the examined application claim is either anticipated by, or would have been obvious over, the reference claim(s). See, e.g., *In re Berg*, 140 F.3d 1428, 46 USPQ2d 1226 (Fed. Cir. 1998); *In re Goodman*, 11 F.3d 1046, 29 USPQ2d 2010 (Fed. Cir. 1993); *In re Longi*, 759 F.2d 887, 225 USPQ 645 (Fed. Cir. 1985); *In re Van Ornum*, 686 F.2d 937, 214 USPQ 761 (CCPA 1982); *In re Vogel*, 422 F.2d 438, 164 USPQ 619 (CCPA 1970); *In re Thorington*, 418 F.2d 528, 163 USPQ 644 (CCPA 1969).

A timely filed terminal disclaimer in compliance with 37 CFR 1.321(c) or 1.321(d) may be used to overcome an actual or provisional rejection based on nonstatutory double patenting provided the reference application or patent either is shown to be commonly owned with the examined application, or claims an invention made as a result of activities undertaken within the scope of a joint research agreement. See MPEP § 717.02 for applications subject to examination under the first inventor to file provisions of the AIA as explained in MPEP § 2159. See MPEP § 2146 *et seq.* for applications not subject to examination under the first inventor to file provisions of the AIA. A terminal disclaimer must be signed in compliance with 37 CFR 1.321(b).

The filing of a terminal disclaimer by itself is not a complete reply to a nonstatutory double patenting (NSDP) rejection. A complete reply requires that the terminal disclaimer be accompanied by a reply requesting reconsideration of the prior Office action. Even where the NSDP rejection is provisional the reply must be complete. See MPEP § 804, subsection I.B.1. For a reply to a non-final Office action, see 37 CFR 1.111(a). For a reply to final Office action, see 37

CFR 1.113(c). A request for reconsideration while not provided for in 37 CFR 1.113(c) may be filed after final for consideration. See MPEP §§ 706.07(e) and 714.13.

The USPTO Internet website contains terminal disclaimer forms which may be used. Please visit www.uspto.gov/patent/patents-forms. The actual filing date of the application in which the form is filed determines what form (e.g., PTO/SB/25, PTO/SB/26, PTO/AIA/25, or PTO/AIA/26) should be used. A web-based eTerminal Disclaimer may be filled out completely online using web-screens. An eTerminal Disclaimer that meets all requirements is auto-processed and approved immediately upon submission. For more information about eTerminal Disclaimers, refer to www.uspto.gov/patents/apply/applying-online/eterminal-disclaimer.

Claims 1-3, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-4, 7-8, 11-14, 17-18, and 21-22 of U.S. Patent No. 10849908 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH. In particular, the patented claims recite treating a human having CAH by administering a therapeutically effective amount of a CRF1 antagonist, resulting in a reduction of the ACTH level by at least 10% from baseline. The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5); although the patented claims don't recite treating menstrual irregularity as required by the instant claims, this patient population would have been prima facie obvious in view of Grigoriadis. Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH

as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method provides a reduction in ACTH levels, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented method by treating menstrual irregularities in a subject with CAH,

and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is overlapping (from 50-1600 mg/day in patented claim 2 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 8). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

8. Claims 1-3, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-4, 7-8, and 11-16 of U.S. Patent No. 11007201 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH. In particular, the patented claims recite treating a human having CAH by administering a therapeutically effective amount of a CRF1 antagonist, resulting in a reduction of the androstenedione (A4) level. The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5); although the patented claims don't recite treating menstrual irregularity as required by the instant claims, this patient population would have been prima facie obvious in view of Grigoriadis. Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract;

p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method of treating CAH by administering a CRF1 antagonist, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented method by

treating menstrual irregularities in a subject with CAH, and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is overlapping (from 50-1600 mg/day in patented claim 2 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 8). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

9. Claims 1-2, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-4, 7-8, and 11-16 of U.S. Patent No. 11311549 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH. In particular, the patented claims recite treating a human having CAH by administering a therapeutically effective amount of a CRF1 antagonist, resulting in a reduction of the androstenedione (A4) level. The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5); although the patented claims don't recite treating menstrual irregularity as required by the instant claims, this patient population would have been prima facie obvious in view of Grigoriadis. Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract;

p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method of treating CAH by administering a CRF1 antagonist, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented method by

treating menstrual irregularities in a subject with CAH, and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is overlapping (from 50-1600 mg/day in patented claim 2 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 8). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

10. Claims 1-3, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-4, 7-8, 11-16, and 19-24 of U.S. Patent No. 11344557 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH. In particular, the patented claims recite treating a human having CAH by administering a therapeutically effective amount of a CRF1 antagonist, resulting in a reduction of the ACTH level by at least 10% from baseline. The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5); although the patented claims don't recite treating menstrual irregularity as required by the instant claims, this patient population would have been prima facie obvious in view of Grigoriadis. Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal

hyperplasia (CAH) (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method provides a reduction in ACTH levels, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented

method by treating menstrual irregularities in a subject with CAH, and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is overlapping (from 50-1600 mg/day in patented claim 2 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 8). Both sets of claims further recite a reduction of 17-OHP levels in the treated subject (see patented claims 1, 19-24; and instant claims 19-21). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

11. Claims 1-2, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-4, 7-8, and 14-19 of U.S. Patent No. 11351177 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH. In particular, the patented claims recite treating a human having CAH by administering a therapeutically effective amount of a CRF1 antagonist, resulting in a reduction of the ACTH level by at least 10% from baseline. The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5); although the patented claims don't recite treating menstrual irregularity as required by the instant claims, this patient population would

have been prima facie obvious in view of Grigoriadis. Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method provides a

reduction in ACTH levels, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented method by treating menstrual irregularities in a subject with CAH, and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is overlapping (from 50-1600 mg/day in patented claim 2 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 8). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

12. Claims 1-3, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-15 and 19 of U.S. Patent No. 11304950 B2 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003; and Grigoriadis, WO 2015112642 A1, publ. 7/30/2015. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH (see patented claims 1-2). The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5). The instant and patented claims differ from each other as the patented claims are drawn to treating ovarian adrenal rest tumors in a

subject with CAH, and not menstrual irregularities. However, Stikkelbroeck teaches females with CAH can have irregular menstrual cycles due to adrenal androgen excess (p. 1030, 1st para). Stikkelbroeck further teaches females with CAH can develop adrenal rest tumors in the ovaries which replace normal ovarian tissue, causing impaired menstrual cycling (p. 1034, right col., last para-p. 1035, left col., top para). Therefore, it would have been prima facie obvious to have applied the instantly claimed method to treat menstrual irregularities in a subject with CAH and ovarian adrenal rest tumors, and have had a reasonable expectation of success.

Furthermore, Grigoriadis teaches corticotropin releasing factor-1 (CRF1) receptor antagonists for the treatment of congenital adrenal hyperplasia (CAH) (title & abstract; p. 1, lines 3-5). Grigoriadis teaches CAH as a group of autosomal recessive genetic disorders that cause little to no cortisol biosynthesis and often 21-hydroxylase deficiency; ACTH production is excessive (p. 1, line 6-p. 2, line 3). CAH affects both males and females (p. 7, lines 22-28). Grigoriadis teaches treatment of CAH typically involves glucocorticoids to establish sufficient cortisol production and for reducing elevated ACTH levels, and suppression of ACTH prevents menstrual disturbances in women (p. 7, line 29-p. 8, line 11). An issue with glucocorticoid therapy is that the doses required to achieve sufficient suppression of excess androgens are usually well above the normal physiologic dose, and over time glucocorticoid therapy can lead to increased cardiovascular risk factors, glucose intolerance, and decreased bone mineral density (p. 9, lines 1-12). Grigoriadis teaches treatment of human and non-human animals; a therapeutically effective dose is taught to range from about 50-1000 mg. (p. 24, line 3-p. 25, line 7). Grigoriadis teaches various routes of administration, including oral (p. 25, lines 15-21). Grigoriadis teaches administration of a CRF1 receptor antagonist as described results in a

clinically significant reduction of both ACTH and 17-OHP levels relative to placebo, including a reduction of at least 25% (p. 28, lines 10-19). Grigoriadis teaches pharmaceutical compositions comprising CRF1 receptor antagonists in the form of tablets or capsules (p. 28, line 20-p. 29, line 16). Grigoriadis further teaches the dose also depends on the patient's condition, stage of disease, severity of symptoms; although bedtime administration is exemplified, the timing of administration can also be adjusted depending on various factors (p. 25, line 22-p. 26, line 20). Therefore, as the patented claims recite the method provides a reduction in ACTH levels, and elevated ACTH levels are associated with menstrual irregularities as taught by Grigoriadis, it would have been prima facie obvious to have modified the patented method by treating menstrual irregularities in a subject with CAH and ovarian adrenal rest tumors, and have had a reasonable expectation of success. Additionally, the amount of CRF1 antagonist administered is the same (from 10-400 mg/day in patented claim 13 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & patented claim 15). Both sets of claims explicitly recite the CRF1 antagonist of instant claims 1-2 (see patented claim 11). As Grigoriadis teaches the dose and timing of administration of the CRF1 receptor antagonist can be adjusted based on various factors, it would have been prima facie obvious to have administered the CRF1 receptor antagonist once per day as recited by instant claim 13, or twice per day as recited by instant claim 14. For these reasons the instant and patented claims are obvious variants of each other and are not patentably distinct.

13. Claims 1-3, and 5-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 72-91 of copending application 18941832 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH (see copending claims 72 and 76). The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5). The instant and copending claims differ from each other as the copending claims are drawn to treating a benign tumor, including ovarian adrenal rest tumors in a subject with CAH (see claims 72 and 75), and not menstrual irregularities. However, Stikkelbroeck teaches females with CAH can have irregular menstrual cycles due to adrenal androgen excess (p. 1030, 1st para). Stikkelbroeck further teaches females with CAH can develop adrenal rest tumors in the ovaries which replace normal ovarian tissue, causing impaired menstrual cycling (p. 1034, right col., last para-p. 1035, left col., top para). Therefore, it would have been prima facie obvious to have applied the instantly claimed method to treat menstrual irregularities in a subject with CAH and ovarian adrenal rest tumors, and have had a reasonable expectation of success.

Additionally, the amount of CRF1 antagonist administered is overlapping (from 5-800 mg/day in copending claim 78 compared to 10-400 mg/day as recited by instant claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & copending claim 91); wherein the treated subject is in a fed state (copending claim 84 and instant claims 14-15); and wherein the CRF1 antagonist is administered twice per day (copending claim 83 & instant claim 14). Both sets of claims explicitly recite the same CRF1

antagonists (see copending claim 72 & instant claim 1); and reduction of the same biomarkers selected from 17-OHP, ACTH, A4, LH, FSH, SHBG, renin, or aldosterone (see instant claim 19 & copending claim 86). As CAH affects females, including women and adolescent females (see Stikkelbroeck, p. 1030, right col., last para-p. 1031, left col., top para), it would have been prima facie obvious to have applied the copending method to treat an adolescent female as recited by instant claim 6, and a female at least 18 years or older as recited by instant claim 7. For these reasons the instant and copending claims are obvious variants of each other and are not patentably distinct.

14. Claims 1-3, and 5-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-5 and 9-21 of copending application 19043275 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH (see copending claims 1 & 5). The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5). The instant and copending claims differ from each other as the copending claims are drawn to treating a benign tumor, including ovarian adrenal rest tumors in a subject with CAH (see copending claim 4), and not menstrual irregularities. However, Stikkelbroeck teaches females with CAH can have irregular menstrual cycles due to adrenal androgen excess (p. 1030, 1st para). Stikkelbroeck further teaches females with CAH can develop adrenal rest tumors in the ovaries which replace normal ovarian tissue, causing impaired menstrual cycling (p. 1034, right col., last para-p. 1035,

left col., top para). Therefore, it would have been prima facie obvious to have applied the instantly claimed method to treat menstrual irregularities in a subject with CAH and ovarian adrenal rest tumors, and have had a reasonable expectation of success.

Additionally, the amount of CRF1 antagonist administered is the same (10-400 mg/day as recited by instant claim 8 and copending claim 11); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & copending claim 21); wherein the treated subject is in a fed state (copending claims 18-19 and instant claims 14-15); and wherein the CRF1 antagonist is administered once per day (copending claim 16 & instant claim 13); and the CRF1 antagonist is administered twice per day (copending claim 17 & instant claim 14). Both sets of claims explicitly recite the same CRF1 antagonists (see copending claim 1 & instant claim 1); treatment of an adolescent (see copending claim 9 & instant claim 6); and treatment of a subject at least 18 years of age (copending claim 10 & instant claim 7). For these reasons the instant and copending claims are obvious variants of each other and are not patentably distinct.

15. Claims 1-3, 5-13, and 17-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-19 of copending application 19043303 in view of Stikkelbroeck et. al., *Fertility & Sterility*, vol. 80(4), pp. 1030-1036, publ. 2003. Although the claims at issue are not identical, they are not patentably distinct from each other because both sets of claims encompass administering a CRF1 antagonist to a subject having CAH (see copending claims 1 & 5). The instant claims similarly recite administering small molecule CRF1 antagonists to treat a subject having CAH (see instant claims 1 and 5). The instant and

compending claims differ from each other as the compending claims are drawn to treating ovarian adrenal rest tumors in a subject with CAH (see compending claims 1 & 5), and not menstrual irregularities. However, Stikkelbroeck teaches females with CAH can have irregular menstrual cycles due to adrenal androgen excess (p. 1030, 1st para). Stikkelbroeck further teaches females with CAH can develop adrenal rest tumors in the ovaries which replace normal ovarian tissue, causing impaired menstrual cycling (p. 1034, right col., last para-p. 1035, left col., top para). Therefore, it would have been prima facie obvious to have applied the instantly claimed method to treat menstrual irregularities in a subject with CAH and ovarian adrenal rest tumors, and have had a reasonable expectation of success.

Additionally, the amount of CRF1 antagonist administered is the same (10-400 mg/day as recited by instant claim 8 and compending claim 8); wherein the CRF1 antagonist is formulated as a capsule or tablet (see instant claim 18 & compending claim 16); wherein the CRF1 antagonist is administered once per day (compending claim 13 & instant claim 13); and the CRF1 antagonist is administered twice per day (compending claim 14 & instant claim 14). Both sets of claims explicitly recite the same CRF1 antagonists (see compending claim 1 & instant claim 1); treatment of an adolescent (see compending claim 9 & instant claim 6); and treatment of a subject at least 18 years of age (compending claim 10 & instant claim 7). As CAH affects females, including women and adolescent females (see Stikkelbroeck, p. 1030, right col., last para-p. 1031, left col., top para), it would have been prima facie obvious to have applied the compending method to treat an adolescent female as recited by instant claim 6, and a female at least 18 years or older as recited by instant claim 7. For these reasons the instant and compending claims are obvious variants of each other and are not patentably distinct.

Information Disclosure Statements

16. The IDS filed on 2/5/25 have been considered.

Conclusion

17. Claims 1-21 are rejected.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SARAH PIHONAK whose telephone number is (571)270-7710. The examiner can normally be reached Monday-Friday 9:00-5:30 EST.

Examiner interviews are available via telephone, in-person, and video conferencing using a USPTO supplied web-based collaboration tool. To schedule an interview, applicant is encouraged to use the USPTO Automated Interview Request (AIR) at <http://www.uspto.gov/interviewpractice>.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Kortney Klinkel can be reached on 571-270-5239. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of published or unpublished applications may be obtained from Patent Center. Unpublished application information in Patent Center is available to registered users. To file and manage patent submissions in Patent Center, visit: <https://patentcenter.uspto.gov>. Visit <https://www.uspto.gov/patents/apply/patent-center> for more information about Patent Center and <https://www.uspto.gov/patents/docx> for

Application/Control Number: 19/043,264
Art Unit: 1627

Page 31

information about filing in DOCX format. For additional questions, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

SARAH . PIHONAK
Primary Examiner
Art Unit 1627

/SARAH PIHONAK/
Primary Examiner, Art Unit 1627

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875				Application or Docket Number 19/043,264		Filing Date 01/31/2025		☐ To be Mailed	
ENTITY: ☒ LARGE ☐ SMALL ☐ MICRO									
APPLICATION AS FILED - PART I									
	(Column 1)	(Column 2)							
FOR	NUMBER FILED	NUMBER EXTRA			RATE (\$)			FEE (\$)	
☐ BASIC FEE (37 CFR 1.16(a), (b), or (c))	N/A	N/A			N/A				
☐ SEARCH FEE (37 CFR 1.16(k), (l), or (m))	N/A	N/A			N/A				
☐ EXAMINATION FEE (37 CFR 1.16(o), (p), or (q))	N/A	N/A			N/A				
TOTAL CLAIMS (37 CFR 1.16(j))	minus 20 =	*			x \$200 =				
INDEPENDENT CLAIMS (37 CFR 1.16(h))	minus 3 =	*			x \$600 =				
☐ APPLICATION SIZE FEE (37 CFR 1.16(s))	If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$310 (\$155 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).								
☐ MULTIPLE DEPENDENT CLAIM PRESENT (37 CFR 1.16(j))									
* If the difference in column 1 is less than zero, enter "0" in column 2.							TOTAL		
APPLICATION AS AMENDED - PART II									
	(Column 1)	(Column 2)	(Column 3)						
AMENDMENT	07/25/2025	CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA		RATE (\$)		ADDITIONAL FEE (\$)	
	Total (37 CFR 1.16(j))	* 21	Minus	** 21	= 0		x \$200 =	0	
	Independent (37 CFR 1.16(h))	* 1	Minus	*** 3	= 0		x \$600 =	0	
	☐ Application Size Fee (37 CFR 1.16(s))								
	☐ FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))								
						TOTAL ADD'L FEE	0		
AMENDMENT		CLAIMS REMAINING AFTER AMENDMENT	HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA		RATE (\$)		ADDITIONAL FEE (\$)	
	Total (37 CFR 1.16(j))	*	Minus	**	=		x \$0 =		
	Independent (37 CFR 1.16(h))	*	Minus	***	=		x \$0 =		
	☐ Application Size Fee (37 CFR 1.16(s))								
	☐ FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))								
						TOTAL ADD'L FEE			
* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.						SLIE			
** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".						/DEIDRE C KING-WOODWARD/			
*** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".									
The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.									

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. **SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.**

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC ACKNOWLEDGEMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 07/25/2025 02:04:48 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Colleen Mensching
PATENT CENTER #	71528969	FILING DATE	01/31/2025
CUSTOMER #	21971	FIRST NAMED INVENTOR	Alexis HOWERTON
CORRESPONDENCE ADDRESS	-	AUTHORIZED BY	Victoria Cheng

Documents

TOTAL DOCUMENTS: 7

DOCUMENT	PAGES	DESCRIPTION	SIZE (KB)
Spruce 50535-710.306 - Internet Comms Auth.pdf	2	Internet Communications Authorization	209 KB
Spruce 50535-710.306 - Office Action Response.pdf	11	-	288 KB
Spruce 50535-710.306 - Office Action Response- A....pdf (1-1)	1	Amendment/Request for Reconsideration-After Non- Final Rejection	108 KB
Spruce 50535-710.306 - Office Action Response- SPEC.pdf (2-2)	1	Specification	107 KB
Spruce 50535-710.306 - Office Action Response- CLM.pdf (3-6)	4	Claims	218 KB

Spruce 50535-710.306 - Office Action Response- REM.pdf	(7-11)	5	Applicant Arguments/Remarks Made in an Amendment	168 KB
Spruce 50535-710.306 - Substitute Specification - Marked.pdf		114	Specification	1120 KB
Spruce 50535-710.306 - Substitute Specification - Clean.pdf		113	Specification	1110 KB

Digest

DOCUMENT	MESSAGE DIGEST(SHA-512)
Spruce 50535-710.306 - Internet Comms Auth.pdf	8642C448A77EABCCD709CB4F70A2C19BFCC9F9ED8F2072A2 161907F56D7B0E600A760C2E9E3D58F737D2018FAC10482053 45BE24A30A8FC4B8CB35900C936CDA
Spruce 50535-710.306 - Office Action Response.pdf	49792DEDF9C13BDFE1F4B945E257AB225FC53C6924E6009F9 95B14EAA4FC3B52C124C52897E76951CC61B37F4933EE5AC FD66479EE89E59A6466A9DC96F2084E
Spruce 50535-710.306 - Office Action Response-A....pdf	365B043151913D460FEFD69F1632EFB9AD221F63061D4A15A E46409580A2AC7CDD72A0F845CB1CE6F070AEAD409DFFF01 05DAB19A62771BF7CEA7983F5F17F13
Spruce 50535-710.306 - Office Action Response-SPEC.pdf	DEEB3D9736BA63247FC1FA470DCC2B683EA3FDE4C4552505 791DB2F20E71EC597432A0C622EE6426FC1D154243A1D30C6 4C3BEE297D8CE237DB922A75E42D747
Spruce 50535-710.306 - Office Action Response-CLM.pdf	E4A80A3D004589A5EB7ABDF5CD760D21EFE695C962EF60E5 AAA62D68116330B4EB8B6BA4C86A1109C705AA3C75986658E 92C8C902860DDC48144EC8B20E50B8E
Spruce 50535-710.306 - Office Action Response-REM.pdf	6CC55C7C2E986B40B927887CA65C2148D1C21509830C84D89 5C58DAEE33B6C69D9A113DED8C0F8B137E092E9D93BEA5B 98541A9E7B8BF8ADFB20373A48CF4D60
Spruce 50535-710.306 - Substitute Specification - Marked.pdf	F3774E93C3F6DF3D8D4A7AC5AA46113F741DCB80BD01AFA1 78ED0942D776CEC63405FAF6EA6FE3D7885AFDF0CBE35AF5 2E1A82E02BA24E2235EA6CD2E3B2AB78
Spruce 50535-710.306 - Substitute Specification - Clean.pdf	2B56BA490E21D995CE7C24472174C33BDE077FDFACE17057 FDD0B5C090C8243454C59BD9EB8BBF87A0EFA9067E6E714B 036BF3B5227F62F1789E61E0DFC21FCE

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

PATENT
Attorney Docket No. 50535-710.306

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of: Inventors: Alexis Howerton et al. Application No.: 19/043,264 Filing Date: January 31, 2025 Title: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS	Confirmation No.: 6961 Art Unit: 1627 Examiner: Pihonak, Sarah Customer No.: 21971 <p style="text-align: center;"><u>Certificate of Electronic Filing</u></p> I hereby certify that this correspondence is being transmitted via the U.S. Patent and Trademark Office's electronic filing system to the USPTO on <u>July 25, 2025</u> . By: _____ / Colleen Mensching / Colleen Mensching
--	---

Mail Stop Amendment
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

AMENDMENT AND RESPONSE TO NON-FINAL OFFICE ACTION

Dear Commissioner:

In response to the Non-Final Office Action mailed April 25, 2025, setting an initial due date of July 25, 2025, Applicant respectfully requests reconsideration of the above-referenced application in view of the following amendments and remarks.

Amendments to the Specification begin on page 2 of this paper.

Amendments to the Claims begin on page 3 of this paper.

Remarks begin on page 7 of this paper.

Conclusion begins on page 11 of this paper.

Application No. 19/043,264
Office Action Mailed April 25, 2025
Amendment dated: July 24, 2025

AMENDMENTS TO THE SPECIFICATION


Applicant provides herewith both a clean copy and a marked copy of a substitute specification, excluding claims, in compliance with 37 C.F.R. §§ 1.52, 1.121, and 1.125. The amendments follow the Certificate of Correction (submitted on March 12, 2025, and accepted by the US Patent Office on June 17, 2025) corresponding to US Pat. No. 11,304,950.


Applicant respectfully requests that the specification of record be replaced with the substitute specification filed herewith.

AMENDMENTS TO THE CLAIMS


Listing of the Claims

1. **(Currently Amended)** A method of treating ovarian adrenal rest tumors (OART) and menstrual irregularity in a subject, comprising administering to the subject a small molecule corticotropin-releasing factor type-1 (CRF₁) antagonist, or a pharmaceutically acceptable salt thereof, wherein the small molecule CRF₁ antagonist is:






2. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is:



3. **(Currently Amended)** The method of claim 1, wherein the small molecule CRF₁ antagonist is:



4. **(Currently Amended)** The method of claim 1, ~~wherein the metabolic parameter is insulin sensitivity~~ further comprising treating ovulatory dysfunction or infertility.
5. (Original) The method of claim 1, wherein the subject has congenital adrenal hyperplasia (CAH).
6. (Original) The method of claim 1, wherein the subject is an adolescent.
7. (Original) The method of claim 1, wherein the subject is at least 18 years or older.
8. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered at a dose of about 10 mg/day to about 400 mg/day to the subject.
9. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 50 mg.
10. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 100 mg.
11. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 200 mg.
12. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 400 mg.
13. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject once per day.
14. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject twice per day.
15. (Original) The method of claim 13, wherein the subject is in a fed state.
16. (Original) The method of claim 14, wherein the subject is in a fed state.
17. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered orally.
18. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is formulated as a capsule or tablet.
19. (Original) The method of claim 1, wherein the administering reduces a plasma concentration of a biomarker comprising 17 α -hydroxyprogesterone (17-OHP), adrenocorticotrophic hormone (ACTH), testosterone, androstenedione (A4), luteinizing hormone (LH), follicle-

stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, or aldosterone, or a combination thereof.

20. (Original) The method of claim 19, wherein the biomarker is selected from the group consisting of 17-OHP, ACTH, and A4.
21. (Original) The method of claim 19, wherein the plasma concentration of the biomarker is reduced by at least 10% from baseline.

REMARKS

Claims 1-21 were pending prior to entry of the amendments presented in this response. Claims 1, 3, and 4 are herein amended, support for which can be found in the as-filed application, at least at ¶¶ [0181] and [0208]. No new matter is added by any of these amendments. Reconsideration and allowance are respectfully requested based on the amendments and the following remarks.

Indefiniteness 35 U.S.C. § 112

Claim 4 is rejected under 35 U.S.C. § 112(b) as allegedly being indefinite. Without conceding to the basis of the rejection and for the purpose of compact prosecution, Applicant herein amends claim 4 and requests withdrawal of the indefiniteness rejection.

Anticipation Under 35 U.S.C. § 102

Claim(s) 1-2 and 5-21 is/are rejected under 35 U.S.C. § 102(a)(1) and § 102(a)(2) as being anticipated by Howerton et. al., WO 2019036503 A1, publ. 2/21/2019, international filing date 8/14/2018, cited in an IDS.

“A claim is anticipated only if each and every element as set forth in the claim is found, either expressly or inherently described, in a single prior art reference.” *Verdegaal Bros. v. Union Oil Co. of California*, 814 F.2d 628, 631 (Fed. Cir. 1987); MPEP § 2131.

Applicant submits amended claim 1 is novel over Howerton et al., which Applicant submits does not describe “treating ovarian adrenal rest tumors (OART).” Therefore, Applicant respectfully requests withdrawal of the anticipation rejection and favorable reconsideration. By virtue of their dependencies, claims 2 and 5-21 are also novel, as they depend from amended claim 1, which is novel for at least the reason above.

Obviousness Under 35 U.S.C. § 103

To establish a *prima facie* case of obviousness, all of the claim limitations must be taught by the prior art. “A prior art reference (or references when combined) must teach or suggest all the claim limitations. The teaching or suggestion to make the claimed combination and the reasonable expectation of success must both be found in the prior art and not based on the

applicant's disclosure." *In re Vaeck*, 947 F.2d 488, 20 USPQ2d 1438 (Fed. Cir. 1991); MPEP § 2141(III).

Grigoriadis

Claim(s) 1, 3-14, and 17-21 is/are rejected under 35 U.S.C. § 103 as being unpatentable over Grigoriadis, WO 2015112642 A1, publ. 7/30/2015, cited in an IDS (hereafter "Grigoriadis").

Applicant submits amended claim 1 is nonobvious over Grigoriadis. For instance, Grigoriadis does not appear to teach or suggest treating "treating ovarian adrenal rest tumors (OART) and menstrual irregularity in a subject" as in amended claim 1. As such, a person of ordinary skill in the art would not find a teaching or suggestion in Grigoriadis to arrive at amended claim 1. Therefore, amended claim 1 is nonobvious over Grigoriadis for at least this reason, and withdrawal of the obviousness rejection is respectfully requested.

By virtue of their dependencies, claims 3-14 and 17-21 are also nonobvious over the cited references, as they depend from amended claim 1, which is nonobvious for the reasons above.

Chen in view of Grigoriadis

Claim(s) 1-2, 5-14, and 17-21 is/are rejected under 35 U.S.C. § 103 as being unpatentable over Chen et. al., WO 2008036579 A1, publ. 3/27/2008, cited in an IDS (hereafter "Chen") in view of Grigoriadis.

Applicant submits amended claim 1 is nonobvious over Chen in view of Grigoriadis. Chen does not appear to teach or suggest treating "treating ovarian adrenal rest tumors (OART) and menstrual irregularity in a subject" as in amended claim 1. As such, a person of ordinary skill in the art would not find a teaching or suggestion in Chen to arrive at amended claim 1. Therefore, amended claim 1 is nonobvious over Chen for at least this reason.

Grigoriadis does not cure the deficiencies of Chen for the same reasons as above. As such, a person of ordinary skill in the art would not find a teaching or suggestion in Chen or Grigoriadis, alone or in combination, to arrive at amended claim 1. Therefore, amended claim 1 is nonobvious over the cited references for at least this reason, and withdrawal of the obviousness rejection is respectfully requested.

By virtue of their dependencies, claims 2, 5-14, and 17-21 are also nonobvious over the cited references, as they depend from amended claim 1, which is nonobvious for the reasons above.

Obviousness-Type Double Patenting

Pending claims 1-3, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over:

- Claims 1-4, 7-8, 11-14, 17-18, and 21-22 of U.S. Patent No. 10,849,908 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015;
- Claims 1-4, 7-8, and 11-16 of U.S. Patent No. 11,007,201 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015;
- Claims 1-4, 7-8, 11-16, and 19-24 of U.S. Patent No. 11,344,557 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015; and
- Claims 1-15 and 19 of U.S. Patent No. 11,304,950 B2 in view of Stikkelbroeck et. al., *Fertility & Sterility*, vol. 80(4), pp. 1030-1036, publ. 2003; and Grigoriadis, WO 2015112642 A1, publ. 7/30/2015.

Pending claims 1-2, 5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over:

- Claims 1-4, 7-8, and 14-19 of U.S. Patent No. 11,351,177 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015.
- Claims 1-4, 7-8, and 11-16 of U.S. Patent No. 11,311,549 B2 in view of Grigoriadis, WO 2015112642 A1, publ. 7/30/2015.

Pending claims 1-3, and 5-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over:

- Pending claims 72-91 of copending application 18941832 in view of Stikkelbroeck et. al., *Fertility & Sterility*, vol. 80(4), pp. 1030-1036, publ. 2003; and
- Pending claims 1-5 and 9-21 of copending application 19043275 in view of Stikkelbroeck et. al., *Fertility & Sterility*, vol. 80(4), pp. 1030-1036, publ. 2003.

Claims 1-3, 5-13, and 17-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-19 of copending application 19043303 in view of Stikkelbroeck et. al., *Fertility & Sterility*, vol. 80(4), pp. 1030-1036, publ. 2003.

Without conceding in the basis of the rejection, Applicant submits the amended claims are patentably distinct from the cited patents, cited patent applications, and cited references above. Accordingly, Applicant respectfully requests the withdrawal of the nonstatutory double patenting rejections.

It shall be understood herein that any instance in which Applicant has addressed certain comments set forth by the Office shall not be construed as a concession to other comments or arguments advanced by the Office. Any circumstance in which Applicant has amended or cancelled a claim also does not mean that Applicant concedes to the arguments or positions advanced by the Office with respect to that claim or other claims pending herein.

Application No. 19/043,264
Office Action Mailed April 25, 2025
Amendment dated: July 24, 2025

CONCLUSION

In light of the remarks set forth above, Applicant believes that the pending claims are under condition for allowance. Applicant respectfully solicits the Examiner to expedite the prosecution of this patent application to issuance. Should the Examiner have any questions, the Examiner is encouraged to telephone the undersigned at (858) 350-2240.

The Commissioner is authorized to charge any underpayment or credit any overpayment to Deposit account No. 23-2415 (Attorney Docket No. 50535-710.306).

Respectfully submitted,

Date: July 25, 2025

By: /Victoria Cheng/
Victoria Cheng, Reg. No. 79,641

WILSON SONSINI GOODRICH & ROSATI
650 Page Mill Road
Palo Alto, CA 94304-1050
Phone: (650) 493-9300
Customer No. 21971

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

CROSS-REFERENCE

[0001] This application is a continuation of U.S. Application No. 18/941,832, filed November 8, 2024, which is a continuation of U.S. Application No. 17/667,285, filed on February 8, 2022, which is a continuation of U.S. Application No. 17/081,694 filed on October 27, 2020, which issued as U.S. Pat. No. 11,304,950 on April 19, 2022, and which is a continuation of International Application No. PCT/US2019/029486 filed on April 26, 2019, which claims the benefit of U.S. Provisional Patent Application No. 62/663,951, filed April 27, 2018, and U.S. Provisional Patent Application No. 62/822,815, filed March 23, 2019, each of which is entirely incorporated herein by reference.

BACKGROUND

[0002] Corticotropes are basophilic cells in the anterior pituitary that produce melanocyte-stimulating hormone, adrenocorticotrophic hormone (ACTH) and lipotropin. The cells produce pro-opiomelanocortin (POMC) which undergoes cleavage to ACTH and β -lipotropin (β -LPH). These cells respond to corticotropin-releasing factor (CRF) and make up about 20% of the cells in the anterior pituitary. CRF is a releasing hormone that belongs to the corticotropin-releasing factor family and is the primary physiological regulator of proopioidmelanocortin (POMC).

[0003] CRF is produced by parvocellular neuroendocrine cells within the paraventricular nucleus of the hypothalamus and is released at the median eminence from neurosecretory terminals of these neurons into the primary capillary plexus of the hypothalamo-hypophyseal portal system. The portal system carries the CRF to the anterior lobe of the pituitary, where it stimulates corticotropes to secrete ACTH and other biologically-active substances. ACTH, in turn, stimulates the synthesis of cortisol, glucocorticoids, mineralocorticoids and DHEA.

[0004] Testicular adrenal rest tumors (TART) and ovarian adrenal rest tumors (OART) are ACTH-responsive lesions of the testes and ovaries derived from adrenal tissue interchelated within these organs during embryogenesis. In response to high ACTH, hyperplasia of this tissue occurs, resulting in single or multiple lesions that may cause pain and infertility. The identification of compounds that modulate CRF function and downstream processes is an ongoing challenge. Currently, there are no non-steroidal, non-surgical solutions to control, shrink, or reduce TART. Disclosed herein, *inter alia*, are solutions to these and other problems in the art.

SUMMARY


[10005] Congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency is a serious genetic disorder characterized by impaired adrenal synthesis of cortisol and consequent overproduction of adrenocorticotrophic hormone (ACTH), 17-hydroxyprogesterone (17OHP), and adrenal androgens such as androstenedione (A4). CAH is an orphan condition with no FDA-approved therapy. To suppress androgen production, patients are often prescribed long-term supraphysiologic doses of glucocorticoid (GC), which has adverse effects and fail to suppress 17OHP and A4 in the majority of patients.

[0006] Compound 1 may be a potent, selective, nonsteroidal, oral corticotropin-releasing factor type-1 (CRF1) receptor antagonist that reduces excess ACTH, 17OHP, and A4 to improve clinical and biochemical sequelae of CAH. This may obviate the need for high dose steroids.

[0007] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0008] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0009] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):




wherein:

R¹ and R² are independently ethyl or n-propyl;

R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R⁴ is hydrogen, Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,




R^a and R^b are independently hydrogen, C₁-C₃alkyl, H₂NCH₂CH₂-, (CH₃)₃COC(O)NHCH₂CH₂-, or CH₃CH₂CH₂NHCH₂CH₂-.


[0010] In some embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl. In some embodiments, R^3 is Cl, Br, or methyl. In some embodiments, R^4 is Br, R^aR^bN- , pyridin-4-yl, morpholin-4-yl,

or . In some embodiments, R^4 is morpholin-4-yl or .


[0011] In some embodiments, R^4 is hydrogen, Br, R^aR^bN- and R^a and R^b are independently C_1 - C_3 alkyl. In some embodiments, the compound is



[0012] In some embodiments, the compound is



[0013] In some embodiments, the compound is



[0014] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0015] In some embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically


acceptable salt thereof, is administered to the subject. In some embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0016] In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 5 mg/day to about 25 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 10 mg/day to about 400 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day is administered to the subject.

[0017] In some embodiments, the compound is administered orally. In some embodiments, the compound is formulated as a capsule or tablet. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state.

[0018] In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

[0019] One aspect described herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I) and a pharmaceutically acceptable excipient:




(I), or a pharmaceutically acceptable salt thereof,

wherein:

R^1 and R^2 are independently ethyl or n-propyl;


R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R^4 is hydrogen, Br, $R^a R^b N-$, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,



R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$.

[0020] In some embodiments, the compound of Formula (I) has the structure:



, or a pharmaceutically acceptable salt thereof.

[0021] In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0022] In some embodiments, the pharmaceutical composition comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265

mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.


[0023] In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0024] In some embodiments, the pharmaceutical composition is administered orally. In some embodiments, the pharmaceutical composition is formulated as capsule or a tablet. In some embodiments, the pharmaceutical composition is administered in the fed state. In some embodiments, the pharmaceutical composition is administered in the fasted state. In some embodiments, the pharmaceutical composition is administered at least once per day.

[0025] In some embodiments, the pharmaceutical composition is administered at bedtime. In some embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 2 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep. In some embodiments, the pharmaceutical composition is administered in the evening. In some embodiments, the pharmaceutical composition is administered at about 10 pm at night.


[0026] In some embodiments, the method further comprises administering to the subject an additional chemotherapeutic agent. In some embodiments, the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In some embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisolone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In some embodiments, the mineralocorticoid is fludrocortisone.

[0027] In some embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist. In some embodiments, the CRF₁ antagonist is



pharmaceutically acceptable salt thereof.

[0028] In some embodiments, the CRF₁ antagonist is



[0029] In some embodiments, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof.

[0030] In some embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof.

[0031] In some embodiments, the surgical resection is prior to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is concurrent with

administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0032] In some embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In some embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0033] In some embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the radiation therapy is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0034] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0035] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0037] FIG. 1A-1B. A schematic representation of intra-patient dose escalation study to evaluate safety, pharmacokinetics, and pharmacodynamics **(A)**. A schematic representation of inhibition of CRF₁ with a CRF₁ antagonist (Compound 1) **(B)**.

[0038] FIG. 2. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0039] FIG. 3. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0040] FIG. 4. Graphic representation of patient-level response (measuring ACTH levels) to a CRF₁ antagonist (Compound 1).

[0041] FIG. 5A-5B. Graphic representation of patient-level response of sample responders (measuring ACTH levels) to a CRF₁ antagonist [Patient 10 **(A)**; Patient 8 **(B)**].

[0042] FIG. 6. Graphic representation of patient-level response (measuring 17-OHP levels) to a CRF₁ antagonist (Compound 1).

[0043] FIG. 7A-7B. Graphic representation of patient-level response of sample responders (measuring 17-OHP levels) to a CRF₁ antagonist [Patient 10 **(A)**; Patient 8 **(B)**].

[0044] FIG. 8. Graphic representation of patient-level response (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1).

[0045] FIG. 9A-9B. Graphic representation of patient-level response of sample responders (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1) [Patient 10 **(A)**; Patient 8 **(B)**].

[0046] FIG. 10A-10B. Images of a patient with TART pre-treatment with a CRF₁ antagonist (Compound 1) **(A)** and 6-weeks post-treatment with a CRF₁ antagonist (Compound 1) **(B)**.

[0047] FIG. 11A-11B. Graphic representation of an *in vitro* binding assay of a CRF₁ antagonist (Compound 1); ¹²⁵I-sauvagine binding to rat cerebellar membranes one hour after oral administration of increasing Compound 1 doses **(A)** and *in vivo* inhibition of CRF-induced activity after treatment with a CRF₁ antagonist; Compound 1 (oral, 10 mg/kg) **(B)**.

[0048] FIG. 12A-12B. Graphic representation of ACTH levels (A) and 17-OHP levels (B) after treatment with a CRF₁ antagonist.

[0049] FIG. 13. Graphic representation of hormone responses after treatment with a CRF₁ antagonist (Compound 1).

[0050] FIG. 14 depicts the proof of concept and escalating QD dosing at 10 pm for Cohort A and the single dose level and BID dosing at 10 am and 10 pm for Cohort B.

[0051] FIG. 15A-15D are graphic representations of changes in androstenedione seen in participants in Cohort A and Cohort B.

[0052] FIG. 16A-16D are graphic representations of changes in 17-hydroxyprogesterone seen in participants in Cohort A and Cohort B.

[0053] FIG. 17A-17D are graphic representations of changes in adrenocorticotrophic hormone seen in participants in Cohort A and Cohort B.

DETAILED DESCRIPTION

[0054] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

I. Definitions

[0055] The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are constructed according to the standard rules of chemical valency known in the chemical arts.

[0056] Where substituent groups are specified by their conventional chemical formulae, written from left to right, they equally encompass the chemically identical substituents that would result from writing the structure from right to left, e.g., $-\text{CH}_2\text{O}-$ is equivalent to $-\text{OCH}_2-$.

[0057] The term “alkyl,” by itself or as part of another substituent, means, unless otherwise stated, a straight (i.e., unbranched) or branched carbon chain (or carbon), or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include mono-, di- and multivalent radicals, having the number of carbon atoms designated (i.e., $\text{C}_1\text{-C}_{10}$ means one to ten carbons). Alkyl is an uncyclized chain. Examples of saturated hydrocarbon radicals include, but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, (cyclohexyl)methyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. An alkoxy is an alkyl attached to the remainder of the molecule via an oxygen linker ($-\text{O}-$).

[0058] The term “alkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyl, as exemplified, but not limited by, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms,

with those groups having 10 or fewer carbon atoms being preferred herein. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms. The term “alkenylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkene.

[0059] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or combinations thereof, including at least one carbon atom and at least one heteroatom (e.g., O, N, P, Si, and S), and wherein the nitrogen and sulfur atoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) (e.g., N, S, Si, or P) may be placed at any interior position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Heteroalkyl is an uncyclized chain. Examples include, but are not limited to: $-\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_3$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{S}(\text{O})-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{S}(\text{O})_2-\text{CH}_3$, $-\text{CH}=\text{CH}-\text{O}-\text{CH}_3$, $-\text{Si}(\text{CH}_3)_3$, $-\text{CH}_2-\text{CH}=\text{N}-\text{OCH}_3$, $-\text{CH}=\text{CH}-\text{N}(\text{CH}_3)-\text{CH}_3$, $-\text{O}-\text{CH}_3$, $-\text{O}-\text{CH}_2-\text{CH}_3$, and $-\text{CN}$. Up to two or three heteroatoms may be consecutive, such as, for example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$ and $-\text{CH}_2-\text{O}-\text{Si}(\text{CH}_3)_3$. A heteroalkyl moiety may include one heteroatom (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include two optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include three optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include four optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include five optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include up to 8 optionally different heteroatoms (e.g., O, N, S, Si, or P).

[0060] Similarly, the term “heteroalkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from heteroalkyl, as exemplified, but not limited by, $-\text{CH}_2-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_2-$ and $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-$. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (e.g., alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied by the direction in which the formula of the linking group is written. For example, the formula $-\text{C}(\text{O})_2\text{R}'$ represents both $-\text{C}(\text{O})_2\text{R}'$ and $-\text{R}'\text{C}(\text{O})_2-$. As described above, heteroalkyl groups, as used herein, include those groups that are attached to the remainder of the molecule through a heteroatom, such as $-\text{C}(\text{O})\text{R}'$, $-\text{C}(\text{O})\text{NR}'$, $-\text{NR}'\text{R}''$, $-\text{OR}'$, $-\text{SR}'$, and/or $-\text{SO}_2\text{R}'$. Where “heteroalkyl” is recited, followed by recitations of specific heteroalkyl groups, such as $-\text{NR}'\text{R}''$ or the like, it will be understood that the terms heteroalkyl and $-\text{NR}'\text{R}''$ are not redundant or mutually exclusive. Rather, the specific heteroalkyl groups are recited to add clarity. Thus, the term “heteroalkyl” should not be interpreted herein as excluding specific heteroalkyl groups, such as $-\text{NR}'\text{R}''$ or the like.

[0061] The terms “cycloalkyl” and “heterocycloalkyl,” by themselves or in combination with other terms, mean, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl,” respectively. Cycloalkyl and heterocycloalkyl are not aromatic. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like. A “cycloalkylene” and a “heterocycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a cycloalkyl and heterocycloalkyl, respectively. “Cycloalkyl” is also meant to refer to bicyclic and polycyclic hydrocarbon rings such as, for example, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, etc.

[0062] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl” are meant to include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” includes, but is not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0063] The term “acyl” means, unless otherwise stated, -C(O)R where R is a substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

[0064] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent, which can be a single ring or multiple rings (preferably from 1 to 3 rings) that are fused together (i.e., a fused ring aryl) or linked covalently. A fused ring aryl refers to multiple rings fused together wherein at least one of the fused rings is an aryl ring. The term “heteroaryl” refers to aryl groups (or rings) that contain at least one heteroatom such as N, O, or S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. Thus, the term “heteroaryl” includes fused ring heteroaryl groups (i.e., multiple rings fused together wherein at least one of the fused rings is a heteroaromatic ring). A 5,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 5 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. Likewise, a 6,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. And a 6,5-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and


the other ring has 5 members, and wherein at least one ring is a heteroaryl ring. A heteroaryl group can be attached to the remainder of the molecule through a carbon or heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, naphthyl, pyrrolyl, pyrazolyl, pyridazinyl, triazinyl, pyrimidinyl, imidazolyl, pyrazinyl, purinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, benzofuran, isobenzofuranyl, indolyl, isoindolyl, benzothiophenyl, isoquinolyl, quinoxalyl, quinolyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below. An “arylene” and a “heteroarylene,” alone or as part of another substituent, mean a divalent radical derived from an aryl and heteroaryl, respectively. A heteroaryl group substituent may be -O- bonded to a ring heteroatom nitrogen.

[0065] Spirocyclic rings are two or more rings wherein adjacent rings are attached through a single atom. The individual rings within spirocyclic rings may be identical or different. Individual rings in spirocyclic rings may be substituted or unsubstituted and may have different substituents from other individual rings within a set of spirocyclic rings. Possible substituents for individual rings within spirocyclic rings are the possible substituents for the same ring when not part of spirocyclic rings (e.g. substituents for cycloalkyl or heterocycloalkyl rings). Spirocyclic rings may be substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkylene, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heterocycloalkylene and individual rings within a spirocyclic ring group may be any of the immediately previous list, including having all rings of one type (e.g. all rings being substituted heterocycloalkylene wherein each ring may be the same or different substituted heterocycloalkylene). When referring to a spirocyclic ring system, heterocyclic spirocyclic rings means a spirocyclic rings wherein at least one ring is a heterocyclic ring and wherein each ring may be a different ring. When referring to a spirocyclic ring system, substituted spirocyclic rings means that at least one ring is substituted and each substituent may optionally be different.

[0066] The symbol “” denotes the point of attachment of a chemical moiety to the remainder of a molecule or chemical formula.

[0067] The term “oxo,” as used herein, means an oxygen that is double bonded to a carbon atom.

[0068] The term “alkylarylene” as an arylene moiety covalently bonded to an alkylene moiety (also referred to herein as an alkylene linker). In embodiments, the alkylarylene group has the formula:



[0069] An alkylarylene moiety may be substituted (e.g. with a substituent group) on the alkylene moiety or the arylene linker (e.g. at carbons 2, 3, 4, or 6) with halogen, oxo, -N₃, -CF₃, -CCl₃, -CBr₃, -Cl₃, -CN, -CHO, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₂CH₃, -SO₃H, -OSO₃H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(O)NHNH₂, substituted or unsubstituted C₁-C₅ alkyl or substituted or unsubstituted 2 to 5 membered heteroalkyl). In embodiments, the alkylarylene is unsubstituted.

[0070] Each of the above terms (e.g., “alkyl,” “heteroalkyl,” “cycloalkyl,” “heterocycloalkyl,” “aryl,” and “heteroaryl”) includes both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

[0071] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be one or more of a variety of groups selected from, but not limited to, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR'C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R''R''')=NR''', -NR-C(NR'R'')=NR'', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -NR'NR''R''', -ONR'R'', -NR'C(O)NR''NR''R''', -CN, -NO₂, -NR'SO₂R'', -NR'C(O)R'', -NR'C(O)-OR'', -NR'OR'', in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such radical. R, R', R'', R''', and R'''' each preferably independently refer to hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl (e.g., aryl substituted with 1-3 halogens), substituted or unsubstituted heteroaryl, substituted or unsubstituted alkyl, alkoxy, or thioalkoxy groups, or arylalkyl groups. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', and R'''' group when more than one of these groups is present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 4-, 5-, 6-, or 7-membered ring. For example, -NR'R'' includes, but is not limited to, 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term “alkyl” is meant to include

groups including carbon atoms bound to groups other than hydrogen groups, such as haloalkyl (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

[0072] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups are varied and are selected from, for example: -OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR'C(O)R', -NR'C(O)NR'R''', -NR'C(O)₂R', -NR-C(NR'R'R''')=NR''', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -NR'NR'R''', -ONR'R'', -NR'C(O)NR''NR''R''', -CN, -NO₂, -R', -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, -NR'SO₂R'', -NR'C(O)R'', -NR'C(O)-OR'', -NR'OR'', in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'', R''', and R'''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', and R'''' groups when more than one of these groups is present.

[0073] Substituents for rings (e.g. cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylene, heterocycloalkylene, arylene, or heteroarylene) may be depicted as substituents on the ring rather than on a specific atom of a ring (commonly referred to as a floating substituent). In such a case, the substituent may be attached to any of the ring atoms (obeying the rules of chemical valency) and in the case of fused rings or spirocyclic rings, a substituent depicted as associated with one member of the fused rings or spirocyclic rings (a floating substituent on a single ring), may be a substituent on any of the fused rings or spirocyclic rings (a floating substituent on multiple rings). When a substituent is attached to a ring, but not a specific atom (a floating substituent), and a subscript for the substituent is an integer greater than one, the multiple substituents may be on the same atom, same ring, different atoms, different fused rings, different spirocyclic rings, and each substituent may optionally be different. Where a point of attachment of a ring to the remainder of a molecule is not limited to a single atom (a floating substituent), the attachment point may be any atom of the ring and in the case of a fused ring or spirocyclic ring, any atom of any of the fused rings or spirocyclic rings while obeying the rules of chemical valency. Where a ring, fused rings, or spirocyclic rings contain one or more ring heteroatoms and the ring, fused rings, or spirocyclic rings are shown with one more floating substituents (including, but not limited to, points of attachment to the remainder of the molecule), the floating substituents may be bonded to the heteroatoms. Where the ring heteroatoms are shown bound to one or more hydrogens (e.g. a ring nitrogen with two bonds to ring atoms and a third bond to a hydrogen) in the structure or formula with the floating substituent, when the

heteroatom is bonded to the floating substituent, the substituent will be understood to replace the hydrogen, while obeying the rules of chemical valency.

[0074] Two or more substituents may optionally be joined to form aryl, heteroaryl, cycloalkyl, or heterocycloalkyl groups. Such so-called ring-forming substituents are typically, though not necessarily, found attached to a cyclic base structure. In one embodiment, the ring-forming substituents are attached to adjacent members of the base structure. For example, two ring-forming substituents attached to adjacent members of a cyclic base structure create a fused ring structure. In another embodiment, the ring-forming substituents are attached to a single member of the base structure. For example, two ring-forming substituents attached to a single member of a cyclic base structure create a spirocyclic structure. In yet another embodiment, the ring-forming substituents are attached to non-adjacent members of the base structure.

[0075] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally form a ring of the formula $-T-C(O)-(CRR')_q-U-$, wherein T and U are independently $-NR'$ -, $-O-$ -, $-CRR'$ -, or a single bond, and q is an integer of from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-A-(CH_2)_r-B-$, wherein A and B are independently $-CRR'$ -, $-O-$ -, $-NR'$ -, $-S-$ -, $-S(O)-$ -, $-S(O)_2-$ -, $-S(O)_2NR'$ -, or a single bond, and r is an integer of from 1 to 4. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-(CRR')_s-X'-(C''R''R''')_d-$, where s and d are independently integers of from 0 to 3, and X' is $-O-$ -, $-NR'$ -, $-S-$ -, $-S(O)-$ -, $-S(O)_2-$ -, or $-S(O)_2NR'$ -. The substituents R, R', R'', and R''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

[0076] As used herein, the terms “heteroatom” or “ring heteroatom” are meant to include oxygen (O), nitrogen (N), sulfur (S), phosphorus (P), and silicon (Si).

[0077] A “substituent group,” as used herein, means a group selected from the following moieties:

(A) oxo, halogen, $-CF_3$ -, $-CN$ -, $-OH$ -, $-NH_2$ -, $-COOH$ -, $-CONH_2$ -, $-NO_2$ -, $-SH$ -, $-SO_3H$ -, $-SO_4H$ -, $-SO_2NH_2$ -, $-NHNH_2$ -, $-ONH_2$ -, $-NHC(=O)NHNH_2$ -, $-NHC(=O)NH_2$ -, $-NHSO_2H$ -, $-NHC(=O)H$ -, $-NHC(O)-OH$ -, $-NHOH$ -, $-OCF_3$ -, $-OCHF_2$ -, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(B) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(i) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NHSO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(ii) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(a) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NHSO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(b) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from: oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NHSO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl.

[0078] A “size-limited substituent” or “size-limited substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl.

[0079] A “lower substituent” or “lower substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each

substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl.

[0080] In some embodiments, each substituted group described in the compounds herein is substituted with at least one substituent group. More specifically, in some embodiments, each substituted alkyl, substituted heteroalkyl, substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl, substituted heteroaryl, substituted alkylene, substituted heteroalkylene, substituted cycloalkylene, substituted heterocycloalkylene, substituted arylene, and/or substituted heteroarylene described in the compounds herein are substituted with at least one substituent group. In other embodiments, at least one or all of these groups are substituted with at least one size-limited substituent group. In other embodiments, at least one or all of these groups are substituted with at least one lower substituent group.

[0081] In other embodiments of the compounds herein, each substituted or unsubstituted alkyl may be a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl. In some embodiments of the compounds herein, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₂₀ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 20 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₈ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a substituted or unsubstituted 3 to 8 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 10 membered heteroarylene.

[0082] In some embodiments, each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl. In some

embodiments, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₈ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 8 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₇ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a substituted or unsubstituted 3 to 7 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 9 membered heteroarylene. In some embodiments, the compound is a chemical species set forth in the Examples section, figures, or tables below.

[0083] Certain compounds of the present invention possess asymmetric carbon atoms (optical or chiral centers) or double bonds; the enantiomers, racemates, diastereomers, tautomers, geometric isomers, stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)- or, as (D)- or (L)- for amino acids, and individual isomers are encompassed within the scope of the present invention. The compounds of the present invention do not include those that are known in art to be too unstable to synthesize and/or isolate. The present invention is meant to include compounds in racemic and optically pure forms. Optically active (R)- and (S)-, or (D)- and (L)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers.

[0084] As used herein, the term “isomers” refers to compounds having the same number and kind of atoms, and hence the same molecular weight, but differing in respect to the structural arrangement or configuration of the atoms.

[0085] The term “tautomer,” as used herein, refers to one of two or more structural isomers which exist in equilibrium and which are readily converted from one isomeric form to another.

[0086] It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention.

[0087] Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention.

[0088] Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a

deuterium or tritium, or the replacement of a carbon by ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention.

[0089] The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I), or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are encompassed within the scope of the present invention.

[0090] It should be noted that throughout the application that alternatives are written in Markush groups, for example, each amino acid position that contains more than one possible amino acid. It is specifically contemplated that each member of the Markush group should be considered separately, thereby comprising another embodiment, and the Markush group is not to be read as a single unit.

[0091] “Analog” or “analogue” is used in accordance with its plain ordinary meaning within Chemistry and Biology and refers to a chemical compound that is structurally similar to another compound (i.e., a so-called “reference” compound) but differs in composition, e.g., in the replacement of one atom by an atom of a different element, or in the presence of a particular functional group, or the replacement of one functional group by another functional group, or the absolute stereochemistry of one or more chiral centers of the reference compound. Accordingly, an analog is a compound that is similar or comparable in function and appearance but not in structure or origin to a reference compound.

[0092] The terms “a” or “an,” as used in herein means one or more. In addition, the phrase “substituted with a[n],” as used herein, means the specified group may be substituted with one or more of any or all of the named substituents. For example, where a group, such as an alkyl or heteroaryl group, is “substituted with an unsubstituted $\text{C}_1\text{-C}_{20}$ alkyl, or unsubstituted 2 to 20 membered heteroalkyl,” the group may contain one or more unsubstituted $\text{C}_1\text{-C}_{20}$ alkyls, and/or one or more unsubstituted 2 to 20 membered heteroalkyls.

[0093] Moreover, where a moiety is substituted with an R substituent, the group may be referred to as “R-substituted.” Where a moiety is R-substituted, the moiety is substituted with at least one R substituent and each R substituent is optionally different. Where a particular R group is present in the description of a chemical genus (such as Formula (I)), a Roman alphabetic symbol may be used to distinguish each appearance of that particular R group. For example, where multiple R^{13} substituents are present, each R^{13} substituent may be distinguished as $\text{R}^{13\text{A}}$, $\text{R}^{13\text{B}}$, $\text{R}^{13\text{C}}$, $\text{R}^{13\text{D}}$, etc., wherein each of $\text{R}^{13\text{A}}$, $\text{R}^{13\text{B}}$, $\text{R}^{13\text{C}}$, $\text{R}^{13\text{D}}$, etc. is defined within the scope of the definition of R^{13} and optionally differently.

[0094] A “detectable moiety” as used herein refers to a moiety that can be covalently or noncovalently attached to a compound or biomolecule that can be detected for instance, using techniques known in the art. In embodiments, the detectable moiety is covalently attached. The detectable moiety may provide for imaging of the attached compound or biomolecule. The detectable moiety may indicate the contacting between two compounds. Exemplary detectable moieties are fluorophores, antibodies, reactive dies, radio-labeled moieties, magnetic contrast agents, and quantum dots. Exemplary fluorophores include fluorescein, rhodamine, GFP, coumarin, FITC, Alexa fluor, Cy3, Cy5, BODIPY, and cyanine dyes. Exemplary radionuclides include Fluorine-18, Gallium-68, and Copper-64. Exemplary magnetic contrast agents include gadolinium, iron oxide and iron platinum, and manganese.

[0095] Descriptions of compounds of the present invention are limited by principles of chemical bonding known to those skilled in the art. Accordingly, where a group may be substituted by one or more of a number of substituents, such substitutions are selected so as to comply with principles of chemical bonding and to give compounds which are not inherently unstable and/or would be known to one of ordinary skill in the art as likely to be unstable under ambient conditions, such as aqueous, neutral, and several known physiological conditions. For example, a heterocycloalkyl or heteroaryl is attached to the remainder of the molecule via a ring heteroatom in compliance with principles of chemical bonding known to those skilled in the art thereby avoiding inherently unstable compounds.

[0096] The term “pharmaceutically acceptable salts” is meant to include salts of the active compounds that are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, oxalic,

methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see*, for example, Berge *et al.*, “Pharmaceutical Salts”, *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0097] Thus, the compounds of the present invention may exist as salts, such as with pharmaceutically acceptable acids. The present invention includes such salts. Non-limiting examples of such salts include hydrochlorides, hydrobromides, phosphates, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, propionates, tartrates (e.g., (+)-tartrates, (-)-tartrates, or mixtures thereof including racemic mixtures), succinates, benzoates, and salts with amino acids such as glutamic acid, and quaternary ammonium salts (e.g. methyl iodide, ethyl iodide, and the like). These salts may be prepared by methods known to those skilled in the art.

[0098] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound may differ from the various salt forms in certain physical properties, such as solubility in polar solvents. In embodiments, compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts. The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in a conventional manner. The parent form of the compounds differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but, unless specifically indicated, the salts disclosed herein are equivalent to the parent form of the compound for the purposes of the present invention.

[0099] In addition to salt forms, the present invention provides compounds, which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Prodrugs of the compounds described herein may be converted *in vivo* after administration. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment, such as, for example, when contacted with a suitable enzyme or chemical reagent.

[0100] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In

general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0101] “Pharmaceutically acceptable excipient” and “pharmaceutically acceptable carrier” refer to a substance that aids the administration of a compound to and absorption by a subject and can be included in the compositions of the present invention without causing a significant adverse toxicological effect on the patient. Non-limiting examples of pharmaceutically acceptable excipients include water, NaCl, normal saline solutions, lactated Ringer’s, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer’s solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with the compounds of the invention. One of skill in the art will recognize that other pharmaceutical excipients are useful in the present invention.

[0102] The term “preparation” is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

[0103] A “CRF1 inhibitor” refers to a compound (e.g., compounds described herein) that reduces the activity of CRF1 when compared to a control, such as absence of the compound or a compound with known inactivity.

[0104] The terms “polypeptide,” “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues, wherein the polymer may optionally be conjugated to a moiety that does not consist of amino acids. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. In embodiments, the terms “polypeptide,” “peptide,” and “protein”, used interchangeably herein, refer to a polymeric form of amino acids of any length, which can include genetically coded and non-genetically coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified polypeptide backbones. The terms include fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence; fusion proteins with heterologous and

homologous leader sequences, with or without N-terminus methionine residues; immunologically tagged proteins; and the like.

[0105] A polypeptide, or a cell is “recombinant” when it is artificial or engineered, or derived from or contains an artificial or engineered protein or nucleic acid (*e.g.* non-natural or not wild type). For example, a polynucleotide that is inserted into a vector or any other heterologous location, *e.g.*, in a genome of a recombinant organism, such that it is not associated with nucleotide sequences that normally flank the polynucleotide as it is found in nature is a recombinant polynucleotide. A protein expressed *in vitro* or *in vivo* from a recombinant polynucleotide is an example of a recombinant polypeptide. Likewise, a polynucleotide sequence that does not appear in nature, for example a variant of a naturally occurring gene, is recombinant.

[0106] “Contacting” is used in accordance with its plain ordinary meaning and refers to the process of allowing at least two distinct species (*e.g.* chemical compounds including biomolecules or cells) to become sufficiently proximal to react, interact or physically touch. It should be appreciated; however, the resulting reaction product can be produced directly from a reaction between the added reagents or from an intermediate from one or more of the added reagents that can be produced in the reaction mixture.

[0107] The term “contacting” may include allowing two species to react, interact, or physically touch, wherein the two species may be a compound as described herein and a protein or enzyme. In some embodiments contacting includes allowing a compound described herein to interact with a protein or enzyme that is involved in a signaling pathway (*e.g.*, MAP kinase pathway).

[0108] As defined herein, the term “activation”, “activate”, “activating” and the like in reference to a protein refers to conversion of a protein into a biologically active derivative from an initial inactive or deactivated state. The terms reference activation, or activating, sensitizing, or up-regulating signal transduction or enzymatic activity or the amount of a protein decreased in a disease.

[0109] The terms “agonist,” “activator,” “upregulator,” etc. refer to a substance capable of detectably increasing the expression or activity of a given gene or protein. The agonist can increase expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the agonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or higher than the expression or activity in the absence of the agonist. In embodiments, an agonist is a molecule that interacts with a target to cause or promote an increase in the activation of the target. In embodiments, activators are molecules that increase, activate, facilitate, enhance activation, sensitize, or up-regulate, *e.g.*, a gene, protein, ligand, receptor, or cell.

[0110] As defined herein, the term “inhibition”, “inhibit”, “inhibiting” and the like in reference to a protein-inhibitor interaction means negatively affecting (e.g. decreasing) the activity or function of the protein relative to the activity or function of the protein in the absence of the inhibitor. In embodiments inhibition means negatively affecting (e.g. decreasing) the concentration or levels of the protein relative to the concentration or level of the protein in the absence of the inhibitor. In embodiments inhibition refers to reduction of a disease or symptoms of disease. In embodiments, inhibition refers to a reduction in the activity of a particular protein target. Thus, inhibition includes, at least in part, partially or totally blocking stimulation, decreasing, preventing, or delaying activation, or inactivating, desensitizing, or down-regulating signal transduction or enzymatic activity or the amount of a protein. In embodiments, inhibition refers to a reduction of activity of a target protein resulting from a direct interaction (e.g. an inhibitor binds to the target protein). In embodiments, inhibition refers to a reduction of activity of a target protein from an indirect interaction (e.g. an inhibitor binds to a protein that activates the target protein, thereby preventing target protein activation).

[0111] The terms “inhibitor,” “repressor” or “antagonist” or “downregulator” interchangeably refer to a substance capable of detectably decreasing the expression or activity of a given gene or protein. The antagonist can decrease expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the antagonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or lower than the expression or activity in the absence of the antagonist. An antagonist prevents, reduces, inhibits, or neutralizes the activity of an agonist, and an antagonist can also prevent, inhibit, or reduce constitutive activity of a target, e.g., a target receptor, even where there is no identified agonist. In embodiments, inhibitors are molecules that decrease, block, prevent, delay activation, inactivate, desensitize, or down-regulate, e.g., a gene, protein, ligand, receptor, or cell. An inhibitor may also be defined as a molecule that reduces, blocks, or inactivates a constitutive activity. An “antagonist” is a molecule that opposes the action(s) of an agonist.

[0112] The terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” refer to a protein (including homologs, isoforms, and functional fragments thereof) and is a receptor for corticotropin releasing-hormone. The terms include any recombinant or naturally-occurring form of CRF₁ variants thereof that maintain CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). The terms include any mutant form of CRF₁ variants (e.g., frameshift mutations) thereof that maintain CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). In embodiments, the

CRF₁ protein encoded by the *CRHR1* gene has the amino acid sequence set forth in or corresponding to Entrez 1233, UniProt P34998.

[0113] The term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion. Expression can be detected using conventional techniques for detecting protein (*e.g.*, ELISA, Western blotting, flow cytometry, immunofluorescence, immunohistochemistry, *etc.*).

[0114] The terms “disease” or “condition” refer to a state of being or health status of a patient or subject capable of being treated with the compounds or methods provided herein. The disease may be a malignant or benign tumor (*e.g.*, testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)). The disease may be cancer. In some further instances, “cancer” refers to human cancers and carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, *etc.*, including solid and lymphoid cancers, kidney, breast, lung, bladder, colon, ovarian, prostate, pancreas, adrenal, stomach, brain, head and neck, skin, uterine, testicular, glioma, esophagus, and liver cancer, including hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin’s lymphomas (*e.g.*, Burkitt’s, Small Cell, and Large Cell lymphomas), Hodgkin’s lymphoma, leukemia (including MDS, AML, ALL, ATLL and CML), or multiple myeloma.

[0115] As used herein, the term “cancer” refers to all types of cancer, neoplasm or malignant tumors found in mammals (*e.g.* humans), including leukemia, carcinomas and sarcomas. Exemplary cancers that may be treated with a compound or method provided herein include adrenal cancer and tumors (*e.g.*, testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) brain cancer, glioma, glioblastoma, neuroblastoma, prostate cancer, colorectal cancer, pancreatic cancer, cervical cancer, gastric cancer, ovarian cancer, lung cancer, and cancer of the head. Exemplary cancers that may be treated with a compound or method provided herein include cancer of the thyroid, endocrine system, brain, breast, cervix, colon, head & neck, liver, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus, Medulloblastoma, colorectal cancer, pancreatic cancer. Additional examples include, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung adenocarcinoma, lung squamous cell carcinoma, Hodgkin’s Disease, Non-Hodgkin’s Lymphoma, multiple myeloma, neuroblastoma, glioma, glioblastoma multiforme, ovarian cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, primary brain tumors, cancer, malignant pancreatic insuloma, malignant carcinoid, urinary bladder cancer,

pre-malignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, endometrial cancer, adrenal cortical cancer, neoplasms of the endocrine or exocrine pancreas, medullary thyroid cancer, medullary thyroid carcinoma, melanoma, colorectal cancer, papillary thyroid cancer, hepatocellular carcinoma, or prostate cancer.

[0116] The term “congenital adrenal hyperplasia” or “CAH” are any of several autosomal recessive diseases resulting from mutations of genes for enzymes mediating the biochemical steps of production of mineralocorticoids, glucocorticoids or sex steroids from cholesterol by the adrenal glands (steroidogenesis). These conditions involve excessive or deficient production of sex steroids and can alter development of primary or secondary sex characteristics in some affected infants, children, or adults.

[0117] “Adrenal rest tumor” refers to a neoplasm of the testis (or ovaries) consisting of adrenal tissue. It may be secondary to hyperplasia of an adrenal gland.

[0118] “Testicular adrenal rest tumor(s)” or “TART” and/or ovarian adrenal rest tumor(s) are a complication of congenital adrenal hyperplasia (CAH), which may develop from ectopic remnants of intra-testicular adrenal tissue stimulated by adrenocorticotrophic hormone (ACTH) hypersecretion. TART lesions are typically located within the rete testis and are bilateral, synchronous, nodular and multiple. TART can lead to testicular structural damage, spermatogenesis disorders, infertility and mass-forming lesions.

[0119] The term “leukemia” refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease-acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number abnormal cells in the blood-leukemic or aleukemic (subleukemic). Exemplary leukemias that may be treated with a compound or method provided herein include, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemetic leukemia, basophilic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic

leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, multiple myeloma, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, or undifferentiated cell leukemia.

[0120] The term “sarcoma” generally refers to a tumor which is made up of a substance like the embryonic connective tissue and is generally composed of closely packed cells embedded in a fibrillar or homogeneous substance. Sarcomas that may be treated with a compound or method provided herein include a chondrosarcoma, fibrosarcoma, lymphosarcoma, melanosarcoma, myxosarcoma, osteosarcoma, Abemethy's sarcoma, adipose sarcoma, liposarcoma, alveolar soft part sarcoma, ameloblastic sarcoma, botryoid sarcoma, chloroma sarcoma, chorio carcinoma, embryonal sarcoma, Wilms' tumor sarcoma, endometrial sarcoma, stromal sarcoma, Ewing's sarcoma, fascial sarcoma, fibroblastic sarcoma, giant cell sarcoma, granulocytic sarcoma, Hodgkin's sarcoma, idiopathic multiple pigmented hemorrhagic sarcoma, immunoblastic sarcoma of B cells, lymphoma, immunoblastic sarcoma of T-cells, Jensen's sarcoma, Kaposi's sarcoma, Kupffer cell sarcoma, angiosarcoma, leukosarcoma, malignant mesenchymoma sarcoma, parosteal sarcoma, reticulocytic sarcoma, Rous sarcoma, serocystic sarcoma, synovial sarcoma, or telangiectatic sarcoma.

[0121] The term “melanoma” is taken to mean a tumor arising from the melanocytic system of the skin and other organs. Melanomas that may be treated with a compound or method provided herein include, for example, acral-lentiginous melanoma, amelanotic melanoma, benign juvenile melanoma, Cloudman's melanoma, S91 melanoma, Harding-Passey melanoma, juvenile melanoma, lentigo maligna melanoma, malignant melanoma, nodular melanoma, subungual melanoma, or superficial spreading melanoma.

[0122] The term “carcinoma” refers to a malignant new growth made up of epithelial cells tending to infiltrate the surrounding tissues and give rise to metastases. Exemplary carcinomas that may be treated with a compound or method provided herein include, for example, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung adenocarcinoma, lung squamous cell carcinoma, medullary thyroid carcinoma, familial medullary thyroid carcinoma, acinar carcinoma, acinous carcinoma, adenocystic carcinoma, adenoid cystic carcinoma, carcinoma adenomatousum, carcinoma of adrenal cortex, alveolar carcinoma, alveolar cell carcinoma, basal cell carcinoma, carcinoma basocellulare, basaloid carcinoma, basosquamous cell carcinoma, bronchioalveolar carcinoma, bronchiolar carcinoma, bronchogenic carcinoma, cerebriiform carcinoma, cholangiocellular carcinoma, chorionic carcinoma, colloid carcinoma,

comedo carcinoma, corpus carcinoma, cribriform carcinoma, carcinoma en cuirasse, carcinoma cutaneum, cylindrical carcinoma, cylindrical cell carcinoma, duct carcinoma, carcinoma durum, embryonal carcinoma, encephaloid carcinoma, epiermoid carcinoma, carcinoma epitheliale adenoides, exophytic carcinoma, carcinoma ex ulcere, carcinoma fibrosum, gelatiniformi carcinoma, gelatinous carcinoma, giant cell carcinoma, carcinoma gigantocellulare, glandular carcinoma, granulosa cell carcinoma, hair-matrix carcinoma, hematoid carcinoma, hepatocellular carcinoma, Hurthle cell carcinoma, hyaline carcinoma, hypernephroid carcinoma, infantile embryonal carcinoma, carcinoma in situ, intraepidermal carcinoma, intraepithelial carcinoma, Krompecher's carcinoma, Kulchitzky-cell carcinoma, large-cell carcinoma, lenticular carcinoma, carcinoma lenticulare, lipomatous carcinoma, lymphoepithelial carcinoma, carcinoma medullare, medullary carcinoma, melanotic carcinoma, carcinoma molle, mucinous carcinoma, carcinoma muciparum, carcinoma mucocellulare, mucoepidermoid carcinoma, carcinoma mucosum, mucous carcinoma, carcinoma myxomatodes, nasopharyngeal carcinoma, oat cell carcinoma, carcinoma ossificans, osteoid carcinoma, papillary carcinoma, periportal carcinoma, preinvasive carcinoma, prickle cell carcinoma, pultaceous carcinoma, renal cell carcinoma of kidney, reserve cell carcinoma, carcinoma sarcomatodes, schneiderian carcinoma, scirrhus carcinoma, carcinoma scroti, signet-ring cell carcinoma, carcinoma simplex, small-cell carcinoma, solanoid carcinoma, spheroidal cell carcinoma, spindle cell carcinoma, carcinoma spongiosum, squamous carcinoma, squamous cell carcinoma, string carcinoma, carcinoma telangiectaticum, carcinoma telangiectodes, transitional cell carcinoma, carcinoma tuberosum, tuberos carcinoma, verrucous carcinoma, or carcinoma villosum.

[0123] The terms “treating” or “treatment” refer to any indicia of success in the therapy or amelioration of an injury, disease, pathology or condition, including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the injury, pathology or condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; improving a patient’s physical or mental well-being. The treatment or amelioration of symptoms can be based on objective or subjective parameters; including the results of a physical examination, neuropsychiatric exams, and/or a psychiatric evaluation. The term “treating” and conjugations thereof, may include prevention of an injury, pathology, condition, or disease. In embodiments, treating is preventing. In embodiments, treating does not include preventing.

[0124] “Treating” or “treatment” as used herein (and as well-understood in the art) also broadly includes any approach for obtaining beneficial or desired results in a subject’s condition, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of the extent

of a disease, stabilizing (*i.e.*, not worsening) the state of disease, prevention of a disease's transmission or spread, delay or slowing of disease progression, amelioration or palliation of the disease state, diminishment of the reoccurrence of disease, and remission, whether partial or total and whether detectable or undetectable. In other words, "treatment" as used herein includes any cure, amelioration, or prevention of a disease. Treatment may prevent the disease from occurring; inhibit the disease's spread; relieve the disease's symptoms (*e.g.*, ocular pain, seeing halos around lights, red eye, very high intraocular pressure), fully or partially remove the disease's underlying cause, shorten a disease's duration, or do a combination of these things.

[0125] "Treating" and "treatment" as used herein include prophylactic treatment. Treatment methods include administering to a subject a therapeutically effective amount of a compound described herein. The administering step may consist of a single administration or may include a series of administrations. The length of the treatment period depends on a variety of factors, such as the severity of the condition, the age of the patient, the concentration of the compound, the activity of the compositions used in the treatment, or a combination thereof. It will also be appreciated that the effective dosage of an agent used for the treatment or prophylaxis may increase or decrease over the course of a particular treatment or prophylaxis regime. Changes in dosage may result and become apparent by standard diagnostic assays known in the art. In some instances, chronic administration may be required. For example, the compositions are administered to the subject in an amount and for a duration sufficient to treat the patient.

[0126] The term "prevent" refers to a decrease in the occurrence of disease symptoms in a patient. As indicated above, the prevention may be complete (no detectable symptoms) or partial, such that fewer symptoms are observed than would likely occur absent treatment. In embodiments, prevent refers to slowing the progression of the disease, disorder or condition or inhibiting progression thereof to a harmful or otherwise undesired state.

[0127] "Patient" or "subject in need thereof" refers to a living organism suffering from or prone to a disease or condition that can be treated by administration of a pharmaceutical composition as provided herein. Non-limiting examples include humans, other mammals, bovines, rats, mice, dogs, monkeys, goat, sheep, cows, deer, and other non-mammalian animals. In some embodiments, a patient is human.

[0128] An "effective amount" is an amount sufficient for a compound to accomplish a stated purpose relative to the absence of the compound (*e.g.*, achieve the effect for which it is administered, treat a disease, reduce enzyme activity, increase enzyme activity, reduce a signaling pathway, or reduce one or more symptoms of a disease or condition). An example of an "effective amount" is an amount sufficient to contribute to the treatment, prevention, or reduction of a symptom or symptoms of a disease, which could also be referred to as a

“therapeutically effective amount.” A “reduction” of a symptom or symptoms (and grammatical equivalents of this phrase) means decreasing of the severity or frequency of the symptom(s), or elimination of the symptom(s). A “prophylactically effective amount” of a drug is an amount of a drug that, when administered to a subject, will have the intended prophylactic effect, e.g., preventing or delaying the onset (or reoccurrence) of an injury, disease, pathology or condition, or reducing the likelihood of the onset (or reoccurrence) of an injury, disease, pathology, or condition, or their symptoms. The full prophylactic effect does not necessarily occur by administration of one dose, and may occur only after administration of a series of doses. Thus, a prophylactically effective amount may be administered in one or more administrations. An “activity decreasing amount,” as used herein, refers to an amount of antagonist required to decrease the activity of an enzyme relative to the absence of the antagonist. A “function disrupting amount,” as used herein, refers to the amount of antagonist required to disrupt the function of an enzyme or protein relative to the absence of the antagonist. The exact amounts will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (*see, e.g.,* Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); and *Remington: The Science and Practice of Pharmacy*, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins). The therapeutically effective amount can be ascertained by measuring relevant physiological effects, and it can be adjusted in connection with the dosing regimen and diagnostic analysis of the subject’s condition, and the like. By way of example, measurement of the serum level of a CRF1 inhibitor (or, e.g., a metabolite thereof) at a particular time post-administration may be indicative of whether a therapeutically effective amount has been administered.

[0129] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0130] Therapeutically effective amounts for use in humans can also be determined from animal models. For example, a dose for humans can be formulated to achieve a concentration that has been found to be effective in animals. The dosage in humans can be adjusted by monitoring compounds effectiveness and adjusting the dosage upwards or downwards, as described above. Adjusting the dose to achieve maximal efficacy in humans based on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan. Adjusting the dose to achieve maximal therapeutic window efficacy or toxicity in humans based

on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan.

[0131] The term “therapeutically effective amount,” as used herein, refers to that amount of the therapeutic agent sufficient to ameliorate the disorder, as described above. For example, for the given parameter, a therapeutically effective amount will show an increase or decrease of at least 5%, 10%, 15%, 20%, 25%, 40%, 50%, 60%, 75%, 80%, 90%, or at least 100%. Therapeutic efficacy can also be expressed as “-fold” increase or decrease. For example, a therapeutically effective amount can have at least a 1.2-fold, 1.5-fold, 2-fold, 5-fold, or more effect over a control.

[0132] The terms “QD,” “qd,” and “q.d.” refer to once a day, i.e., a drug or therapeutic agent is administered once a day for the duration of the therapeutic regimen.

[0133] The terms “BD,” “bd,” and “b.i.d.” refer to twice a day, i.e., a drug or therapeutic agent is administered two times a day for the duration of the therapeutic regimen.

[0134] The terms “QID,” “qid,” and “q.i.d.” refer to three times a day, i.e., a drug or therapeutic agent is administered three times a day for the duration of the therapeutic regimen.

[0135] Dosages may be varied depending upon the requirements of the patient and the compound being employed. The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial therapeutic response in the patient over time. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects. Determination of the proper dosage for a particular situation is within the skill of the practitioner. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. Dosage amounts and intervals can be adjusted individually to provide levels of the administered compound effective for the particular clinical indication being treated. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease state.

[0136] As used herein, the term “administering” means oral administration, administration as a suppository, topical contact, intravenous, parenteral, intraperitoneal, intramuscular, intralesional, intrathecal, intracranial, intranasal or subcutaneous administration, or the implantation of a slow-release device, *e.g.*, a mini-osmotic pump, to a subject. Administration is by any route, including parenteral and transmucosal (*e.g.*, buccal, sublingual, palatal, gingival, nasal, vaginal, rectal, or transdermal). Parenteral administration includes, *e.g.*, intravenous, intramuscular, intra-arteriole, intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial. Other modes of delivery include, but are not limited to, the use of liposomal formulations, intravenous infusion, transdermal patches, *etc.* By “co-administer” it is meant that a composition described herein is

administered at the same time, just prior to, or just after the administration of one or more additional therapies (e.g. anti-cancer agent, chemotherapeutic, or treatment for a neurodegenerative disease). The compound of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compound individually or in combination (more than one compound or agent). Thus, the preparations can also be combined, when desired, with other active substances (e.g. to reduce metabolic degradation). The compositions of the present invention can be delivered by transdermally, by a topical route, formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols. Oral preparations include tablets, pills, powder, dragees, capsules, liquids, lozenges, cachets, gels, syrups, slurries, suspensions, etc., suitable for ingestion by the patient. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. The compositions of the present invention may additionally include components to provide sustained release and/or comfort. Such components include high molecular weight, anionic mucomimetic polymers, gelling polysaccharides and finely-divided drug carrier substrates. These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841; 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by reference in their entirety for all purposes. The compositions of the present invention can also be delivered as microspheres for slow release in the body. For example, microspheres can be administered via intradermal injection of drug-containing microspheres, which slowly release subcutaneously (see Rao, *J. Biomater Sci. Polym. Ed.* 7:623-645, 1995; as biodegradable and injectable gel formulations (see, e.g., Gao *Pharm. Res.* 12:857-863, 1995); or, as microspheres for oral administration (see, e.g., Eyles, *J. Pharm. Pharmacol.* 49:669-674, 1997). In another embodiment, the formulations of the compositions of the present invention can be delivered by the use of liposomes which fuse with the cellular membrane or are endocytosed, *i.e.*, by employing receptor ligands attached to the liposome, that bind to surface membrane protein receptors of the cell resulting in endocytosis. By using liposomes, particularly where the liposome surface carries receptor ligands specific for target cells, or are otherwise preferentially directed to a specific organ, one can focus the delivery of the compositions of the present invention into the target cells *in vivo*. (See, *e.g.*, Al-Muhammed, *J. Microencapsul.* 13:293-306, 1996; Chonn, *Curr. Opin. Biotechnol.* 6:698-708, 1995; Ostro, *Am. J. Hosp. Pharm.* 46:1576-1587, 1989). The compositions of the present invention can also be delivered as nanoparticles.

[0137] By “co-administer” it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies. The compounds of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compounds individually or in combination (more than one compound). The compositions of the present invention can be delivered transdermally, by a topical route, or formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols.

[0138] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0139] Utilizing the teachings provided herein, an effective prophylactic or therapeutic treatment regimen can be planned that does not cause substantial toxicity and yet is effective to treat the clinical symptoms demonstrated by the particular patient. This planning should involve the careful choice of active compound by considering factors such as compound potency, relative bioavailability, patient body weight, presence and severity of adverse side effects, preferred mode of administration and the toxicity profile of the selected agent.

[0140] The compounds described herein can be used in combination with one another, with other active agents known to be useful in treating cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0141] In some embodiments, co-administration includes administering one active agent within 0.5, 1, 2, 4, 6, 8, 10, 12, 16, 20, 24 hours, 2 days, 4 days, 1 week or 1 month of a second active agent. Co-administration includes administering two active agents simultaneously, approximately simultaneously (e.g., within about 1, 5, 10, 15, 20, or 30 minutes of each other), or sequentially in any order. In some embodiments, co-administration can be accomplished by co-formulation, i.e., preparing a single pharmaceutical composition including both active agents. In other embodiments, the active agents can be formulated separately. In another embodiment, the active and/or adjunctive agents may be linked or conjugated to one another. In some embodiments, the compounds described herein may be combined with treatments for cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0142] The compounds described herein can be administered to treat a fertility disease or disorder and/or cancer. In this regard, the compounds disclosed herein may be administered

either alone to treat such diseases or disorders or may be co-administered with another therapeutic agent to treat such diseases or disorders.

[0143] The compounds disclosed herein may be co-administered with an agonist or antagonist of CRF₁ function. The compounds described herein can be co-administered with conventional anti-cancer/chemotherapeutic.

[0144] “Anti-cancer agent” is used in accordance with its plain ordinary meaning and refers to a composition (e.g. compound, drug, antagonist, inhibitor, modulator) having antineoplastic properties or the ability to inhibit the growth or proliferation of cells. In some embodiments, an anti-cancer agent is a chemotherapeutic. In some embodiments, an anti-cancer agent is an agent identified herein having utility in methods of treating cancer. In some embodiments, an anti-cancer agent is an agent approved by the FDA or similar regulatory agency of a country other than the USA, for treating cancer. Examples of anti-cancer agents include, but are not limited to, MEK (e.g. MEK1, MEK2, or MEK1 and MEK2) inhibitors (e.g. XL518, CI-1040, PD035901, selumetinib/ AZD6244, GSK1120212/ trametinib, GDC-0973, ARRY-162, ARRY-300, AZD8330, PD0325901, U0126, PD98059, TAK-733, PD318088, AS703026, BAY 869766), alkylating agents (e.g., cyclophosphamide, ifosfamide, chlorambucil, busulfan, melphalan, mechlorethamine, uramustine, thiotepa, nitrosoureas, nitrogen mustards (e.g., mechlorethamine, cyclophosphamide, chlorambucil, melphalan), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomustine, semustine, streptozocin), triazenes (decarbazine)), anti-metabolites (e.g., 5-azathioprine, leucovorin, capecitabine, fludarabine, gemcitabine, pemetrexed, raltitrexed, folic acid analog (e.g., methotrexate), or pyrimidine analogs (e.g., fluorouracil, floxouridine, Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin), *etc.*), plant alkaloids (e.g., vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, paclitaxel, docetaxel, *etc.*), topoisomerase inhibitors (e.g., irinotecan, topotecan, amsacrine, etoposide (VP16), etoposide phosphate, teniposide, *etc.*), antitumor antibiotics (e.g., doxorubicin, adriamycin, daunorubicin, epirubicin, actinomycin, bleomycin, mitomycin, mitoxantrone, plicamycin, *etc.*), platinum-based compounds (e.g. cisplatin, oxaloplatin, carboplatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant (e.g., mitotane, aminoglutethimide), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), inhibitors of mitogen-activated protein kinase signaling (e.g. U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-9006, wortmannin, or LY294002, Syk inhibitors, mTOR inhibitors, antibodies (e.g., rituxan), gossypol, genasense, polyphenol E, Chlorofusin, all trans-retinoic acid (ATRA), bryostatin,

tumor necrosis factor-related apoptosis-inducing ligand (TRAIL), 5-aza-2'-deoxycytidine, all trans retinoic acid, doxorubicin, vincristine, etoposide, gemcitabine, imatinib (Gleevec.RTM.), geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), flavopiridol, LY294002, bortezomib, trastuzumab, BAY 11-7082, PKC412, PD184352, 20-epi-1, 25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimestine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatan; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrididemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; 9-dioxamycin; diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilomofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins;

iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole;
isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide;
leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor;
leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarazole;
linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds;
lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin
A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors;
menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone;
miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol;
mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone;
mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin;
monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene
inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B;
mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides;
nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin;
nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide
modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone;
oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer;
ormaplatin; osaterone; oxaliplatin; oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic
acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan
polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol;
phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride;
pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum
complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin;
prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based
immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein
tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins;
pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists;
raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP
inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII
retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol;
saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence
derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction
modulators; single chain antigen-binding protein; sizofuran; sobuzoxane; sodium borocaptate;

sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; tricyribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; zinostatin stimalamer, Adriamycin, Dactinomycin, Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqune; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; iimofosine; interleukin II (including recombinant interleukin II, or rIL.sub.2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1; interferon alfa-n3; interferon beta-1a; interferon gamma-1b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan;

menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedapa;
mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane;
mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin;
oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide;
pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium;
porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride;
pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene;
sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin;
streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone
hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine;
thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricitabine phosphate;
trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard;
uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine
sulfate; vinepidine sulfate; vinylicinate sulfate; vinleurosine sulfate; vinorelbine tartrate;
vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride,
agents that arrest cells in the G2-M phases and/or modulate the formation or stability of
microtubules, (e.g. Taxol.TM (i.e. paclitaxel), Taxotere.TM, compounds comprising the taxane
skeleton, Eribulin (i.e. R-55104), Dolastatin 10 (i.e. DLS-10 and NSC-376128), Mivobulin
isethionate (i.e. as CI-980), Vincristine, NSC-639829, Discodermolide (i.e. as NVP-XX-A-296),
ABT-751 (Abbott, i.e. E-7010), Altorhyrtins (e.g. Altorhyrtin A and Altorhyrtin C),
Spongistatins (e.g. Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5,
Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride
(i.e. LU-103793 and NSC-D-669356), Epothilones (e.g. Epothilone A, Epothilone B, Epothilone
C (i.e. desoxyepothilone A or dEpoA), Epothilone D (i.e. KOS-862, dEpoB, and
desoxyepothilone B), Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide,
16-aza-epothilone B, 21-aminoepothilone B (i.e. BMS-310705), 21-hydroxyepothilone D (i.e.
Desoxyepothilone F and dEpoF), 26-fluoroepothilone, Auristatin PE (i.e. NSC-654663),
Soblidotin (i.e. TZT-1027), LS-4559-P (Pharmacia, i.e. LS-4577), LS-4578 (Pharmacia, i.e. LS-
477-P), LS-4477 (Pharmacia), LS-4559 (Pharmacia), RPR-112378 (Aventis), Vincristine sulfate,
DZ-3358 (Daiichi), FR-182877 (Fujisawa, i.e. WS-9885B), GS-164 (Takeda), GS-198 (Takeda),
KAR-2 (Hungarian Academy of Sciences), BSF-223651 (BASF, i.e. ILX-651 and LU-223651),
SAH-49960 (Lilly/Novartis), SDZ-268970 (Lilly/Novartis), AM-97 (Armad/Kyowa Hakko),
AM-132 (Armad), AM-138 (Armad/Kyowa Hakko), IDN-5005 (Indena), Cryptophycin 52 (i.e.
LY-355703), AC-7739 (Ajinomoto, i.e. AVE-8063A and CS-39.HCl), AC-7700 (Ajinomoto, i.e.
AVE-8062, AVE-8062A, CS-39-L-Ser.HCl, and RPR-258062A), Vitilevuamide, Tubulysin A,

Canadensol, Centaureidin (i.e. NSC-106969), T-138067 (Tularik, i.e. T-67, TL-138067 and TI-138067), COBRA-1 (Parker Hughes Institute, i.e. DDE-261 and WHI-261), H10 (Kansas State University), H16 (Kansas State University), Oncocidin A1 (i.e. BTO-956 and DIME), DDE-313 (Parker Hughes Institute), Fijianolide B, Laulimalide, SPA-2 (Parker Hughes Institute), SPA-1 (Parker Hughes Institute, i.e. SPIKET-P), 3-IAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-569), Narcosine (also known as NSC-5366), Nascapine, D-24851 (Asta Medica), A-105972 (Abbott), Hemiasterlin, 3-BAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-191), TMPN (Arizona State University), Vanadocene acetylacetonate, T-138026 (Tularik), Monsatrol, Inanocine (i.e. NSC-698666), 3-IAABE (Cytoskeleton/Mt. Sinai School of Medicine), A-204197 (Abbott), T-607 (Tularik, i.e. T-900607), RPR-115781 (Aventis), Eleutherobins (such as Desmethyleleutherobin, Desaetyeleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, D-64131 (Asta Medica), D-68144 (Asta Medica), Diazonamide A, A-293620 (Abbott), NPI-2350 (Nereus), Taccalonolide A, TUB-245 (Aventis), A-259754 (Abbott), Diozostatin, (-)-Phenylahistin (i.e. NSCL-96F037), D-68838 (Asta Medica), D-68836 (Asta Medica), Myoseverin B, D-43411 (Zentaris, i.e. D-81862), A-289099 (Abbott), A-318315 (Abbott), HTI-286 (i.e. SPA-110, trifluoroacetate salt) (Wyeth), D-82317 (Zentaris), D-82318 (Zentaris), SC-12983 (NCI), Resverastatin phosphate sodium, BPR-OY-007 (National Health Research Institutes), and SSR-250411 (Sanofi)), steroids (*e.g.*, dexamethasone), finasteride, aromatase inhibitors, gonadotropin-releasing hormone agonists (GnRH) such as goserelin or leuprolide, adrenocorticosteroids (*e.g.*, prednisone), progestins (*e.g.*, hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (*e.g.*, diethylstilbestrol, ethinyl estradiol), antiestrogen (*e.g.*, tamoxifen), androgens (*e.g.*, testosterone propionate, fluoxymesterone), antiandrogen (*e.g.*, flutamide), immunostimulants (*e.g.*, Bacillus Calmette-Guérin (BCG), levamisole, interleukin-2, alpha-interferon, *etc.*), monoclonal antibodies (*e.g.*, anti-CD20, anti-HER2, anti-CD52, anti-HLA-DR, and anti-VEGF monoclonal antibodies), immunotoxins (*e.g.*, anti-CD33 monoclonal antibody-calicheamicin conjugate, anti-CD22 monoclonal antibody-pseudomonas exotoxin conjugate, *etc.*), radioimmunotherapy (*e.g.*, anti-CD20 monoclonal antibody conjugated to ¹¹¹In, ⁹⁰Y, or ¹³¹I, *etc.*), triptolide, homoharringtonine, dactinomycin, doxorubicin, epirubicin, topotecan, itraconazole, vindesine, cerivastatin, vincristine, deoxyadenosine, sertraline, pitavastatin, irinotecan, clofazimine, 5-nonyloxytryptamine, vemurafenib, dabrafenib, erlotinib, gefitinib, EGFR inhibitors, epidermal growth factor receptor (EGFR)-targeted therapy or therapeutic (*e.g.* gefitinib (Iressa™), erlotinib (Tarceva™), cetuximab (Erbix™), lapatinib (Tykerb™), panitumumab (Vectibix™), vandetanib (Caprelsa™), afatinib/BIBW2992, CI-1033/canertinib, neratinib/HKI-272, CP-724714, TAK-285, AST-1306, ARRY334543, ARRY-

380, AG-1478, dacomitinib/PF299804, OSI-420/desmethyl erlotinib, AZD8931, AEE788, pelitinib/EKB-569, CUDC-101, WZ8040, WZ4002, WZ3146, AG-490, XL647, PD153035, BMS-599626), sorafenib, imatinib, sunitinib, dasatinib, or the like.

[0145] “Chemotherapeutic” or “chemotherapeutic agent” is used in accordance with its plain ordinary meaning and refers to a chemical composition or compound having antineoplastic properties or the ability to inhibit the growth or proliferation of cells.

[0146] The compounds disclosed herein may be co-administered with an antiproliferative/antineoplastic drug or a combination thereof, as used in medical oncology, such as an alkylating agent (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan or a nitrosourea); an antimetabolite (for example an antifolate such as a fluoropyrimidine like 5-fluorouracil or tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, gemcitabine or paclitaxel); an antitumour antibiotic (for example an anthracycline such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin or mithramycin); an antimitotic agent (for example a vinca alkaloid such as vincristine, vinblastine, vindesine or vinorelbine, or a taxoid such as taxol or taxotere); or a topoisomerase inhibitor (for example an epipodophyllotoxin such as etoposide, teniposide, amsacrine, topotecan or a camptothecin); (ii) a cytostatic agent such as an antioestrogen (for example tamoxifen, toremifene, raloxifene, droloxifene or iodoxyfene), an oestrogen receptor down regulator (for example fulvestrant), an antiandrogen (for example bicalutamide, flutamide, nilutamide or cyproterone acetate), a LHRH antagonist or LHRH agonist (for example goserelin, leuprorelin or buserelin), a progestogen (for example megestrol acetate), an aromatase inhibitor (for example as anastrozole, letrozole, vorazole or exemestane) or an inhibitor of 5.alpha.-reductase such as finasteride; (iii) an agent which inhibits cancer cell invasion (for example a metalloproteinase inhibitor like marimastat or an inhibitor of urokinase plasminogen activator receptor function); (iv) an inhibitor of growth factor function, for example: a growth factor antibody (for example the anti-erbB2 antibody trastuzumab, or the anti-erbB1 antibody cetuximab [C225]), a farnesyl transferase inhibitor, a tyrosine kinase inhibitor or a serine/threonine kinase inhibitor, an inhibitor of the epidermal growth factor family (for example an EGFR family tyrosine kinase inhibitor such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD 1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) or 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), an inhibitor of the platelet-derived growth factor family, or an inhibitor of the hepatocyte growth factor family; (v) an antiangiogenic agent such as one which inhibits the effects of vascular endothelial growth factor (for example the anti-vascular endothelial cell growth factor antibody bevacizumab, a

compound disclosed in WO 97/22596, WO 97/30035, WO 97/32856 or WO 98/13354), or a compound that works by another mechanism (for example linomide, an inhibitor of integrin .alpha.v.beta.3 function or an angiostatin); (vi) a vascular damaging agent such as combretastatin A4, or a compound disclosed in WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 or WO 02/08213; (vii) an agent used in antisense therapy, for example one directed to one of the targets listed above, such as ISIS 2503, an anti-ras antisense; (viii) an agent used in a gene therapy approach, for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; or (ix) an agent used in an immunotherapeutic approach, for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

[0147] In embodiments, the compounds disclosed herein can be co-administered with an antibody, such as a monoclonal antibody targeting B-Lymphocytes (such as CD20 (rituximab), MRA-aIL16R and T-Lymphocytes, CTLA4-Ig, HuMax IL-15) or antibody modulating Ig function such as anti-IgE (for example omalizumab).

[0148] In embodiments, treatment of cancer includes administration of an effective amount of at least two of the following: a CRF1 inhibitor, an inhibitor of the PD-L1/PD-1 pathway, an inhibitor of CTLA-4, an agonistic antibody of CD137 (4-1BB). In some embodiments, the method may comprise the use of two or more combinations.

[0149] In a further embodiment, the compounds described herein can be co-administered with conventional radiotherapeutic agents including, but not limited to, radionuclides such as ⁴⁷Sc, ⁶⁴Cu, ⁶⁷Cu, ⁸⁹Sr, ⁸⁶Y, ⁸⁷Y, ⁹⁰Y, ¹⁰⁵Rh, ¹¹¹Ag, ¹¹¹In, ^{117m}Sn, ¹⁴⁹Pm, ¹⁵³Sm, ¹⁶⁶Ho, ¹⁷⁷Lu, ¹⁸⁶Re, ¹⁸⁸Re, ²¹¹At, and ²¹²Bi, optionally conjugated to antibodies directed against tumor antigens.

[0150] In addition, a CRF1 inhibitor may be combined with the therapeutic administration of immune cells, sometimes referred to as adoptive cell transfer. These cells may be cells from the patient, a genetically related or unrelated donor, they may be genetically modified (e.g. CAR-T cells, NK cells, etc), cell lines, genetically modified cell lines and live or dead versions of the above. CRF1 inhibitors may also be combined with vaccines of any kind (e.g. protein/peptide, viral, bacterial, cellular) to stimulate immune responses to cancer.

[0151] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramusculay, intranasal, intraocularal, intrapericardial, intraperitoneal, intrapleural, intraprostatical, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenousl, intravesicullar, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0152] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramusculay, intranasal, intraocularal, intrapericardial, intraperitoneal, intrapleural, intraprostatical, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenousl, intravesicullar, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0153] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered once daily until study reached endpoint. The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered at least three times but in some studies four or more times depending on the length of the study and/or the design of the study.

[0154] The methods disclosed herein may be used in combination with additional cancer therapy. In some embodiments, the distinct cancer therapy comprises surgery, radiotherapy, chemotherapy, toxin therapy, immunotherapy, cryotherapy or gene therapy. In some embodiments, the cancer is a chemotherapy-resistant or radio-resistant cancer.

[0155] A “cell” as used herein, refers to a cell carrying out metabolic or other function sufficient to preserve or replicate its genomic DNA. A cell can be identified by well-known methods in the art including, for example, presence of an intact membrane, staining by a particular dye, ability to produce progeny or, in the case of a gamete, ability to combine with a second gamete to produce a viable offspring. Cells may include prokaryotic and eukaryotic cells. Prokaryotic cells include but are not limited to bacteria. Eukaryotic cells include but are not limited to yeast cells and cells derived from plants and animals, for example mammalian, insect (*e.g.*, spodoptera) and human cells. Cells may be useful when they are naturally nonadherent or have been treated not to adhere to surfaces, for example by trypsinization.

[0156] “Control” or “control experiment” is used in accordance with its plain ordinary meaning and refers to an experiment in which the subjects or reagents of the experiment are treated as in a parallel experiment except for omission of a procedure, reagent, or variable of the experiment. In some instances, the control is used as a standard of comparison in evaluating experimental effects. In some embodiments, a control is the measurement of the activity of a protein in the absence of a compound as described herein (including embodiments and examples).

[0157] The term “modulator” refers to a composition that increases or decreases the level of a target molecule or the function of a target molecule or the physical state of the target of the molecule. In some embodiments, a CRF₁ associated disease modulator is a compound that reduces the severity of one or more symptoms of a disease associated with CRF₁ (e.g. an endocrine disorder, a cancer including but not limited to TART or OART). A CRF₁ modulator is a compound that increases or decreases the activity or function or level of activity or level of function of CRF₁. A modulator may act alone, or it may use a cofactor, e.g., a protein, metal ion, or small molecule. Examples of modulators include small molecule compounds and other bioorganic molecules. Numerous libraries of small molecule compounds (e.g., combinatorial libraries) are commercially available and can serve as a starting point for identifying a modulator. The skilled artisan is able to develop one or more assays (e.g., biochemical or cell-based assays) in which such compound libraries can be screened in order to identify one or more compounds having the desired properties; thereafter, the skilled medicinal chemist is able to optimize such one or more compounds by, for example, synthesizing and evaluating analogs and derivatives thereof. Synthetic and/or molecular modeling studies can also be utilized in the identification of an activator.

[0158] The term “modulate” is used in accordance with its plain ordinary meaning and refers to the act of changing or varying one or more properties. “Modulation” refers to the process of changing or varying one or more properties. For example, as applied to the effects of a modulator on a target protein, to modulate means to change by increasing or decreasing a property or function of the target molecule or the amount of the target molecule. In embodiments, the terms “modulate”, “modulation” and the like refer to the ability of a molecule (e.g., an activator or an inhibitor) to increase or decrease the function or activity of CRF₁, either directly or indirectly, relative to the absence of the molecule.

[0159] The term “associated” or “associated with” in the context of a substance or substance activity or function associated with a disease (e.g. a protein associated disease, an endocrine disorder, a disease associated with CRF₁ activity, a CRF₁ associated disease (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART))) means that the disease (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) is caused by (in

whole or in part), or a symptom of the disease is caused by (in whole or in part) the substance or substance activity or function. For example, an endocrine disorder associated with CRF₁ activity or function may be an endocrine disorder that results (entirely or partially) from aberrant CRF₁ function (e.g. enzyme activity, protein-protein interaction, signaling pathway) or an endocrine disorder wherein a particular symptom of the disease is caused (entirely or partially) by aberrant CRF₁ activity or function. As used herein, what is described as being associated with a disease, if a causative agent, could be a target for treatment of the disease. For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated disease (e.g., an endocrine disorder including but not limited to TART or OART), may be treated with a compound described herein (e.g., CRF₁ modulator, CRF₁ inhibitor, or CRF₁ antagonist), in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease (e.g., an endocrine disorder such as TART or OART). For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated endocrine disorder, may be treated with an CRF₁ modulator or CRF₁ inhibitor, in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease.

[0160] The term “aberrant” as used herein refers to different from normal. When used to describe enzymatic activity or protein function, aberrant refers to activity or function that is greater or less than a normal control or the average of normal non-diseased control samples. Aberrant activity may refer to an amount of activity that results in a disease, wherein returning the aberrant activity to a normal or non-disease-associated amount (e.g. by administering a compound or using a method as described herein), results in reduction of the disease or one or more disease symptoms.

[0161] The term “signaling pathway” as used herein refers to a series of interactions between cellular and optionally extra-cellular components (e.g. proteins, nucleic acids, small molecules, ions, lipids) that conveys a change in one component to one or more other components, which in turn may convey a change to additional components, which is optionally propagated to other signaling pathway components. For example, binding of a CRF₁ with a compound as described herein may reduce the level of a product of the CRF₁ catalyzed reaction or the level of a downstream derivative of the product or binding may reduce the interactions between the CRF₁ or a reaction product and downstream effectors or signaling pathway components (e.g., MAP kinase pathway), resulting in changes in cell growth, proliferation, or survival.

[0162] As used herein, the terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” and all other related art-accepted terms, many of which are set forth below, refer to a compound capable of modulating, either directly or indirectly, the CRF₁ receptor in an *in vitro* assay, an *in vivo* model, and/or other

means indicative of therapeutic efficacy. The terms also refer to a compound that exhibits at least some therapeutic benefit in a human subject.

[0163] The phrase “in a sufficient amount to effect a change” means that there is a detectable difference between a level of an indicator measured before (e.g., a baseline level) and after administration of a particular therapy. Indicators include any objective parameter (e.g., serum concentration) or subjective parameter (e.g., a subject’s feeling of well-being).

[0164] The “activity” of a molecule may describe or refer to the binding of the molecule to a ligand or to a receptor; to catalytic activity; to the ability to stimulate gene expression or cell signaling, differentiation, or maturation; to antigenic activity; to the modulation of activities of other molecules; and the like. The term “proliferative activity” encompasses an activity that promotes, that is necessary for, or that is specifically associated with, for example, normal cell division, as well as cancer, tumors, dysplasia, cell transformation, metastasis, and angiogenesis.

[0165] “Substantially pure” indicates that a component makes up greater than about 50% of the total content of the composition, and typically greater than about 60% of the total polypeptide content. More typically, “substantially pure” refers to compositions in which at least 75%, at least 85%, at least 90% or more of the total composition is the component of interest. In some cases, the polypeptide will make up greater than about 90%, or greater than about 95% of the total content of the composition (percentage in a weight per weight basis).

[0166] The terms “specifically binds” and “selectively binds,” when referring to a ligand/receptor, antibody/antigen, or other binding pair, indicates a binding reaction which is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated conditions, a specified ligand binds to a particular receptor and does not bind in a significant amount to other proteins present in the sample. The antibody, or binding composition derived from the antigen-binding site of an antibody, of the contemplated method binds to its antigen, or a variant or mutein thereof, with an affinity that is at least two-fold greater, at least 10-times greater, at least 20-times greater, or at least 100-times greater than the affinity with any other antibody, or binding composition derived therefrom. In embodiments, the antibody will have an affinity that is greater than about 10^9 liters/mol, as determined by, e.g., Scatchard analysis (Munsen, et al. (1980) *Analyt. Biochem.* 107:220-239).


[0167] The terms “DNA,” “nucleic acid,” “nucleic acid molecule,” “polynucleotide” and the like are used interchangeably herein to refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Non-limiting examples of polynucleotides include linear and circular nucleic acids, messenger RNA (mRNA), complementary DNA (cDNA), recombinant polynucleotides, vectors, probes, primers and the like.

[0168] As used herein, the terms “variants” and “homologs” are used interchangeably to refer to amino acid or nucleic acid sequences that are similar to reference amino acid or nucleic acid sequences, respectively. The term encompasses naturally-occurring variants and non-naturally-occurring variants. Naturally-occurring variants include homologs (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one species to another), and allelic variants (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one individual to another within a species). Thus, variants and homologs encompass naturally occurring amino acid and nucleic acid sequences encoded thereby and their isoforms, as well as splice variants of a protein or gene. The terms also encompass nucleic acid sequences that vary in one or more bases from a naturally-occurring nucleic acid sequence but still translate into an amino acid sequence that corresponds to the naturally-occurring protein due to degeneracy of the genetic code. Non-naturally-occurring variants and homologs include polypeptides and nucleic acids that comprise a change in amino acid or nucleotide sequence, respectively, where the change in sequence is artificially introduced (e.g., muteins); for example, the change is generated in the laboratory by human intervention (“hand of man”). Therefore, non-naturally occurring variants and homologs may also refer to those that differ from the naturally-occurring sequences by one or more conservative substitutions and/or tags and/or conjugates.

[0169] The term “muteins” as used herein refers broadly to mutated recombinant proteins. These proteins usually carry single or multiple amino acid substitutions and are frequently derived from cloned genes that have been subjected to site-directed or random mutagenesis, or from completely synthetic genes.





II. Compounds

[0170] In an aspect provided herein, is a compound of Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 and R^2 are independently ethyl or n-propyl. R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R^4 is hydrogen,

Br, R^aR^bN- , methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl,




. R^a and R^b are independently hydrogen, C_1 - C_3 alkyl,

$\text{H}_2\text{NCH}_2\text{CH}_2-$, $(\text{CH}_3)_3\text{COC}(\text{O})\text{NHCH}_2\text{CH}_2-$, or $\text{CH}_3\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2-$ In embodiments, the inhibitor forms an irreversible bond with the amino acid residue. In embodiments, the bond is a covalent bond.


[0171] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0172] In certain embodiments, R^3 is Cl, Br, or methyl.

[0173] In certain embodiments, R^4 is Br, $\text{R}^a\text{R}^b\text{N}-$, pyridin-4-yl, morpholin-4-yl, or .


[0174] In certain embodiments, R^4 is morpholin-4-yl or .

[0175] In certain embodiments, R^4 is hydrogen, Br, $\text{R}^a\text{R}^b\text{N}-$ and R^a and R^b are independently C_1 - C_3 alkyl.




[0176] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0177] In certain embodiments, the compound is



[0178] In certain embodiments, the compound is



[0179] In one aspect described herein, there is a method of treating a subject in need thereof comprising administering an astressin or a pharmaceutically acceptable salt or solvate thereof.

An astressin generally refers to a nonselective corticotropin releasing hormone antagonist that reduces the synthesis of ACTH and cortisol.

[0180] In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, the method comprising: (i) measuring a hormone level in the subject in need thereof; (ii) administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof; and repeating steps (i) and (ii) until the hormone level reaches a pre-determined range followed by a maintenance therapy of a daily dosing of said astressin.

[0181] In one aspect described herein, there is a method of improving hyperandrogenic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating menstrual irregularity, ovulatory dysfunction or infertility, in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of improving metabolic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto; or a pharmaceutically acceptable salt or solvate thereof.

[0182] In one aspect described herein, there is a method of improving quality of life in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

III. Pharmaceutical Compositions

[0183] In an aspect, there is provided a pharmaceutical composition, including a compound as described herein, including embodiments, or the structural Formula (I), and a pharmaceutically acceptable excipient.

[0184] The compounds (e.g., CRF₁ inhibitors/antagonists) of the present disclosure may be in the form of compositions suitable for administration to a subject. In general, such compositions are “pharmaceutical compositions” comprising a compound (e.g., CRF₁ inhibitor(s)/antagonist(s)) and one or more pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients. In certain embodiments, the compound (e.g., CRF₁ inhibitor/antagonist) are present in a therapeutically acceptable amount. The pharmaceutical compositions may be used in the methods of the present disclosure; thus, for example, the

pharmaceutical compositions can be administered *ex vivo* or *in vivo* to a subject in order to practice the therapeutic and prophylactic methods and uses described herein.

[0185] In one aspect described herein, there is a pharmaceutical composition in the form of a capsule comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a pharmaceutical composition in the form of a tablet comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

[0186] The pharmaceutical compositions of the present disclosure can be formulated to be compatible with the intended method or route of administration; exemplary routes of administration are set forth herein.

[0187] The pharmaceutical compositions containing the active ingredient (e.g., an inhibitor of CRF₁ function) may be in a form suitable for oral use, for example, as tablets, capsules, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups, solutions, microbeads or elixirs. Pharmaceutical compositions intended for oral use may be prepared for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents such as, for example, sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets, capsules and the like contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture thereof. These excipients may be, for example, diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc.

[0188] The tablets, capsules and the like suitable for oral administration may be uncoated or coated to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action. For example, a time-delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated to form osmotic therapeutic tablets for controlled release. Additional agents include biodegradable or biocompatible particles or a polymeric substance such as polyesters, polyamine acids, hydrogel, polyvinyl pyrrolidone, polyanhydrides, polyglycolic acid, ethylene-vinylacetate, methylcellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymers, polylactide/glycolide copolymers, or ethylenevinylacetate copolymers in order to control delivery of an administered composition. For example, the oral agent can be entrapped in microcapsules prepared by coacervation techniques or by interfacial polymerization, by the use

of hydroxymethylcellulose or gelatin-microcapsules or poly(methylmethacrolate) microcapsules, respectively, or in a colloid drug delivery system. Colloidal dispersion systems include macromolecule complexes, nano-capsules, microspheres, microbeads, and lipid-based systems, including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Methods for the preparation of the above-mentioned formulations will be apparent to those skilled in the art.

[0189] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate, kaolin or microcrystalline cellulose, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

[0190] Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture thereof. Such excipients can be suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents, for example a naturally-occurring phosphatide (e.g., lecithin), or condensation products of an alkylene oxide with fatty acids (e.g., polyoxy-ethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols (e.g., for heptadecaethyleneoxycetanol), or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol (e.g., polyoxyethylene sorbitol monooleate), or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides (e.g., polyethylene sorbitan monooleate). The aqueous suspensions may also contain one or more preservatives.

[0191] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

[0192] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, and optionally one or more suspending agents and/or preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified herein.

[0193] The pharmaceutical compositions of the present disclosure may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example, liquid paraffin, or mixtures of these. Suitable emulsifying agents may be naturally occurring gums, for example, gum acacia or gum tragacanth; naturally occurring phosphatides, for example, soy bean, lecithin, and esters or partial esters derived from

fatty acids; hexitol anhydrides, for example, sorbitan monooleate; and condensation products of partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate.

[0194] The pharmaceutical compositions typically comprise a therapeutically effective amount of a CRF₁ inhibitor/antagonist contemplated by the present disclosure and one or more pharmaceutically and physiologically acceptable formulation agents. Suitable pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients include, but are not limited to, antioxidants (e.g., ascorbic acid and sodium bisulfate), preservatives (e.g., benzyl alcohol, methyl parabens, ethyl or n-propyl, p-hydroxybenzoate), emulsifying agents, suspending agents, dispersing agents, solvents, fillers, bulking agents, detergents, buffers, vehicles, diluents, and/or adjuvants. For example, a suitable vehicle may be physiological saline solution or citrate-buffered saline, possibly supplemented with other materials common in pharmaceutical compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Those skilled in the art will readily recognize a variety of buffers that can be used in the pharmaceutical compositions and dosage forms contemplated herein. Typical buffers include, but are not limited to, pharmaceutically acceptable weak acids, weak bases, or mixtures thereof. As an example, the buffer components can be water soluble materials such as phosphoric acid, tartaric acids, lactic acid, succinic acid, citric acid, acetic acid, ascorbic acid, aspartic acid, glutamic acid, and salts thereof. Acceptable buffering agents include, for example, a Tris buffer; N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) (HEPES); 2-(N-Morpholino)ethanesulfonic acid (MES); 2-(N-Morpholino)ethanesulfonic acid sodium salt (MES); 3-(N-Morpholino)propanesulfonic acid (MOPS); and N-tris[Hydroxymethyl]methyl-3-aminopropanesulfonic acid (TAPS).

[0195] After a pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form, a lyophilized form requiring reconstitution prior to use, a liquid form requiring dilution prior to use, or other acceptable form. In some embodiments, the pharmaceutical composition is provided in a single-use container (e.g., a single-use vial, ampule, syringe, or autoinjector (similar to, e.g., an EpiPen®)), whereas a multi-use container (e.g., a multi-use vial) is provided in other embodiments.

[0196] Formulations can also include carriers to protect the composition against rapid degradation or elimination from the body, such as a controlled release formulation, including liposomes, hydrogels, prodrugs and microencapsulated delivery systems. For example, a time-delay material such as glyceryl monostearate or glyceryl stearate alone, or in combination with a wax, may be employed. Any drug delivery apparatus may be used to deliver a CRF₁

inhibitor/antagonist, including implants (e.g., implantable pumps) and catheter systems, slow injection pumps and devices, all of which are well known to the skilled artisan.

[0197] Depot injections, which are generally administered subcutaneously or intramuscularly, may also be utilized to release the compound (e.g., CRF₁ inhibitor/antagonist) disclosed herein over a defined period of time. Depot injections are usually either solid- or oil-based and generally comprise at least one of the formulation components set forth herein. One of ordinary skill in the art is familiar with possible formulations and uses of depot injections.

[0198] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents mentioned herein. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Acceptable diluents, solvents and dispersion media that may be employed include water, Ringer's solution, isotonic sodium chloride solution, Cremophor® EL (BASF, Parsippany, NJ) or phosphate buffered saline (PBS), ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. In addition, sterile fixed oils are conventionally employed as a solvent or suspending medium; for this purpose, any bland fixed oil may be employed, including synthetic mono- or diglycerides. Moreover, fatty acids, such as oleic acid, find use in the preparation of injectables. Prolonged absorption of particular injectable formulations can be achieved by including an agent that delays absorption (e.g., aluminum monostearate or gelatin).

[0199] The present disclosure contemplates the administration of the compound (e.g., CRF₁ inhibitor/antagonist) in the form of suppositories for rectal administration. The suppositories can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include, but are not limited to, cocoa butter and polyethylene glycols.

[0200] The compound (e.g., CRF₁ inhibitor/antagonist) contemplated by the present disclosure may be in the form of any other suitable pharmaceutical composition (e.g., sprays for nasal or inhalation use) currently known or developed in the future.

IV. Methods

[0201] In certain embodiments, provided herein are methods for treating or preventing a disease comprising administering an effective amount of a compound of Formula (I), or an isotopic variant thereof or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0202] In one embodiment, provided herein are methods for treating or preventing a tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a benign tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a malignant tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal rest tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0203] In one embodiment, provided herein are methods for treating or preventing infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing male infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm count, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm motility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm morphology, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing female infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0204] In one embodiment, provided herein are methods for treating or preventing a proliferative or hyperproliferative disease or disorder, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, the proliferative disease is cancer. In certain embodiments, the proliferative disease is a cancer of the breast, skin, prostate, cervix,

uterus, ovary, testes, bladder, lung, liver, larynx, oral cavity, colon and gastrointestinal tract (e.g., esophagus, stomach, pancreas), brain, thyroid, blood, and lymphatic system.

[0205] In certain embodiments, the cancer treatable with the methods provided herein includes, but is not limited to, (1) leukemias, including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome or a symptom thereof (such as anemia, thrombocytopenia, neutropenia, bicytopenia or pancytopenia), refractory anemia (RA), RA with ringed sideroblasts (RARS), RA with excess blasts (RAEB), RAEB in transformation (RAEB-T), preleukemia, and chronic myelomonocytic leukemia (CMML), (2) chronic leukemias, including, but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, and hairy cell leukemia; (3) polycythemia vera; (4) lymphomas, including, but not limited to, Hodgkin's disease and non-Hodgkin's disease; (5) multiple myelomas, including, but not limited to, smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma, and extramedullary plasmacytoma; (6) Waldenstrom's macroglobulinemia; (7) monoclonal gammopathy of undetermined significance; (8) benign monoclonal gammopathy; (9) heavy chain disease; (10) bone and connective tissue sarcomas, including, but not limited to, bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, metastatic cancers, neurilemmoma, rhabdomyosarcoma, and synovial sarcoma; (11) brain tumors, including, but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, and primary brain lymphoma; (12) breast cancer, including, but not limited to, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, primary cancers, Paget's disease, and inflammatory breast cancer; (13) adrenal cancer, including, but not limited to, pheochromocytoma and adrenocortical carcinoma; (14) thyroid cancer, including, but not limited to, papillary or follicular thyroid cancer, medullary thyroid cancer, and anaplastic thyroid cancer; (15) pancreatic cancer, including, but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; (16) pituitary cancer, including, but limited to, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; (17) eye cancer, including, but not limited, to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; (18) vaginal

cancer, including, but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; (19) vulvar cancer, including, but not limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; (20) cervical cancers, including, but not limited to, squamous cell carcinoma, and adenocarcinoma; (21) uterine cancer, including, but not limited to, endometrial carcinoma and uterine sarcoma; (22) ovarian cancer, including, but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; (23) esophageal cancer, including, but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; (24) stomach cancer, including, but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; (25) colon cancer; (26) rectal cancer; (27) liver cancer, including, but not limited to, hepatocellular carcinoma and hepatoblastoma; (28) gallbladder cancer, including, but not limited to, adenocarcinoma; (29) cholangiocarcinomas, including, but not limited to, papillary, nodular, and diffuse; (30) lung cancer, including, but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma, and small-cell lung cancer; (31) testicular cancer, including, but not limited to, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, and choriocarcinoma (yolk-sac tumor); (32) prostate cancer, including, but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; (33) penile cancer; (34) oral cancer, including, but not limited to, squamous cell carcinoma; (35) basal cancer; (36) salivary gland cancer, including, but not limited to, adenocarcinoma, mucoepidermoid carcinoma, and adenoicystic carcinoma; (37) pharynx cancer, including, but not limited to, squamous cell cancer and verrucous; (38) skin cancer, including, but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, and acral lentiginous melanoma; (39) kidney cancer, including, but not limited to, renal cell cancer, adenocarcinoma, hypernephroma, fibrosarcoma, and transitional cell cancer (renal pelvis and/or ureter); (40) Wilms' tumor; (41) bladder cancer, including, but not limited to, transitional cell carcinoma, squamous cell cancer, adenocarcinoma, and carcinosarcoma; and other cancer, including, not limited to, myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangio-endotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, and papillary adenocarcinomas (*See Fishman et al.*, 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy *et al.*, 1997, *Informed Decisions*:

The Complete Book of Cancer Diagnosis, Treatment, and Recovery, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

[0206] Depending on the disorder, disease, or condition to be treated, and the subject's condition, the compounds or pharmaceutical compositions provided herein can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), inhalation, nasal, vaginal, rectal, sublingual, or topical (e.g., transdermal or local) routes of administration and can be formulated, alone or together, in suitable dosage unit with pharmaceutically acceptable excipients, carriers, adjuvants, and vehicles appropriate for each route of administration as described elsewhere herein.

[0207] The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. Recommended routes of administration for the second active agents are known to those of ordinary skill in the art. See, e.g., Physicians' Desk Reference, 1755-1760 (56th ed., 2002).


[0208] In one embodiment, provided herein are methods for treating or preventing testicular adrenal rest tumors (TART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).

[0209] In one embodiment, provided herein are methods for treating or preventing ovarian adrenal rest tumors (OART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).

[0210] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.





[0211] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0212] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a compound of structural Formula (I):




, or a pharmaceutically acceptable salt thereof. R^1 and R^2 are independently ethyl or n-propyl. R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R^4 is hydrogen,

Br, $R^a R^b N-$, methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl,




. R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$.

[0213] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments.

[0214] In an aspect provided herein, is a method of decreasing or reducing adrenocorticotrophic hormone (ACTH) levels in a subject in need thereof, comprising administering to the subject contacting the ACTH with a compound of structural Formula (I):




, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 60% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 70% of the subjects demonstrate reduction in

ACTH. In certain embodiments, at least about 80% of the subjects demonstrate reduction in ACTH. In embodiments, at least about 10% of subjects were in the standard or normal range for ACTH after treatment.


[0215] In embodiments, at least about 20% of subjects were in the standard or normal range for ACTH after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for ACTH after treatment. In certain embodiments, ACTH levels are decreased or reduced within about 30 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 1 hour post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 90 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 2 hours post dosing.

[0216] In an aspect provided herein, is a method of decreasing or reducing 17 α -hydroxyprogesterone (17-OHP) levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R¹, R², R³ and R⁴ are as described herein, including embodiments. R¹, R², R³ and R⁴ are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 60% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 70% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 80% of the subjects demonstrate reduction in 17-OHP. In embodiments, at least about 30% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 40% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 60% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 60% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment.

[0217] In an aspect provided herein, is a method of decreasing or reducing androstenedione levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 60% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 70% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 80% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 90% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least 100% of the subjects demonstrate reduction in androstenedione. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least 60% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 40% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range for androstenedione after treatment.

[0218] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0219] In certain embodiments, R^3 is Cl, Br, or methyl.


[0220] In certain embodiments, R^4 is Br, $R^a R^b N-$, pyridin-4-yl, morpholin-4-yl, or



[0221] In certain embodiments, R^4 is morpholin-4-yl or




[0222] In certain embodiments, R^4 is hydrogen, Br, $R^a R^b N-$ and R^a and R^b are independently C_1-C_3 alkyl.




[0223] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0224] In certain embodiments, the compound is



[0225] In certain embodiments, the compound is



[0226] In certain embodiments, a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is cyclically administered to a patient. Cycling therapy involves the administration of an active agent or a combination of active agents for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

[0227] In certain embodiments, in the treatment, prevention, or amelioration of one or more symptoms of the disorders, diseases, or conditions described herein (e.g., treatment or prevention of testicular adrenal rest tumors (TART) or treatment or prevention of ovarian adrenal rest tumors (OART)), an appropriate dosage level of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof generally is ranging from about 1 mg to 3000 mg, from about 1 mg to 2000 mg, from about 1 mg to 1000 mg, from about 1 mg to about 500 mg, from about 5 mg to about 500 mg,

from about 5 mg to about 400 mg, from about 5 mg to about 300 mg, from about 5 mg to about 250 mg, from about 5 mg to about 200 mg, from about 50 mg to about 200 mg, from about 10 mg to about 400 mg, from about 5 mg to about 25 mg, from about 10 mg to about 25 mg, or from about 25 to about 200 mg, which can be administered in single or multiple doses. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg/day.

[0228] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 270 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is

administered in an amount of about 600 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg.

[0229] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, or about 1000 mg.

[0230] For oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing from about 1.0 mg to about 1,000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about

350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 980 mg, about 990 mg, about 990 mg, or about 1000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof for the symptomatic adjustment of the dosage to the patient to be treated. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 50 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 200 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. The pharmaceutical compositions can be administered on a regimen of one (1) to four (4) times per day, including once, twice, three times, and four times per day. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about

310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 990 mg, about 990 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg,

about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered three times per day.

[0231] In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of one day. In some embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of two days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of three days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of four days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of five days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of six days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of seven days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of eight days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of nine days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of ten days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 11 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 12 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 13 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 14 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of 15 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 16 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 17 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 18 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 19 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 20 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 21 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 22 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 23 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 24 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 25 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 26 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 27 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 28 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 29 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of 30 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 31 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 32 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 33 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate,

hydrate, or prodrug thereof is administered over a period of 34 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 35 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 36 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 37 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 38 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 39 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 40 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 41 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 42 days or more.

[0232] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days.

[0233] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least

about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 14 days.

[0234] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily

for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily

for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 14 days.

[0235] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330

mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 28 days.

[0236] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 220 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula

(I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 28 days.

[0237] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 255 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230

mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 42 days.

[0238] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 5 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula

(I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an

amount of about 370 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 42 days.

[0239] In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, or about 400 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200

mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, or about 800 mg of the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered at least once per day.

[0240] In certain embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0241] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered orally.

[0242] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is formulated as a capsule or tablet.

[0243] In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.


[0244] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0245] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day.

[0246] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 6 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 5 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 3 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 1 hour before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep.

[0247] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

[0248] In another aspect is provided a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I): and a pharmaceutically acceptable excipient comprising with a compound as described herein, including embodiments, or the structural Formulae (I) and a pharmaceutically acceptable excipient:



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 , and R^4 are as described herein, including embodiments.

[0249] In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments,

the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0250] In certain embodiments, the pharmaceutical composition comprises about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0251] In certain embodiments, the pharmaceutical composition is administered orally.

[0252] In certain embodiments, the pharmaceutical composition is formulated as a capsule or tablet. In certain embodiments, the capsule or tablet comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, or about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 50 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 200 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0253] In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically

acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0254] In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0255] In certain embodiments, the pharmaceutical composition is administered in the fed state. In certain embodiments, the pharmaceutical composition is administered in the fasted state. In certain embodiments, the pharmaceutical composition is administered at least once per day.

[0256] In certain embodiments, the pharmaceutical composition is administered at bedtime. In certain embodiments, the pharmaceutical composition is administered less than about 6 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 5 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 3 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 2 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 1 hour before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep.


[0257] In certain embodiments, the pharmaceutical composition is administered in the evening. In certain embodiments, the pharmaceutical composition is administered at about 10 pm at night.

[0258] In certain embodiments, the subject has congenital adrenal hyperplasia (CAH).


[0259] In certain embodiments, the size and/or number of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced.

[0260] In certain embodiments of the methods disclosed herein, the method further comprises administering to the subject an additional chemotherapeutic agent.

[0261] In certain embodiments, the the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In certain embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In certain embodiments, the mineralocorticoid is fludrocortisone. In certain embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist. In certain embodiments, the CRF₁ antagonist is:



pharmaceutically acceptable salt thereof. In certain embodiments, the CRF₁ antagonist is



[0262] In certain embodiments of the methods disclosed herein, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof. In certain embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the surgical resection is prior

to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0263] In certain embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0264] In one embodiment, the pharmaceutical compositions are provided in a dosage form for oral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. The pharmaceutical compositions provided herein that are formulated for oral administration may be in tablet, capsule, powder, or liquid form. In some embodiments, a tablet comprises a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil, or synthetic oil. Physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol, or polyethylene glycol may be included. In some embodiments, a capsule comprises a solid carrier such as gelatin.

[0265] In another embodiment, the pharmaceutical compositions are provided in a dosage form for parenteral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. Where pharmaceutical compositions may be formulated for intravenous, cutaneous or subcutaneous injection, the active ingredient will be in the form of a parenterally acceptable aqueous solution, which is pyrogen-free and has a suitable pH, isotonicity, and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles, such as Sodium Chloride injection, Ringer's injection, or Lactated Ringer's injection. In some embodiments, preservatives, stabilisers, buffers, antioxidants, and/or other additives are included as required.

[0266] In yet another embodiment, the pharmaceutical compositions are provided in a dosage form for topical administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers.

[0267] The pharmaceutical compositions can also be formulated as modified release dosage forms, including delayed-, extended-, prolonged-, sustained-, pulsatile-, controlled-, accelerated-, fast-, targeted-, and programmed-release, and gastric retention dosage forms. These dosage forms can be prepared according to conventional methods and techniques known to those skilled in the art (*see, Remington: The Science and Practice of Pharmacy, supra; Modified-Release Drug Delivery Technology*, 2nd Edition, Rathbone *et al.*, Eds., Marcel Dekker, Inc.: New York, NY, 2008).

[0268] The pharmaceutical compositions provided herein can be provided in a unit-dosage form or multiple-dosage form. A unit-dosage form, as used herein, refers to physically discrete a unit suitable for administration to a human and animal subject, and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of an active ingredient(s) sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carriers or excipients. Examples of a unit-dosage form include an ampoule, syringe, and individually packaged tablet and capsule. A unit-dosage form may be administered in fractions or multiples thereof. A multiple-dosage form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dosage form. Examples of a multiple-dosage form include a vial, bottle of tablets or capsules, or bottle of pints or gallons.

[0269] The pharmaceutical compositions provided herein can be administered at once, or multiple times at intervals of time. It is understood that the precise dosage and duration of treatment may vary with the age, weight, and condition of the patient being treated, and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test or diagnostic data. It is further understood that for any particular individual, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations.

[0270] In certain embodiments, the pharmaceutical compositions provided herein further comprise one or more chemotherapeutic agents as defined herein.

A. Oral Administration

[0271] The pharmaceutical compositions provided herein for oral administration can be provided in solid, semisolid, or liquid dosage forms for oral administration. As used herein, oral administration also includes buccal, lingual, and sublingual administration. Suitable oral dosage

forms include, but are not limited to, tablets, fastmelts, chewable tablets, capsules, pills, strips, troches, lozenges, pastilles, cachets, pellets, medicated chewing gum, bulk powders, effervescent or non-effervescent powders or granules, oral mists, solutions, emulsions, suspensions, wafers, sprinkles, elixirs, and syrups. In addition to the active ingredient(s), the pharmaceutical compositions can contain one or more pharmaceutically acceptable carriers or excipients, including, but not limited to, binders, fillers, diluents, disintegrants, wetting agents, lubricants, glidants, coloring agents, dye-migration inhibitors, sweetening agents, flavoring agents, emulsifying agents, suspending and dispersing agents, preservatives, solvents, non-aqueous liquids, organic acids, and sources of carbon dioxide.

[0272] Binders or granulators impart cohesiveness to a tablet to ensure the tablet remaining intact after compression. Suitable binders or granulators include, but are not limited to, starches, such as corn starch, potato starch, and pre-gelatinized starch (e.g., STARCH 1500); gelatin; sugars, such as sucrose, glucose, dextrose, molasses, and lactose; natural and synthetic gums, such as acacia, alginic acid, alginates, extract of Irish moss, panwar gum, ghatti gum, mucilage of isabgol husks, carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone (PVP), Veegum, larch arabogalactan, powdered tragacanth, and guar gum; celluloses, such as ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose, methyl cellulose, hydroxyethylcellulose (HEC), hydroxypropylcellulose (HPC), hydroxypropyl methyl cellulose (HPMC); microcrystalline celluloses, such as AVICEL-PH-101, AVICEL-PH-103, AVICEL RC-581, AVICEL-PH-105 (FMC Corp., Marcus Hook, PA); and mixtures thereof. Suitable fillers include, but are not limited to, talc, calcium carbonate, microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The amount of a binder or filler in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The binder or filler may be present from about 50 to about 99% by weight in the pharmaceutical compositions provided herein.

[0273] Suitable diluents include, but are not limited to, dicalcium phosphate, calcium sulfate, lactose, sorbitol, sucrose, inositol, cellulose, kaolin, mannitol, sodium chloride, dry starch, and powdered sugar. Certain diluents, such as mannitol, lactose, sorbitol, sucrose, and inositol, when present in sufficient quantity, can impart properties to some compressed tablets that permit disintegration in the mouth by chewing. Such compressed tablets can be used as chewable tablets. The amount of a diluent in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art.

[0274] Suitable disintegrants include, but are not limited to, agar; bentonite; celluloses, such as methylcellulose and carboxymethylcellulose; wood products; natural sponge; cation-exchange

resins; alginic acid; gums, such as guar gum and Veegum HV; citrus pulp; cross-linked celluloses, such as croscarmellose; cross-linked polymers, such as crospovidone; cross-linked starches; calcium carbonate; microcrystalline cellulose, such as sodium starch glycolate; polyacrilin potassium; starches, such as corn starch, potato starch, tapioca starch, and pre-gelatinized starch; clays; aligns; and mixtures thereof. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The pharmaceutical compositions provided herein may contain from about 0.5 to about 15% or from about 1 to about 5% by weight of a disintegrant.

[0275] Suitable lubricants include, but are not limited to, calcium stearate; magnesium stearate; mineral oil; light mineral oil; glycerin; sorbitol; mannitol; glycols, such as glycerol behenate and polyethylene glycol (PEG); stearic acid; sodium lauryl sulfate; talc; hydrogenated vegetable oil, including peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil; zinc stearate; ethyl oleate; ethyl laureate; agar; starch; lycopodium; silica or silica gels, such as AEROSIL® 200 (W.R. Grace Co., Baltimore, MD) and CAB-O-SIL® (Cabot Co. of Boston, MA); and mixtures thereof. The pharmaceutical compositions provided herein may contain about 0.1 to about 5% by weight of a lubricant.

[0276] Suitable glidants include, but are not limited to, colloidal silicon dioxide, CAB-O-SIL® (Cabot Co. of Boston, MA), and asbestos-free talc. Suitable coloring agents include, but are not limited to, any of the approved, certified, water soluble FD&C dyes, and water insoluble FD&C dyes suspended on alumina hydrate, and color lakes and mixtures thereof. A color lake is the combination by adsorption of a water-soluble dye to a hydrous oxide of a heavy metal, resulting in an insoluble form of the dye. Suitable flavoring agents include, but are not limited to, natural flavors extracted from plants, such as fruits, and synthetic blends of compounds which produce a pleasant taste sensation, such as peppermint and methyl salicylate. Suitable sweetening agents include, but are not limited to, sucrose, lactose, mannitol, syrups, glycerin, and artificial sweeteners, such as saccharin and aspartame. Suitable emulsifying agents include, but are not limited to, gelatin, acacia, tragacanth, bentonite, and surfactants, such as polyoxyethylene sorbitan monooleate (TWEEN® 20), polyoxyethylene sorbitan monooleate 80 (TWEEN® 80), and triethanolamine oleate. Suitable suspending and dispersing agents include, but are not limited to, sodium carboxymethylcellulose, pectin, tragacanth, Veegum, acacia, sodium carbomethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable preservatives include, but are not limited to, glycerin, methyl and propylparaben, benzoic add,

sodium benzoate and alcohol. Suitable wetting agents include, but are not limited to, propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate, and polyoxyethylene lauryl ether. Suitable solvents include, but are not limited to, glycerin, sorbitol, ethyl alcohol, and syrup. Suitable non-aqueous liquids utilized in emulsions include, but are not limited to, mineral oil and cottonseed oil. Suitable organic acids include, but are not limited to, citric and tartaric acid. Suitable sources of carbon dioxide include, but are not limited to, sodium bicarbonate and sodium carbonate.

[0277] It should be understood that many carriers and excipients may serve several functions, even within the same formulation.

[0278] The pharmaceutical compositions provided herein for oral administration can be provided as compressed tablets, tablet triturates, chewable lozenges, rapidly dissolving tablets, multiple compressed tablets, or enteric-coating tablets, sugar-coated, or film-coated tablets. Enteric-coated tablets are compressed tablets coated with substances that resist the action of stomach acid but dissolve or disintegrate in the intestine, thus protecting the active ingredients from the acidic environment of the stomach. Enteric-coatings include, but are not limited to, fatty acids, fats, phenyl salicylate, waxes, shellac, ammoniated shellac, and cellulose acetate phthalates. Sugar-coated tablets are compressed tablets surrounded by a sugar coating, which may be beneficial in covering up objectionable tastes or odors and in protecting the tablets from oxidation. Film-coated tablets are compressed tablets that are covered with a thin layer or film of a water-soluble material. Film coatings include, but are not limited to, hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000, and cellulose acetate phthalate. Film coating imparts the same general characteristics as sugar coating. Multiple compressed tablets are compressed tablets made by more than one compression cycle, including layered tablets, and press-coated or dry-coated tablets.

[0279] The tablet dosage forms can be prepared from the active ingredient in powdered, crystalline, or granular forms, alone or in combination with one or more carriers or excipients described herein, including binders, disintegrants, controlled-release polymers, lubricants, diluents, and/or colorants. Flavoring and sweetening agents are especially useful in the formation of chewable tablets and lozenges.

[0280] The pharmaceutical compositions provided herein for oral administration can be provided as soft or hard capsules, which can be made from gelatin, methylcellulose, starch, or calcium alginate. The hard gelatin capsule, also known as the dry-filled capsule (DFC), consists of two sections, one slipping over the other, thus completely enclosing the active ingredient. The soft elastic capsule (SEC) is a soft, globular shell, such as a gelatin shell, which is plasticized by the addition of glycerin, sorbitol, or a similar polyol. The soft gelatin shells may contain a

preservative to prevent the growth of microorganisms. Suitable preservatives are those as described herein, including methyl- and propyl-parabens, and sorbic acid. The liquid, semisolid, and solid dosage forms provided herein may be encapsulated in a capsule. Suitable liquid and semisolid dosage forms include solutions and suspensions in propylene carbonate, vegetable oils, or triglycerides. Capsules containing such solutions can be prepared as described in U.S. Pat. Nos. 4,328,245; 4,409,239; and 4,410,545. The capsules may also be coated as known by those of skill in the art in order to modify or sustain dissolution of the active ingredient.

[0281] Coloring and flavoring agents can be used in all of the above dosage forms.

[0282] The pharmaceutical compositions provided herein for oral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

B. Parenteral Administration

[0283] The pharmaceutical compositions provided herein can be administered parenterally by injection, infusion, or implantation, for local or systemic administration. Parenteral administration, as used herein, include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular, intrasynovial, intravesical, and subcutaneous administration.

[0284] The pharmaceutical compositions provided herein for parenteral administration can be formulated in any dosage forms that are suitable for parenteral administration, including solutions, suspensions, emulsions, micelles, liposomes, microspheres, nanosystems, and solid forms suitable for solutions or suspensions in liquid prior to injection. Such dosage forms can be prepared according to conventional methods known to those skilled in the art of pharmaceutical science (*see, Remington: The Science and Practice of Pharmacy, supra*).

[0285] The pharmaceutical compositions intended for parenteral administration can include one or more pharmaceutically acceptable carriers and excipients, including, but not limited to, aqueous vehicles, water-miscible vehicles, non-aqueous vehicles, antimicrobial agents or preservatives against the growth of microorganisms, stabilizers, solubility enhancers, isotonic agents, buffering agents, antioxidants, local anesthetics, suspending and dispersing agents, wetting or emulsifying agents, complexing agents, sequestering or chelating agents, cryoprotectants, lyoprotectants, thickening agents, pH adjusting agents, and inert gases.

[0286] Suitable aqueous vehicles include, but are not limited to, water, saline, physiological saline or phosphate buffered saline (PBS), sodium chloride injection, Ringers injection, isotonic dextrose injection, sterile water injection, dextrose and lactated Ringers injection. Suitable non-aqueous vehicles include, but are not limited to, fixed oils of vegetable origin, castor oil, corn oil, cottonseed oil, olive oil, peanut oil, peppermint oil, safflower oil, sesame oil, soybean oil,

hydrogenated vegetable oils, hydrogenated soybean oil, and medium-chain triglycerides of coconut oil, and palm seed oil. Suitable water-miscible vehicles include, but are not limited to, ethanol, 1,3-butanediol, liquid polyethylene glycol (e.g., polyethylene glycol 300 and polyethylene glycol 400), propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, *N,N*-dimethylacetamide, and dimethyl sulfoxide.

[0287] Suitable antimicrobial agents or preservatives include, but are not limited to, phenols, cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl *p*-hydroxybenzoates, thimerosal, benzalkonium chloride (e.g., benzethonium chloride), methyl- and propyl-parabens, and sorbic acid. Suitable isotonic agents include, but are not limited to, sodium chloride, glycerin, and dextrose. Suitable buffering agents include, but are not limited to, phosphate and citrate. Suitable antioxidants are those as described herein, including bisulfite and sodium metabisulfite. Suitable local anesthetics include, but are not limited to, procaine hydrochloride. Suitable suspending and dispersing agents are those as described herein, including sodium carboxymethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable emulsifying agents are those described herein, including polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate 80, and triethanolamine oleate. Suitable sequestering or chelating agents include, but are not limited to EDTA. Suitable pH adjusting agents include, but are not limited to, sodium hydroxide, hydrochloric acid, citric acid, and lactic acid. Suitable complexing agents include, but are not limited to, cyclodextrins, including α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, sulfobutylether- β -cyclodextrin, and sulfobutylether 7- β -cyclodextrin (CAPTISOL®, CyDex, Lenexa, KS).

[0288] When the pharmaceutical compositions provided herein are formulated for multiple dosage administration, the multiple dosage parenteral formulations must contain an antimicrobial agent at bacteriostatic or fungi static concentrations. All parenteral formulations must be sterile, as known and practiced in the art.

[0289] In one embodiment, the pharmaceutical compositions for parenteral administration are provided as ready-to-use sterile solutions. In another embodiment, the pharmaceutical compositions are provided as sterile dry soluble products, including lyophilized powders and hypodermic tablets, to be reconstituted with a vehicle prior to use. In yet another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile suspensions. In yet another embodiment, the pharmaceutical compositions are provided as sterile dry insoluble products to be reconstituted with a vehicle prior to use. In still another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile emulsions.

[0290] The pharmaceutical compositions provided herein for parenteral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

[0291] The pharmaceutical compositions provided herein for parenteral administration can be formulated as a suspension, solid, semi-solid, or thixotropic liquid, for administration as an implanted depot. In one embodiment, the pharmaceutical compositions provided herein are dispersed in a solid inner matrix, which is surrounded by an outer polymeric membrane that is insoluble in body fluids but allows the active ingredient in the pharmaceutical compositions diffuse through.

[0292] Suitable inner matrixes include, but are not limited to, polymethylmethacrylate, polybutyl-methacrylate, plasticized or unplasticized polyvinylchloride, plasticized nylon, plasticized polyethylene terephthalate, natural rubber, polyisoprene, polyisobutylene, polybutadiene, polyethylene, ethylene-vinyl acetate copolymers, silicone rubbers, polydimethylsiloxanes, silicone carbonate copolymers, hydrophilic polymers, such as hydrogels of esters of acrylic and methacrylic acid, collagen, cross-linked polyvinyl alcohol, and cross-linked partially hydrolyzed polyvinyl acetate.

[0293] Suitable outer polymeric membranes include but are not limited to, polyethylene, polypropylene, ethylene/propylene copolymers, ethylene/ethyl acrylate copolymers, ethylene/vinyl acetate copolymers, silicone rubbers, polydimethyl siloxanes, neoprene rubber, chlorinated polyethylene, polyvinylchloride, vinyl chloride copolymers with vinyl acetate, vinylidene chloride, ethylene and propylene, ionomer polyethylene terephthalate, butyl rubber epichlorohydrin rubbers, ethylene/vinyl alcohol copolymer, ethylene/vinyl acetate/vinyl alcohol terpolymer, and ethylene/vinyloxyethanol copolymer.

V. Examples

[0294] While preferred embodiments of the present disclosure have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.


Example 1. A Phase 2, Open-Label Study to Evaluate the Efficacy and Safety of Compound 1 in Subjects with Testicular Adrenal Rest Tumors

[0295] Testicular adrenal rest tumors (TART) is a common complication in congenital adrenal hyperplasia (CAH) that is caused by adrenocorticotropin hormone (ACTH)–driven

overstimulation of aberrant adrenal cells within the testes and results in impaired spermatogenesis and infertility. TARTS can be found anywhere on the testes, though a typical (very common) location is within the mediastinum. This is a particularly problematic location. Because of the central location of TARTs being in the mediastinum, compression of the nearby seminiferous tubules is common and can lead to obstructive azoospermia, irreversible damage to surrounding tissue, and permanent infertility in males.

[0296] Current treatment for TART consists of chronic high-dose glucocorticoids, a problematic therapy with significant side effects and a narrow therapeutic window. Given the serious nature of TART and the limitations and risks of chronic steroid therapy, new treatment modalities are needed for patients with TART. Compound 1, a high-affinity and selective small-molecule antagonist of the corticotropin-releasing factor type-1 (CRF₁) receptor, may be effective in blocking ACTH synthesis, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of glucocorticoids. This mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-hydroxyprogesterone (17-OHP) in subjects with CAH (FIG. 12A-12B). Compound 1 has been shown to have an acceptable safety and tolerability profile in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. These promising results provide a rationale for investigating the use of Compound 1 in the treatment of TART, in combination with replacement glucocorticoids and mineralocorticoids.

[0297] The structure of Compound 1 is:



receive active study drug. Subjects will serve as their own controls in that their outcome measures after 12 weeks of study drug dosing will be compared to their baseline measures.

[0299] The schedule of study visits (every 2 weeks through Week 4, then every 4 weeks for the remainder of the study) provides an appropriate balance between minimizing subject burden and closely monitoring subject safety and response to therapy. The 4-week post-treatment washout period will allow for adequate safety follow-up and analysis of any HPA axis rebound after the last dose of study drug.

[0300] The specification of study drug dosing at 10pm and after consumption of a snack is based on information from the Phase 1 studies in healthy volunteers. The plasma C_{max} (maximum concentration) for Compound 1 in the Phase 1 studies occurred approximately 5 hours after dosing. Therefore, Compound 1 will be dosed daily at 10pm in this study in order to achieve peak plasma study drug concentration at the time of day when most adrenal androgens are produced. Subjects will consume a standardized snack before taking study drug because the Phase 1 studies showed minimal study drug absorption when subjects were dosed in a fasted state. Specific dose reduction criteria, strict individual treatment stopping criteria, and appropriate laboratory and medical assessments have been established to ensure subject safety. **(FIGS. 1A-1B).**

Objectives	Endpoints
Primary	
<ul style="list-style-type: none"> To assess the efficacy of Compound 1 in subjects with TART 	<ul style="list-style-type: none"> Reduction in the size/volume of TART on testicular ultrasound from screening to Week 12
Secondary	
<ul style="list-style-type: none"> To evaluate the safety of Compound 1 in subjects with TART 	<ul style="list-style-type: none"> Adverse events (AEs), serious adverse events (SAEs), AEs leading to withdrawal, AEs of special interest (AESIs) Physical examination findings Vital signs Changes in electrocardiogram (ECG) Changes in clinical laboratory values
<ul style="list-style-type: none"> To assess the pharmacodynamic (PD) biomarker effects of Compound 1 	<ul style="list-style-type: none"> Changes in plasma concentrations of key PD biomarkers, including ACTH, 17-OHP, testosterone, androstenedione, luteinizing hormone (LH),

follicle-stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, and aldosterone from baseline to Week 12

- To assess changes on semen analysis
 - To assess changes in metabolism
 - To evaluate changes in quality of life (QoL) and mood
- Changes in sperm count, morphology, and motility from baseline to Week 12
 - Changes in metabolic parameters, including fasting glucose, HbA1c, insulin sensitivity, and lipid profile (including LDL cholesterol, HDL cholesterol, and triglycerides), from baseline to Week 12
 - Changes in body weight, body mass index (BMI), waist circumference, and blood pressure
 - Changes in QoL measured using the Short Form 36 (SF-36)
 - Changes in mood measured using the Hospital Anxiety and Depression Scale (HADS)

Study Population

[0301] Approximately 12 male subjects with TART will be enrolled at approximately 6 investigative sites in the United States, United Kingdom, and possibly other countries. Up to 12 additional subjects may be added to further evaluate specific safety, PD, or efficacy findings.

Eligibility Criteria

Inclusion Criteria

- Adult male subjects aged 18 to 30, inclusive
- Documented historical diagnosis of CAH based on documented genetic mutation or historical documentation of elevated 17-OHP
- Diagnosis of TART via testicular ultrasound and/or magnetic resonance imaging (MRI) at screening with ≥ 1 lesion(s) measuring ≥ 2 cm in diameter
- On a stable regimen of GC replacement for ≥ 30 days before baseline
- Agree to follow contraception guidelines and refrain from donating sperm throughout the treatment period and for 90 days after the last dose of study drug
- Subject is able to understand all study procedures and risks involved and provides written informed consent indicating willingness to comply with all aspects of the protocol

Exclusion Criteria

- Any clinically significant unstable medical condition, medically significant illness, or chronic disease within 30 days of screening, including but not limited to:

- A malignancy or <3 years of remission history from any malignancy, other than successfully treated localized skin cancer
- Presence of clinically significant renal disease, as evidenced by an estimated glomerular filtration rate (eGFR) of <60 mL/min/1.73m²
- Current or chronic history of liver disease or known hepatic or biliary abnormalities (with the exception of Gilbert's syndrome or asymptomatic gallstones)
- Confirmed positive test at screening for active hepatitis B, hepatitis C, or human immunodeficiency virus (HIV)
- Any clinically significant psychiatric disorders, including any current major depressive episode, bipolar disorder, schizophrenia, schizoaffective disorder, major depressive disorder with psychotic features, or any other psychotic disorder within the preceding 6 months
- HADS score >12 for either depression or anxiety
- At increased risk of suicide on the basis of the Investigator's judgment or the results of the Columbia-Suicide Severity Rating Scale (C-SSRS) conducted at screening and baseline (eg, C-SSRS Type 3, 4, or 5 ideation during the preceding 6 months or any suicidal behavior within the past 12 months)
- Clinically significant and abnormal clinical or laboratory assessments must be discussed with the Medical Monitor to determine eligibility for this study. Abnormal assessments that must be reviewed to determine eligibility include but are not limited to:
 - Clinically meaningful abnormal ECG results, in the opinion of the Investigator
 - Fridericia-corrected QT interval (QTcF) >450 msec
 - Alanine aminotransferase (ALT) >2x upper limit of normal (ULN)
 - Bilirubin >1.5x ULN (isolated bilirubin >1.5x ULN is acceptable if bilirubin is fractionated and direct bilirubin is <35%)
- A history that includes bilateral adrenalectomy or hypopituitarism
- Use of any other investigational drug within 30 days or 5 half-lives, whichever is longer, before initial screening
- Require use of prohibited concomitant medications, including strong inhibitors and/or inducers of CYP3A4 (with the exception of glucocorticoids, hydrocortisone, and dexamethasone) within 30 days or 5 half-lives (whichever is longer) of first dose of study drug. Medications that are sensitive substrates or substrates with narrow therapeutic ranges (metabolized by CYP3A4, 2C8, 2C9, or 2C19) should be discussed on a case-by-case basis with the Medical Monitor to determine whether the medication should be discontinued or may be continued with caution. If washout is feasible, then the medication should be

withdrawn at least 30 days or 5 half-lives (whichever is longer) before the first dose of study drug.

- Donation of blood within 60 days before first dose of study drug, or donation of platelets, white blood cells, or plasma within 15 days before first dose of study drug.

Study Duration

[0302] The expected duration of study participation for each subject is up to approximately 20 weeks. This includes a screening period of ≤ 4 weeks, a treatment period of 12 weeks, and a safety follow-up period of 4 weeks.

Study Design

[0303] This is a Phase 2, open-label efficacy and safety study of Compound 1 for the treatment of TART. All subjects will receive a daily oral dose of Compound 1 for 12 weeks. In-clinic study visits will occur for screening and at Day 1 (baseline) and Weeks 2, 4, 8, 12 (end of treatment), and 16 (follow-up).

Dose, Route, Regimen

[0304] All subjects will receive a daily oral dose of Compound 1 for 12 weeks. Subjects will receive 600 mg/day during the first 2 weeks and 1000 mg/day during the remaining 10 weeks of the treatment period. Study drug will be taken daily at 10pm (or bedtime, if earlier), 5 to 15 minutes after consumption of a standardized snack.

[0305] On mornings of study visits with 8am blood draws for laboratory assessments, subjects should take their morning glucocorticoid replacement after the blood draws. For all other days during the study, subjects may take their morning glucocorticoid replacement at their usual time (generally between 6am and 8am).

[0306] If a subject experiences a Grade 3 or higher TEAE considered at least possibly related to study drug, the Investigator may reduce the subject's daily dose of Compound 1 down in increments of 200 mg. Study drug will be discontinued in subjects who experience clinically significant liver chemistry, QTcF, or other individual treatment stopping criteria.

Investigational Drug

[0307] Compound 1 is a small-molecule CRF₁ receptor antagonist and will be supplied as 200-mg, size 1, white, hard gelatin capsules.

Example 2.

[0308] Since elevated ACTH is the primary driver for TART growth, blocking ACTH synthesis using Compound 1, a small-molecule antagonist of the CRF₁ receptor, may be effective in reducing ACTH levels, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of GCs. In CAH, this mechanism of action has been

validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH (FIG. 12A-12B). The safety profile of Compound 1 has been established in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. A Phase 2 study in subjects with CAH is currently ongoing.

Nonclinical Toxicology of Compound 1

[0309] The nonclinical safety profile of Compound 1 has been evaluated in repeat-dose range-finding and definitive toxicology studies of up to 91 days in duration in rats and dogs using the clinically relevant route (oral) and schedule of administration (once daily). In rats and beagle dogs administered oral Compound 1 once daily for 91 consecutive days at doses up to 2000 mg/kg/day, toxicologic findings were all mild and limited to the male reproductive tract, and all Compound 1-related findings were reversible following a 6-week treatment-free recovery period.

[0310] Additionally, a fertility toxicology study was conducted in rats. In this study, male rats were administered oral Compound 1 once daily at 0, 5, 20, or 1000 mg/kg for 85 to 87 days (70 days before mating, during mating, and ≤4 days between mating and necropsy). Male reproductive performance (mean day of mating, mating and fertility indices, and conception rates) was unaffected by Compound 1 treatment. Untreated females mated with the Compound 1-treated males had no ovarian and uterine effects.

[0311] In addition, a complete battery of standard genotoxicity tests consisting of in vitro bacterial reverse mutation and chromosome aberration assays and an in vivo (rat) bone marrow micronucleus assay has been conducted. Exploratory and definitive embryo-fetal development studies in rabbits and a bovine corneal opacity and permeability assay have also been conducted.

Clinical Experience with Compound 1

[0312] To date, a total of 53 subjects, all healthy adults aged 21 to 65 years, have been treated with at least 1 dose of Compound 1 in 2 Phase 1 studies. The first-in-human study of Compound 1 (Study I3C-FW-BLAA) evaluated single ascending doses of 2 mg to 800 mg. In this study, a small number of subjects experienced diarrhea and headache of mild or moderate severity. A subsequent study (Study I3C-FW-BLAB) evaluated repeat doses of 50 mg, 100 mg, or 200 mg Compound 1 once daily for 14 consecutive days. A small number of subjects in the repeat-dose study experienced dyspnea, rhinorrhea, palpitations, and headache. The Phase 1 studies showed no clinically significant alterations in safety laboratory values, vital signs, or electrocardiograms (ECGs). Compound 1 has been well tolerated in healthy human subjects, with no SAEs, AEs leading to discontinuation, or dose-limiting toxicities (DLTs). A maximum tolerated dose has not been reached in humans.

[0313] An ongoing Phase 2 open-label study (Study Compound 2) is currently evaluating the safety and efficacy of Compound 1 dosed once daily for 6 weeks in adults with classic CAH. In the initial Cohort A of this study, all subjects are dosed starting at 200 mg/day for 14 days, escalating for each subject to 600 mg/day for 14 days, then 1000 mg/day for 14 days if the subject experiences no DLTs at the previous dose level. (**FIGS. 2-3**).

Known Potential Risks

[0314] Because the reversible testicular effects (degeneration/atrophy) observed in rats and dogs treated with high-dose Compound 1 (2,000 mg/kg/day) are likely associated with on-target (mechanism-driven) effects, these effects are likely to be relevant to human risk. However, male reproductive studies in rodents showed no impairment in fertility at the highest dose tested (1,000 mg/kg/day). Additionally, there were no effects on clinical laboratory markers of testicular function (LH and FSH) when Compound 1 was administered to healthy male volunteers at doses up to 200 mg per day for 14 consecutive days. Clinical laboratory measures of testicular function (LH, FSH, inhibin B, and SHBG) will be assessed throughout this study to monitor the risk of testicular injury. Testicular ultrasounds and semen analysis will also be performed at screening/baseline and the end of the treatment period.

[0315] Laboratory abnormalities in liver function tests (LFTs) have been observed at significantly higher doses in nonclinical studies. However, these events were not observed in the Phase 1 clinical studies. Clinical laboratory measures of liver chemistry will be assessed throughout this study, and strict individual treatment stopping criteria for liver chemistry findings are in place.

[0316] Compound 1-induced effects on the thyroid gland were noted in dogs after 91 days of high-dose (2,000 mg/kg/day) treatment but not in any other nonclinical studies. Clinical pathology parameters of thyroid function (eg, T4, T3, thyroid-stimulating hormone [TSH]) were not evaluated in the Phase 1 studies but will be monitored throughout this study.

Known Potential Benefits

[0317] Early clinical studies suggests that Compound 1 is effective as a CRF₁ receptor antagonist, a mechanism of action that supports the clinical investigation of Compound 1 for treatment of TART. In CAH, this mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH (FIG. 12A-12B).

Assessment of Potential Risks and Benefits

[0318] The mild and reversible toxicologic effects of Compound 1 on the male reproductive tract in nonclinical studies occurred in the context of animals with normal, rather than elevated, baseline hormone function. As a CRF₁ receptor antagonist, Compound 1 is intended to reduce

abnormally elevated ACTH levels, which in TART drives overstimulation of aberrant adrenal cells within the testes. In nonclinical models where baseline hormone levels are within normal ranges, clinical benefits are not expected, and indeed, toxicities may be expected as a result of androgen reduction outside of the normal range. In TART, where baseline ACTH and androgen levels are pathologically elevated, androgen reduction would be considered a positive, therapeutic effect.

[0319] Given the serious nature of TART and the limitations (and risks) of chronic steroid therapy, new treatment modalities are needed for patients with TART. Given the acceptable safety and tolerability profile of Compound 1 in healthy volunteers from the Phase 1 program, the low probability of demonstrating meaningful PD marker changes in healthy subjects, and the lack of any clear benefit to exposing more healthy subjects to study drug, the Sponsor believes the benefit-to-risk profile favors the entry of Compound 1 into Phase 2 in subjects with TART.

Example 3. Congenital Adrenal Hyperplasia (CAH)

[0320] CAH results from a genetic defect in the 21-hydroxylase enzyme. CAH patients are thus unable to produce cortisol. Without cortisol to provide negative feedback, patients over-produce ACTH. High ACTH promotes the overproduction of potent androgens (male sex hormones). Without wishing to be bound by theory, Compound 1 blocks CRF₁ receptor to prevent ACTH release by the pituitary and associated drive on the hypothalamic-pituitary-adrenal axis. Replacement of cortisol to physiological levels will correct cortisol deficiency, however, will not lower ACTH, resulting in hyperplasia of adrenal gland and associated overproduction of adrenal androgens.

[0321] As shown in **FIGS. 4** and **5A-5B**, the attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism. 80% of subjects demonstrated reduction in ACTH. 70% of subjects demonstrated > 25% reduction in ACTH. 30% of subjects were in the normal range (*) after treatment with a CRF₁ receptor antagonist.

[0322] **FIGS. 6** and **7A-7B** demonstrate “control” of disease based on standard guidelines, which would then allow steroid taper. 80% of subjects demonstrated reduction in 17-OHP. 50% of subjects demonstrated > 25% reduction in ACTH. 50% of subjects were in the guideline range (*; 1200 ng/dL) after treatment with a CRF₁ receptor antagonist.

[0323] **FIGS. 8** and **9A-9B** demonstrate attenuation of morning rise in A4 upon treatment with a CRF₁ antagonist, indicative of its ability to control excess androgen production and associated symptoms. 100% of subjects demonstrated reduction in androstenedione (at various doses). 60% of subjects demonstrated > 25% reduction in androstenedione. 50% of subjects were within the normal reference range after treatment with a CRF₁ receptor antagonist.

[0324] FIGS. 10A-10B show approximately 25% shrinkage in TART in a 25 year old male with classic congenital adrenal hyperplasia following 6 weeks treatment with a CRF₁ receptor antagonist.

[0325] FIGS. 11A-11B demonstrate Compound 1 is a potent, selective, and functional CRF₁ receptor antagonist. In in vitro and ex vivo studies, Compound 1 was determined to be a potent and selective CRF₁ receptor antagonist with high affinity and selective for the hCRF₁ receptor. Compound 1 was found to have functional potency to inhibit adenylate cyclase activity (CRF₁ receptor-mediated downstream cascade), demonstrating in vitro efficacy as a CRF₁ receptor antagonist. Compound 1 was found to significantly inhibit CRF-induced locomotor behavior in response to i.c.v. administration of CRF (**FIG. 11B**).

[0326] FIGS. 12A-12B demonstrate single pill of CRF₁ receptor antagonist (i.e., as described in US 2017/0020877 A1) improves ACTH and 17-OHP in CAH patients.

[0327] FIG. 13 shows the subject level hormone changes (ACTH, 17-OHP, and androstenedione) by dose of a CRF₁ receptor antagonist.

Example 4. Assess reduction in excess ACTH and adrenal steroid production

[0328] The objectives of the study are: to assess the ability of Compound 1 to reduce excess ACTH and adrenal steroid production in participants with classic CAH on a stable glucocorticoid regimen; to evaluate the safety and tolerability of tildacerfont in participants with classic CAH; and to evaluate the pharmacokinetics (PK) of tildacerfont.

ELIGIBILITY CRITERIA

[0329] Inclusion Criteria:

- Adult males and females 18 years or older
- Diagnosis of classic CAH due to 21-hydroxylase deficiency
- Morning 17OHP $\geq$ 800 ng/dL, before taking any morning GC dose
- On a stable regimen of glucocorticoid replacement for $\geq$ 30 days before enrollment

[0330] Exclusion Criteria:

- Clinically significant unstable medical condition, illness, or chronic disease within 30 days of screening
- Clinically significant psychiatric disorder
- History of bilateral adrenalectomy or hypopituitarism

STUDY DESIGN

[0331] FIG. 14 depicts the proof of concept and escalating QD dosing at 10 PM for Cohort A and the single dose level and BID dosing at 10 AM and 10 pm for Cohort B.

BASELINE CHARACTERISTICS

[0332] Table 1 provides a description of the baseline characteristics displayed by Cohort A and Cohort B. In Cohort A, 10 participants were enrolled and received at least 1 dose of Compound 1. One participant withdrew early because of an occupation-related scheduling conflict. In Cohort B, 9 participants were enrolled and received at least one dose of Compound 1. One participant was withdrawn upon discovery of a major protocol violation at Baseline and did not have postbaseline efficacy/PK data. Thus, baseline characteristics are calculated for n=8 in Cohort B. One participant in Cohort B was taking both HC and prednisone.

Table 1 – Baseline characteristics of each cohort

Parameter	Cohort A (n=10)	Cohort B (n=8)
Demographics		
Age (years), mean (range)	39.3 (24-66)	36.8 (19-66)
Sex	5M, 5F	1M, 7F
Bmi (kg/m ²), mean $\pm$ SD	31.5 $\pm$ 12.0	31.1 $\pm$ 6.7
Daily GC dose (mg HC equivalents), mean range	28 (13-40)	35 (20-50)
Types of GCs	4 JC, 3 pred, 3 dex	2HC, 4 pred, 3 dex
17OHP at screening ^a (ng/dl), mean $\pm$ SD	5,985 $\pm$ 6,347	4,592 $\pm$ 4,624

Abbreviations: 17OHP: 17-hydroxyprogesterone; BMI: body mass index; dex: dexamethasone; F: female; GC: glucocorticoid; HC: hydrocortisone; M: male; pred: prednisone; SD: standard deviation.

^a 17OHP reference ranges: 27-199 ng/dL males, 15-290 ng/dL females.

SAFETY

[0333] Across both cohorts, Compound 1 was generally well tolerated. The most common type of adverse event (AE) was gastrointestinal disorders. There were no serious AEs and no AEs leading to withdrawal.

PHARMACOKINETICS

[0334] Over a wide range of doses (200 to 1000 mg QD), Compound 1 exhibited a predictable, dose-dependent PK profile.

EFFICACY

[0335] FIG. 15A-15D show the changes in levels of Androstenedione (A4) across both cohorts. Levels of A4 improved at ≥ 1 dose level for 10 out of 10 participants in Cohort A and 6 out of 8 participants in Cohort B. Excess A4 is associated with clinical sequelae in CAH. Attenuation of A4 demonstrates control of adrenal androgen.

[0336] FIG. 16A-16D show the changes in levels of 17-hydroxyprogesterone (17-OHP) across both cohorts. Levels of 17OHP improved at ≥ 1 dose level for 8 out of 10 participants in Cohort A and 6 out of 7 participants in Cohort B. 17OHP is a critical hormone in CAH diagnosis and management. Attenuation of 17OHP demonstrates control of adrenal steroid production.

[0337] FIG. 17A-17D show the changes in levels of adrenocorticotrophic hormone (ACTH) across both cohorts. Levels of ACTH improved at ≥ 1 dose level for 8 out of 10 participants in

Cohort A and 5 out of 7 participants in Cohort B. Attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism.

Table 2 – Percentage change from baseline in ACTH, 17OHP and A4.

	Cohort A			Cohort B
	200 mg QD (Week 2)	600 mg QD (Week 4)	1000 mg QD (Week 6)	200 mg BID (Week 2)
ACTH	-25.7 (±51.6)	-60.1 (±37.5)	-32.7 (±47.0)	-36.5 (±58.0)
17OHP	+22.2 (±172.3)	-23.7 (±64.2)	-13.8 (±74.7)	-47.8 (±63.3)
A4	-12.1 (±47.4)	-33.2 (±36.5)	-28.6 (±44.6)	-16.0 (±81.4)

CONCLUSIONS

[0338] Compound 1, a potent and selective CRF₁ receptor antagonist, offers a novel nonsteroidal oral treatment approach for CAH. Compound 1 was well tolerated and efficacious in reducing excess ACTH, 17OHP, and androstenedione in patients with classic CAH. Compound 1 exhibited a predictable, dose-dependent PK profile suitable for the target population.

[0339] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

ABSTRACT OF THE DISCLOSURE

Provided herein are compounds and pharmaceutical compositions for the prevention and treatment of testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART).

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

CROSS-REFERENCE

[0001] This application is a continuation of U.S. Application No. 18/941,832, filed November 8, 2024, which is a continuation of U.S. Application No. 17/667,285, filed on February 8, 2022, which is a continuation of U.S. Application No. 17/081,694 filed on October 27, 2020, which issued as U.S. Pat. No. 11,304,950 on April 19, 2022, and which is a continuation of International Application No. PCT/US2019/029486 filed on April 26, 2019, which claims the benefit of U.S. Provisional Patent Application No. 62/663,951, filed April 27, 2018, and U.S. Provisional Patent Application No. 62/822,815, filed March 23, 2019, each of which is entirely incorporated herein by reference.

BACKGROUND

[0002] Corticotropes are basophilic cells in the anterior pituitary that produce melanocyte-stimulating hormone, adrenocorticotrophic hormone (ACTH) and lipotropin. The cells produce pro-opiomelanocortin (POMC) which undergoes cleavage to ACTH and β -lipotropin (β -LPH). These cells respond to corticotropin-releasing factor (CRF) and make up about 20% of the cells in the anterior pituitary. CRF is a releasing hormone that belongs to the corticotropin-releasing factor family and is the primary physiological regulator of proopioidmelanocortin (POMC).

[0003] CRF is produced by parvocellular neuroendocrine cells within the paraventricular nucleus of the hypothalamus and is released at the median eminence from neurosecretory terminals of these neurons into the primary capillary plexus of the hypothalamo-hypophyseal portal system. The portal system carries the CRF to the anterior lobe of the pituitary, where it stimulates corticotropes to secrete ACTH and other biologically-active substances. ACTH, in turn, stimulates the synthesis of cortisol, glucocorticoids, mineralocorticoids and DHEA.

[0004] Testicular adrenal rest tumors (TART) and ovarian adrenal rest tumors (OART) are ACTH-responsive lesions of the testes and ovaries derived from adrenal tissue interchelated within these organs during embryogenesis. In response to high ACTH, hyperplasia of this tissue occurs, resulting in single or multiple lesions that may cause pain and infertility. The identification of compounds that modulate CRF function and downstream processes is an ongoing challenge. Currently, there are no non-steroidal, non-surgical solutions to control, shrink, or reduce TART. Disclosed herein, *inter alia*, are solutions to these and other problems in the art.

SUMMARY


[0005] Congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency is a serious genetic disorder characterized by impaired adrenal synthesis of cortisol and consequent overproduction of adrenocorticotrophic hormone (ACTH), 17-hydroxyprogesterone (17OHP), and adrenal androgens such as androstenedione (A4). CAH is an orphan condition with no FDA-approved therapy. To suppress androgen production, patients are often prescribed long-term supraphysiologic doses of glucocorticoid (GC), which has adverse effects and fail to suppress 17OHP and A4 in the majority of patients.

[0006] Compound 1 may be a potent, selective, nonsteroidal, oral corticotropin-releasing factor type-1 (CRF1) receptor antagonist that reduces excess ACTH, 17OHP, and A4 to improve clinical and biochemical sequelae of CAH. This may obviate the need for high dose steroids.

[0007] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0008] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0009] One aspect provided herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):




wherein:

R¹ and R² are independently ethyl or n-propyl;



R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R⁴ is hydrogen, Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,




R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$.


[0010] In some embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl. In some embodiments, R^3 is Cl, Br, or methyl. In some embodiments, R^4 is Br, R^aR^bN- , pyridin-4-yl, morpholin-4-yl,

or . In some embodiments, R^4 is morpholin-4-yl or .


[0011] In some embodiments, R^4 is hydrogen, Br, R^aR^bN- and R^a and R^b are independently C_1 - C_3 alkyl. In some embodiments, the compound is



[0012] In some embodiments, the compound is



[0013] In some embodiments, the compound is



[0014] In some embodiments, the subject has congenital adrenal hyperplasia (CAH). In some embodiments, the size and/or number of the tumors is decreased or reduced.

[0015] In some embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a


pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In some embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0016] In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 5 mg/day to about 25 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose from about 10 mg/day to about 400 mg/day is administered to the subject. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day is administered to the subject.

[0017] In some embodiments, the compound is administered orally. In some embodiments, the compound is formulated as a capsule or tablet. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state.

[0018] In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

[0019] One aspect described herein is a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I) and a pharmaceutically acceptable excipient:




(I), or a pharmaceutically acceptable salt thereof,

wherein:

R^1 and R^2 are independently ethyl or n-propyl;


R^3 is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy; and

R^4 is hydrogen, Br, $R^a R^b N-$, methoxymethyl, n-butyl, acetamido, pyridin-4-yl,



R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$.

[0020] In some embodiments, the compound of Formula (I) has the structure:



, or a pharmaceutically acceptable salt thereof.

[0021] In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0022] In some embodiments, the pharmaceutical composition comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0023] In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day. In some embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.


[0024] In some embodiments, the pharmaceutical composition is administered orally. In some embodiments, the pharmaceutical composition is formulated as capsule or a tablet. In some embodiments, the pharmaceutical composition is administered in the fed state. In some embodiments, the pharmaceutical composition is administered in the fasted state. In some embodiments, the pharmaceutical composition is administered at least once per day.

[0025] In some embodiments, the pharmaceutical composition is administered at bedtime. In some embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 2 hours before sleep. In some embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep. In some embodiments, the pharmaceutical composition is administered in the evening. In some embodiments, the pharmaceutical composition is administered at about 10 pm at night.


[0026] In some embodiments, the method further comprises administering to the subject an additional chemotherapeutic agent. In some embodiments, the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In some embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In some embodiments, the mineralocorticoid is fludrocortisone.

[0027] In some embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist.

In some embodiments, the CRF₁ antagonist is



[0028] In some embodiments, the CRF₁ antagonist is



[0029] In some embodiments, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof.

[0030] In some embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural

Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof.

[0031] In some embodiments, the surgical resection is prior to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the surgical resection is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0032] In some embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In some embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0033] In some embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In some embodiments, the radiation therapy is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0034] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure.

Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0035] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings (also “Figure” and “FIG.” herein), of which:

[0037] FIG. 1A-1B. A schematic representation of intra-patient dose escalation study to evaluate safety, pharmacokinetics, and pharmacodynamics **(A)**. A schematic representation of inhibition of CRF₁ with a CRF₁ antagonist (Compound 1) **(B)**.

[0038] FIG. 2. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0039] FIG. 3. Graphic representation of pharmacokinetics (dose-proportional, adequate exposure at safe doses) in patients with CAH following 14-days of once daily dosing at each dose level with a CRF₁ antagonist (Compound 1).

[0040] FIG. 4. Graphic representation of patient-level response (measuring ACTH levels) to a CRF₁ antagonist (Compound 1).

[0041] FIG. 5A-5B. Graphic representation of patient-level response of sample responders (measuring ACTH levels) to a CRF₁ antagonist [~~Subject 003001~~ Patient 10 **(A)**; ~~Subject 007002~~ Patient 8 **(B)**].

[0042] FIG. 6. Graphic representation of patient-level response (measuring 17-OHP levels) to a CRF₁ antagonist (Compound 1).

[0043] FIG. 7A-7B. Graphic representation of patient-level response of sample responders (measuring 17-OHP levels) to a CRF₁ antagonist [~~Subject 003001~~ Patient 10 **(A)**; ~~Subject 007002~~ Patient 8 **(B)**].

[0044] FIG. 8. Graphic representation of patient-level response (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1).

[0045] FIG. 9A-9B. Graphic representation of patient-level response of sample responders (measuring androstenedione levels) to a CRF₁ antagonist (Compound 1) [~~Subject 003001~~ Patient 10 **(A)**; ~~Subject 007002~~ Patient 8 **(B)**].

[0046] FIG. 10A-10B. Images of a patient with TART pre-treatment with a CRF₁ antagonist (Compound 1) **(A)** and 6-weeks post-treatment with a CRF₁ antagonist (Compound 1) **(B)**.

[0047] FIG. 11A-11B. Graphic representation of an *in vitro* binding assay of a CRF₁ antagonist (Compound 1); ¹²⁵I-sauvagine binding to rat cerebellar membranes one hour after oral

administration of increasing Compound 1 doses (A) and *in vivo* inhibition of CRF-induced activity after treatment with a CRF₁ antagonist; Compound 1 (oral, 10 mg/kg) (B).

[0048] FIG. 12A-12B. ~~Graphic representation of ACTH levels (A) and 17-OHP levels (B) after treatment with a CRF₁ antagonist (Compound 1)~~ Graphic representation of ACTH levels (A) and 17-OHP levels (B) after treatment with a CRF₁ antagonist.

[0049] FIG. 13. Graphic representation of hormone responses after treatment with a CRF₁ antagonist (Compound 1).

[0050] FIG. 14 depicts the proof of concept and escalating QD dosing at 10 pm for Cohort A and the single dose level and BID dosing at 10 am and 10 pm for Cohort B.

[0051] FIG. 15A-15D are graphic representations of changes in androstenedione seen in participants in Cohort A and Cohort B.

[0052] FIG. 16A-16D are graphic representations of changes in 17-hydroxyprogesterone seen in participants in Cohort A and Cohort B.

[0053] FIG. 17A-17D are graphic representations of changes in adrenocorticotrophic hormone seen in participants in Cohort A and Cohort B.

DETAILED DESCRIPTION

[0054] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

I. Definitions

[0055] The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are constructed according to the standard rules of chemical valency known in the chemical arts.

[0056] Where substituent groups are specified by their conventional chemical formulae, written from left to right, they equally encompass the chemically identical substituents that would result from writing the structure from right to left, e.g., -CH₂O- is equivalent to -OCH₂-.

[0057] The term “alkyl,” by itself or as part of another substituent, means, unless otherwise stated, a straight (i.e., unbranched) or branched carbon chain (or carbon), or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include mono-, di- and multivalent radicals, having the number of carbon atoms designated (i.e., C₁-C₁₀ means one to ten carbons). Alkyl is an uncyclized chain. Examples of saturated hydrocarbon radicals include, but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl,

sec-butyl, (cyclohexyl)methyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. An alkoxy is an alkyl attached to the remainder of the molecule via an oxygen linker (-O-).

[0058] The term “alkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyl, as exemplified, but not limited by, -CH₂CH₂CH₂CH₂-. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred herein. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms. The term “alkenylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkene.

[0059] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or combinations thereof, including at least one carbon atom and at least one heteroatom (e.g., O, N, P, Si, and S), and wherein the nitrogen and sulfur atoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) (e.g., N, S, Si, or P) may be placed at any interior position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Heteroalkyl is an uncyclized chain. Examples include, but are not limited to: -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃, -O-CH₃, -O-CH₂-CH₃, and -CN. Up to two or three heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. A heteroalkyl moiety may include one heteroatom (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include two optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include three optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include four optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include five optionally different heteroatoms (e.g., O, N, S, Si, or P). A heteroalkyl moiety may include up to 8 optionally different heteroatoms (e.g., O, N, S, Si, or P).

[0060] Similarly, the term “heteroalkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from heteroalkyl, as exemplified, but not limited by, -CH₂-CH₂-S-CH₂-CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (e.g., alkyleneoxy, alkyleneedioxy, alkyleneamino, alkylene-diamino, and the like). Still further, for alkylene and

heteroalkylene linking groups, no orientation of the linking group is implied by the direction in which the formula of the linking group is written. For example, the formula $-C(O)_2R'$ represents both $-C(O)_2R'$ and $-R'C(O)_2-$. As described above, heteroalkyl groups, as used herein, include those groups that are attached to the remainder of the molecule through a heteroatom, such as $-C(O)R'$, $-C(O)NR'$, $-NR'R''$, $-OR'$, $-SR'$, and/or $-SO_2R'$. Where “heteroalkyl” is recited, followed by recitations of specific heteroalkyl groups, such as $-NR'R''$ or the like, it will be understood that the terms heteroalkyl and $-NR'R''$ are not redundant or mutually exclusive. Rather, the specific heteroalkyl groups are recited to add clarity. Thus, the term “heteroalkyl” should not be interpreted herein as excluding specific heteroalkyl groups, such as $-NR'R''$ or the like.

[0061] The terms “cycloalkyl” and “heterocycloalkyl,” by themselves or in combination with other terms, mean, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl,” respectively. Cycloalkyl and heterocycloalkyl are not aromatic. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like. A “cycloalkylene” and a “heterocycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a cycloalkyl and heterocycloalkyl, respectively. “Cycloalkyl” is also meant to refer to bicyclic and polycyclic hydrocarbon rings such as, for example, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, etc.

[0062] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl” are meant to include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” includes, but is not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0063] The term “acyl” means, unless otherwise stated, $-C(O)R$ where R is a substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

[0064] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent, which can be a single ring or multiple rings (preferably from 1 to 3 rings) that are fused together (i.e., a fused ring aryl) or linked covalently. A fused ring aryl refers to multiple rings fused together wherein at least one of the fused rings is an aryl ring. The term

“heteroaryl” refers to aryl groups (or rings) that contain at least one heteroatom such as N, O, or S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. Thus, the term “heteroaryl” includes fused ring heteroaryl groups (i.e., multiple rings fused together wherein at least one of the fused rings is a heteroaromatic ring). A 5,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 5 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. Likewise, a 6,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. And a 6,5-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 5 members, and wherein at least one ring is a heteroaryl ring. A heteroaryl group can be attached to the remainder of the molecule through a carbon or heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, naphthyl, pyrrolyl, pyrazolyl, pyridazinyl, triazinyl, pyrimidinyl, imidazolyl, pyrazinyl, purinyl, oxazolyl, isoxazolyl, thiazolyl, furyl, thienyl, pyridyl, pyrimidyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, benzofuran, isobenzofuranyl, indolyl, isoindolyl, benzothiophenyl, isoquinolyl, quinoxalyl, quinolyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below. An “arylene” and a “heteroarylene,” alone or as part of another substituent, mean a divalent radical derived from an aryl and heteroaryl, respectively. A heteroaryl group substituent may be -O- bonded to a ring heteroatom nitrogen.


[0065] Spirocyclic rings are two or more rings wherein adjacent rings are attached through a single atom. The individual rings within spirocyclic rings may be identical or different. Individual rings in spirocyclic rings may be substituted or unsubstituted and may have different substituents from other individual rings within a set of spirocyclic rings. Possible substituents for individual rings within spirocyclic rings are the possible substituents for the same ring when not part of spirocyclic rings (e.g. substituents for cycloalkyl or heterocycloalkyl rings). Spirocyclic rings may be substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkylene, substituted or unsubstituted heterocycloalkyl or substituted or unsubstituted heterocycloalkylene and individual rings within a spirocyclic ring group may be any of the immediately previous list, including having all rings of one type (e.g. all rings being substituted

heterocycloalkylene wherein each ring may be the same or different substituted heterocycloalkylene). When referring to a spirocyclic ring system, heterocyclic spirocyclic rings means a spirocyclic rings wherein at least one ring is a heterocyclic ring and wherein each ring may be a different ring. When referring to a spirocyclic ring system, substituted spirocyclic rings means that at least one ring is substituted and each substituent may optionally be different.

[0066] The symbol “” denotes the point of attachment of a chemical moiety to the remainder of a molecule or chemical formula.

[0067] The term “oxo,” as used herein, means an oxygen that is double bonded to a carbon atom.

[0068] The term “alkylarylene” as an arylene moiety covalently bonded to an alkylene moiety (also referred to herein as an alkylene linker). In embodiments, the alkylarylene group has the formula:



[0069] An alkylarylene moiety may be substituted (e.g. with a substituent group) on the alkylene moiety or the arylene linker (e.g. at carbons 2, 3, 4, or 6) with halogen, oxo, -N₃, -CF₃, -CCl₃, -CBr₃, -Cl₃, -CN, -CHO, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₂CH₃, -SO₃H, -OSO₃H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(O)NHNH₂, substituted or unsubstituted C₁-C₅ alkyl or substituted or unsubstituted 2 to 5 membered heteroalkyl). In embodiments, the alkylarylene is unsubstituted.

[0070] Each of the above terms (e.g., “alkyl,” “heteroalkyl,” “cycloalkyl,” “heterocycloalkyl,” “aryl,” and “heteroaryl”) includes both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

[0071] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be one or more of a variety of groups selected from, but not limited to, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR'C(O)R', -NR'-C(O)NR''R''', -NR'C(O)₂R', -NR-C(NR'R''R''')=NR''', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -NR'NR''R''', -ONR'R'', -NR'C(O)NR''NR''R''', -CN, -NO₂, -NR'SO₂R'', -NR'C(O)R'', -NR'C(O)-OR'', -NR'OR'', in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such radical. R, R', R'', R''', and R'''' each preferably independently refer to hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl (e.g., aryl substituted with 1-3 halogens), substituted or unsubstituted heteroaryl, substituted or

unsubstituted alkyl, alkoxy, or thioalkoxy groups, or arylalkyl groups. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', and R'''' group when more than one of these groups is present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 4-, 5-, 6-, or 7-membered ring. For example, -NR'R'' includes, but is not limited to, 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term “alkyl” is meant to include groups including carbon atoms bound to groups other than hydrogen groups, such as haloalkyl (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

[0072] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups are varied and are selected from, for example: -OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR'C(O)R', -NR'C(O)NR'R''', -NR'C(O)₂R', -NR-C(NR'R''R''')=NR''', -NR-C(NR'R'')=NR'', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -NR'NR'R''', -ONR'R'', -NR'C(O)NR''NR''''R''', -CN, -NO₂, -R', -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, -NR'SO₂R'', -NR'C(O)R'', -NR'C(O)-OR'', -NR'OR'', in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'', R''', and R'''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl. When a compound described herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''', and R'''' groups when more than one of these groups is present.

[0073] Substituents for rings (e.g. cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylene, heterocycloalkylene, arylene, or heteroarylene) may be depicted as substituents on the ring rather than on a specific atom of a ring (commonly referred to as a floating substituent). In such a case, the substituent may be attached to any of the ring atoms (obeying the rules of chemical valency) and in the case of fused rings or spirocyclic rings, a substituent depicted as associated with one member of the fused rings or spirocyclic rings (a floating substituent on a single ring), may be a substituent on any of the fused rings or spirocyclic rings (a floating substituent on multiple rings). When a substituent is attached to a ring, but not a specific atom (a floating substituent), and a subscript for the substituent is an integer greater than one, the multiple substituents may be on the same atom, same ring, different atoms, different fused rings, different spirocyclic rings, and each substituent may optionally be different. Where a point of attachment of a ring to the remainder of a molecule is not limited to a single atom (a floating substituent), the attachment point may be any atom of the ring and in the case of a fused ring or spirocyclic

ring, any atom of any of the fused rings or spirocyclic rings while obeying the rules of chemical valency. Where a ring, fused rings, or spirocyclic rings contain one or more ring heteroatoms and the ring, fused rings, or spirocyclic rings are shown with one more floating substituents (including, but not limited to, points of attachment to the remainder of the molecule), the floating substituents may be bonded to the heteroatoms. Where the ring heteroatoms are shown bound to one or more hydrogens (e.g. a ring nitrogen with two bonds to ring atoms and a third bond to a hydrogen) in the structure or formula with the floating substituent, when the heteroatom is bonded to the floating substituent, the substituent will be understood to replace the hydrogen, while obeying the rules of chemical valency.

[0074] Two or more substituents may optionally be joined to form aryl, heteroaryl, cycloalkyl, or heterocycloalkyl groups. Such so-called ring-forming substituents are typically, though not necessarily, found attached to a cyclic base structure. In one embodiment, the ring-forming substituents are attached to adjacent members of the base structure. For example, two ring-forming substituents attached to adjacent members of a cyclic base structure create a fused ring structure. In another embodiment, the ring-forming substituents are attached to a single member of the base structure. For example, two ring-forming substituents attached to a single member of a cyclic base structure create a spirocyclic structure. In yet another embodiment, the ring-forming substituents are attached to non-adjacent members of the base structure.

[0075] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally form a ring of the formula $-T-C(O)-(CRR')_q-U-$, wherein T and U are independently $-NR-$, $-O-$, $-CRR'-$, or a single bond, and q is an integer of from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-A-(CH_2)_r-B-$, wherein A and B are independently $-CRR'-$, $-O-$, $-NR-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-S(O)_2NR'-$, or a single bond, and r is an integer of from 1 to 4. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-(CRR')_s-X'-(C''R''R''')_d-$, where s and d are independently integers of from 0 to 3, and X' is $-O-$, $-NR'-$, $-S-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)_2NR'-$. The substituents R, R', R'', and R''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

[0076] As used herein, the terms “heteroatom” or “ring heteroatom” are meant to include oxygen (O), nitrogen (N), sulfur (S), phosphorus (P), and silicon (Si).

[0077] A “substituent group,” as used herein, means a group selected from the following moieties:

(A) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NH₂SO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(B) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(i) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NH₂SO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(ii) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from:

(a) oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NH₂SO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(b) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, substituted with at least one substituent selected from: oxo, halogen, -CF₃, -CN, -OH, -NH₂, -COOH, -CONH₂, -NO₂, -SH, -SO₃H, -SO₄H, -SO₂NH₂, -NHNH₂, -ONH₂, -NHC(=O)NHNH₂, -NHC(=O)NH₂, -NH₂SO₂H, -NHC(=O)H, -NHC(O)-OH, -NHOH, -OCF₃, -OCHF₂, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl.

[0078] A “size-limited substituent” or “size-limited substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or

unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl.

[0079] A “lower substituent” or “ lower substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl.

[0080] In some embodiments, each substituted group described in the compounds herein is substituted with at least one substituent group. More specifically, in some embodiments, each substituted alkyl, substituted heteroalkyl, substituted cycloalkyl, substituted heterocycloalkyl, substituted aryl, substituted heteroaryl, substituted alkylene, substituted heteroalkylene, substituted cycloalkylene, substituted heterocycloalkylene, substituted arylene, and/or substituted heteroarylene described in the compounds herein are substituted with at least one substituent group. In other embodiments, at least one or all of these groups are substituted with at least one size-limited substituent group. In other embodiments, at least one or all of these groups are substituted with at least one lower substituent group.

[0081] In other embodiments of the compounds herein, each substituted or unsubstituted alkyl may be a substituted or unsubstituted C₁-C₂₀ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 20 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₈ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 8 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 10 membered heteroaryl. In some embodiments of the compounds herein, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₂₀ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 20 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₈ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a substituted or unsubstituted 3 to 8 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 10 membered heteroarylene.

[0082] In some embodiments, each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2 to 8 membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₃-C₇ cycloalkyl, each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 3 to 7 membered heterocycloalkyl, each substituted or unsubstituted aryl is a substituted or unsubstituted C₆-C₁₀ aryl, and/or each substituted or unsubstituted heteroaryl is a substituted or unsubstituted 5 to 9 membered heteroaryl. In some embodiments, each substituted or unsubstituted alkylene is a substituted or unsubstituted C₁-C₈ alkylene, each substituted or unsubstituted heteroalkylene is a substituted or unsubstituted 2 to 8 membered heteroalkylene, each substituted or unsubstituted cycloalkylene is a substituted or unsubstituted C₃-C₇ cycloalkylene, each substituted or unsubstituted heterocycloalkylene is a substituted or unsubstituted 3 to 7 membered heterocycloalkylene, each substituted or unsubstituted arylene is a substituted or unsubstituted C₆-C₁₀ arylene, and/or each substituted or unsubstituted heteroarylene is a substituted or unsubstituted 5 to 9 membered heteroarylene. In some embodiments, the compound is a chemical species set forth in the Examples section, figures, or tables below.

[0083] Certain compounds of the present invention possess asymmetric carbon atoms (optical or chiral centers) or double bonds; the enantiomers, racemates, diastereomers, tautomers, geometric isomers, stereoisometric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)- or, as (D)- or (L)- for amino acids, and individual isomers are encompassed within the scope of the present invention. The compounds of the present invention do not include those that are known in art to be too unstable to synthesize and/or isolate. The present invention is meant to include compounds in racemic and optically pure forms. Optically active (R)- and (S)-, or (D)- and (L)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers.

[0084] As used herein, the term “isomers” refers to compounds having the same number and kind of atoms, and hence the same molecular weight, but differing in respect to the structural arrangement or configuration of the atoms.

[0085] The term “tautomer,” as used herein, refers to one of two or more structural isomers which exist in equilibrium and which are readily converted from one isomeric form to another.

[0086] It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention.

[0087] Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention.

[0088] Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention.

[0089] The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I), or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are encompassed within the scope of the present invention.

[0090] It should be noted that throughout the application that alternatives are written in Markush groups, for example, each amino acid position that contains more than one possible amino acid. It is specifically contemplated that each member of the Markush group should be considered separately, thereby comprising another embodiment, and the Markush group is not to be read as a single unit.

[0091] “Analog” or “analogue” is used in accordance with its plain ordinary meaning within Chemistry and Biology and refers to a chemical compound that is structurally similar to another compound (i.e., a so-called “reference” compound) but differs in composition, e.g., in the replacement of one atom by an atom of a different element, or in the presence of a particular functional group, or the replacement of one functional group by another functional group, or the absolute stereochemistry of one or more chiral centers of the reference compound. Accordingly, an analog is a compound that is similar or comparable in function and appearance but not in structure or origin to a reference compound.

[0092] The terms “a” or “an,” as used in herein means one or more. In addition, the phrase “substituted with a[n],” as used herein, means the specified group may be substituted with one or more of any or all of the named substituents. For example, where a group, such as an alkyl or heteroaryl group, is “substituted with an unsubstituted $\text{C}_1\text{-C}_{20}$ alkyl, or unsubstituted 2 to 20 membered heteroalkyl,” the group may contain one or more unsubstituted $\text{C}_1\text{-C}_{20}$ alkyls, and/or one or more unsubstituted 2 to 20 membered heteroalkyls.

[0093] Moreover, where a moiety is substituted with an R substituent, the group may be referred to as “R-substituted.” Where a moiety is R-substituted, the moiety is substituted with at least one

R substituent and each R substituent is optionally different. Where a particular R group is present in the description of a chemical genus (such as Formula (I)), a Roman alphabetic symbol may be used to distinguish each appearance of that particular R group. For example, where multiple R^{13} substituents are present, each R^{13} substituent may be distinguished as R^{13A} , R^{13B} , R^{13C} , R^{13D} , etc., wherein each of R^{13A} , R^{13B} , R^{13C} , R^{13D} , etc. is defined within the scope of the definition of R^{13} and optionally differently.

[0094] A “detectable moiety” as used herein refers to a moiety that can be covalently or noncovalently attached to a compound or biomolecule that can be detected for instance, using techniques known in the art. In embodiments, the detectable moiety is covalently attached. The detectable moiety may provide for imaging of the attached compound or biomolecule. The detectable moiety may indicate the contacting between two compounds. Exemplary detectable moieties are fluorophores, antibodies, reactive dies, radio-labeled moieties, magnetic contrast agents, and quantum dots. Exemplary fluorophores include fluorescein, rhodamine, GFP, coumarin, FITC, Alexa fluor, Cy3, Cy5, BODIPY, and cyanine dyes. Exemplary radionuclides include Fluorine-18, Gallium-68, and Copper-64. Exemplary magnetic contrast agents include gadolinium, iron oxide and iron platinum, and manganese.

[0095] Descriptions of compounds of the present invention are limited by principles of chemical bonding known to those skilled in the art. Accordingly, where a group may be substituted by one or more of a number of substituents, such substitutions are selected so as to comply with principles of chemical bonding and to give compounds which are not inherently unstable and/or would be known to one of ordinary skill in the art as likely to be unstable under ambient conditions, such as aqueous, neutral, and several known physiological conditions. For example, a heterocycloalkyl or heteroaryl is attached to the remainder of the molecule via a ring heteroatom in compliance with principles of chemical bonding known to those skilled in the art thereby avoiding inherently unstable compounds.

[0096] The term “pharmaceutically acceptable salts” is meant to include salts of the active compounds that are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of

pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, oxalic, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see*, for example, Berge *et al.*, “Pharmaceutical Salts”, *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0097] Thus, the compounds of the present invention may exist as salts, such as with pharmaceutically acceptable acids. The present invention includes such salts. Non-limiting examples of such salts include hydrochlorides, hydrobromides, phosphates, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, propionates, tartrates (e.g., (+)-tartrates, (-)-tartrates, or mixtures thereof including racemic mixtures), succinates, benzoates, and salts with amino acids such as glutamic acid, and quaternary ammonium salts (e.g. methyl iodide, ethyl iodide, and the like). These salts may be prepared by methods known to those skilled in the art.

[0098] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound may differ from the various salt forms in certain physical properties, such as solubility in polar solvents. In embodiments, compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts. The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in a conventional manner. The parent form of the compounds differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but, unless specifically indicated, the salts disclosed herein are equivalent to the parent form of the compound for the purposes of the present invention.

[0099] In addition to salt forms, the present invention provides compounds, which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Prodrugs of the compounds described herein may be converted *in vivo* after administration. Additionally, prodrugs can be converted to the compounds of the present

invention by chemical or biochemical methods in an *ex vivo* environment, such as, for example, when contacted with a suitable enzyme or chemical reagent.

[0100] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0101] “Pharmaceutically acceptable excipient” and “pharmaceutically acceptable carrier” refer to a substance that aids the administration of a compound to and absorption by a subject and can be included in the compositions of the present invention without causing a significant adverse toxicological effect on the patient. Non-limiting examples of pharmaceutically acceptable excipients include water, NaCl, normal saline solutions, lactated Ringer’s, normal sucrose, normal glucose, binders, fillers, disintegrants, lubricants, coatings, sweeteners, flavors, salt solutions (such as Ringer’s solution), alcohols, oils, gelatins, carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, and colors, and the like. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with the compounds of the invention. One of skill in the art will recognize that other pharmaceutical excipients are useful in the present invention.

[0102] The term “preparation” is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

[0103] A “CRF1 inhibitor” refers to a compound (e.g., compounds described herein) that reduces the activity of CRF1 when compared to a control, such as absence of the compound or a compound with known inactivity.

[0104] The terms “polypeptide,” “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues, wherein the polymer may optionally be conjugated to a moiety that does not consist of amino acids. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. In embodiments, the terms “polypeptide,” “peptide,” and

“protein”, used interchangeably herein, refer to a polymeric form of amino acids of any length, which can include genetically coded and non-genetically coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified polypeptide backbones. The terms include fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence; fusion proteins with heterologous and homologous leader sequences, with or without N-terminus methionine residues; immunologically tagged proteins; and the like.

[0105] A polypeptide, or a cell is “recombinant” when it is artificial or engineered, or derived from or contains an artificial or engineered protein or nucleic acid (*e.g.* non-natural or not wild type). For example, a polynucleotide that is inserted into a vector or any other heterologous location, *e.g.*, in a genome of a recombinant organism, such that it is not associated with nucleotide sequences that normally flank the polynucleotide as it is found in nature is a recombinant polynucleotide. A protein expressed *in vitro* or *in vivo* from a recombinant polynucleotide is an example of a recombinant polypeptide. Likewise, a polynucleotide sequence that does not appear in nature, for example a variant of a naturally occurring gene, is recombinant.

[0106] “Contacting” is used in accordance with its plain ordinary meaning and refers to the process of allowing at least two distinct species (*e.g.* chemical compounds including biomolecules or cells) to become sufficiently proximal to react, interact or physically touch. It should be appreciated; however, the resulting reaction product can be produced directly from a reaction between the added reagents or from an intermediate from one or more of the added reagents that can be produced in the reaction mixture.

[0107] The term “contacting” may include allowing two species to react, interact, or physically touch, wherein the two species may be a compound as described herein and a protein or enzyme. In some embodiments contacting includes allowing a compound described herein to interact with a protein or enzyme that is involved in a signaling pathway (*e.g.*, MAP kinase pathway).

[0108] As defined herein, the term “activation”, “activate”, “activating” and the like in reference to a protein refers to conversion of a protein into a biologically active derivative from an initial inactive or deactivated state. The terms reference activation, or activating, sensitizing, or up-regulating signal transduction or enzymatic activity or the amount of a protein decreased in a disease.

[0109] The terms “agonist,” “activator,” “upregulator,” etc. refer to a substance capable of detectably increasing the expression or activity of a given gene or protein. The agonist can increase expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the agonist. In certain instances, expression or activity

is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or higher than the expression or activity in the absence of the agonist. In embodiments, an agonist is a molecule that interacts with a target to cause or promote an increase in the activation of the target. In embodiments, activators are molecules that increase, activate, facilitate, enhance activation, sensitize, or up-regulate, e.g., a gene, protein, ligand, receptor, or cell.

[0110] As defined herein, the term “inhibition”, “inhibit”, “inhibiting” and the like in reference to a protein-inhibitor interaction means negatively affecting (e.g. decreasing) the activity or function of the protein relative to the activity or function of the protein in the absence of the inhibitor. In embodiments inhibition means negatively affecting (e.g. decreasing) the concentration or levels of the protein relative to the concentration or level of the protein in the absence of the inhibitor. In embodiments inhibition refers to reduction of a disease or symptoms of disease. In embodiments, inhibition refers to a reduction in the activity of a particular protein target. Thus, inhibition includes, at least in part, partially or totally blocking stimulation, decreasing, preventing, or delaying activation, or inactivating, desensitizing, or down-regulating signal transduction or enzymatic activity or the amount of a protein. In embodiments, inhibition refers to a reduction of activity of a target protein resulting from a direct interaction (e.g. an inhibitor binds to the target protein). In embodiments, inhibition refers to a reduction of activity of a target protein from an indirect interaction (e.g. an inhibitor binds to a protein that activates the target protein, thereby preventing target protein activation).

[0111] The terms “inhibitor,” “repressor” or “antagonist” or “downregulator” interchangeably refer to a substance capable of detectably decreasing the expression or activity of a given gene or protein. The antagonist can decrease expression or activity 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control in the absence of the antagonist. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or lower than the expression or activity in the absence of the antagonist. An antagonist prevents, reduces, inhibits, or neutralizes the activity of an agonist, and an antagonist can also prevent, inhibit, or reduce constitutive activity of a target, e.g., a target receptor, even where there is no identified agonist. In embodiments, inhibitors are molecules that decrease, block, prevent, delay activation, inactivate, desensitize, or down-regulate, e.g., a gene, protein, ligand, receptor, or cell. An inhibitor may also be defined as a molecule that reduces, blocks, or inactivates a constitutive activity. An “antagonist” is a molecule that opposes the action(s) of an agonist.

[0112] The terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” refer to a protein (including homologs, isoforms, and functional fragments thereof) and is a receptor for corticotropin releasing-hormone. The terms include any recombinant or naturally-occurring form of CRF₁ variants thereof that maintain

CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). The terms include any mutant form of CRF₁ variants (e.g., frameshift mutations) thereof that maintain CRF₁ activity (e.g. within at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% activity compared to wildtype CRF₁). In embodiments, the CRF₁ protein encoded by the *CRHR1* gene has the amino acid sequence set forth in or corresponding to Entrez 1233, UniProt P34998.

[0113] The term “expression” includes any step involved in the production of the polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion. Expression can be detected using conventional techniques for detecting protein (e.g., ELISA, Western blotting, flow cytometry, immunofluorescence, immunohistochemistry, *etc.*).

[0114] The terms “disease” or “condition” refer to a state of being or health status of a patient or subject capable of being treated with the compounds or methods provided herein. The disease may be a malignant or benign tumor (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)). The disease may be cancer. In some further instances, “cancer” refers to human cancers and carcinomas, sarcomas, adenocarcinomas, lymphomas, leukemias, etc., including solid and lymphoid cancers, kidney, breast, lung, bladder, colon, ovarian, prostate, pancreas, adrenal, stomach, brain, head and neck, skin, uterine, testicular, glioma, esophagus, and liver cancer, including hepatocarcinoma, lymphoma, including B-acute lymphoblastic lymphoma, non-Hodgkin’s lymphomas (e.g., Burkitt’s, Small Cell, and Large Cell lymphomas), Hodgkin’s lymphoma, leukemia (including MDS, AML, ALL, ATLL and CML), or multiple myeloma.

[0115] As used herein, the term “cancer” refers to all types of cancer, neoplasm or malignant tumors found in mammals (e.g. humans), including leukemia, carcinomas and sarcomas. Exemplary cancers that may be treated with a compound or method provided herein include adrenal cancer and tumors (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) brain cancer, glioma, glioblastoma, neuroblastoma, prostate cancer, colorectal cancer, pancreatic cancer, cervical cancer, gastric cancer, ovarian cancer, lung cancer, and cancer of the head. Exemplary cancers that may be treated with a compound or method provided herein include cancer of the thyroid, endocrine system, brain, breast, cervix, colon, head & neck, liver, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus, Medulloblastoma, colorectal cancer, pancreatic cancer. Additional examples include, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung

adenocarcinoma, lung squamous cell carcinoma, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, glioma, glioblastoma multiforme, ovarian cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, primary brain tumors, cancer, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, endometrial cancer, adrenal cortical cancer, neoplasms of the endocrine or exocrine pancreas, medullary thyroid cancer, medullary thyroid carcinoma, melanoma, colorectal cancer, papillary thyroid cancer, hepatocellular carcinoma, or prostate cancer.

[0116] The term “congenital adrenal hyperplasia” or “CAH” are any of several autosomal recessive diseases resulting from mutations of genes for enzymes mediating the biochemical steps of production of mineralocorticoids, glucocorticoids or sex steroids from cholesterol by the adrenal glands (steroidogenesis). These conditions involve excessive or deficient production of sex steroids and can alter development of primary or secondary sex characteristics in some affected infants, children, or adults.

[0117] “Adrenal rest tumor” refers to a neoplasm of the testis (or ovaries) consisting of adrenal tissue. It may be secondary to hyperplasia of an adrenal gland.

[0118] “Testicular adrenal rest tumor(s)” or “TART” and/or ovarian adrenal rest tumor(s) are a complication of congenital adrenal hyperplasia (CAH), which may develop from ectopic remnants of intra-testicular adrenal tissue stimulated by adrenocorticotrophic hormone (ACTH) hypersecretion. TART lesions are typically located within the rete testis and are bilateral, synchronous, nodular and multiple. TART can lead to testicular structural damage, spermatogenesis disorders, infertility and mass-forming lesions.

[0119] The term “leukemia” refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease-acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number abnormal cells in the blood-leukemic or aleukemic (subleukemic). Exemplary leukemias that may be treated with a compound or method provided herein include, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemc leukemia, basophylic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia,

hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, multiple myeloma, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, or undifferentiated cell leukemia.

[0120] The term “sarcoma” generally refers to a tumor which is made up of a substance like the embryonic connective tissue and is generally composed of closely packed cells embedded in a fibrillar or homogeneous substance. Sarcomas that may be treated with a compound or method provided herein include a chondrosarcoma, fibrosarcoma, lymphosarcoma, melanomasarcoma, myxosarcoma, osteosarcoma, Abemethy's sarcoma, adipose sarcoma, liposarcoma, alveolar soft part sarcoma, ameloblastic sarcoma, botryoid sarcoma, chloroma sarcoma, chorio carcinoma, embryonal sarcoma, Wilms' tumor sarcoma, endometrial sarcoma, stromal sarcoma, Ewing's sarcoma, fascial sarcoma, fibroblastic sarcoma, giant cell sarcoma, granulocytic sarcoma, Hodgkin's sarcoma, idiopathic multiple pigmented hemorrhagic sarcoma, immunoblastic sarcoma of B cells, lymphoma, immunoblastic sarcoma of T-cells, Jensen's sarcoma, Kaposi's sarcoma, Kupffer cell sarcoma, angiosarcoma, leukosarcoma, malignant mesenchymoma sarcoma, parosteal sarcoma, reticulocytic sarcoma, Rous sarcoma, serocystic sarcoma, synovial sarcoma, or telangiectaltic sarcoma.

[0121] The term “melanoma” is taken to mean a tumor arising from the melanocytic system of the skin and other organs. Melanomas that may be treated with a compound or method provided herein include, for example, acral-lentiginous melanoma, amelanotic melanoma, benign juvenile melanoma, Cloudman's melanoma, S91 melanoma, Harding-Passey melanoma, juvenile melanoma, lentigo maligna melanoma, malignant melanoma, nodular melanoma, subungal melanoma, or superficial spreading melanoma.

[0122] The term “carcinoma” refers to a malignant new growth made up of epithelial cells tending to infiltrate the surrounding tissues and give rise to metastases. Exemplary carcinomas that may be treated with a compound or method provided herein include, for example, thyroid carcinoma, cholangiocarcinoma, pancreatic adenocarcinoma, skin cutaneous melanoma, colon adenocarcinoma, rectum adenocarcinoma, stomach adenocarcinoma, esophageal carcinoma, head and neck squamous cell carcinoma, breast invasive carcinoma, lung adenocarcinoma, lung squamous cell carcinoma, medullary thyroid carcinoma, familial medullary thyroid carcinoma, acinar carcinoma, acinous carcinoma, adenocystic carcinoma, adenoid cystic carcinoma,

carcinoma adenomatosum, carcinoma of adrenal cortex, alveolar carcinoma, alveolar cell carcinoma, basal cell carcinoma, carcinoma basocellulare, basaloid carcinoma, basosquamous cell carcinoma, bronchioalveolar carcinoma, bronchiolar carcinoma, bronchogenic carcinoma, cerebriiform carcinoma, cholangiocellular carcinoma, chorionic carcinoma, colloid carcinoma, comedo carcinoma, corpus carcinoma, cribriform carcinoma, carcinoma en cuirasse, carcinoma cutaneum, cylindrical carcinoma, cylindrical cell carcinoma, duct carcinoma, carcinoma durum, embryonal carcinoma, encephaloid carcinoma, epiermoid carcinoma, carcinoma epitheliale adenoides, exophytic carcinoma, carcinoma ex ulcere, carcinoma fibrosum, gelatiniformi carcinoma, gelatinous carcinoma, giant cell carcinoma, carcinoma gigantocellulare, glandular carcinoma, granulosa cell carcinoma, hair-matrix carcinoma, hematoid carcinoma, hepatocellular carcinoma, Hurthle cell carcinoma, hyaline carcinoma, hypernephroid carcinoma, infantile embryonal carcinoma, carcinoma in situ, intraepidermal carcinoma, intraepithelial carcinoma, Krompecher's carcinoma, Kulchitzky-cell carcinoma, large-cell carcinoma, lenticular carcinoma, carcinoma lenticulare, lipomatous carcinoma, lymphoepithelial carcinoma, carcinoma medullare, medullary carcinoma, melanotic carcinoma, carcinoma molle, mucinous carcinoma, carcinoma muciparum, carcinoma mucocellulare, mucoepidermoid carcinoma, carcinoma mucosum, mucous carcinoma, carcinoma myxomatodes, nasopharyngeal carcinoma, oat cell carcinoma, carcinoma ossificans, osteoid carcinoma, papillary carcinoma, periportal carcinoma, preinvasive carcinoma, prickle cell carcinoma, pultaceous carcinoma, renal cell carcinoma of kidney, reserve cell carcinoma, carcinoma sarcomatodes, schneiderian carcinoma, scirrhus carcinoma, carcinoma scroti, signet-ring cell carcinoma, carcinoma simplex, small-cell carcinoma, solanoid carcinoma, spheroidal cell carcinoma, spindle cell carcinoma, carcinoma spongiosum, squamous carcinoma, squamous cell carcinoma, string carcinoma, carcinoma telangiectaticum, carcinoma telangiectodes, transitional cell carcinoma, carcinoma tuberosum, tuberos carcinoma, verrucous carcinoma, or carcinoma villosum.

[0123] The terms “treating” or “treatment” refer to any indicia of success in the therapy or amelioration of an injury, disease, pathology or condition, including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the injury, pathology or condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; improving a patient’s physical or mental well-being. The treatment or amelioration of symptoms can be based on objective or subjective parameters; including the results of a physical examination, neuropsychiatric exams, and/or a psychiatric evaluation. The term “treating” and conjugations thereof, may include prevention of an injury, pathology, condition, or disease. In embodiments, treating is preventing. In embodiments, treating does not include preventing.

[0124] “Treating” or “treatment” as used herein (and as well-understood in the art) also broadly includes any approach for obtaining beneficial or desired results in a subject’s condition, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of the extent of a disease, stabilizing (*i.e.*, not worsening) the state of disease, prevention of a disease’s transmission or spread, delay or slowing of disease progression, amelioration or palliation of the disease state, diminishment of the reoccurrence of disease, and remission, whether partial or total and whether detectable or undetectable. In other words, “treatment” as used herein includes any cure, amelioration, or prevention of a disease. Treatment may prevent the disease from occurring; inhibit the disease’s spread; relieve the disease’s symptoms (*e.g.*, ocular pain, seeing halos around lights, red eye, very high intraocular pressure), fully or partially remove the disease’s underlying cause, shorten a disease’s duration, or do a combination of these things.

[0125] “Treating” and “treatment” as used herein include prophylactic treatment. Treatment methods include administering to a subject a therapeutically effective amount of a compound described herein. The administering step may consist of a single administration or may include a series of administrations. The length of the treatment period depends on a variety of factors, such as the severity of the condition, the age of the patient, the concentration of the compound, the activity of the compositions used in the treatment, or a combination thereof. It will also be appreciated that the effective dosage of an agent used for the treatment or prophylaxis may increase or decrease over the course of a particular treatment or prophylaxis regime. Changes in dosage may result and become apparent by standard diagnostic assays known in the art. In some instances, chronic administration may be required. For example, the compositions are administered to the subject in an amount and for a duration sufficient to treat the patient.

[0126] The term “prevent” refers to a decrease in the occurrence of disease symptoms in a patient. As indicated above, the prevention may be complete (no detectable symptoms) or partial, such that fewer symptoms are observed than would likely occur absent treatment. In embodiments, prevent refers to slowing the progression of the disease, disorder or condition or inhibiting progression thereof to a harmful or otherwise undesired state.

[0127] “Patient” or “subject in need thereof” refers to a living organism suffering from or prone to a disease or condition that can be treated by administration of a pharmaceutical composition as provided herein. Non-limiting examples include humans, other mammals, bovines, rats, mice, dogs, monkeys, goat, sheep, cows, deer, and other non-mammalian animals. In some embodiments, a patient is human.

[0128] An “effective amount” is an amount sufficient for a compound to accomplish a stated purpose relative to the absence of the compound (*e.g.*, achieve the effect for which it is

administered, treat a disease, reduce enzyme activity, increase enzyme activity, reduce a signaling pathway, or reduce one or more symptoms of a disease or condition). An example of an “effective amount” is an amount sufficient to contribute to the treatment, prevention, or reduction of a symptom or symptoms of a disease, which could also be referred to as a “therapeutically effective amount.” A “reduction” of a symptom or symptoms (and grammatical equivalents of this phrase) means decreasing of the severity or frequency of the symptom(s), or elimination of the symptom(s). A “prophylactically effective amount” of a drug is an amount of a drug that, when administered to a subject, will have the intended prophylactic effect, e.g., preventing or delaying the onset (or reoccurrence) of an injury, disease, pathology or condition, or reducing the likelihood of the onset (or reoccurrence) of an injury, disease, pathology, or condition, or their symptoms. The full prophylactic effect does not necessarily occur by administration of one dose, and may occur only after administration of a series of doses. Thus, a prophylactically effective amount may be administered in one or more administrations. An “activity decreasing amount,” as used herein, refers to an amount of antagonist required to decrease the activity of an enzyme relative to the absence of the antagonist. A “function disrupting amount,” as used herein, refers to the amount of antagonist required to disrupt the function of an enzyme or protein relative to the absence of the antagonist. The exact amounts will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (*see, e.g.,* Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); and *Remington: The Science and Practice of Pharmacy*, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins). The therapeutically effective amount can be ascertained by measuring relevant physiological effects, and it can be adjusted in connection with the dosing regimen and diagnostic analysis of the subject’s condition, and the like. By way of example, measurement of the serum level of a CRF1 inhibitor (or, e.g., a metabolite thereof) at a particular time post-administration may be indicative of whether a therapeutically effective amount has been administered.

[0129] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0130] Therapeutically effective amounts for use in humans can also be determined from animal models. For example, a dose for humans can be formulated to achieve a concentration that has been found to be effective in animals. The dosage in humans can be adjusted by monitoring compounds effectiveness and adjusting the dosage upwards or downwards, as described above.

Adjusting the dose to achieve maximal efficacy in humans based on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan. Adjusting the dose to achieve maximal therapeutic window efficacy or toxicity in humans based on the methods described above and other methods is well within the capabilities of the ordinarily skilled artisan.

[0131] The term “therapeutically effective amount,” as used herein, refers to that amount of the therapeutic agent sufficient to ameliorate the disorder, as described above. For example, for the given parameter, a therapeutically effective amount will show an increase or decrease of at least 5%, 10%, 15%, 20%, 25%, 40%, 50%, 60%, 75%, 80%, 90%, or at least 100%. Therapeutic efficacy can also be expressed as “-fold” increase or decrease. For example, a therapeutically effective amount can have at least a 1.2-fold, 1.5-fold, 2-fold, 5-fold, or more effect over a control.

[0132] The terms “QD,” “qd,” and “q.d.” refer to once a day, i.e., a drug or therapeutic agent is administered once a day for the duration of the therapeutic regimen.

[0133] The terms “BD,” “bd,” and “b.i.d.” refer to twice a day, i.e., a drug or therapeutic agent is administered two times a day for the duration of the therapeutic regimen.

[0134] The terms “QID,” “qid,” and “q.i.d.” refer to three times a day, i.e., a drug or therapeutic agent is administered three times a day for the duration of the therapeutic regimen.

[0135] Dosages may be varied depending upon the requirements of the patient and the compound being employed. The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial therapeutic response in the patient over time. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects. Determination of the proper dosage for a particular situation is within the skill of the practitioner. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. Dosage amounts and intervals can be adjusted individually to provide levels of the administered compound effective for the particular clinical indication being treated. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease state.

[0136] As used herein, the term “administering” means oral administration, administration as a suppository, topical contact, intravenous, parenteral, intraperitoneal, intramuscular, intralesional, intrathecal, intracranial, intranasal or subcutaneous administration, or the implantation of a slow-release device, *e.g.*, a mini-osmotic pump, to a subject. Administration is by any route, including parenteral and transmucosal (*e.g.*, buccal, sublingual, palatal, gingival, nasal, vaginal, rectal, or transdermal). Parenteral administration includes, *e.g.*, intravenous, intramuscular, intra-arteriole,

intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial. Other modes of delivery include, but are not limited to, the use of liposomal formulations, intravenous infusion, transdermal patches, *etc.* By “co-administer” it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies (e.g. anti-cancer agent, chemotherapeutic, or treatment for a neurodegenerative disease). The compound of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compound individually or in combination (more than one compound or agent). Thus, the preparations can also be combined, when desired, with other active substances (e.g. to reduce metabolic degradation). The compositions of the present invention can be delivered by transdermally, by a topical route, formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols. Oral preparations include tablets, pills, powder, dragees, capsules, liquids, lozenges, cachets, gels, syrups, slurries, suspensions, *etc.*, suitable for ingestion by the patient. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. The compositions of the present invention may additionally include components to provide sustained release and/or comfort. Such components include high molecular weight, anionic mucomimetic polymers, gelling polysaccharides and finely-divided drug carrier substrates. These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841; 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by reference in their entirety for all purposes. The compositions of the present invention can also be delivered as microspheres for slow release in the body. For example, microspheres can be administered via intradermal injection of drug-containing microspheres, which slowly release subcutaneously (see Rao, *J. Biomater Sci. Polym. Ed.* 7:623-645, 1995; as biodegradable and injectable gel formulations (see, e.g., Gao *Pharm. Res.* 12:857-863, 1995); or, as microspheres for oral administration (see, e.g., Eyles, *J. Pharm. Pharmacol.* 49:669-674, 1997). In another embodiment, the formulations of the compositions of the present invention can be delivered by the use of liposomes which fuse with the cellular membrane or are endocytosed, *i.e.*, by employing receptor ligands attached to the liposome, that bind to surface membrane protein receptors of the cell resulting in endocytosis. By using liposomes, particularly where the liposome surface carries receptor ligands specific for target cells, or are otherwise preferentially directed to a specific organ, one can focus the delivery of the compositions of the present invention into the target cells *in vivo*. (See, e.g., Al-Muhammed, *J. Microencapsul.* 13:293-306, 1996; Chonn, *Curr. Opin. Biotechnol.* 6:698-708, 1995; Ostro, *Am. J. Hosp. Pharm.*

46:1576-1587, 1989). The compositions of the present invention can also be delivered as nanoparticles.

[0137] By “co-administer” it is meant that a composition described herein is administered at the same time, just prior to, or just after the administration of one or more additional therapies. The compounds of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the compounds individually or in combination (more than one compound). The compositions of the present invention can be delivered transdermally, by a topical route, or formulated as applicator sticks, solutions, suspensions, emulsions, gels, creams, ointments, pastes, jellies, paints, powders, and aerosols.

[0138] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of achieving the methods described herein, as measured using the methods described herein or known in the art.

[0139] Utilizing the teachings provided herein, an effective prophylactic or therapeutic treatment regimen can be planned that does not cause substantial toxicity and yet is effective to treat the clinical symptoms demonstrated by the particular patient. This planning should involve the careful choice of active compound by considering factors such as compound potency, relative bioavailability, patient body weight, presence and severity of adverse side effects, preferred mode of administration and the toxicity profile of the selected agent.

[0140] The compounds described herein can be used in combination with one another, with other active agents known to be useful in treating cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0141] In some embodiments, co-administration includes administering one active agent within 0.5, 1, 2, 4, 6, 8, 10, 12, 16, 20, 24 hours, 2 days, 4 days, 1 week or 1 month of a second active agent. Co-administration includes administering two active agents simultaneously, approximately simultaneously (e.g., within about 1, 5, 10, 15, 20, or 30 minutes of each other), or sequentially in any order. In some embodiments, co-administration can be accomplished by co-formulation, i.e., preparing a single pharmaceutical composition including both active agents. In other embodiments, the active agents can be formulated separately. In another embodiment, the active and/or adjunctive agents may be linked or conjugated to one another. In some embodiments, the compounds described herein may be combined with treatments for cancer (e.g. colon cancer), cardiovascular disease, metabolic disease, immune or inflammatory disease or disorder.

[0142] The compounds described herein can be administered to treat a fertility disease or disorder and/or cancer. In this regard, the compounds disclosed herein may be administered either alone to treat such diseases or disorders or may be co-administered with another therapeutic agent to treat such diseases or disorders.

[0143] The compounds disclosed herein may be co-administered with an agonist or antagonist of CRF₁ function. The compounds described herein can be co-administered with conventional anti-cancer/chemotherapeutic.

[0144] “Anti-cancer agent” is used in accordance with its plain ordinary meaning and refers to a composition (e.g. compound, drug, antagonist, inhibitor, modulator) having antineoplastic properties or the ability to inhibit the growth or proliferation of cells. In some embodiments, an anti-cancer agent is a chemotherapeutic. In some embodiments, an anti-cancer agent is an agent identified herein having utility in methods of treating cancer. In some embodiments, an anti-cancer agent is an agent approved by the FDA or similar regulatory agency of a country other than the USA, for treating cancer. Examples of anti-cancer agents include, but are not limited to, MEK (e.g. MEK1, MEK2, or MEK1 and MEK2) inhibitors (e.g. XL518, CI-1040, PD035901, selumetinib/ AZD6244, GSK1120212/ trametinib, GDC-0973, ARRY-162, ARRY-300, AZD8330, PD0325901, U0126, PD98059, TAK-733, PD318088, AS703026, BAY 869766), alkylating agents (e.g., cyclophosphamide, ifosfamide, chlorambucil, busulfan, melphalan, mechlorethamine, uramustine, thiotepa, nitrosoureas, nitrogen mustards (e.g., mechlorethamine, cyclophosphamide, chlorambucil, melphalan), ethylenimine and methylmelamines (e.g., hexamethylmelamine, thiotepa), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine, lomustine, semustine, streptozocin), triazenes (decarbazine)), anti-metabolites (e.g., 5-azathioprine, leucovorin, capecitabine, fludarabine, gemcitabine, pemetrexed, raltitrexed, folic acid analog (e.g., methotrexate), or pyrimidine analogs (e.g., fluorouracil, floxouridine, Cytarabine), purine analogs (e.g., mercaptopurine, thioguanine, pentostatin), *etc.*), plant alkaloids (e.g., vincristine, vinblastine, vinorelbine, vindesine, podophyllotoxin, paclitaxel, docetaxel, *etc.*), topoisomerase inhibitors (e.g., irinotecan, topotecan, amsacrine, etoposide (VP16), etoposide phosphate, teniposide, *etc.*), antitumor antibiotics (e.g., doxorubicin, adriamycin, daunorubicin, epirubicin, actinomycin, bleomycin, mitomycin, mitoxantrone, plicamycin, *etc.*), platinum-based compounds (e.g. cisplatin, oxaloplatin, carboplatin), anthracenedione (e.g., mitoxantrone), substituted urea (e.g., hydroxyurea), methyl hydrazine derivative (e.g., procarbazine), adrenocortical suppressant (e.g., mitotane, aminoglutethimide), epipodophyllotoxins (e.g., etoposide), antibiotics (e.g., daunorubicin, doxorubicin, bleomycin), enzymes (e.g., L-asparaginase), inhibitors of mitogen-activated protein kinase signaling (e.g. U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-

9006, wortmannin, or LY294002, Syk inhibitors, mTOR inhibitors, antibodies (e.g., rituxan), gossypol, genasense, polyphenol E, Chlorofusin, all trans-retinoic acid (ATRA), bryostatins, tumor necrosis factor-related apoptosis-inducing ligand (TRAIL), 5-aza-2'-deoxycytidine, all trans retinoic acid, doxorubicin, vincristine, etoposide, gemcitabine, imatinib (Gleevec.RTM.), geldanamycin, 17-N-Allylamino-17-Demethoxygeldanamycin (17-AAG), flavopiridol, LY294002, bortezomib, trastuzumab, BAY 11-7082, PKC412, PD184352, 20-epi-1, 25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; broprimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydridemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziqune; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; 9-dioxamycin; diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene;

idramantone; ilmofofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides;
 insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins;
 iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole;
 isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide;
 leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor;
 leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarazole;
 linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds;
 lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin;
 loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin
 A; marimastat; masoprocot; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors;
 menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone;
 miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol;
 mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone;
 mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin;
 monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene
 inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B;
 mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides;
 nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin;
 nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide
 modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone;
 oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer;
 ormaplatin; osaterone; oxaliplatin; oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic
 acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan
 polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol;
 phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride;
 pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum
 complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin;
 prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based
 immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein
 tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins;
 pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists;
 raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP
 inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII
 retinamide; roglitimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol;
 saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence

derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen-binding protein; sizofuran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; tricyribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; zinostatin stimalamer, Adriamycin, Dactinomycin, Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqune; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; iimofosine; interleukin II (including recombinant interleukin II, or rIL.sub.2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1; interferon alfa-n3; interferon beta-1a; interferon gamma-1b; iuprolatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole

hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedapa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; pipsulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricirbine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulazole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride, agents that arrest cells in the G2-M phases and/or modulate the formation or stability of microtubules, (e.g. Taxol.TM (i.e. paclitaxel), Taxotere.TM, compounds comprising the taxane skeleton, Erbulozole (i.e. R-55104), Dolastatin 10 (i.e. DLS-10 and NSC-376128), Mivobulin isethionate (i.e. as CI-980), Vincristine, NSC-639829, Discodermolide (i.e. as NVP-XX-A-296), ABT-751 (Abbott, i.e. E-7010), Altorhyrtins (e.g. Altorhyrtin A and Altorhyrtin C), Spongistatins (e.g. Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5, Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride (i.e. LU-103793 and NSC-D-669356), Epothilones (e.g. Epothilone A, Epothilone B, Epothilone C (i.e. desoxyepothilone A or dEpoA), Epothilone D (i.e. KOS-862, dEpoB, and desoxyepothilone B), Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide, 16-aza-epothilone B, 21-aminoepothilone B (i.e. BMS-310705), 21-hydroxyepothilone D (i.e. Desoxyepothilone F and dEpoF), 26-fluoroepothilone, Auristatin PE (i.e. NSC-654663), Soblidotin (i.e. TZT-1027), LS-4559-P (Pharmacia, i.e. LS-4577), LS-4578 (Pharmacia, i.e. LS-477-P), LS-4477 (Pharmacia), LS-4559 (Pharmacia), RPR-112378 (Aventis), Vincristine sulfate, DZ-3358 (Daiichi), FR-182877 (Fujisawa, i.e. WS-9885B), GS-164 (Takeda), GS-198 (Takeda), KAR-2 (Hungarian Academy of Sciences), BSF-223651 (BASF, i.e. ILX-651 and LU-223651), SAH-49960 (Lilly/Novartis), SDZ-268970 (Lilly/Novartis), AM-97 (Armad/Kyowa Hakko), AM-132 (Armad), AM-138 (Armad/Kyowa Hakko), IDN-5005 (Indena), Cryptophycin 52 (i.e.

LY-355703), AC-7739 (Ajinomoto, i.e. AVE-8063A and CS-39.HCl), AC-7700 (Ajinomoto, i.e. AVE-8062, AVE-8062A, CS-39-L-Ser.HCl, and RPR-258062A), Vitilevuamide, Tubulysin A, Canadensol, Centaureidin (i.e. NSC-106969), T-138067 (Tularik, i.e. T-67, TL-138067 and TI-138067), COBRA-1 (Parker Hughes Institute, i.e. DDE-261 and WHI-261), H10 (Kansas State University), H16 (Kansas State University), Oncocidin A1 (i.e. BTO-956 and DIME), DDE-313 (Parker Hughes Institute), Fijianolide B, Laulimalide, SPA-2 (Parker Hughes Institute), SPA-1 (Parker Hughes Institute, i.e. SPIKET-P), 3-IAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-569), Narcosine (also known as NSC-5366), Nascapine, D-24851 (Asta Medica), A-105972 (Abbott), Hemiasterlin, 3-BAABU (Cytoskeleton/Mt. Sinai School of Medicine, i.e. MF-191), TMPN (Arizona State University), Vanadocene acetylacetonate, T-138026 (Tularik), Monsatrol, Inanocine (i.e. NSC-698666), 3-IAABE (Cytoskeleton/Mt. Sinai School of Medicine), A-204197 (Abbott), T-607 (Tularik, i.e. T-900607), RPR-115781 (Aventis), Eleutherobins (such as Desmethyleleutherobin, Desaetyeleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, D-64131 (Asta Medica), D-68144 (Asta Medica), Diazonamide A, A-293620 (Abbott), NPI-2350 (Nereus), Taccalonolide A, TUB-245 (Aventis), A-259754 (Abbott), Diozostatin, (-)-Phenylahistin (i.e. NSCL-96F037), D-68838 (Asta Medica), D-68836 (Asta Medica), Myoseverin B, D-43411 (Zentaris, i.e. D-81862), A-289099 (Abbott), A-318315 (Abbott), HTI-286 (i.e. SPA-110, trifluoroacetate salt) (Wyeth), D-82317 (Zentaris), D-82318 (Zentaris), SC-12983 (NCI), Resverastatin phosphate sodium, BPR-OY-007 (National Health Research Institutes), and SSR-250411 (Sanofi)), steroids (*e.g.*, dexamethasone), finasteride, aromatase inhibitors, gonadotropin-releasing hormone agonists (GnRH) such as goserelin or leuprolide, adrenocorticosteroids (*e.g.*, prednisone), progestins (*e.g.*, hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (*e.g.*, diethylstilbestrol, ethinyl estradiol), antiestrogen (*e.g.*, tamoxifen), androgens (*e.g.*, testosterone propionate, fluoxymesterone), antiandrogen (*e.g.*, flutamide), immunostimulants (*e.g.*, Bacillus Calmette-Guérin (BCG), levamisole, interleukin-2, alpha-interferon, *etc.*), monoclonal antibodies (*e.g.*, anti-CD20, anti-HER2, anti-CD52, anti-HLA-DR, and anti-VEGF monoclonal antibodies), immunotoxins (*e.g.*, anti-CD33 monoclonal antibody-calicheamicin conjugate, anti-CD22 monoclonal antibody-pseudomonas exotoxin conjugate, *etc.*), radioimmunotherapy (*e.g.*, anti-CD20 monoclonal antibody conjugated to ¹¹¹In, ⁹⁰Y, or ¹³¹I, *etc.*), triptolide, homoharringtonine, dactinomycin, doxorubicin, epirubicin, topotecan, itraconazole, vindesine, cerivastatin, vincristine, deoxyadenosine, sertraline, pitavastatin, irinotecan, clofazimine, 5-nonyloxytryptamine, vemurafenib, dabrafenib, erlotinib, gefitinib, EGFR inhibitors, epidermal growth factor receptor (EGFR)-targeted therapy or therapeutic (*e.g.* gefitinib (Iressa™), erlotinib (Tarceva™), cetuximab (Erbix™), lapatinib

(Tykerb™), panitumumab (Vectibix™), vandetanib (Caprelsa™), afatinib/BIBW2992, CI-1033/canertinib, neratinib/HKI-272, CP-724714, TAK-285, AST-1306, ARRY334543, ARRY-380, AG-1478, dacomitinib/PF299804, OSI-420/desmethyl erlotinib, AZD8931, AEE788, pelitinib/EKB-569, CUDC-101, WZ8040, WZ4002, WZ3146, AG-490, XL647, PD153035, BMS-599626), sorafenib, imatinib, sunitinib, dasatinib, or the like.

[0145] “Chemotherapeutic” or “chemotherapeutic agent” is used in accordance with its plain ordinary meaning and refers to a chemical composition or compound having antineoplastic properties or the ability to inhibit the growth or proliferation of cells.

[0146] The compounds disclosed herein may be co-administered with an antiproliferative/antineoplastic drug or a combination thereof, as used in medical oncology, such as an alkylating agent (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan or a nitrosourea); an antimetabolite (for example an antifolate such as a fluoropyrimidine like 5-fluorouracil or tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, gemcitabine or paclitaxel); an antitumour antibiotic (for example an anthracycline such as adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin or mithramycin); an antimitotic agent (for example a vinca alkaloid such as vincristine, vinblastine, vindesine or vinorelbine, or a taxoid such as taxol or taxotere); or a topoisomerase inhibitor (for example an epipodophyllotoxin such as etoposide, teniposide, amsacrine, topotecan or a camptothecin); (ii) a cytostatic agent such as an antioestrogen (for example tamoxifen, toremifene, raloxifene, droloxifene or iodoxyfene), an oestrogen receptor down regulator (for example fulvestrant), an antiandrogen (for example bicalutamide, flutamide, nilutamide or cyproterone acetate), a LHRH antagonist or LHRH agonist (for example goserelin, leuprorelin or buserelin), a progestogen (for example megestrol acetate), an aromatase inhibitor (for example as anastrozole, letrozole, vorazole or exemestane) or an inhibitor of 5.alpha.-reductase such as finasteride; (iii) an agent which inhibits cancer cell invasion (for example a metalloproteinase inhibitor like marimastat or an inhibitor of urokinase plasminogen activator receptor function); (iv) an inhibitor of growth factor function, for example: a growth factor antibody (for example the anti-erbB2 antibody trastuzumab, or the anti-erbB1 antibody cetuximab [C225]), a farnesyl transferase inhibitor, a tyrosine kinase inhibitor or a serine/threonine kinase inhibitor, an inhibitor of the epidermal growth factor family (for example an EGFR family tyrosine kinase inhibitor such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD 1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) or 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), an inhibitor of the platelet-derived growth factor family, or an inhibitor of the hepatocyte growth factor family; (v)

an antiangiogenic agent such as one which inhibits the effects of vascular endothelial growth factor (for example the anti-vascular endothelial cell growth factor antibody bevacizumab, a compound disclosed in WO 97/22596, WO 97/30035, WO 97/32856 or WO 98/13354), or a compound that works by another mechanism (for example linomide, an inhibitor of integrin .alpha.v.beta.3 function or an angiostatin); (vi) a vascular damaging agent such as combretastatin A4, or a compound disclosed in WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 or WO 02/08213; (vii) an agent used in antisense therapy, for example one directed to one of the targets listed above, such as ISIS 2503, an anti-ras antisense; (viii) an agent used in a gene therapy approach, for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; or (ix) an agent used in an immunotherapeutic approach, for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

[0147] In embodiments, the compounds disclosed herein can be co-administered with an antibody, such as a monoclonal antibody targeting B-Lymphocytes (such as CD20 (rituximab), MRA-aIL16R and T-Lymphocytes, CTLA4-Ig, HuMax IL-15) or antibody modulating Ig function such as anti-IgE (for example omalizumab).

[0148] In embodiments, treatment of cancer includes administration of an effective amount of at least two of the following: a CRF1 inhibitor, an inhibitor of the PD-L1/PD-1 pathway, an inhibitor of CTLA-4, an agonistic antibody of CD137 (4-1BB). In some embodiments, the method may comprise the use of two or more combinations.

[0149] In a further embodiment, the compounds described herein can be co-administered with conventional radiotherapeutic agents including, but not limited to, radionuclides such as ⁴⁷Sc, ⁶⁴Cu, ⁶⁷Cu, ⁸⁹Sr, ⁸⁶Y, ⁸⁷Y, ⁹⁰Y, ¹⁰⁵Rh, ¹¹¹Ag, ¹¹¹In, ^{117m}Sn, ¹⁴⁹Pm, ¹⁵³Sm, ¹⁶⁶Ho, ¹⁷⁷Lu, ¹⁸⁶Re, ¹⁸⁸Re, ²¹¹At, and ²¹²Bi, optionally conjugated to antibodies directed against tumor antigens.

[0150] In addition, a CRF₁ inhibitor may be combined with the therapeutic administration of immune cells, sometimes referred to as adoptive cell transfer. These cells may be cells from the patient, a genetically related or unrelated donor, they may be genetically modified (e.g. CAR-T cells, NK cells, etc), cell lines, genetically modified cell lines and live or dead versions of the

above. CRF₁ inhibitors may also be combined with vaccines of any kind (e.g. protein/peptide, viral, bacterial, cellular) to stimulate immune responses to cancer.

[0151] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramusculay, intranasal, intraocularal, intrapericardial, intraperitoneal, intrapleural, intraprostatical, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenousl, intravesicullar, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0152] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein can be administered by any acceptable route, such oral, intraadiposal, intraarterial, intraarticular, intracranial, intradermal, intralesional, intramusculay, intranasal, intraocularal, intrapericardial, intraperitoneal, intrapleural, intraprostatical, intrarectal, intrathecal, intratracheal, intratumoral, intraumbilical, intravaginal, intravenousl, intravesicullar, intravitreal, liposomal, local, mucosal, parenteral, rectal, subconjunctival, subcutaneous, sublingual, topical, transbuccal, transdermal, vaginal, in cremes, in lipid compositions, via a catheter, via a lavage, via continuous infusion, via infusion, via inhalation, via injection, via local delivery, via localized perfusion, bathing target cells directly, or any combination thereof.

[0153] The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered once daily until study reached endpoint. The CRF₁ inhibitors/CRF₁ antagonists disclosed herein may be administered at least three times but in some studies four or more times depending on the length of the study and/or the design of the study.

[0154] The methods disclosed herein may be used in combination with additional cancer therapy. In some embodiments, the distinct cancer therapy comprises surgery, radiotherapy, chemotherapy, toxin therapy, immunotherapy, cryotherapy or gene therapy. In some embodiments, the cancer is a chemotherapy-resistant or radio-resistant cancer.

[0155] A “cell” as used herein, refers to a cell carrying out metabolic or other function sufficient to preserve or replicate its genomic DNA. A cell can be identified by well-known methods in the art including, for example, presence of an intact membrane, staining by a particular dye, ability to produce progeny or, in the case of a gamete, ability to combine with a second gamete to produce a viable offspring. Cells may include prokaryotic and eukaryotic cells. Prokaryotic cells include but are not limited to bacteria. Eukaryotic cells include but are not limited to yeast cells and cells derived from plants and animals, for example mammalian, insect (e.g., spodoptera) and

human cells. Cells may be useful when they are naturally nonadherent or have been treated not to adhere to surfaces, for example by trypsinization.

[0156] “Control” or “control experiment” is used in accordance with its plain ordinary meaning and refers to an experiment in which the subjects or reagents of the experiment are treated as in a parallel experiment except for omission of a procedure, reagent, or variable of the experiment. In some instances, the control is used as a standard of comparison in evaluating experimental effects. In some embodiments, a control is the measurement of the activity of a protein in the absence of a compound as described herein (including embodiments and examples).

[0157] The term “modulator” refers to a composition that increases or decreases the level of a target molecule or the function of a target molecule or the physical state of the target of the molecule. In some embodiments, a CRF₁ associated disease modulator is a compound that reduces the severity of one or more symptoms of a disease associated with CRF₁ (e.g. an endocrine disorder, a cancer including but not limited to TART or OART). A CRF₁ modulator is a compound that increases or decreases the activity or function or level of activity or level of function of CRF₁. A modulator may act alone, or it may use a cofactor, e.g., a protein, metal ion, or small molecule. Examples of modulators include small molecule compounds and other bioorganic molecules. Numerous libraries of small molecule compounds (e.g., combinatorial libraries) are commercially available and can serve as a starting point for identifying a modulator. The skilled artisan is able to develop one or more assays (e.g., biochemical or cell-based assays) in which such compound libraries can be screened in order to identify one or more compounds having the desired properties; thereafter, the skilled medicinal chemist is able to optimize such one or more compounds by, for example, synthesizing and evaluating analogs and derivatives thereof. Synthetic and/or molecular modeling studies can also be utilized in the identification of an activator.

[0158] The term “modulate” is used in accordance with its plain ordinary meaning and refers to the act of changing or varying one or more properties. “Modulation” refers to the process of changing or varying one or more properties. For example, as applied to the effects of a modulator on a target protein, to modulate means to change by increasing or decreasing a property or function of the target molecule or the amount of the target molecule. In embodiments, the terms “modulate”, “modulation” and the like refer to the ability of a molecule (e.g., an activator or an inhibitor) to increase or decrease the function or activity of CRF₁, either directly or indirectly, relative to the absence of the molecule.

[0159] The term “associated” or “associated with” in the context of a substance or substance activity or function associated with a disease (e.g. a protein associated disease, an endocrine disorder, a disease associated with CRF₁ activity, a CRF₁ associated disease (e.g., testicular

adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) means that the disease (e.g., testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART)) is caused by (in whole or in part), or a symptom of the disease is caused by (in whole or in part) the substance or substance activity or function. For example, an endocrine disorder associated with CRF₁ activity or function may be an endocrine disorder that results (entirely or partially) from aberrant CRF₁ function (e.g. enzyme activity, protein-protein interaction, signaling pathway) or an endocrine disorder wherein a particular symptom of the disease is caused (entirely or partially) by aberrant CRF₁ activity or function. As used herein, what is described as being associated with a disease, if a causative agent, could be a target for treatment of the disease. For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated disease (e.g., an endocrine disorder including but not limited to TART or OART), may be treated with a compound described herein (e.g., CRF₁ modulator, CRF₁ inhibitor, or CRF₁ antagonist), in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease (e.g., an endocrine disorder such as TART or OART). For example, an endocrine disorder associated with CRF₁ activity or function or a CRF₁ associated endocrine disorder, may be treated with an CRF₁ modulator or CRF₁ inhibitor, in the instance where increased CRF₁ activity or function (e.g. signaling pathway activity) causes the disease.

[0160] The term “aberrant” as used herein refers to different from normal. When used to describe enzymatic activity or protein function, aberrant refers to activity or function that is greater or less than a normal control or the average of normal non-diseased control samples. Aberrant activity may refer to an amount of activity that results in a disease, wherein returning the aberrant activity to a normal or non-disease-associated amount (e.g. by administering a compound or using a method as described herein), results in reduction of the disease or one or more disease symptoms.

[0161] The term “signaling pathway” as used herein refers to a series of interactions between cellular and optionally extra-cellular components (e.g. proteins, nucleic acids, small molecules, ions, lipids) that conveys a change in one component to one or more other components, which in turn may convey a change to additional components, which is optionally propagated to other signaling pathway components. For example, binding of a CRF₁ with a compound as described herein may reduce the level of a product of the CRF₁ catalyzed reaction or the level of a downstream derivative of the product or binding may reduce the interactions between the CRF₁ or a reaction product and downstream effectors or signaling pathway components (e.g., MAP kinase pathway), resulting in changes in cell growth, proliferation, or survival.

[0162] As used herein, the terms “corticotropin-releasing factor type-1,” “CRF₁,” “corticotropin-releasing hormone receptor 1,” and “CHHR1” and all other related art-accepted

terms, many of which are set forth below, refer to a compound capable of modulating, either directly or indirectly, the CRF₁ receptor in an *in vitro* assay, an *in vivo* model, and/or other means indicative of therapeutic efficacy. The terms also refer to a compound that exhibits at least some therapeutic benefit in a human subject.

[0163] The phrase “in a sufficient amount to effect a change” means that there is a detectable difference between a level of an indicator measured before (e.g., a baseline level) and after administration of a particular therapy. Indicators include any objective parameter (e.g., serum concentration) or subjective parameter (e.g., a subject’s feeling of well-being).

[0164] The “activity” of a molecule may describe or refer to the binding of the molecule to a ligand or to a receptor; to catalytic activity; to the ability to stimulate gene expression or cell signaling, differentiation, or maturation; to antigenic activity; to the modulation of activities of other molecules; and the like. The term “proliferative activity” encompasses an activity that promotes, that is necessary for, or that is specifically associated with, for example, normal cell division, as well as cancer, tumors, dysplasia, cell transformation, metastasis, and angiogenesis.

[0165] “Substantially pure” indicates that a component makes up greater than about 50% of the total content of the composition, and typically greater than about 60% of the total polypeptide content. More typically, “substantially pure” refers to compositions in which at least 75%, at least 85%, at least 90% or more of the total composition is the component of interest. In some cases, the polypeptide will make up greater than about 90%, or greater than about 95% of the total content of the composition (percentage in a weight per weight basis).

[0166] The terms “specifically binds” and “selectively binds,” when referring to a ligand/receptor, antibody/antigen, or other binding pair, indicates a binding reaction which is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated conditions, a specified ligand binds to a particular receptor and does not bind in a significant amount to other proteins present in the sample. The antibody, or binding composition derived from the antigen-binding site of an antibody, of the contemplated method binds to its antigen, or a variant or mutein thereof, with an affinity that is at least two-fold greater, at least 10-times greater, at least 20-times greater, or at least 100-times greater than the affinity with any other antibody, or binding composition derived therefrom. In embodiments, the antibody will have an affinity that is greater than about 10⁹ liters/mol, as determined by, e.g., Scatchard analysis (Munsen, et al. (1980) *Analyt. Biochem.* 107:220-239).

[0167] The terms “DNA,” “nucleic acid,” “nucleic acid molecule,” “polynucleotide” and the like are used interchangeably herein to refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Non-limiting examples of polynucleotides include linear and circular nucleic acids, messenger RNA (mRNA),


complementary DNA (cDNA), recombinant polynucleotides, vectors, probes, primers and the like.

[0168] As used herein, the terms “variants” and “homologs” are used interchangeably to refer to amino acid or nucleic acid sequences that are similar to reference amino acid or nucleic acid sequences, respectively. The term encompasses naturally-occurring variants and non-naturally-occurring variants. Naturally-occurring variants include homologs (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one species to another), and allelic variants (polypeptides and nucleic acids that differ in amino acid or nucleotide sequence, respectively, from one individual to another within a species). Thus, variants and homologs encompass naturally occurring amino acid and nucleic acid sequences encoded thereby and their isoforms, as well as splice variants of a protein or gene. The terms also encompass nucleic acid sequences that vary in one or more bases from a naturally-occurring nucleic acid sequence but still translate into an amino acid sequence that corresponds to the naturally-occurring protein due to degeneracy of the genetic code. Non-naturally-occurring variants and homologs include polypeptides and nucleic acids that comprise a change in amino acid or nucleotide sequence, respectively, where the change in sequence is artificially introduced (e.g., muteins); for example, the change is generated in the laboratory by human intervention (“hand of man”). Therefore, non-naturally occurring variants and homologs may also refer to those that differ from the naturally-occurring sequences by one or more conservative substitutions and/or tags and/or conjugates.


[0169] The term “muteins” as used herein refers broadly to mutated recombinant proteins. These proteins usually carry single or multiple amino acid substitutions and are frequently derived from cloned genes that have been subjected to site-directed or random mutagenesis, or from completely synthetic genes.


II. Compounds

[0170] In an aspect provided herein, is a compound of Formula (I):



, or a pharmaceutically acceptable salt thereof. R¹ and R² are independently ethyl or n-propyl. R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R⁴ is hydrogen,

Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl, ,





R^a and R^b are independently hydrogen, C_1 - C_3 alkyl, $H_2NCH_2CH_2-$, $(CH_3)_3COC(O)NHCH_2CH_2-$, or $CH_3CH_2CH_2NHCH_2CH_2-$. In embodiments, the inhibitor forms an irreversible bond with the amino acid residue. In embodiments, the bond is a covalent bond.


[0171] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0172] In certain embodiments, R^3 is Cl, Br, or methyl.


[0173] In certain embodiments, R^4 is Br, R^aR^bN- , pyridin-4-yl, morpholin-4-yl, or



[0174] In certain embodiments, R^4 is morpholin-4-yl or




[0175] In certain embodiments, R^4 is hydrogen, Br, R^aR^bN- and R^a and R^b are independently C_1 - C_3 alkyl.




[0176] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0177] In certain embodiments, the compound is



, or a pharmaceutically acceptable salt thereof.

[0178] In certain embodiments, the compound is



, or a pharmaceutically acceptable salt thereof.

[0179] In one aspect described herein, there is a method of treating a subject in need thereof comprising administering an astressin or a pharmaceutically acceptable salt or solvate thereof. An astressin generally refers to a nonselective corticotropin releasing hormone antagonist that reduces the synthesis of ACTH and cortisol.

[0180] In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating congenital adrenal hyperplasia (CAH) in a subject in need thereof, the method comprising: (i) measuring a hormone level in the subject in need thereof; (ii) administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof; and repeating steps (i) and (ii) until the hormone level reaches a pre-determined range followed by a maintenance therapy of a daily dosing of said astressin.

[0181] In one aspect described herein, there is a method of improving hyperandrogenic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of treating menstrual irregularity, ovulatory dysfunction or infertility, in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a method of improving metabolic symptoms in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto; or a pharmaceutically acceptable salt or solvate thereof.

[0182] In one aspect described herein, there is a method of improving quality of life in a subject in need thereof, comprising administering an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

III. Pharmaceutical Compositions

[0183] In an aspect, there is provided a pharmaceutical composition, including a compound as described herein, including embodiments, or the structural Formula (I), and a pharmaceutically acceptable excipient.

[0184] The compounds (e.g., CRF₁ inhibitors/antagonists) of the present disclosure may be in the form of compositions suitable for administration to a subject. In general, such compositions are “pharmaceutical compositions” comprising a compound (e.g., CRF₁ inhibitor(s)/antagonist(s)) and one or more pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients. In certain embodiments, the compound (e.g., CRF₁ inhibitor/antagonist) are present in a therapeutically acceptable amount. The pharmaceutical compositions may be used in the methods of the present disclosure; thus, for example, the pharmaceutical compositions can be administered *ex vivo* or *in vivo* to a subject in order to practice the therapeutic and prophylactic methods and uses described herein.

[0185] In one aspect described herein, there is a pharmaceutical composition in the form of a capsule comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof. In one aspect described herein, there is a pharmaceutical composition in the form of a tablet comprising an astressin or a variant having at least about 70% identity thereto, or a pharmaceutically acceptable salt or solvate thereof.

[0186] The pharmaceutical compositions of the present disclosure can be formulated to be compatible with the intended method or route of administration; exemplary routes of administration are set forth herein.

[0187] The pharmaceutical compositions containing the active ingredient (e.g., an inhibitor of CRF₁ function) may be in a form suitable for oral use, for example, as tablets, capsules, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups, solutions, microbeads or elixirs. Pharmaceutical compositions intended for oral use may be prepared for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents such as, for example, sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets, capsules and the like contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture thereof. These excipients may be, for example, diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc.

[0188] The tablets, capsules and the like suitable for oral administration may be uncoated or coated to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action. For example, a time-delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated to form osmotic therapeutic tablets for

controlled release. Additional agents include biodegradable or biocompatible particles or a polymeric substance such as polyesters, polyamine acids, hydrogel, polyvinyl pyrrolidone, polyanhydrides, polyglycolic acid, ethylene-vinylacetate, methylcellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymers, polylactide/glycolide copolymers, or ethylenevinylacetate copolymers in order to control delivery of an administered composition. For example, the oral agent can be entrapped in microcapsules prepared by coacervation techniques or by interfacial polymerization, by the use of hydroxymethylcellulose or gelatin-microcapsules or poly(methylmethacrolate) microcapsules, respectively, or in a colloid drug delivery system. Colloidal dispersion systems include macromolecule complexes, nano-capsules, microspheres, microbeads, and lipid-based systems, including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Methods for the preparation of the above-mentioned formulations will be apparent to those skilled in the art.

[0189] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate, kaolin or microcrystalline cellulose, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

[0190] Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture thereof. Such excipients can be suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents, for example a naturally-occurring phosphatide (e.g., lecithin), or condensation products of an alkylene oxide with fatty acids (e.g., polyoxy-ethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols (e.g., for heptadecaethyleneoxycetanol), or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol (e.g., polyoxyethylene sorbitol monooleate), or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides (e.g., polyethylene sorbitan monooleate). The aqueous suspensions may also contain one or more preservatives.

[0191] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

[0192] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting

agent, and optionally one or more suspending agents and/or preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified herein.

[0193] The pharmaceutical compositions of the present disclosure may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example, liquid paraffin, or mixtures of these. Suitable emulsifying agents may be naturally occurring gums, for example, gum acacia or gum tragacanth; naturally occurring phosphatides, for example, soy bean, lecithin, and esters or partial esters derived from fatty acids; hexitol anhydrides, for example, sorbitan monooleate; and condensation products of partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate.

[0194] The pharmaceutical compositions typically comprise a therapeutically effective amount of a CRF₁ inhibitor/antagonist contemplated by the present disclosure and one or more pharmaceutically and physiologically acceptable formulation agents. Suitable pharmaceutically acceptable or physiologically acceptable diluents, carriers or excipients include, but are not limited to, antioxidants (e.g., ascorbic acid and sodium bisulfate), preservatives (e.g., benzyl alcohol, methyl parabens, ethyl or n-propyl, p-hydroxybenzoate), emulsifying agents, suspending agents, dispersing agents, solvents, fillers, bulking agents, detergents, buffers, vehicles, diluents, and/or adjuvants. For example, a suitable vehicle may be physiological saline solution or citrate-buffered saline, possibly supplemented with other materials common in pharmaceutical compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Those skilled in the art will readily recognize a variety of buffers that can be used in the pharmaceutical compositions and dosage forms contemplated herein. Typical buffers include, but are not limited to, pharmaceutically acceptable weak acids, weak bases, or mixtures thereof. As an example, the buffer components can be water soluble materials such as phosphoric acid, tartaric acids, lactic acid, succinic acid, citric acid, acetic acid, ascorbic acid, aspartic acid, glutamic acid, and salts thereof. Acceptable buffering agents include, for example, a Tris buffer; N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) (HEPES); 2-(N-Morpholino)ethanesulfonic acid (MES); 2-(N-Morpholino)ethanesulfonic acid sodium salt (MES); 3-(N-Morpholino)propanesulfonic acid (MOPS); and N-tris[Hydroxymethyl]methyl-3-aminopropanesulfonic acid (TAPS).

[0195] After a pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form, a lyophilized form requiring reconstitution prior to use, a liquid form requiring dilution prior to use, or other acceptable form. In some embodiments, the pharmaceutical composition is provided in a single-use container

(e.g., a single-use vial, ampule, syringe, or autoinjector (similar to, e.g., an EpiPen®)), whereas a multi-use container (e.g., a multi-use vial) is provided in other embodiments.

[0196] Formulations can also include carriers to protect the composition against rapid degradation or elimination from the body, such as a controlled release formulation, including liposomes, hydrogels, prodrugs and microencapsulated delivery systems. For example, a time-delay material such as glyceryl monostearate or glyceryl stearate alone, or in combination with a wax, may be employed. Any drug delivery apparatus may be used to deliver a CRF₁ inhibitor/antagonist, including implants (e.g., implantable pumps) and catheter systems, slow injection pumps and devices, all of which are well known to the skilled artisan.

[0197] Depot injections, which are generally administered subcutaneously or intramuscularly, may also be utilized to release the compound (e.g., CRF₁ inhibitor/antagonist) disclosed herein over a defined period of time. Depot injections are usually either solid- or oil-based and generally comprise at least one of the formulation components set forth herein. One of ordinary skill in the art is familiar with possible formulations and uses of depot injections.

[0198] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents mentioned herein. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Acceptable diluents, solvents and dispersion media that may be employed include water, Ringer's solution, isotonic sodium chloride solution, Cremophor® EL (BASF, Parsippany, NJ) or phosphate buffered saline (PBS), ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. In addition, sterile fixed oils are conventionally employed as a solvent or suspending medium; for this purpose, any bland fixed oil may be employed, including synthetic mono- or diglycerides. Moreover, fatty acids, such as oleic acid, find use in the preparation of injectables. Prolonged absorption of particular injectable formulations can be achieved by including an agent that delays absorption (e.g., aluminum monostearate or gelatin).

[0199] The present disclosure contemplates the administration of the compound (e.g., CRF₁ inhibitor/antagonist) in the form of suppositories for rectal administration. The suppositories can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include, but are not limited to, cocoa butter and polyethylene glycols.

[0200] The compound (e.g., CRF₁ inhibitor/antagonist) contemplated by the present disclosure may be in the form of any other suitable pharmaceutical composition (e.g., sprays for nasal or inhalation use) currently known or developed in the future.

IV. Methods

[0201] In certain embodiments, provided herein are methods for treating or preventing a disease comprising administering an effective amount of a compound of Formula (I), or an isotopic variant thereof or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0202] In one embodiment, provided herein are methods for treating or preventing a tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a benign tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing a malignant tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing an adrenal rest tumor, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0203] In one embodiment, provided herein are methods for treating or preventing infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for treating or preventing male infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm count, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm motility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are methods for improving or increasing sperm morphology, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In one embodiment, provided herein are

methods for treating or preventing female infertility, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof.

[0204] In one embodiment, provided herein are methods for treating or preventing a proliferative or hyperproliferative disease or disorder, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, the proliferative disease is cancer. In certain embodiments, the proliferative disease is a cancer of the breast, skin, prostate, cervix, uterus, ovary, testes, bladder, lung, liver, larynx, oral cavity, colon and gastrointestinal tract (e.g., esophagus, stomach, pancreas), brain, thyroid, blood, and lymphatic system.

[0205] In certain embodiments, the cancer treatable with the methods provided herein includes, but is not limited to, (1) leukemias, including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome or a symptom thereof (such as anemia, thrombocytopenia, neutropenia, bicytopenia or pancytopenia), refractory anemia (RA), RA with ringed sideroblasts (RARS), RA with excess blasts (RAEB), RAEB in transformation (RAEB-T), preleukemia, and chronic myelomonocytic leukemia (CMML), (2) chronic leukemias, including, but not limited to, chronic myelocytic (granulocytic) leukemia, chronic lymphocytic leukemia, and hairy cell leukemia; (3) polycythemia vera; (4) lymphomas, including, but not limited to, Hodgkin's disease and non-Hodgkin's disease; (5) multiple myelomas, including, but not limited to, smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma, and extramedullary plasmacytoma; (6) Waldenstrom's macroglobulinemia; (7) monoclonal gammopathy of undetermined significance; (8) benign monoclonal gammopathy; (9) heavy chain disease; (10) bone and connective tissue sarcomas, including, but not limited to, bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, metastatic cancers, neurilemmoma, rhabdomyosarcoma, and synovial sarcoma; (11) brain tumors, including, but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendroglioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, and primary brain lymphoma; (12) breast cancer, including, but not limited to, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, primary cancers, Paget's disease, and

inflammatory breast cancer; (13) adrenal cancer, including, but not limited to, pheochromocytom and adrenocortical carcinoma; (14) thyroid cancer, including, but not limited to, papillary or follicular thyroid cancer, medullary thyroid cancer, and anaplastic thyroid cancer; (15) pancreatic cancer, including, but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatin-secreting tumor, and carcinoid or islet cell tumor; (16) pituitary cancer, including, but limited to, Cushing's disease, prol actin-secreting tumor, acromegaly, and diabetes insipius; (17) eye cancer, including, but not limited, to ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; (18) vaginal cancer, including, but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; (19) vulvar cancer, including, but not limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; (20) cervical cancers, including, but not limited to, squamous cell carcinoma, and adenocarcinoma; (21) uterine cancer, including, but not limited to, endometrial carcinoma and uterine sarcoma; (22) ovarian cancer, including, but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; (23) esophageal cancer, including, but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; (24) stomach cancer, including, but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; (25) colon cancer; (26) rectal cancer; (27) liver cancer, including, but not limited to, hepatocellular carcinoma and hepatoblastoma; (28) gallbladder cancer, including, but not limited to, adenocarcinoma; (29) cholangiocarcinomas, including, but not limited to, pappillary, nodular, and diffuse; (30) lung cancer, including, but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma, and small-cell lung cancer; (31) testicular cancer, including, but not limited to, germinal tumor, seminoma, anaplastic, classic (typical), spermatocytic, nonseminoma, embryonal carcinoma, teratoma carcinoma, and choriocarcinoma (yolk-sac tumor); (32) prostate cancer, including, but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; (33) penal cancer; (34) oral cancer, including, but not limited to, squamous cell carcinoma; (35) basal cancer; (36) salivary gland cancer, including, but not limited to, adenocarcinoma, mucoepidermoid carcinoma, and adenoidcystic carcinoma; (37) pharynx cancer, including, but not limited to, squamous cell cancer and verrucous; (38) skin cancer, including, but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, and acral lentiginous melanoma; (39) kidney cancer, including, but not limited to, renal cell

cancer, adenocarcinoma, hypernephroma, fibrosarcoma, and transitional cell cancer (renal pelvis and/or uterine); (40) Wilms' tumor; (41) bladder cancer, including, but not limited to, transitional cell carcinoma, squamous cell cancer, adenocarcinoma, and carcinosarcoma; and other cancer, including, not limited to, myxosarcoma, osteogenic sarcoma, endotheliosarcoma, lymphangio-endotheliosarcoma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, and papillary adenocarcinomas (*See Fishman et al.*, 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and *Murphy et al.*, 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

[0206] Depending on the disorder, disease, or condition to be treated, and the subject's condition, the compounds or pharmaceutical compositions provided herein can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), inhalation, nasal, vaginal, rectal, sublingual, or topical (e.g., transdermal or local) routes of administration and can be formulated, alone or together, in suitable dosage unit with pharmaceutically acceptable excipients, carriers, adjuvants, and vehicles appropriate for each route of administration as described elsewhere herein.

[0207] The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. Recommended routes of administration for the second active agents are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).


[0208] In one embodiment, provided herein are methods for treating or preventing testicular adrenal rest tumors (TART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).

[0209] In one embodiment, provided herein are methods for treating or preventing ovarian adrenal rest tumors (OART) in a subject, comprising administering a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof, wherein the subject has congenital adrenal hyperplasia (CAH).


[0210] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0211] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a corticotropin-releasing factor type-1 (CRF₁) antagonist or a pharmaceutically acceptable salt thereof.

[0212] In an aspect provided herein, is a method of treating or preventing testicular adrenal rest tumors (TART), comprising administering to a subject in need thereof a compound of structural Formula (I):




, or a pharmaceutically acceptable salt thereof. R¹ and R² are independently ethyl or n-propyl. R³ is hydrogen, Cl, Br, methyl, trifluoromethyl, or methoxy. R⁴ is hydrogen,




Br, R^aR^bN-, methoxymethyl, n-butyl, acetamido, pyridin-4-yl, morpholin-4-yl, R^a and R^b are independently hydrogen, C₁-C₃alkyl, H₂NCH₂CH₂-, (CH₃)₃COC(O)NHCH₂CH₂-, or CH₃CH₂CH₂NHCH₂CH₂-.

[0213] In an aspect provided herein, is a method of treating or preventing ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R¹, R², R³ and R⁴ are as described herein, including embodiments.


[0214] In an aspect provided herein, is a method of decreasing or reducing adrenocorticotrophic hormone (ACTH) levels in a subject in need thereof, comprising administering to the subject contacting the ACTH with a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 60% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 70% of the subjects demonstrate reduction in ACTH. In certain embodiments, at least about 80% of the subjects demonstrate reduction in ACTH. In embodiments, at least about 10% of subjects were in the standard or normal range for ACTH after treatment.

[0215] In embodiments, at least about 20% of subjects were in the standard or normal range for ACTH after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for ACTH after treatment. In certain embodiments, ACTH levels are decreased or reduced within about 30 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 1 hour post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 90 minutes post dosing. In certain embodiments, ACTH levels are decreased or reduced within about 2 hours post dosing.


[0216] In an aspect provided herein, is a method of decreasing or reducing 17 α -hydroxyprogesterone (17-OHP) levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. R^1 , R^2 , R^3 and R^4 are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 60% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 70% of the subjects demonstrate reduction in 17-OHP. In certain embodiments, at least about 80% of the subjects demonstrate reduction in 17-OHP. In embodiments, at least about 30% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 40% of subjects were in the standard

or normal range of 1200 ng/dL after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 60% of subjects were in the standard or normal range of 1200 ng/dL after treatment. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment. In embodiments, at least about 60% of subjects demonstrated an at least about 25% reduction in 17-OHP after treatment.

[0217] In an aspect provided herein, is a method of decreasing or reducing androstenedione levels in a subject in need thereof, comprising administering to the subject a compound of structural Formula (I):





, or a pharmaceutically acceptable salt thereof. R¹, R², R³ and R⁴ are as described herein, including embodiments. R¹, R², R³ and R⁴ are as described herein, including embodiments. In certain embodiments, at least about 50% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 60% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 70% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 80% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least about 90% of the subjects demonstrate reduction in androstenedione. In certain embodiments, at least 100% of the subjects demonstrate reduction in androstenedione. In embodiments, at least about 30% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 40% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 50% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least 60% of subjects demonstrated an at least about 25% reduction in androstenedione after treatment. In embodiments, at least about 30% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 40% of subjects were in the standard or normal range for androstenedione after treatment. In embodiments, at least about 50% of subjects were in the standard or normal range for androstenedione after treatment.


[0218] In certain embodiments, R^3 is Cl, Br, methyl, or trifluoromethyl.

[0219] In certain embodiments, R^3 is Cl, Br, or methyl.


[0220] In certain embodiments, R^4 is Br, $R^a R^b N-$, pyridin-4-yl, morpholin-4-yl, or



[0221] In certain embodiments, R^4 is morpholin-4-yl or




[0222] In certain embodiments, R^4 is hydrogen, Br, $R^a R^b N-$ and R^a and R^b are independently C_1 - C_3 alkyl.




[0223] In certain embodiments, the compound is pharmaceutically acceptable salt thereof.

[0224] In certain embodiments, the compound is



, or a pharmaceutically acceptable salt thereof.

[0225] In certain embodiments, the compound is



, or a pharmaceutically acceptable salt thereof.

[0226] In certain embodiments, a compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is cyclically administered

to a patient. Cycling therapy involves the administration of an active agent or a combination of active agents for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

[0227] In certain embodiments, in the treatment, prevention, or amelioration of one or more symptoms of the disorders, diseases, or conditions described herein (e.g., treatment or prevention of testicular adrenal rest tumors (TART) or treatment or prevention of ovarian adrenal rest tumors (OART)), an appropriate dosage level of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof generally is ranging from about 1 mg to 3000 mg, from about 1 mg to 2000 mg, from about 1 mg to 1000 mg, from about 1 mg to about 500 mg, from about 5 mg to about 500 mg, from about 5 mg to about 400 mg, from about 5 mg to about 300 mg, from about 5 mg to about 250 mg, from about 5 mg to about 200 mg, from about 50 mg to about 200 mg, from about 10 mg to about 400 mg, from about 5 mg to about 25 mg, from about 10 mg to about 25 mg, or from about 25 to about 200 mg, which can be administered in single or multiple doses. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55

mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 450 mg, about 500 mg, about 600 mg, about 800 mg, or about 1000 mg/day.

[0228] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg. In certain embodiments, the compound of Formula (I), or an isotopic

variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg.

[0229] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, or about 1000 mg.

[0230] For oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing from about 1.0 mg to about 1,000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In certain embodiments, for oral administration, the

pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 1 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg, about 100 mg, about 105 mg, about 110 mg, about 115 mg, about 120 mg, about 125 mg, about 130 mg, about 135 mg, about 140 mg, about 145 mg, about 150 mg, about 155 mg, about 160 mg, about 165 mg, about 170 mg, about 175 mg, about 180 mg, about 185 mg, about 190 mg, about 195 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 980 mg, about 990 mg, about 990 mg, or about 1000 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof for the symptomatic adjustment of the dosage to the patient to be treated. In certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 50 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. In

certain embodiments, for oral administration, the pharmaceutical compositions provided herein can be formulated in the form of tablets or capsules containing about 200 mg of a compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof. The pharmaceutical compositions can be administered on a regimen of one (1) to four (4) times per day, including once, twice, three times, and four times per day. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 810 mg, about 820 mg, about 830 mg, about 840 mg, about 850 mg, about 860 mg, about 870 mg, about 880 mg, about 890 mg, about 900 mg, about 910 mg, about 920 mg, about 930 mg, about 940 mg, about 950 mg, about 960 mg, about 970 mg, about 980 mg, about 990 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered once per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg,

about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg, about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 405 mg, about 410 mg, about 415 mg, about 420 mg, about 425 mg, about 430 mg, about 435 mg, about 440 mg, about 445 mg, about 450 mg, about 455 mg, about 460 mg, about 465 mg, about 470 mg, about 475 mg, about 480 mg, about 485 mg, about 490 mg, about 495 mg, about 500 mg, about 505 mg, about 510 mg, about 515 mg, about 520 mg, about 525 mg, about 530 mg, about 535 mg, about 540 mg, about 545 mg, about 550 mg, about 555 mg, about 560 mg, about 565 mg, about 570 mg, about 575 mg, about 580 mg, about 585 mg, about 590 mg, about 595 mg, about 600 mg, about 605 mg,

about 610 mg, about 615 mg, about 620 mg, about 625 mg, about 630 mg, about 635 mg, about 640 mg, about 645 mg, about 650 mg, about 655 mg, about 660 mg, about 665 mg, about 670 mg, about 675 mg, about 680 mg, about 685 mg, about 690 mg, about 695 mg, about 700 mg, about 705 mg, about 710 mg, about 715 mg, about 720 mg, about 725 mg, about 730 mg, about 735 mg, about 740 mg, about 745 mg, about 750 mg, about 755 mg, about 760 mg, about 765 mg, about 770 mg, about 775 mg, about 780 mg, about 785 mg, about 790 mg, about 795 mg, about 800 mg, about 850 mg, about 900 mg, about 950 mg, or about 1000 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered three times per day.

[0231] In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of one day. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of two days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of three days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of four days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of five days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of six days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of seven days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of eight days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of nine days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of ten days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 11 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 12 days. In

some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 13 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 14 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of 15 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 16 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 17 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 18 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 19 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 20 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 21 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 22 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 23 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 24 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 25 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 26 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 27 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 28 days. In some embodiments, the

compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 29 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered for a period of 30 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 31 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 32 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 33 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 34 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 35 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 36 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 37 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 38 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 39 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 40 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 41 days. In some embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered over a period of 42 days or more.

[0232] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days.

[0233] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is

administered in an amount of about 260 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 14 days. In certain embodiments, the

compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 14 days.

[0234] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug

thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 14 days. In certain

embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 140 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or

prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 14 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 14 days.

[0235] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 28 days.

In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt,

solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 28

days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 28 days.

[0236] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof, or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 280 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at

least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically

acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 28 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 28 days.

[0237] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 255 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg once daily for

a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a

pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg once daily for

a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg once daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg once daily for a period of up to at least about 42 days.

[0238] In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 5 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 10 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 15 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 20 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 25 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 50 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 100 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 150 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 200 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 210 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 220 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 230 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 240 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 250 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 260 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 270 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 420 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 290 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 300 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 310 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 320 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 330

mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 340 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 350 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 360 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 370 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 380 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 390 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 400 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 450 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 500 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 600 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 700 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 800 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant

thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 900 mg twice daily for a period of up to at least about 42 days. In certain embodiments, the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered in an amount of about 1000 mg twice daily for a period of up to at least about 42 days.

[0239] In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, or about 400 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered twice per day. In some embodiments, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 700 mg, or about 800 mg of the compound of Formula (I), or an isotopic variant thereof; or a pharmaceutically acceptable salt, solvate, hydrate, or prodrug thereof is administered at least once per day.

[0240] In certain embodiments, from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0241] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered orally.

[0242] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is formulated as a capsule or tablet.

[0243] In certain embodiments, from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0244] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the compound of


Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0245] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fed state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the fasted state. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at least once per day.

[0246] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at bedtime. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 6 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 5 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 4 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 3 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 2 hours before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 1 hour before sleep. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered less than about 30 minutes before sleep.

[0247] In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered in the evening. In certain embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered at about 10 pm at night.

[0248] In another aspect is provided a method of treating or preventing testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART), comprising administering to a subject in need thereof a pharmaceutical composition, comprising a compound of structural Formula (I): and a pharmaceutically acceptable excipient comprising with a compound as described herein, including embodiments, or the structural Formulae (I) and a pharmaceutically acceptable excipient:



, or a pharmaceutically acceptable salt thereof. R^1 , R^2 , R^3 , and R^4 are as described herein, including embodiments.

[0249] In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the pharmaceutical composition comprises from about 100 mg to about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0250] In certain embodiments, the pharmaceutical composition comprises about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360 mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, about 400 mg, about 500 mg, or about 600 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0251] In certain embodiments, the pharmaceutical composition is administered orally.

[0252] In certain embodiments, the pharmaceutical composition is formulated as a capsule or tablet. In certain embodiments, the capsule or tablet comprises about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 205 mg, about 210 mg, about 215 mg, about 220 mg, about 225 mg, about 230 mg, about 235 mg, about 240 mg, about 245 mg, about 250 mg, about 255 mg, about 260 mg, about 265 mg, about 270 mg, about 275 mg, about 280 mg, about 285 mg, about 290 mg, about 295 mg, about 300 mg, about 305 mg, about 310 mg, about 315 mg, about 320 mg, about 325 mg, about 330 mg, about 335 mg, about 340 mg, about 345 mg, about 350 mg, about 355 mg, about 360

mg, about 365 mg, about 370 mg, about 375 mg, about 380 mg, about 385 mg, about 390 mg, about 395 mg, or about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 50 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In certain embodiments, the capsule or tablet comprises about 200 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

[0253] In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg to about 1000 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg to about 25 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg to about 400 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the subject.

[0254] In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 25 mg/day to about 1000 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 5 mg/day to about 25 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 10 mg/day to about 400 mg/day is administered to the subject. In certain embodiments, the pharmaceutical composition is administered at a dose of from about 100 mg/day to about 600 mg/day.

[0255] In certain embodiments, the pharmaceutical composition is administered in the fed state. In certain embodiments, the pharmaceutical composition is administered in the fasted state. In certain embodiments, the pharmaceutical composition is administered at least once per day.

[0256] In certain embodiments, the pharmaceutical composition is administered at bedtime. In certain embodiments, the pharmaceutical composition is administered less than about 6 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 5 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 4 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 3 hours before sleep. In certain embodiments, the

pharmaceutical composition is administered less than about 2 hours before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 1 hour before sleep. In certain embodiments, the pharmaceutical composition is administered less than about 30 minutes before sleep.


[0257] In certain embodiments, the pharmaceutical composition is administered in the evening. In certain embodiments, the pharmaceutical composition is administered at about 10 pm at night.


[0258] In certain embodiments, the subject has congenital adrenal hyperplasia (CAH).

[0259] In certain embodiments, the size and/or number of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced. In certain embodiments, the size of the tumors is decreased or reduced.


[0260] In certain embodiments of the methods disclosed herein, the method further comprises administering to the subject an additional chemotherapeutic agent.

[0261] In certain embodiments, the the additional chemotherapeutic agent is a glucocorticoid, a mineralocorticoid, an ACAT1 inhibitor, or an anti-androgen. In certain embodiments, the glucocorticoid is beclomethasone, betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, or triamcinolone. In certain embodiments, the mineralocorticoid is fludrocortisone. In certain embodiments, the additional chemotherapeutic agent is another CRF₁ antagonist. In certain embodiments, the CRF₁ antagonist is:





, or a pharmaceutically acceptable salt thereof. In certain embodiments, the



CRF₁ antagonist is

, or a pharmaceutically acceptable salt thereof.

[0262] In certain embodiments of the methods disclosed herein, the method further comprises an additional treatment selected from surgical resection of the tumors and radiation therapy, or a combination thereof. In certain embodiments, the additional therapy is surgical resection and the surgical resection is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the surgical resection is prior to administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the surgical resection is concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0263] In certain embodiments, the additional treatment is radiation therapy and the radiation therapy is prior to, after, and/or concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition, or any combination thereof. In certain embodiments, the radiation therapy is prior to administration of the CRF₁ antagonist of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is after administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition. In certain embodiments, the radiation therapy is

concurrent with administration of the compound of structural Formula (I), or a pharmaceutically acceptable salt thereof or the pharmaceutical composition.

[0264] In one embodiment, the pharmaceutical compositions are provided in a dosage form for oral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. The pharmaceutical compositions provided herein that are formulated for oral administration may be in tablet, capsule, powder, or liquid form. In some embodiments, a tablet comprises a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil, or synthetic oil. Physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol, or polyethylene glycol may be included. In some embodiments, a capsule comprises a solid carrier such as gelatin.

[0265] In another embodiment, the pharmaceutical compositions are provided in a dosage form for parenteral administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers. Where pharmaceutical compositions may be formulated for intravenous, cutaneous or subcutaneous injection, the active ingredient will be in the form of a parenterally acceptable aqueous solution, which is pyrogen-free and has a suitable pH, isotonicity, and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles, such as Sodium Chloride injection, Ringer's injection, or Lactated Ringer's injection. In some embodiments, preservatives, stabilisers, buffers, antioxidants, and/or other additives are included as required.

[0266] In yet another embodiment, the pharmaceutical compositions are provided in a dosage form for topical administration, which comprise a compound provided herein, and one or more pharmaceutically acceptable excipients or carriers.

[0267] The pharmaceutical compositions can also be formulated as modified release dosage forms, including delayed-, extended-, prolonged-, sustained-, pulsatile-, controlled-, accelerated-, fast-, targeted-, and programmed-release, and gastric retention dosage forms. These dosage forms can be prepared according to conventional methods and techniques known to those skilled in the art (*see, Remington: The Science and Practice of Pharmacy, supra; Modified-Release Drug Delivery Technology*, 2nd Edition, Rathbone *et al.*, Eds., Marcel Dekker, Inc.: New York, NY, 2008).

[0268] The pharmaceutical compositions provided herein can be provided in a unit-dosage form or multiple-dosage form. A unit-dosage form, as used herein, refers to physically discrete a unit suitable for administration to a human and animal subject, and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of an active ingredient(s)

sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carriers or excipients. Examples of a unit-dosage form include an ampoule, syringe, and individually packaged tablet and capsule. A unit-dosage form may be administered in fractions or multiples thereof. A multiple-dosage form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dosage form. Examples of a multiple-dosage form include a vial, bottle of tablets or capsules, or bottle of pints or gallons.

[0269] The pharmaceutical compositions provided herein can be administered at once, or multiple times at intervals of time. It is understood that the precise dosage and duration of treatment may vary with the age, weight, and condition of the patient being treated, and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test or diagnostic data. It is further understood that for any particular individual, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations.

[0270] In certain embodiments, the pharmaceutical compositions provided herein further comprise one or more chemotherapeutic agents as defined herein.

A. Oral Administration

[0271] The pharmaceutical compositions provided herein for oral administration can be provided in solid, semisolid, or liquid dosage forms for oral administration. As used herein, oral administration also includes buccal, lingual, and sublingual administration. Suitable oral dosage forms include, but are not limited to, tablets, fastmelts, chewable tablets, capsules, pills, strips, troches, lozenges, pastilles, cachets, pellets, medicated chewing gum, bulk powders, effervescent or non-effervescent powders or granules, oral mists, solutions, emulsions, suspensions, wafers, sprinkles, elixirs, and syrups. In addition to the active ingredient(s), the pharmaceutical compositions can contain one or more pharmaceutically acceptable carriers or excipients, including, but not limited to, binders, fillers, diluents, disintegrants, wetting agents, lubricants, glidants, coloring agents, dye-migration inhibitors, sweetening agents, flavoring agents, emulsifying agents, suspending and dispersing agents, preservatives, solvents, non-aqueous liquids, organic acids, and sources of carbon dioxide.

[0272] Binders or granulators impart cohesiveness to a tablet to ensure the tablet remaining intact after compression. Suitable binders or granulators include, but are not limited to, starches, such as corn starch, potato starch, and pre-gelatinized starch (e.g., STARCH 1500); gelatin; sugars, such as sucrose, glucose, dextrose, molasses, and lactose; natural and synthetic gums, such as acacia, alginic acid, alginates, extract of Irish moss, panwar gum, ghatti gum, mucilage

of isabgol husks, carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone (PVP), Veegum, larch arabogalactan, powdered tragacanth, and guar gum; celluloses, such as ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose, methyl cellulose, hydroxyethylcellulose (HEC), hydroxypropylcellulose (HPC), hydroxypropyl methyl cellulose (HPMC); microcrystalline celluloses, such as AVICEL-PH-101, AVICEL-PH-103, AVICEL RC-581, AVICEL-PH-105 (FMC Corp., Marcus Hook, PA); and mixtures thereof. Suitable fillers include, but are not limited to, talc, calcium carbonate, microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The amount of a binder or filler in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The binder or filler may be present from about 50 to about 99% by weight in the pharmaceutical compositions provided herein.

[0273] Suitable diluents include, but are not limited to, dicalcium phosphate, calcium sulfate, lactose, sorbitol, sucrose, inositol, cellulose, kaolin, mannitol, sodium chloride, dry starch, and powdered sugar. Certain diluents, such as mannitol, lactose, sorbitol, sucrose, and inositol, when present in sufficient quantity, can impart properties to some compressed tablets that permit disintegration in the mouth by chewing. Such compressed tablets can be used as chewable tablets. The amount of a diluent in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art.

[0274] Suitable disintegrants include, but are not limited to, agar; bentonite; celluloses, such as methylcellulose and carboxymethylcellulose; wood products; natural sponge; cation-exchange resins; alginic acid; gums, such as guar gum and Veegum HV; citrus pulp; cross-linked celluloses, such as croscarmellose; cross-linked polymers, such as crospovidone; cross-linked starches; calcium carbonate; microcrystalline cellulose, such as sodium starch glycolate; polacrillin potassium; starches, such as corn starch, potato starch, tapioca starch, and pre-gelatinized starch; clays; aligns; and mixtures thereof. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The amount of a disintegrant in the pharmaceutical compositions provided herein varies upon the type of formulation, and is readily discernible to those of ordinary skill in the art. The pharmaceutical compositions provided herein may contain from about 0.5 to about 15% or from about 1 to about 5% by weight of a disintegrant.

[0275] Suitable lubricants include, but are not limited to, calcium stearate; magnesium stearate; mineral oil; light mineral oil; glycerin; sorbitol; mannitol; glycols, such as glycerol behenate and polyethylene glycol (PEG); stearic acid; sodium lauryl sulfate; talc; hydrogenated vegetable oil,

including peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil; zinc stearate; ethyl oleate; ethyl laureate; agar; starch; lycopodium; silica or silica gels, such as AEROSIL® 200 (W.R. Grace Co., Baltimore, MD) and CAB-O-SIL® (Cabot Co. of Boston, MA); and mixtures thereof. The pharmaceutical compositions provided herein may contain about 0.1 to about 5% by weight of a lubricant.

[0276] Suitable glidants include, but are not limited to, colloidal silicon dioxide, CAB-O-SIL® (Cabot Co. of Boston, MA), and asbestos-free talc. Suitable coloring agents include, but are not limited to, any of the approved, certified, water soluble FD&C dyes, and water insoluble FD&C dyes suspended on alumina hydrate, and color lakes and mixtures thereof. A color lake is the combination by adsorption of a water-soluble dye to a hydrous oxide of a heavy metal, resulting in an insoluble form of the dye. Suitable flavoring agents include, but are not limited to, natural flavors extracted from plants, such as fruits, and synthetic blends of compounds which produce a pleasant taste sensation, such as peppermint and methyl salicylate. Suitable sweetening agents include, but are not limited to, sucrose, lactose, mannitol, syrups, glycerin, and artificial sweeteners, such as saccharin and aspartame. Suitable emulsifying agents include, but are not limited to, gelatin, acacia, tragacanth, bentonite, and surfactants, such as polyoxyethylene sorbitan monooleate (TWEEN® 20), polyoxyethylene sorbitan monooleate 80 (TWEEN® 80), and triethanolamine oleate. Suitable suspending and dispersing agents include, but are not limited to, sodium carboxymethylcellulose, pectin, tragacanth, Veegum, acacia, sodium carbomethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable preservatives include, but are not limited to, glycerin, methyl and propylparaben, benzoic acid, sodium benzoate and alcohol. Suitable wetting agents include, but are not limited to, propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate, and polyoxyethylene lauryl ether. Suitable solvents include, but are not limited to, glycerin, sorbitol, ethyl alcohol, and syrup. Suitable non-aqueous liquids utilized in emulsions include, but are not limited to, mineral oil and cottonseed oil. Suitable organic acids include, but are not limited to, citric and tartaric acid. Suitable sources of carbon dioxide include, but are not limited to, sodium bicarbonate and sodium carbonate.

[0277] It should be understood that many carriers and excipients may serve several functions, even within the same formulation.

[0278] The pharmaceutical compositions provided herein for oral administration can be provided as compressed tablets, tablet triturates, chewable lozenges, rapidly dissolving tablets, multiple compressed tablets, or enteric-coating tablets, sugar-coated, or film-coated tablets. Enteric-coated tablets are compressed tablets coated with substances that resist the action of stomach acid but dissolve or disintegrate in the intestine, thus protecting the active ingredients

from the acidic environment of the stomach. Enteric-coatings include, but are not limited to, fatty acids, fats, phenyl salicylate, waxes, shellac, ammoniated shellac, and cellulose acetate phthalates. Sugar-coated tablets are compressed tablets surrounded by a sugar coating, which may be beneficial in covering up objectionable tastes or odors and in protecting the tablets from oxidation. Film-coated tablets are compressed tablets that are covered with a thin layer or film of a water-soluble material. Film coatings include, but are not limited to, hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000, and cellulose acetate phthalate. Film coating imparts the same general characteristics as sugar coating. Multiple compressed tablets are compressed tablets made by more than one compression cycle, including layered tablets, and press-coated or dry-coated tablets.

[0279] The tablet dosage forms can be prepared from the active ingredient in powdered, crystalline, or granular forms, alone or in combination with one or more carriers or excipients described herein, including binders, disintegrants, controlled-release polymers, lubricants, diluents, and/or colorants. Flavoring and sweetening agents are especially useful in the formation of chewable tablets and lozenges.

[0280] The pharmaceutical compositions provided herein for oral administration can be provided as soft or hard capsules, which can be made from gelatin, methylcellulose, starch, or calcium alginate. The hard gelatin capsule, also known as the dry-filled capsule (DFC), consists of two sections, one slipping over the other, thus completely enclosing the active ingredient. The soft elastic capsule (SEC) is a soft, globular shell, such as a gelatin shell, which is plasticized by the addition of glycerin, sorbitol, or a similar polyol. The soft gelatin shells may contain a preservative to prevent the growth of microorganisms. Suitable preservatives are those as described herein, including methyl- and propyl-parabens, and sorbic acid. The liquid, semisolid, and solid dosage forms provided herein may be encapsulated in a capsule. Suitable liquid and semisolid dosage forms include solutions and suspensions in propylene carbonate, vegetable oils, or triglycerides. Capsules containing such solutions can be prepared as described in U.S. Pat. Nos. 4,328,245; 4,409,239; and 4,410,545. The capsules may also be coated as known by those of skill in the art in order to modify or sustain dissolution of the active ingredient.

[0281] Coloring and flavoring agents can be used in all of the above dosage forms.

[0282] The pharmaceutical compositions provided herein for oral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

B. Parenteral Administration

[0283] The pharmaceutical compositions provided herein can be administered parenterally by injection, infusion, or implantation, for local or systemic administration. Parenteral

administration, as used herein, include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular, intrasynovial, intravesical, and subcutaneous administration.

[0284] The pharmaceutical compositions provided herein for parenteral administration can be formulated in any dosage forms that are suitable for parenteral administration, including solutions, suspensions, emulsions, micelles, liposomes, microspheres, nanosystems, and solid forms suitable for solutions or suspensions in liquid prior to injection. Such dosage forms can be prepared according to conventional methods known to those skilled in the art of pharmaceutical science (*see, Remington: The Science and Practice of Pharmacy, supra*).

[0285] The pharmaceutical compositions intended for parenteral administration can include one or more pharmaceutically acceptable carriers and excipients, including, but not limited to, aqueous vehicles, water-miscible vehicles, non-aqueous vehicles, antimicrobial agents or preservatives against the growth of microorganisms, stabilizers, solubility enhancers, isotonic agents, buffering agents, antioxidants, local anesthetics, suspending and dispersing agents, wetting or emulsifying agents, complexing agents, sequestering or chelating agents, cryoprotectants, lyoprotectants, thickening agents, pH adjusting agents, and inert gases.

[0286] Suitable aqueous vehicles include, but are not limited to, water, saline, physiological saline or phosphate buffered saline (PBS), sodium chloride injection, Ringers injection, isotonic dextrose injection, sterile water injection, dextrose and lactated Ringers injection. Suitable non-aqueous vehicles include, but are not limited to, fixed oils of vegetable origin, castor oil, corn oil, cottonseed oil, olive oil, peanut oil, peppermint oil, safflower oil, sesame oil, soybean oil, hydrogenated vegetable oils, hydrogenated soybean oil, and medium-chain triglycerides of coconut oil, and palm seed oil. Suitable water-miscible vehicles include, but are not limited to, ethanol, 1,3-butanediol, liquid polyethylene glycol (e.g., polyethylene glycol 300 and polyethylene glycol 400), propylene glycol, glycerin, *N*-methyl-2-pyrrolidone, *N,N*-dimethylacetamide, and dimethyl sulfoxide.

[0287] Suitable antimicrobial agents or preservatives include, but are not limited to, phenols, cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl *p*-hydroxybenzoates, thimerosal, benzalkonium chloride (e.g., benzethonium chloride), methyl-and propyl-parabens, and sorbic acid. Suitable isotonic agents include, but are not limited to, sodium chloride, glycerin, and dextrose. Suitable buffering agents include, but are not limited to, phosphate and citrate. Suitable antioxidants are those as described herein, including bisulfite and sodium metabisulfite. Suitable local anesthetics include, but are not limited to, procaine hydrochloride. Suitable suspending and dispersing agents are those as described herein, including sodium carboxymethylcellulose, hydroxypropyl methylcellulose, and polyvinylpyrrolidone. Suitable

emulsifying agents are those described herein, including polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate 80, and triethanolamine oleate. Suitable sequestering or chelating agents include, but are not limited to EDTA. Suitable pH adjusting agents include, but are not limited to, sodium hydroxide, hydrochloric acid, citric acid, and lactic acid. Suitable complexing agents include, but are not limited to, cyclodextrins, including α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, sulfobutylether- β -cyclodextrin, and sulfobutylether 7- β -cyclodextrin (CAPTISOL®, CyDex, Lenexa, KS).

[0288] When the pharmaceutical compositions provided herein are formulated for multiple dosage administration, the multiple dosage parenteral formulations must contain an antimicrobial agent at bacteriostatic or fungi static concentrations. All parenteral formulations must be sterile, as known and practiced in the art.

[0289] In one embodiment, the pharmaceutical compositions for parenteral administration are provided as ready-to-use sterile solutions. In another embodiment, the pharmaceutical compositions are provided as sterile dry soluble products, including lyophilized powders and hypodermic tablets, to be reconstituted with a vehicle prior to use. In yet another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile suspensions. In yet another embodiment, the pharmaceutical compositions are provided as sterile dry insoluble products to be reconstituted with a vehicle prior to use. In still another embodiment, the pharmaceutical compositions are provided as ready-to-use sterile emulsions.

[0290] The pharmaceutical compositions provided herein for parenteral administration can be formulated as immediate or modified release dosage forms, including delayed-, sustained, pulsed-, controlled, targeted-, and programmed-release forms.

[0291] The pharmaceutical compositions provided herein for parenteral administration can be formulated as a suspension, solid, semi-solid, or thixotropic liquid, for administration as an implanted depot. In one embodiment, the pharmaceutical compositions provided herein are dispersed in a solid inner matrix, which is surrounded by an outer polymeric membrane that is insoluble in body fluids but allows the active ingredient in the pharmaceutical compositions diffuse through.

[0292] Suitable inner matrixes include, but are not limited to, polymethylmethacrylate, polybutyl-methacrylate, plasticized or unplasticized polyvinylchloride, plasticized nylon, plasticized polyethylene terephthalate, natural rubber, polyisoprene, polyisobutylene, polybutadiene, polyethylene, ethylene-vinyl acetate copolymers, silicone rubbers, polydimethylsiloxanes, silicone carbonate copolymers, hydrophilic polymers, such as hydrogels of esters of acrylic and metliacrylic acid, collagen, cross-linked polyvinyl alcohol, and cross-linked partially hydrolyzed polyvinyl acetate.

[0293] Suitable outer polymeric membranes include but are not limited to, polyethylene, polypropylene, ethylene/propylene copolymers, ethylene/ethyl acrylate copolymers, ethylene/vinyl acetate copolymers, silicone rubbers, polydimethyl siloxanes, neoprene rubber, chlorinated polyethylene, polyvinylchloride, vinyl chloride copolymers with vinyl acetate, vinylidene chloride, ethylene and propylene, ionomer polyethylene terephthalate, butyl rubber epichlorohydrin rubbers, ethylene/vinyl alcohol copolymer, ethylene/vinyl acetate/vinyl alcohol terpolymer, and ethylene/vinyloxyethanol copolymer.

V. Examples

[0294] While preferred embodiments of the present disclosure have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.


Example 1. A Phase 2, Open-Label Study to Evaluate the Efficacy and Safety of Compound 1 in Subjects with Testicular Adrenal Rest Tumors

[0295] Testicular adrenal rest tumors (TART) is a common complication in congenital adrenal hyperplasia (CAH) that is caused by adrenocorticotropin hormone (ACTH)–driven overstimulation of aberrant adrenal cells within the testes and results in impaired spermatogenesis and infertility. TARTs can be found anywhere on the testes, though a typical (very common) location is within the mediastinum. This is a particularly problematic location. Because of the central location of TARTs being in the mediastinum, compression of the nearby seminiferous tubules is common and can lead to obstructive azoospermia, irreversible damage to surrounding tissue, and permanent infertility in males.

[0296] Current treatment for TART consists of chronic high-dose glucocorticoids, a problematic therapy with significant side effects and a narrow therapeutic window. Given the serious nature of TART and the limitations and risks of chronic steroid therapy, new treatment modalities are needed for patients with TART. Compound 1, a high-affinity and selective small-molecule antagonist of the corticotropin-releasing factor type-1 (CRF₁) receptor, may be effective in blocking ACTH synthesis, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of glucocorticoids. This mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-hydroxyprogesterone (17-OHP) in subjects with CAH (FIG. 12A-12B). Compound 1 has been

shown to have an acceptable safety and tolerability profile in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. These promising results provide a rationale for investigating the use of Compound 1 in the treatment of TART, in combination with replacement glucocorticoids and mineralocorticoids.

[0297] The structure of Compound 1 is:



[0298] Building upon the 2 completed Phase 1 studies of Compound 1 in healthy volunteers and the ongoing Phase 2 proof-of-concept study of Compound 1 in subjects with CAH, this Phase 2 will study Compound 1 specifically in male CAH subjects with TART for the first time. A sample size of approximately 12 subjects was selected to provide sufficient data for an initial evaluation of efficacy and safety in subjects with TART while maintaining feasible subject recruitment goals for this rare condition. Given the rarity of the disease and thus the small sample size that can be recruited, this study will be open label, and all enrolled subjects will receive active study drug. Subjects will serve as their own controls in that their outcome measures after 12 weeks of study drug dosing will be compared to their baseline measures.

[0299] The schedule of study visits (every 2 weeks through Week 4, then every 4 weeks for the remainder of the study) provides an appropriate balance between minimizing subject burden and closely monitoring subject safety and response to therapy. The 4-week post-treatment washout period will allow for adequate safety follow-up and analysis of any HPA axis rebound after the last dose of study drug.

[0300] The specification of study drug dosing at 10pm and after consumption of a snack is based on information from the Phase 1 studies in healthy volunteers. The plasma C_{max} (maximum concentration) for Compound 1 in the Phase 1 studies occurred approximately 5 hours after dosing. Therefore, Compound 1 will be dosed daily at 10pm in this study in order to achieve peak plasma study drug concentration at the time of day when most adrenal androgens are produced. Subjects will consume a standardized snack before taking study drug because the Phase 1 studies showed minimal study drug absorption when subjects were dosed in a fasted state. Specific dose reduction criteria, strict individual treatment stopping criteria, and appropriate laboratory and medical assessments have been established to ensure subject safety. (FIGS. 1A-1B).

Objectives	Endpoints
Primary	
<ul style="list-style-type: none"> To assess the efficacy of Compound 1 in subjects with TART 	<ul style="list-style-type: none"> Reduction in the size/volume of TART on testicular ultrasound from screening to Week 12
Secondary	
<ul style="list-style-type: none"> To evaluate the safety of Compound 1 in subjects with TART 	<ul style="list-style-type: none"> Adverse events (AEs), serious adverse events (SAEs), AEs leading to withdrawal, AEs of special interest (AESIs) Physical examination findings Vital signs Changes in electrocardiogram (ECG) Changes in clinical laboratory values
<ul style="list-style-type: none"> To assess the pharmacodynamic (PD) biomarker effects of Compound 1 	<ul style="list-style-type: none"> Changes in plasma concentrations of key PD biomarkers, including ACTH, 17-OHP, testosterone, androstenedione, luteinizing hormone (LH), follicle-stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, and aldosterone from baseline to Week 12
<ul style="list-style-type: none"> To assess changes on semen analysis 	<ul style="list-style-type: none"> Changes in sperm count, morphology, and motility from baseline to Week 12
<ul style="list-style-type: none"> To assess changes in metabolism 	<ul style="list-style-type: none"> Changes in metabolic parameters, including fasting glucose, HbA1c, insulin sensitivity, and lipid profile (including LDL cholesterol, HDL cholesterol, and triglycerides), from baseline to Week 12 Changes in body weight, body mass index (BMI), waist circumference, and blood pressure
<ul style="list-style-type: none"> To evaluate changes in quality of life (QoL) and mood 	<ul style="list-style-type: none"> Changes in QoL measured using the Short Form 36 (SF-36) Changes in mood measured using the Hospital Anxiety and Depression Scale (HADS)

Study Population

[0301] Approximately 12 male subjects with TART will be enrolled at approximately 6 investigative sites in the United States, United Kingdom, and possibly other countries. Up to 12 additional subjects may be added to further evaluate specific safety, PD, or efficacy findings.

Eligibility Criteria**Inclusion Criteria**

- Adult male subjects aged 18 to 30, inclusive
- Documented historical diagnosis of CAH based on documented genetic mutation or historical documentation of elevated 17-OHP
- Diagnosis of TART via testicular ultrasound and/or magnetic resonance imaging (MRI) at screening with ≥ 1 lesion(s) measuring ≥ 2 cm in diameter
- On a stable regimen of GC replacement for ≥ 30 days before baseline
- Agree to follow contraception guidelines and refrain from donating sperm throughout the treatment period and for 90 days after the last dose of study drug
- Subject is able to understand all study procedures and risks involved and provides written informed consent indicating willingness to comply with all aspects of the protocol

Exclusion Criteria

- Any clinically significant unstable medical condition, medically significant illness, or chronic disease within 30 days of screening, including but not limited to:
 - A malignancy or < 3 years of remission history from any malignancy, other than successfully treated localized skin cancer
 - Presence of clinically significant renal disease, as evidenced by an estimated glomerular filtration rate (eGFR) of < 60 mL/min/1.73m²
 - Current or chronic history of liver disease or known hepatic or biliary abnormalities (with the exception of Gilbert's syndrome or asymptomatic gallstones)
 - Confirmed positive test at screening for active hepatitis B, hepatitis C, or human immunodeficiency virus (HIV)
- Any clinically significant psychiatric disorders, including any current major depressive episode, bipolar disorder, schizophrenia, schizoaffective disorder, major depressive disorder with psychotic features, or any other psychotic disorder within the preceding 6 months
- HADS score > 12 for either depression or anxiety
- At increased risk of suicide on the basis of the Investigator's judgment or the results of the Columbia-Suicide Severity Rating Scale (C-SSRS) conducted at screening and baseline (eg,

C-SSRS Type 3, 4, or 5 ideation during the preceding 6 months or any suicidal behavior within the past 12 months)

- Clinically significant and abnormal clinical or laboratory assessments must be discussed with the Medical Monitor to determine eligibility for this study. Abnormal assessments that must be reviewed to determine eligibility include but are not limited to:
 - Clinically meaningful abnormal ECG results, in the opinion of the Investigator
 - Fridericia-corrected QT interval (QTcF) >450 msec
 - Alanine aminotransferase (ALT) >2x upper limit of normal (ULN)
 - Bilirubin >1.5x ULN (isolated bilirubin >1.5x ULN is acceptable if bilirubin is fractionated and direct bilirubin is <35%)
- A history that includes bilateral adrenalectomy or hypopituitarism
- Use of any other investigational drug within 30 days or 5 half-lives, whichever is longer, before initial screening
- Require use of prohibited concomitant medications, including strong inhibitors and/or inducers of CYP3A4 (with the exception of glucocorticoids, hydrocortisone, and dexamethasone) within 30 days or 5 half-lives (whichever is longer) of first dose of study drug. Medications that are sensitive substrates or substrates with narrow therapeutic ranges (metabolized by CYP3A4, 2C8, 2C9, or 2C19) should be discussed on a case-by-case basis with the Medical Monitor to determine whether the medication should be discontinued or may be continued with caution. If washout is feasible, then the medication should be withdrawn at least 30 days or 5 half-lives (whichever is longer) before the first dose of study drug.
- Donation of blood within 60 days before first dose of study drug, or donation of platelets, white blood cells, or plasma within 15 days before first dose of study drug.

Study Duration

[0302] The expected duration of study participation for each subject is up to approximately 20 weeks. This includes a screening period of ≤4 weeks, a treatment period of 12 weeks, and a safety follow-up period of 4 weeks.

Study Design

[0303] This is a Phase 2, open-label efficacy and safety study of Compound 1 for the treatment of TART. All subjects will receive a daily oral dose of Compound 1 for 12 weeks. In-clinic study visits will occur for screening and at Day 1 (baseline) and Weeks 2, 4, 8, 12 (end of treatment), and 16 (follow-up).

Dose, Route, Regimen

[0304] All subjects will receive a daily oral dose of Compound 1 for 12 weeks. Subjects will receive 600 mg/day during the first 2 weeks and 1000 mg/day during the remaining 10 weeks of the treatment period. Study drug will be taken daily at 10pm (or bedtime, if earlier), 5 to 15 minutes after consumption of a standardized snack.

[0305] On mornings of study visits with 8am blood draws for laboratory assessments, subjects should take their morning glucocorticoid replacement after the blood draws. For all other days during the study, subjects may take their morning glucocorticoid replacement at their usual time (generally between 6am and 8am).

[0306] If a subject experiences a Grade 3 or higher TEAE considered at least possibly related to study drug, the Investigator may reduce the subject's daily dose of Compound 1 down in increments of 200 mg. Study drug will be discontinued in subjects who experience clinically significant liver chemistry, QTcF, or other individual treatment stopping criteria.

Investigational Drug

[0307] Compound 1 is a small-molecule CRF₁ receptor antagonist and will be supplied as 200-mg, size 1, white, hard gelatin capsules.

Example 2.

[0308] Since elevated ACTH is the primary driver for TART growth, blocking ACTH synthesis using Compound 1, a small-molecule antagonist of the CRF₁ receptor, may be effective in reducing ACTH levels, thereby reducing the size of testicular lesions in TART and reducing the need for chronic supraphysiologic doses of GCs. In CAH, this mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH (FIG. 12A-12B). The safety profile of Compound 1 has been established in nonclinical toxicology studies and in 2 completed Phase 1 clinical studies in healthy volunteers. A Phase 2 study in subjects with CAH is currently ongoing.

Nonclinical Toxicology of Compound 1

[0309] The nonclinical safety profile of Compound 1 has been evaluated in repeat-dose range-finding and definitive toxicology studies of up to 91 days in duration in rats and dogs using the clinically relevant route (oral) and schedule of administration (once daily). In rats and beagle dogs administered oral Compound 1 once daily for 91 consecutive days at doses up to 2000 mg/kg/day, toxicologic findings were all mild and limited to the male reproductive tract, and all Compound 1-related findings were reversible following a 6-week treatment-free recovery period.

[0310] Additionally, a fertility toxicology study was conducted in rats. In this study, male rats were administered oral Compound 1 once daily at 0, 5, 20, or 1000 mg/kg for 85 to 87 days (70

days before mating, during mating, and ≤ 4 days between mating and necropsy). Male reproductive performance (mean day of mating, mating and fertility indices, and conception rates) was unaffected by Compound 1 treatment. Untreated females mated with the Compound 1-treated males had no ovarian and uterine effects.

[0311] In addition, a complete battery of standard genotoxicity tests consisting of in vitro bacterial reverse mutation and chromosome aberration assays and an in vivo (rat) bone marrow micronucleus assay has been conducted. Exploratory and definitive embryo-fetal development studies in rabbits and a bovine corneal opacity and permeability assay have also been conducted.

Clinical Experience with Compound 1

[0312] To date, a total of 53 subjects, all healthy adults aged 21 to 65 years, have been treated with at least 1 dose of Compound 1 in 2 Phase 1 studies. The first-in-human study of Compound 1 (Study I3C-FW-BLAA) evaluated single ascending doses of 2 mg to 800 mg. In this study, a small number of subjects experienced diarrhea and headache of mild or moderate severity. A subsequent study (Study I3C-FW-BLAB) evaluated repeat doses of 50 mg, 100 mg, or 200 mg Compound 1 once daily for 14 consecutive days. A small number of subjects in the repeat-dose study experienced dyspnea, rhinorrhea, palpitations, and headache. The Phase 1 studies showed no clinically significant alterations in safety laboratory values, vital signs, or electrocardiograms (ECGs). Compound 1 has been well tolerated in healthy human subjects, with no SAEs, AEs leading to discontinuation, or dose-limiting toxicities (DLTs). A maximum tolerated dose has not been reached in humans.

[0313] An ongoing Phase 2 open-label study (Study Compound 2) is currently evaluating the safety and efficacy of Compound 1 dosed once daily for 6 weeks in adults with classic CAH. In the initial Cohort A of this study, all subjects are dosed starting at 200 mg/day for 14 days, escalating for each subject to 600 mg/day for 14 days, then 1000 mg/day for 14 days if the subject experiences no DLTs at the previous dose level. **(FIGS. 2-3).**

Known Potential Risks

[0314] Because the reversible testicular effects (degeneration/atrophy) observed in rats and dogs treated with high-dose Compound 1 (2,000 mg/kg/day) are likely associated with on-target (mechanism-driven) effects, these effects are likely to be relevant to human risk. However, male reproductive studies in rodents showed no impairment in fertility at the highest dose tested (1,000 mg/kg/day). Additionally, there were no effects on clinical laboratory markers of testicular function (LH and FSH) when Compound 1 was administered to healthy male volunteers at doses up to 200 mg per day for 14 consecutive days. Clinical laboratory measures of testicular function (LH, FSH, inhibin B, and SHBG) will be assessed throughout this study to

monitor the risk of testicular injury. Testicular ultrasounds and semen analysis will also be performed at screening/baseline and the end of the treatment period.

[0315] Laboratory abnormalities in liver function tests (LFTs) have been observed at significantly higher doses in nonclinical studies. However, these events were not observed in the Phase 1 clinical studies. Clinical laboratory measures of liver chemistry will be assessed throughout this study, and strict individual treatment stopping criteria for liver chemistry findings are in place.

[0316] Compound 1-induced effects on the thyroid gland were noted in dogs after 91 days of high-dose (2,000 mg/kg/day) treatment but not in any other nonclinical studies. Clinical pathology parameters of thyroid function (eg, T4, T3, thyroid-stimulating hormone [TSH]) were not evaluated in the Phase 1 studies but will be monitored throughout this study.

Known Potential Benefits

[0317] Early clinical studies suggests that Compound 1 is effective as a CRF₁ receptor antagonist, a mechanism of action that supports the clinical investigation of Compound 1 for treatment of TART. In CAH, this mechanism of action has been validated in a previous proof-of-concept study using a similar small-molecule CRF₁ receptor antagonist, which produced dose-dependent reductions in ACTH and 17-OHP in subjects with CAH (FIG. 12A-12B).

Assessment of Potential Risks and Benefits

[0318] The mild and reversible toxicologic effects of Compound 1 on the male reproductive tract in nonclinical studies occurred in the context of animals with normal, rather than elevated, baseline hormone function. As a CRF₁ receptor antagonist, Compound 1 is intended to reduce abnormally elevated ACTH levels, which in TART drives overstimulation of aberrant adrenal cells within the testes. In nonclinical models where baseline hormone levels are within normal ranges, clinical benefits are not expected, and indeed, toxicities may be expected as a result of androgen reduction outside of the normal range. In TART, where baseline ACTH and androgen levels are pathologically elevated, androgen reduction would be considered a positive, therapeutic effect.

[0319] Given the serious nature of TART and the limitations (and risks) of chronic steroid therapy, new treatment modalities are needed for patients with TART. Given the acceptable safety and tolerability profile of Compound 1 in healthy volunteers from the Phase 1 program, the low probability of demonstrating meaningful PD marker changes in healthy subjects, and the lack of any clear benefit to exposing more healthy subjects to study drug, the Sponsor believes the benefit-to-risk profile favors the entry of Compound 1 into Phase 2 in subjects with TART.

Example 3. Congenital Adrenal Hyperplasia (CAH)

[0320] CAH results from a genetic defect in the 21-hydroxylase enzyme. CAH patients are thus unable to produce cortisol. Without cortisol to provide negative feedback, patients over-produce ACTH. High ACTH promotes the overproduction of potent androgens (male sex hormones). Without wishing to be bound by theory, Compound 1 blocks CRF₁ receptor to prevent ACTH release by the pituitary and associated drive on the hypothalamic-pituitary-adrenal axis. Replacement of cortisol to physiological levels will correct cortisol deficiency, however, will not lower ACTH, resulting in hyperplasia of adrenal gland and associated overproduction of adrenal androgens.

[0321] As shown in **FIGS. 4** and **5A-5B**, the attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism. 80% of subjects demonstrated reduction in ACTH. 70% of subjects demonstrated > 25% reduction in ACTH. 30% of subjects were in the normal range (*) after treatment with a CRF₁ receptor antagonist.

[0322] **FIGS. 6** and **7A-7B** demonstrate “control” of disease based on standard guidelines, which would then allow steroid taper. 80% of subjects demonstrated reduction in 17-OHP. 50% of subjects demonstrated > 25% reduction in ACTH. 50% of subjects were in the guideline range (*; 1200 ng/dL) after treatment with a CRF₁ receptor antagonist.

[0323] **FIGS. 8** and **9A-9B** demonstrate attenuation of morning rise in A4 upon treatment with a CRF₁ antagonist, indicative of its ability to control excess androgen production and associated symptoms. 100% of subjects demonstrated reduction in androstenedione (at various doses). 60% of subjects demonstrated > 25% reduction in androstenedione. 50% of subjects were within the normal reference range after treatment with a CRF₁ receptor antagonist.

[0324] **FIGS. 10A-10B** show approximately 25% shrinkage in TART in a 25 year old male with classic congenital adrenal hyperplasia following 6 weeks treatment with a CRF₁ receptor antagonist.

[0325] **FIGS. 11A-11B** demonstrate Compound 1 is a potent, selective, and functional CRF₁ receptor antagonist. In in vitro and ex vivo studies, Compound 1 was determined to be a potent and selective CRF₁ receptor antagonist with high affinity and selective for the hCRF₁ receptor. Compound 1 was found to have functional potency to inhibit adenylate cyclase activity (CRF₁ receptor-mediated downstream cascade), demonstrating in vitro efficacy as a CRF₁ receptor antagonist. Compound 1 was found to significantly inhibit CRF-induced locomotor behavior in response to i.c.v. administration of CRF (**FIG. 11B**).

[0326] ~~**FIGS. 12A-12B** demonstrate single pill of CRF₁ receptor antagonist improves ACTH and 17-OHP in CAH patients.~~ **FIGS. 12A-12B** demonstrate single pill of CRF₁ receptor antagonist (i.e., as described in US 2017/0020877 A1) improves ACTH and 17-OHP in CAH patients.

[0327] **FIG. 13** shows the subject level hormone changes (ACTH, 17-OHP, and androstenedione) by dose of a CRF₁ receptor antagonist.

Example 4. Assess reduction in excess ACTH and adrenal steroid production

[0328] The objectives of the study are: to assess the ability of Compound 1 to reduce excess ACTH and adrenal steroid production in participants with classic CAH on a stable glucocorticoid regimen; to evaluate the safety and tolerability of tildacerfont in participants with classic CAH; and to evaluate the pharmacokinetics (PK) of tildacerfont.

ELIGIBILITY CRITERIA

[0329] Inclusion Criteria:

- Adult males and females 18 years or older
- Diagnosis of classic CAH due to 21-hydroxylase deficiency
- Morning 17OHP ≥ 800 ng/dL, before taking any morning GC dose
- On a stable regimen of glucocorticoid replacement for ≥ 30 days before enrollment

[0330] Exclusion Criteria:

- Clinically significant unstable medical condition, illness, or chronic disease within 30 days of screening
- Clinically significant psychiatric disorder
- History of bilateral adrenalectomy or hypopituitarism

STUDY DESIGN

[0331] **FIG. 14** depicts the proof of concept and escalating QD dosing at 10 PM for Cohort A and the single dose level and BID dosing at 10 AM and 10 pm for Cohort B.

BASELINE CHARACTERISTICS

[0332] Table 1 provides a description of the baseline characteristics displayed by Cohort A and Cohort B. In Cohort A, 10 participants were enrolled and received at least 1 dose of Compound 1. One participant withdrew early because of an occupation-related scheduling conflict. In Cohort B, 9 participants were enrolled and received at least one dose of Compound 1. One participant was withdrawn upon discovery of a major protocol violation at Baseline and did not have postbaseline efficacy/PK data. Thus, baseline characteristics are calculated for n=8 in Cohort B. One participant in Cohort B was taking both HC and prednisone.

Table 1 – Baseline characteristics of each cohort

Parameter	Cohort A (n=10)	Cohort B (n=8)
Demographics		
Age (years), mean (range)	39.3 (24-66)	36.8 (19-66)
Sex	5M, 5F	1M, 7F
Bmi (kg/m ²), mean $\pm$ SD	31.5 $\pm$ 12.0	31.1 $\pm$ 6.7

Daily GC dose (mg HC equivalents), mean range	28 (13-40)	35 (20-50)
Types of GCs	4 JC, 3 pred, 3 dex	2HC, 4 pred, 3 dex
17OHP at screening ^a (ng/dL), mean $\pm$ SD	5,985 $\pm$ 6,347	4,592 $\pm$ 4,624

Abbreviations: 17OHP: 17-hydroxyprogesterone; BMI: body mass index; dex: dexamethasone; F: female; GC: glucocorticoid; HC: hydrocortisone; M: male; pred: prednisone; SD: standard deviation.
^a 17OHP reference ranges: 27-199 ng/dL males, 15-290 ng/dL females.

SAFETY

[0333] Across both cohorts, Compound 1 was generally well tolerated. The most common type of adverse event (AE) was gastrointestinal disorders. There were no serious AEs and no AEs leading to withdrawal.

PHARMACOKINETICS

[0334] Over a wide range of doses (200 to 1000 mg QD), Compound 1 exhibited a predictable, dose-dependent PK profile.

EFFICACY

[0335] FIG. 15A-15D show the changes in levels of Androstenedione (A4) across both cohorts. Levels of A4 improved at ≥ 1 dose level for 10 out of 10 participants in Cohort A and 6 out of 8 participants in Cohort B. Excess A4 is associated with clinical sequelae in CAH. Attenuation of A4 demonstrates control of adrenal androgen.

[0336] FIG. 16A-16D show the changes in levels of 17-hydroxyprogesterone (17-OHP) across both cohorts. Levels of 17OHP improved at ≥ 1 dose level for 8 out of 10 participants in Cohort A and 6 out of 7 participants in Cohort B. 17OHP is a critical hormone in CAH diagnosis and management. Attenuation of 17OHP demonstrates control of adrenal steroid production.

[0337] FIG. 17A-17D show the changes in levels of adrenocorticotrophic hormone (ACTH) across both cohorts. Levels of ACTH improved at ≥ 1 dose level for 8 out of 10 participants in Cohort A and 5 out of 7 participants in Cohort B. Attenuation of ACTH demonstrates target engagement and functional CRF₁ receptor antagonism.

Table 2 – Percentage change from baseline in ACTH, 17OHP and A4.

	Cohort A			Cohort B
	200 mg QD (Week 2)	600 mg QD (Week 4)	1000 mg QD (Week 6)	200 mg BID (Week 2)
ACTH	-25.7 ($\pm$ 51.6)	-60.1 ($\pm$ 37.5)	-32.7 ($\pm$ 47.0)	-36.5 ($\pm$ 58.0)
17OHP	+22.2 ($\pm$ 172.3)	-23.7 ($\pm$ 64.2)	-13.8 ($\pm$ 74.7)	-47.8 ($\pm$ 63.3)
A4	-12.1 ($\pm$ 47.4)	-33.2 ($\pm$ 36.5)	-28.6 ($\pm$ 44.6)	-16.0 ($\pm$ 81.4)

CONCLUSIONS

[0338] Compound 1, a potent and selective CRF₁ receptor antagonist, offers a novel nonsteroidal oral treatment approach for CAH. Compound 1 was well tolerated and efficacious in reducing excess ACTH, 17OHP, and androstenedione in patients with classic CAH.

Compound 1 exhibited a predictable, dose-dependent PK profile suitable for the target population.

[0339] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

ABSTRACT OF THE DISCLOSURE

Provided herein are compounds and pharmaceutical compositions for the prevention and treatment of testicular adrenal rest tumors (TART) or ovarian adrenal rest tumors (OART).

Doc Code: ECOMM.AUTH/ECOMM.WTDW

Doc Description: Internet Communications Authorization/Internet Communications Authorization Withdrawal

PTO/SB/439 (11-15)

**AUTHORIZATION FOR INTERNET
COMMUNICATIONS IN A PATENT
APPLICATION OR REQUEST TO
WITHDRAW AUTHORIZATION FOR
INTERNET COMMUNICATIONS**

Application No.	19/043,264
Filing Date	January 31, 2025
First Named Inventor	Alexis Howerton
Art Unit	1627
Examiner Name	Pihonak, Sarah
Practitioner Docket No.	50535-710.306

To: Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

I. To authorize permission for Internet Communications.

☒ Recognizing that Internet communications are not secure, I hereby authorize the USPTO to communicate with the undersigned and practitioners in accordance with 37 CFR 1.33 and 37 CFR 1.34 concerning any subject matter of this application via video conferencing, instant messaging, or electronic mail. I understand that a copy of these communications will be made of record in the application file. (MPEP 502.03)

II. To withdraw authorization for Internet Communications.

☐ The authorization given on _____, to the USPTO to communicate with the undersigned and any practitioner in accordance with 37 CFR 1.33 and 37 CFR 1.34 concerning any subject matter of this application via Internet communications such as video conferencing, instant messaging, or electronic mail is hereby withdrawn. I understand that the withdrawal is effective when approved rather than when received.

I am the

- ☐ applicant.
- ☐ attorney or agent of record. Registration number _____.
- ☒ attorney or agent acting under 37 CFR 1.34. Registration number 79,641.

/Victoria Cheng/

07/24/2025

Signature

Date

Victoria Cheng

(858) 350-2240

Typed or printed name

Telephone Number

NOTE: This form must be signed in accordance with 37 CFR 1.33. See 37 CFR 1.4 for signature requirements and certifications. Juristic entities must be represented by a patent practitioner (see 37 CFR 1.31, which is applicable to any paper filed on or after September 16, 2012, that is presented on behalf of a juristic entity, regardless of application filing date). Submit multiple forms if more than one signature is required, see below*.

☒ * Total of 1 forms are submitted.

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. The United States Patent and Trademark Office (USPTO) collects the information in this record under authority of 35 U.S.C. 2. The USPTO's system of records is used to manage all applicant and owner information including name, citizenship, residence, post office address, and other information with respect to inventors and their legal representatives pertaining to the applicant's/owner's activities in connection with the invention for which a patent is sought or has been granted. The applicable Privacy Act System of Records Notice for the information collected in this form is COMMERCE/PAT-TM-7 Patent Application Files, available in the Federal Register at 78 FR 19243 (March 29, 2013), <https://www.govinfo.gov/content/pkg/FR-2013-03-29/pdf/2013-07341.pdf>.

Routine uses of the information in this record may include disclosure to: 1) law enforcement, in the event that the system of records indicates a violation or potential violation of law; 2) a federal, state, local, or international agency, in response to its request; 3) a contractor of the USPTO having need for the information in order to perform a contract; 4) the Department of Justice for determination of whether the Freedom of Information Act (FOIA) requires disclosure of the record; 5) a Member of Congress submitting a request involving an individual to whom the record pertains, when the individual has requested the Member's assistance with respect to the subject matter of the record; 6) a court, magistrate, or administrative tribunal, in the course of presenting evidence, including disclosures to opposing counsel in the course of settlement negotiations; 7) the Administrator, General Services Administration (GSA), or their designee, during an inspection of records conducted by GSA under authority of 44 U.S.C. 2904 and 2906, in accordance with the GSA regulations and any other relevant (i.e., GSA or Commerce) directive, where such disclosure shall not be used to make determinations about individuals; 8) another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)); 9) the Office of Personnel Management (OPM) for personnel research purposes; and 10) the Office of Management and Budget (OMB) for legislative coordination and clearance.

If you do not furnish the information requested on this form, the USPTO may not be able to process and/or examine your submission, which may result in termination of proceedings, abandonment of the application, and/or expiration of the patent.

PATENT APPLICATION FEE DETERMINATION RECORD Substitute for Form PTO-875				Application or Docket Number 19/043,264		Filing Date 01/31/2025		☐ To be Mailed		
ENTITY: ☒ LARGE ☐ SMALL ☐ MICRO										
APPLICATION AS FILED - PART I										
		(Column 1)	(Column 2)							
FOR		NUMBER FILED	NUMBER EXTRA	RATE (\$)		FEE (\$)				
☐ BASIC FEE (37 CFR 1.16(a), (b), or (c))		N/A	N/A	N/A						
☐ SEARCH FEE (37 CFR 1.16(k), (l), or (m))		N/A	N/A	N/A						
☐ EXAMINATION FEE (37 CFR 1.16(o), (p), or (q))		N/A	N/A	N/A						
TOTAL CLAIMS (37 CFR 1.16(j))		minus 20 =	*	x \$200 =						
INDEPENDENT CLAIMS (37 CFR 1.16(h))		minus 3 =	*	x \$600 =						
☐ APPLICATION SIZE FEE (37 CFR 1.16(s))		If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$310 (\$155 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).								
☐ MULTIPLE DEPENDENT CLAIM PRESENT (37 CFR 1.16(j))										
* If the difference in column 1 is less than zero, enter "0" in column 2.				TOTAL						
APPLICATION AS AMENDED - PART II										
		(Column 1)	(Column 2)	(Column 3)						
AMENDMENT	10/27/2025	CLAIMS REMAINING AFTER AMENDMENT		HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)		ADDITIONAL FEE (\$)		
	Total (37 CFR 1.16(i))	* 21	Minus	** 21	= 0	x \$200 =		0		
	Independent (37 CFR 1.16(h))	* 1	Minus	*** 3	= 0	x \$600 =		0		
	☐ Application Size Fee (37 CFR 1.16(s))									
	☐ FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))									
						TOTAL ADD'L FEE		0		
AMENDMENT		(Column 1)	(Column 2)	(Column 3)						
		CLAIMS REMAINING AFTER AMENDMENT		HIGHEST NUMBER PREVIOUSLY PAID FOR	PRESENT EXTRA	RATE (\$)		ADDITIONAL FEE (\$)		
	Total (37 CFR 1.16(i))	*	Minus	**	=	x \$0 =				
	Independent (37 CFR 1.16(h))	*	Minus	***	=	x \$0 =				
	☐ Application Size Fee (37 CFR 1.16(s))									
☐ FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM (37 CFR 1.16(j))										
						TOTAL ADD'L FEE				
* If the entry in column 1 is less than the entry in column 2, write "0" in column 3.						SLIE				
** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 20, enter "20".						/TONYA Q MCBRIDE/				
*** If the "Highest Number Previously Paid For" IN THIS SPACE is less than 3, enter "3".										
The "Highest Number Previously Paid For" (Total or Independent) is the highest number found in the appropriate box in column 1.										

This collection of information is required by 37 CFR 1.16. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.



P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

TERMINAL DISCLAIMER TO OBVIATE A DOUBLE PATENTING REJECTION OVER A PRIOR PATENT

APPLICATION #	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET #
19043264	01/31/2025	Alexis HOWERTON	50535-710.306

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS



Filing of terminal disclaimer does not obviate requirement for response under 37 CFR 1.111 to outstanding Office Action



This electronic Terminal Disclaimer is not being used for a Joint Research Agreement.

Owner	Percent interest
Spruce Biosciences, Inc.	100%
Total	100%

The owner(s) of percent interest listed above in the instant application hereby disclaims, except as provided below, the terminal part of the statutory term of any patent granted on the instant application which would extend beyond the expiration date of the full statutory term of any patent granted on pending reference Application Number(s)

Application #	Filing Date
18941832	11/08/2024

19043275	01/31/2025
19043303	01/31/2025

as the term of any patent granted on said reference application may be shortened by any terminal disclaimer filed prior to the grant of any patent on the pending reference application. The owner hereby agrees that any patent so granted on the instant application shall be enforceable only for and during such period that it and any patent granted on the reference application are commonly owned. This agreement runs with any patent granted on the instant application and is binding upon the grantee, its successors or assigns.

In making the above disclaimer, the owner does not disclaim the terminal part of any patent granted on the instant application that would extend to the expiration date of the full statutory term of any patent granted on said reference application, "as the term of any patent granted on said reference application may be shortened by any terminal disclaimer filed prior to the grant of any patent on the pending reference application," in the event that any such patent granted on the pending reference application: expires for failure to pay a maintenance fee, is held unenforceable, is found invalid by a court of competent jurisdiction, is statutorily disclaimed in whole or terminally disclaimed under 37 CFR 1.321, has all claims canceled by a reexamination certificate, is reissued, or is in any manner terminated prior to the expiration of its full statutory term as shortened by any terminal disclaimer filed prior to its grant.

The owner(s) of percent interest listed above in the instant application hereby disclaims, except as provided below, the terminal part of the statutory term of any patent granted on the instant application which would extend beyond the expiration date of the full statutory term of prior patent number(s)

Patent #
11304950

as the term of said prior patent is presently shortened by any terminal disclaimer. The owner hereby agrees that any patent so granted on the instant application shall be enforceable only for and during such period that it and the prior patent are commonly owned. This agreement runs with any patent granted on the instant application and is binding upon the grantee, its successors or assigns.

In making the above disclaimer, the owner does not disclaim the terminal part of the term of any patent granted on the instant application that would extend to the expiration date of the full statutory term of the prior patent, "as the term of said prior patent is presently shortened by any terminal disclaimer," in the event that said prior patent later:

- expires for failure to pay a maintenance fee;
- is held unenforceable;
- is found invalid by a court of competent jurisdiction;
- is statutorily disclaimed in whole or terminally disclaimed under 37 CFR 1.321;
- has all claims canceled by a reexamination certificate;
- is reissued; or
- is in any manner terminated prior to the expiration of its full statutory term as presently shortened by any terminal

disclaimer.



Terminal disclaimer fee under 37 CFR 1.20(d) included with Electronic Terminal Disclaimer request.

Applicant claims the following entity status:

Regular Undiscounted

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

I certify, in accordance with 37 CFR 1.4(d)(4) that I am: An attorney or agent registered to practice before the Patent and Trademark Office who is of record in this application

Signature	Name	Registration #
/Victoria Cheng/	Victoria Cheng	79641

* Statement under 37 CFR 3.73(b) is required if terminal disclaimer is signed by the assignee (owner). Form PTO/SB/96 may be used for making this certification. See MPEP 324.



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC ACKNOWLEDGEMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 10/27/2025 02:35:03 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Victoria Cheng
PATENT CENTER #	72905171	FILING DATE	01/31/2025
CUSTOMER #	21971	FIRST NAMED INVENTOR	Alexis HOWERTON
CORRESPONDENCE ADDRESS	-	AUTHORIZED BY	Victoria Cheng

Documents

TOTAL DOCUMENTS: 2

DOCUMENT	PAGES	DESCRIPTION	SIZE (KB)
petition-request.pdf	3	Terminal Disclaimer-Filed (Electronic)	46 KB
grantLetter.pdf	1	Terminal Disclaimer-Electronic- Approved	18 KB

Digest

DOCUMENT	MESSAGE DIGEST(SHA-512)
petition-request.pdf	1C004DD234919BC4EA7B50A38352244BA91B130A3F7BD54D BDA0492F3B3503F4B3A8933F7538278DD9F0009EBD4D7DC27 B07D74FDBF366AD92ACE42F47B05055
grantLetter.pdf	367F4AFD2AF4D825CA4BD66BD3B8C943516AF833CAE48B58

2485AA924EE39D742B0A86498232CBC5FC95126364D208F50
B734B133E3C7CFBFA51F0B45B3C36DA

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



**UNITED STATES
PATENT AND TRADEMARK OFFICE**

Commissioner for Patents
United States Patent and Trademark Office
P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

APPROVAL LETTER

APPLICATION #
19/043,264

FILING DATE
01/31/2025

APPLICANT/PATENT UNDER REEXAMINATION
Alexis HOWERTON

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Electronic terminal disclaimer filed on 10/27/2025



Approved

This patent is subject to a Terminal Disclaimer

Approved / Disapproved by: Electronic Terminal Disclaimer automatically approved



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC PAYMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 10/27/2025 02:35:03 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Victoria Cheng
PATENT CENTER #	72905171	AUTHORIZED BY	Victoria Cheng
CUSTOMER #	21971	FILING DATE	01/31/2025
CORRESPONDENCE ADDRESS	-	FIRST NAMED INVENTOR	Alexis HOWERTON

Payment Information

PAYMENT METHOD DA / 232415	PAYMENT TRANSACTION ID E20250QE39118524	PAYMENT AUTHORIZED BY Victoria Cheng
--------------------------------------	---	--

FEE CODE	DESCRIPTION	ITEM PRICE(\$)	QUANTITY	ITEM TOTAL(\$)
1814	STATUTORY DISCLAIMER, INCLUDING TERMINAL DISCLAIMER	183.00	1	183.00
TOTAL AMOUNT:				\$183.00

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C.

371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.

PATENT
Attorney Docket No. 50535-710.306

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:	Confirmation No.: 6961
Inventors: Alexis Howerton et al.	Art Unit: 1627
Application No.: 19/043,264	Examiner: Pihonak, Sarah
Filing Date: January 31, 2025	Customer No.: 21971
Title: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS	<p style="text-align: center;"><u>Certificate of Electronic Filing</u></p> <p>I hereby certify that this correspondence is being transmitted via the U.S. Patent and Trademark Office's electronic filing system to the USPTO on <u>October 27, 2025</u>.</p> <p>By: _____ / Colleen Mensching / Colleen Mensching</p>

Mail Stop AF/RCE
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

AMENDMENT AND RESPONSE TO FINAL OFFICE ACTION

Dear Commissioner:

In response to the Final Office Action mailed August 18, 2025, setting an initial due date of November 18, 2025, Applicant respectfully requests reconsideration of the above-referenced application in view of the following amendments and remarks.

A Request for Continued Examination pursuant to 37 CFR § 1.114 is filed herewith.

Amendments to the Claims begin on page 2 of this paper.


Remarks begin on page 5 of this paper.

Conclusion begins on page 6 of this paper.


AMENDMENTS TO THE CLAIMS

Listing of the Claims

1. (Previously Presented) A method of treating ovarian adrenal rest tumors (OART) and menstrual irregularity in a subject, comprising administering to the subject a small molecule corticotropin-releasing factor type-1 (CRF₁) antagonist, or a pharmaceutically acceptable salt thereof, wherein the small molecule CRF₁ antagonist is:




2. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is:



, or a pharmaceutically acceptable salt thereof.

3. (Previously Presented) The method of claim 1, wherein the small molecule CRF₁ antagonist is:



, or a pharmaceutically acceptable salt thereof.

4. (Previously Presented) The method of claim 1, further comprising treating ovulatory dysfunction or infertility.
5. (Original) The method of claim 1, wherein the subject has congenital adrenal hyperplasia (CAH).
6. (Original) The method of claim 1, wherein the subject is an adolescent.
7. (Original) The method of claim 1, wherein the subject is at least 18 years or older.
8. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered at a dose of about 10 mg/day to about 400 mg/day to the subject.
9. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 50 mg.
10. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 100 mg.
11. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 200 mg.
12. (Original) The method of claim 8, wherein the small molecule CRF₁ antagonist is administered at a dose of about 400 mg.

13. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject once per day.
14. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered to the subject twice per day.
15. (Original) The method of claim 13, wherein the subject is in a fed state.
16. (Original) The method of claim 14, wherein the subject is in a fed state.
17. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is administered orally.
18. (Original) The method of claim 1, wherein the small molecule CRF₁ antagonist is formulated as a capsule or tablet.
19. (Original) The method of claim 1, wherein the administering reduces a plasma concentration of a biomarker comprising 17 α -hydroxyprogesterone (17-OHP), adrenocorticotrophic hormone (ACTH), testosterone, androstenedione (A4), luteinizing hormone (LH), follicle-stimulating hormone (FSH), inhibin B, sex hormone-binding globulin (SHBG), renin, or aldosterone, or a combination thereof.
20. (Original) The method of claim 19, wherein the biomarker is selected from the group consisting of 17-OHP, ACTH, and A4.
21. (Original) The method of claim 19, wherein the plasma concentration of the biomarker is reduced by at least 10% from baseline.

REMARKS

No claims are amended herein. Accordingly, claims 1-21 are currently pending. Allowance is respectfully requested in view of the remarks below.

Double Patenting

Claims 1-5, 8-14, and 17-21 are rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-15 and 19 of U.S. Patent No. 11,304,950 B2 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003; and Grigoriadis, WO 2015112642 A1, publ. 7/30/2015.

Claims 1-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over:

- Claims 72-91 of copending application 18941832 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003;
- Claims 1-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-5 and 9-21 of copending application 19043275 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003; and
- Claims 1-13 and 17-21 are provisionally rejected on the ground of nonstatutory double patenting as being unpatentable over claims 1-19 of copending application 19043303 in view of Stikkelbroeck et. al., Fertility & Sterility, vol. 80(4), pp. 1030-1036, publ. 2003, collectively the "Cited References".

Without acquiescing to the basis of the rejection and for the purpose of compact prosecution, Applicant files herein a terminal disclaimer over U.S. Patent No. 11,304,950, and U.S. Application Nos. 18/941,832, 19/043,275, and 19/043,303. Therefore, Applicant respectfully requests withdrawal of the rejections above and allowance of the pending claims.

Application No. 19/043,264
Office Action Mailed August 18, 2025
Amendment dated: October 27, 2025

CONCLUSION

In light of the remarks set forth above, Applicant believes that the pending claims are under condition for allowance. Applicant respectfully solicits the Examiner to expedite the prosecution of this patent application to issuance. Should the Examiner have any questions, the Examiner is encouraged to telephone the undersigned at (858) 350-2240.

The Commissioner is authorized to charge any underpayment or credit any overpayment to Deposit account No. 23-2415 (Attorney Docket No. 50535-710.306).

Respectfully submitted,

Date: October 27, 2025

By: /Victoria Cheng/
Victoria Cheng, Reg. No. 79,641

WILSON SONSINI GOODRICH & ROSATI
650 Page Mill Road
Palo Alto, CA 94304-1050
Phone: (650) 493-9300
Customer No. 21971

Doc code: RCEX

Doc description: Request for Continued Examination (RCE)

PTO/SB/30EFS (01-22)

Approved for use through 05/31/2024. OMB 0651-0031

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**REQUEST FOR CONTINUED EXAMINATION(RCE)TRANSMITTAL
(Submitted Only via EFS-Web)**

Application Number	19/043,264	Filing Date	2025-01-31	Docket Number (if applicable)	50535-710.306	Art Unit	1627
First Named Inventor	Spruce Biosciences, Inc.			Examiner Name	SARAH PIHONAK		

This is a Request for Continued Examination (RCE) under 37 CFR 1.114 of the above-identified application.Request for Continued Examination (RCE) practice under 37 CFR 1.114 does not apply to any utility or plant application filed prior to June 8, 1995, or to any design application. The Instruction Sheet for this form is located at WWW.USPTO.GOV**SUBMISSION REQUIRED UNDER 37 CFR 1.114**

Note: If the RCE is proper, any previously filed unentered amendments and amendments enclosed with the RCE will be entered in the order in which they were filed unless applicant instructs otherwise. If applicant does not wish to have any previously filed unentered amendment(s) entered, applicant must request non-entry of such amendment(s).

☐ Previously submitted. If a final Office action is outstanding, any amendments filed after the final Office action may be considered as a submission even if this box is not checked.☐ Consider the arguments in the Appeal Brief or Reply Brief previously filed on _____☐ Other _____☒ Enclosed☒ Amendment/Reply☐ Information Disclosure Statement (IDS)☐ Affidavit(s)/ Declaration(s)☐ Other _____**MISCELLANEOUS**☐ Suspension of action on the above-identified application is requested under 37 CFR 1.103(c) for a period of months _____. (Period of suspension shall not exceed 3 months; Fee under 37 CFR 1.17(i) required)☐ Other _____**FEES****The RCE fee under 37 CFR 1.17(e) is required by 37 CFR 1.114 when the RCE is filed.**☒ The Director is hereby authorized to charge any underpayment of fees, or credit any overpayments, to Deposit Account No. 232415**SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED**☒ Patent Practitioner Signature☐ Applicant Signature

Doc code: RCEX

Doc description: Request for Continued Examination (RCE)

PTO/SB/30EFS (01-22)

Approved for use through 05/31/2024. OMB 0651-0031

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Signature of Registered U.S. Patent Practitioner			
Signature	/Victoria Cheng/	Date (YYYY-MM-DD)	2025-10-27
Name	VICTORIA CHENG	Registration Number	79641

This collection of information is required by 37 CFR 1.114. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. Accordingly, pursuant to the requirements of the Act, please be advised that: (1) the general authority for the collection of this information is 35 U.S.C. 2(b)(2); (2) furnishing of the information solicited is voluntary; and (3) the principal purpose for which the information is used by the U.S. Patent and Trademark Office is to process and/or examine your submission related to a patent application or patent. If you do not furnish the requested information, the U.S. Patent and Trademark Office may not be able to process and/or examine your submission, which may result in termination of proceedings or abandonment of the application or expiration of the patent.

The information provided by you in this form will be subject to the following routine uses:

1. The information on this form will be treated confidentially to the extent allowed under the Freedom of Information Act (5 U.S.C. 552) and the Privacy Act (5 U.S.C. 552a). Records from this system of records may be disclosed to the Department of Justice to determine whether the Freedom of Information Act requires disclosure of these records.
2. A record from this system of records may be disclosed, as a routine use, in the course of presenting evidence to a court, magistrate, or administrative tribunal, including disclosures to opposing counsel in the course of settlement negotiations.
3. A record in this system of records may be disclosed, as a routine use, to a Member of Congress submitting a request involving an individual, to whom the record pertains, when the individual has requested assistance from the Member with respect to the subject matter of the record.
4. A record in this system of records may be disclosed, as a routine use, to a contractor of the Agency having need for the information in order to perform a contract. Recipients of information shall be required to comply with the requirements of the Privacy Act of 1974, as amended, pursuant to 5 U.S.C. 552a(m).
5. A record related to an International Application filed under the Patent Cooperation Treaty in this system of records may be disclosed, as a routine use, to the International Bureau of the World Intellectual Property Organization, pursuant to the Patent Cooperation Treaty.
6. A record in this system of records may be disclosed, as a routine use, to another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c)).
7. A record from this system of records may be disclosed, as a routine use, to the Administrator, General Services, or his/her designee, during an inspection of records conducted by GSA as part of that agency's responsibility to recommend improvements in records management practices and programs, under authority of 44 U.S.C. 2904 and 2906. Such disclosure shall be made in accordance with the GSA regulations governing inspection of records for this purpose, and any other relevant (i.e., GSA or Commerce) directive. Such disclosure shall not be used to make determinations about individuals.
8. A record from this system of records may be disclosed, as a routine use, to the public after either publication of the application pursuant to 35 U.S.C. 122(b) or issuance of a patent pursuant to 35 U.S.C. 151. Further, a record may be disclosed, subject to the limitations of 37 CFR 1.14, as a routine use, to the public if the record was filed in an application which became abandoned or in which the proceedings were terminated and which application is referenced by either a published application, an application open to public inspections or an issued patent.
9. A record from this system of records may be disclosed, as a routine use, to a Federal, State, or local law enforcement agency, if the USPTO becomes aware of a violation or potential violation of law or regulation.



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC PAYMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 10/27/2025 04:02:51 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Colleen Mensching
PATENT CENTER #	72918266	AUTHORIZED BY	Victoria Cheng
CUSTOMER #	21971	FILING DATE	01/31/2025
CORRESPONDENCE ADDRESS	-	FIRST NAMED INVENTOR	Alexis HOWERTON

Payment Information

PAYMENT METHOD DA / 232415	PAYMENT TRANSACTION ID E20250QG03539961	PAYMENT AUTHORIZED BY Colleen Mensching
--------------------------------------	---	---

FEE CODE	DESCRIPTION	ITEM PRICE(\$)	QUANTITY	ITEM TOTAL(\$)
1801	REQUEST FOR CONTINUED EXAMINATION (RCE) - 1ST REQUEST (SEE 37 CFR 1.114)	1500.00	1	1500.00
TOTAL AMOUNT:				\$1,500.00

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C.

371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



UNITED STATES
PATENT AND TRADEMARK OFFICE

P.O. Box 1450
Alexandria, VA 22313 - 1450
www.uspto.gov

ELECTRONIC ACKNOWLEDGEMENT RECEIPT

APPLICATION # 19/043,264	RECEIPT DATE / TIME 10/27/2025 04:02:51 PM Z ET	ATTORNEY DOCKET # 50535-710.306
------------------------------------	---	---

Title of Invention

METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

Application Information

APPLICATION TYPE	Utility - Nonprovisional Application under 35 USC 111(a)	PATENT #	-
CONFIRMATION #	6961	FILED BY	Colleen Mensching
PATENT CENTER #	72918266	FILING DATE	01/31/2025
CUSTOMER #	21971	FIRST NAMED INVENTOR	Alexis HOWERTON
CORRESPONDENCE ADDRESS	-	AUTHORIZED BY	Victoria Cheng

Documents

TOTAL DOCUMENTS: 4

DOCUMENT	PAGES	DESCRIPTION	SIZE (KB)
Spruce 50535-710.306 - Response Final.pdf	6	-	222 KB
Spruce 50535-710.306 - Response Final-AMSB.pdf	(1-1) 1	Amendment Submitted/Entered with Filing of Continued Prosecution Application (CPA)/Request for Continued Examination(RCE)	106 KB
Spruce 50535-710.306 - Response Final-CLM.pdf	(2-4) 3	Claims	200 KB
Spruce 50535-710.306 - Response Final-REM.pdf	(5-6) 2	Applicant Arguments/Remarks Made in an Amendment	119 KB
Spruce 50535-710.306 - RCE Final.pdf	3	Request for Continued Examination (RCE)	1631 KB

Digest

DOCUMENT	MESSAGE DIGEST(SHA-512)
Spruce 50535-710.306 - Response Final.pdf	988E9F9091ED57FE9789D3BE6B553EC195176699737B63E0A 18A4647EE671F126A62B165675E832F136C0594B48B6EB8CB0 9B408F82505698C90F067D46FFBED
Spruce 50535-710.306 - Response Final-AMSB.pdf	D5A8D498AFF759FD6453DD9BF3E5A98CB61602D0BBAE4779 BEA897D439F8F701B6C6861F5C882ABD89DB1E37284BEFCD E4246593088D2E5FF40D6A71EE2E71F7
Spruce 50535-710.306 - Response Final-CLM.pdf	9507F20848740F1E90B5337022A8F93ABB78C99B34E2110D6A 4B2E34D130A2B8FF58D104910AB4626A80ECB29103B8F4D82 953013611DE4E9F41409F7C3C4734
Spruce 50535-710.306 - Response Final-REM.pdf	E5F0AD4CA96A234C5FDF0CC403EE19BC414734156E82342C 958F965B20FDE183F759174525450CD27584FD2B936745046B 786A3D8ECF78FC470057B7CA96E2E6
Spruce 50535-710.306 - RCE Final.pdf	9AA69C8A363175AD2AAE811F6D8D383267DA9DD86E37A0B9 0C99489947399C24737C92743A462AFFB171477718BFD8568A 332E12C957EDA8F074163771F8052A

This Acknowledgement Receipt evidences receipt on the noted date by the USPTO of the indicated documents, characterized by the applicant, and including page counts, where applicable. It serves as evidence of receipt similar to a Post Card, as described in MPEP 503.

New Applications Under 35 U.S.C. 111

If a new application is being filed and the application includes the necessary components for filing date (see 37 CFR 1.53(b)-(d) and MPEP 506), a Filing Receipt (37 CFR 1.54) will be issued in due course and the date shown on this Acknowledgement Receipt will establish the filing date of the application

National Stage of an International Application under 35 U.S.C. 371

If a timely submission to enter the national stage of an international application is compliant with the conditions of 35 U.S.C. 371 and other applicable requirements a Form PCT/DO/EO/903 indicating acceptance of the application as a national stage submission under 35 U.S.C. 371 will be issued in addition to the Filing Receipt, in due course.

New International Application Filed with the USPTO as a Receiving Office

If a new international application is being filed and the international application includes the necessary components for an international filing date (see PCT Article 11 and MPEP 1810), a Notification of the International Application Number and of the International Filing Date (Form PCT/RO/105) will be issued in due course, subject to prescriptions concerning national security, and the date shown on this Acknowledgement Receipt will establish the international filing date of the application.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

NOTICE OF ALLOWANCE AND FEE(S) DUE

21971 7590 11/17/2025
WILSON SONSINI GOODRICH & ROSATI
650 PAGE MILL ROAD
PALO ALTO, CA 94304-1050

EXAMINER

PIHONAK, SARAH

ART UNIT

PAPER NUMBER

1627

DATE MAILED: 11/17/2025

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961

TITLE OF INVENTION: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

APPLN. TYPE	ENTITY STATUS	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
nonprovisional	UNDISCOUNTED	\$1290	\$0.00	\$0.00	\$1290	02/17/2026

THE APPLICATION IDENTIFIED ABOVE HAS BEEN EXAMINED AND IS ALLOWED FOR ISSUANCE AS A PATENT. PROSECUTION ON THE MERITS IS CLOSED. THIS NOTICE OF ALLOWANCE IS NOT A GRANT OF PATENT RIGHTS. THIS APPLICATION IS SUBJECT TO WITHDRAWAL FROM ISSUE AT THE INITIATIVE OF THE OFFICE OR UPON PETITION BY THE APPLICANT. SEE 37 CFR 1.313 AND MPEP 1308.

THE ISSUE FEE AND PUBLICATION FEE (IF REQUIRED) MUST BE PAID WITHIN THREE MONTHS FROM THE MAILING DATE OF THIS NOTICE OR THIS APPLICATION SHALL BE REGARDED AS ABANDONED. THIS STATUTORY PERIOD CANNOT BE EXTENDED. SEE 35 U.S.C. 151. THE ISSUE FEE DUE INDICATED ABOVE DOES NOT REFLECT A CREDIT FOR ANY PREVIOUSLY PAID ISSUE FEE IN THIS APPLICATION. IF AN ISSUE FEE HAS PREVIOUSLY BEEN PAID IN THIS APPLICATION (AS SHOWN ABOVE), THE RETURN OF PART B OF THIS FORM WILL BE CONSIDERED A REQUEST TO REAPPLY THE PREVIOUSLY PAID ISSUE FEE TOWARD THE ISSUE FEE NOW DUE.

HOW TO REPLY TO THIS NOTICE:

I. Review the ENTITY STATUS shown above. If the ENTITY STATUS is shown as SMALL or MICRO, verify whether entitlement to that entity status still applies.

If the ENTITY STATUS is the same as shown above, pay the TOTAL FEE(S) DUE shown above.

If the ENTITY STATUS is changed from that shown above, on PART B - FEE(S) TRANSMITTAL, complete section number 5 titled "Change in Entity Status (from status indicated above)".

For purposes of this notice, small entity fees are 40% the amount of undiscounted fees, and micro entity fees are 20% the amount of undiscounted fees.

II. PART B - FEE(S) TRANSMITTAL, or its equivalent, must be completed and returned to the United States Patent and Trademark Office (USPTO) with your ISSUE FEE and PUBLICATION FEE (if required). If you are charging the fee(s) to your deposit account, section "4b" of Part B - Fee(s) Transmittal should be completed. If an equivalent of Part B is filed, a request to reapply a previously paid issue fee must be clearly made, and delays in processing may occur due to the difficulty in recognizing the paper as an equivalent of Part B.

III. All communications regarding this application must give the application number. Please direct all communications prior to issuance to Mail Stop ISSUE FEE unless advised to the contrary.

IMPORTANT REMINDER: Maintenance fees are due in utility patents issuing on applications filed on or after Dec. 12, 1980. It is patentee's responsibility to ensure timely payment of maintenance fees when due. More information is available at www.uspto.gov/PatentMaintenanceFees.

PART B - FEE(S) TRANSMITTAL

Complete and send this form, together with applicable fee(s), by mail or fax, or via the USPTO patent electronic filing system.

By mail, send to: Mail Stop ISSUE FEE
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

By fax, send to: (571)-273-2885

INSTRUCTIONS: This form should be used for transmitting the ISSUE FEE and PUBLICATION FEE (if required). Blocks 1 through 5 should be completed where appropriate. All further correspondence will be mailed to the current correspondence address as indicated unless corrected below or directed otherwise in Block 1, by (a) specifying a new correspondence address; and/or (b) indicating a separate "FEE ADDRESS" for maintenance fee notifications. **Because electronic patent issuance may occur shortly after issue fee payment, any desired continuing application should preferably be filed prior to payment of this issue fee in order not to jeopardize copendency.**

CURRENT CORRESPONDENCE ADDRESS (Note: Use Block 1 for any change of address)

21971 7590 11/17/2025
WILSON SONSINI GOODRICH & ROSATI
650 PAGE MILL ROAD
PALO ALTO, CA 94304-1050

Note: A certificate of mailing can only be used for domestic mailings of the Fee(s) Transmittal. This certificate cannot be used for any other accompanying papers. Each additional paper, such as an assignment or formal drawing, must have its own certificate of mailing or transmission.

Certificate of Mailing or Transmission

I hereby certify that this Fee(s) Transmittal is being deposited with the United States Postal Service with sufficient postage for first class mail in an envelope addressed to the Mail Stop ISSUE FEE address above, or being transmitted to the USPTO via the USPTO patent electronic filing system or by facsimile to (571) 273-2885, on the date below.

(Typed or printed name)
(Signature)
(Date)

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961

TITLE OF INVENTION: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

APPLN. TYPE	ENTITY STATUS	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
nonprovisional	UNDISCOUNTED	\$1290	\$0.00	\$0.00	\$1290	02/17/2026

EXAMINER	ART UNIT	CLASS-SUBCLASS
PIHONAK, SARAH	1627	514-233200

1. Change of correspondence address or indication of "Fee Address" (37 CFR 1.363).

☐ Change of correspondence address (or Change of Correspondence Address form PTO/AIA/122 or PTO/SB/122) attached.

☐ "Fee Address" indication (or "Fee Address" Indication form PTO/AIA/47 or PTO/SB/47; Rev 03-02 or more recent) attached. **Use of a Customer Number is required.**

2. For printing on the patent front page, list

(1) The names of up to 3 registered patent attorneys or agents OR, alternatively,

(2) The name of a single firm (having as a member a registered attorney or agent) and the names of up to 2 registered patent attorneys or agents. If no name is listed, no name will be printed.

1 _____
2 _____
3 _____

3. ASSIGNEE NAME AND RESIDENCE DATA TO BE PRINTED ON THE PATENT (print or type)

PLEASE NOTE: Unless an assignee is identified below, no assignee data will appear on the patent. If an assignee is identified below, the document must have been previously recorded, or filed for recordation, as set forth in 37 CFR 3.11 and 37 CFR 3.81(a). Completion of this form is NOT a substitute for filing an assignment.

(A) NAME OF ASSIGNEE

(B) RESIDENCE: (CITY and STATE OR COUNTRY)

Please check the appropriate assignee category or categories (will not be printed on the patent): ☐ Individual ☐ Corporation or other private group entity ☐ Government

4a. Fees submitted: ☐ Issue Fee ☐ Publication Fee (if required)

4b. Method of Payment: (Please first reapply any previously paid fee shown above)

☐ Electronic Payment via the USPTO patent electronic filing system ☐ Enclosed check ☐ Non-electronic payment by credit card (Attach form PTO-2038)

☐ The Director is hereby authorized to charge the required fee(s), any deficiency, or credit any overpayment to Deposit Account No. _____

5. **Change in Entity Status** (from status indicated above)

☐ Applicant certifying micro entity status. See 37 CFR 1.29

☐ Applicant asserting small entity status. See 37 CFR 1.27

☐ Applicant changing to regular undiscounted fee status.

NOTE: Absent a valid certification of Micro Entity Status (see forms PTO/SB/15A and 15B), issue fee payment in the micro entity amount will not be accepted at the risk of application abandonment.

NOTE: If the application was previously under micro entity status, checking this box will be taken to be a notification of loss of entitlement to micro entity status.

NOTE: Checking this box will be taken to be a notification of loss of entitlement to small or micro entity status, as applicable.

NOTE: This form must be signed in accordance with 37 CFR 1.31 and 1.33. See 37 CFR 1.4 for signature requirements and certifications.

Authorized Signature _____

Date _____

Typed or printed name _____

Registration No. _____



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961
21971 7590 11/17/2025 WILSON SONSINI GOODRICH & ROSATI 650 PAGE MILL ROAD PALO ALTO, CA 94304-1050			EXAMINER PIHONAK, SARAH	
			ART UNIT 1627	PAPER NUMBER
DATE MAILED: 11/17/2025				

Determination of Patent Term Adjustment under 35 U.S.C. 154 (b) (Applications filed on or after May 29, 2000)

The Office has discontinued providing a Patent Term Adjustment (PTA) calculation with the Notice of Allowance.

Section 1(h)(2) of the AIA Technical Corrections Act amended 35 U.S.C. 154(b)(3)(B)(i) to eliminate the requirement that the Office provide a patent term adjustment determination with the notice of allowance. See Revisions to Patent Term Adjustment, 78 Fed. Reg. 19416, 19417 (Apr. 1, 2013). Therefore, the Office is no longer providing an initial patent term adjustment determination with the notice of allowance. The Office will continue to provide a patent term adjustment determination with the Issue Notification Letter that is mailed to applicant approximately three weeks prior to the issue date of the patent, and will include the patent term adjustment on the patent. Any request for reconsideration of the patent term adjustment determination (or reinstatement of patent term adjustment) should follow the process outlined in 37 CFR 1.705.

Any questions regarding the Patent Term Extension or Adjustment determination should be directed to the Office of Patent Legal Administration at (571)-272-7702. Questions relating to issue and publication fee payments should be directed to the Customer Service Center of the Office of Patent Publication at 1-(888)-786-0101 or (571)-272-4200.

<i>Notice Requiring Inventor's Oath or Declaration</i>	Application No. 19/043,264	Applicant(s) Alexis HOWERTON	
	Examiner PIHONAK, SARAH	Art Unit 1627	

This notice is an attachment to the Notice of Allowability (PTOL-37), or the Notice of Allowability For A Design Application (PTOL-37D).

An inventor's oath or declaration in compliance with 37 CFR 1.63 or 1.64 executed by or with respect to each inventor has not yet been submitted.

An oath or declaration in compliance with 37 CFR 1.63, or a substitute statement in compliance with 37 CFR 1.64, executed by or with respect to each inventor (for any inventor for which a compliant oath, declaration, or substitute statement has not yet been submitted) **MUST** be filed no later than the date on which the issue fee is paid. See 35 U.S.C. 115(f). Failure to timely comply will result in ABANDONMENT of this application.

A properly executed inventor's oath to declaration has not been received for the following inventor(s):

If applicant previously filed one or more oaths, declarations, or substitute statements, applicant may have received an informational notice regarding deficiencies therein.

The following deficiencies are noted:

INFORMAL ACTION PROBLEMS

- A properly executed inventor's oath or declaration has not been received for the following inventor(s): **Alexis HOWERTON** and **Hal GERBER**.

Applicant may submit the inventor's oath or declaration at any time before the Notice of Allowance and Fee(s) Due, PTOL-85, is mailed.

Questions relating to this Notice should be directed to the Application Assistance Unit at 571-272-4200.

OMB Clearance and PRA Burden Statement for PTOL-85 Part B

The Paperwork Reduction Act (PRA) of 1995 requires Federal agencies to obtain Office of Management and Budget approval before requesting most types of information from the public. When OMB approves an agency request to collect information from the public, OMB (i) provides a valid OMB Control Number and expiration date for the agency to display on the instrument that will be used to collect the information and (ii) requires the agency to inform the public about the OMB Control Number's legal significance in accordance with 5 CFR 1320.5(b).

The information collected by PTOL-85 Part B is required by 37 CFR 1.311. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 30 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, Virginia 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450. Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. The United States Patent and Trademark Office (USPTO) collects the information in this record under authority of 35 U.S.C. 2. The USPTO's system of records is used to manage all applicant and owner information including name, citizenship, residence, post office address, and other information with respect to inventors and their legal representatives pertaining to the applicant's/owner's activities in connection with the invention for which a patent is sought or has been granted. The applicable Privacy Act System of Records Notice for the information collected in this form is COMMERCE/PAT-TM-7 Patent Application Files, available in the Federal Register at 78 FR 19243 (March 29, 2013).

<https://www.govinfo.gov/content/pkg/FR-2013-03-29/pdf/2013-07341.pdf>

Routine uses of the information in this record may include disclosure to:

- 1) law enforcement, in the event that the system of records indicates a violation or potential violation of law;
- 2) a federal, state, local, or international agency, in response to its request;
- 3) a contractor of the USPTO having need for the information in order to perform a contract;
- 4) the Department of Justice for determination of whether the Freedom of Information Act (FOIA) requires disclosure of the record;
- 5) a Member of Congress submitting a request involving an individual to whom the record pertains, when the individual has requested the Member's assistance with respect to the subject matter of the record;
- 6) a court, magistrate, or administrative tribunal, in the course of presenting evidence, including disclosures to opposing counsel in the course of settlement negotiations;
- 7) the Administrator, General Services Administration (GSA), or their designee, during an inspection of records conducted by GSA under authority of 44 U.S.C. 2904 and 2906, in accordance with the GSA regulations and any other relevant (i.e., GSA or Commerce) directive, where such disclosure shall not be used to make determinations about individuals;
- 8) another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c));
- 9) the Office of Personnel Management (OPM) for personnel research purposes; and
- 10) the Office of Management and Budget (OMB) for legislative coordination and clearance.

If you do not furnish the information requested on this form, the USPTO may not be able to process and/or examine your submission, which may result in termination of proceedings, abandonment of the application, and/or expiration of the patent.

Notice of Allowability	Application No. 19/043,264	Applicant(s) HOWERTON et al.	
	Examiner SARAH PIHONAK	Art Unit 1627	AIA (First Inventor to File) Status Yes

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address--

All claims being allowable, PROSECUTION ON THE MERITS IS (OR REMAINS) CLOSED in this application. If not included herewith (or previously mailed), a Notice of Allowance (PTOL-85) or other appropriate communication will be mailed in due course. **THIS NOTICE OF ALLOWABILITY IS NOT A GRANT OF PATENT RIGHTS.** This application is subject to withdrawal from issue at the initiative of the Office or upon petition by the applicant. See 37 CFR 1.313 and MPEP 1308.

1. ☒ This communication is responsive to 10/27/25.
☐ A declaration(s)/affidavit(s) under **37 CFR 1.130(b)** was/were filed on _____.

2. ☐ An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.

3. ☒ The allowed claim(s) is/are 1-21. As a result of the allowed claim(s), you may be eligible to benefit from the **Patent Prosecution Highway** program at a participating intellectual property office for the corresponding application. For more information, please see http://www.uspto.gov/patents/init_events/pph/index.jsp or send an inquiry to PPHfeedback@uspto.gov.

4. ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
Certified copies:
a) ☐ All b) ☐ Some* c) ☐ None of the:
1. ☐ Certified copies of the priority documents have been received.
2. ☐ Certified copies of the priority documents have been received in Application No. _____.
3. ☐ Copies of the certified copies of the priority documents have been received in this national stage application from the International Bureau (PCT Rule 17.2(a)).
* Certified copies not received: _____.

Applicant has THREE MONTHS FROM THE "MAILING DATE" of this communication to file a reply complying with the requirements noted below. Failure to timely comply will result in ABANDONMENT of this application.
THIS THREE-MONTH PERIOD IS NOT EXTENDABLE.

5. ☐ CORRECTED DRAWINGS (as "replacement sheets") must be submitted.
☐ including changes required by the attached Examiner's Amendment / Comment or in the Office action of Paper No./Mail Date _____.

Identifying indicia such as the application number (see 37 CFR 1.84(c)) should be written on the drawings in the front (not the back) of each sheet. Replacement sheet(s) should be labeled as such in the header according to 37 CFR 1.121(d).

6. ☐ DEPOSIT OF and/or INFORMATION about the deposit of BIOLOGICAL MATERIAL must be submitted. Note the attached Examiner's comment regarding REQUIREMENT FOR THE DEPOSIT OF BIOLOGICAL MATERIAL.

Attachment(s)

1. ☐ Notice of References Cited (PTO-892)	5. ☐ Examiner's Amendment/Comment
2. ☒ Information Disclosure Statements (PTO/SB/08), Paper No./Mail Date _____.	6. ☒ Examiner's Statement of Reasons for Allowance
3. ☐ Examiner's Comment Regarding Requirement for Deposit of Biological Material _____.	7. ☐ Other _____.
4. ☐ Interview Summary (PTO-413), Paper No./Mail Date. _____.	

/SARAH PIHONAK/ Primary Examiner, Art Unit 1627	
--	--

Notice of Pre-AIA or AIA Status

The present application, filed on or after March 16, 2013, is being examined under the first inventor to file provisions of the AIA.

A request for continued examination under 37 CFR 1.114, including the fee set forth in 37 CFR 1.17(e), was filed in this application after final rejection. Since this application is eligible for continued examination under 37 CFR 1.114, and the fee set forth in 37 CFR 1.17(e) has been timely paid, the finality of the previous Office action has been withdrawn pursuant to 37 CFR 1.114. Applicant's submission filed on 10/27/25 has been entered.

Status of Claims

1. Claims 1-21 are pending as of the response filed on 10/27/25.

The rejections for nonstatutory double patenting over the claims of US 11304950, 18941832, 19043275, and 19043303 are withdrawn in consideration of the acceptance of the terminal disclaimer filed on 10/27/25.

2. Claims 1-21 are allowed.

Reasons for Allowance

3. The following is an examiner's statement of reasons for allowance: Grigoriadis, WO 2015112642 A1, of record, represents the closest prior art. Grigoriadis teaches CRF1 antagonists that include those recited in the instant claims for the treatment of congenital

adrenal hyperplasia, however, treatment of a subject having an ovarian adrenal rest tumor is not taught or suggested.

Any comments considered necessary by applicant must be submitted no later than the payment of the issue fee and, to avoid processing delays, should preferably accompany the issue fee. Such submissions should be clearly labeled "Comments on Statement of Reasons for Allowance."

Information Disclosure Statement

4. The IDS filed on 10/27/25 has been considered.

Correspondence

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SARAH PIHONAK whose telephone number is (571)270-7710. The examiner can normally be reached Monday-Friday 9:00-5:30 EST.

Examiner interviews are available via telephone, in-person, and video conferencing using a USPTO supplied web-based collaboration tool. To schedule an interview, applicant is encouraged to use the USPTO Automated Interview Request (AIR) at <http://www.uspto.gov/interviewpractice>.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Kortney Klinkel can be reached at 571-270-5239. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of published or unpublished applications may be obtained from Patent Center. Unpublished application information in Patent Center is available to registered users. To file and manage patent submissions in Patent Center, visit: <https://patentcenter.uspto.gov>. Visit <https://www.uspto.gov/patents/apply/patent-center> for more information about Patent Center and <https://www.uspto.gov/patents/docx> for information about filing in DOCX format. For additional questions, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

SARAH . PIHONAK
Primary Examiner
Art Unit 1627

/SARAH PIHONAK/
Primary Examiner, Art Unit 1627



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

NOTICE OF ALLOWANCE AND FEE(S) DUE

21971 7590 01/06/2026
WILSON SONSINI GOODRICH & ROSATI
650 PAGE MILL ROAD
PALO ALTO, CA 94304-1050

EXAMINER

PIHONAK, SARAH

ART UNIT

PAPER NUMBER

1627

DATE MAILED: 01/06/2026

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961

TITLE OF INVENTION: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

APPLN. TYPE	ENTITY STATUS	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
nonprovisional	UNDISCOUNTED	\$1290	\$0.00	\$0.00	\$1290	04/06/2026

THE APPLICATION IDENTIFIED ABOVE HAS BEEN EXAMINED AND IS ALLOWED FOR ISSUANCE AS A PATENT. PROSECUTION ON THE MERITS IS CLOSED. THIS NOTICE OF ALLOWANCE IS NOT A GRANT OF PATENT RIGHTS. THIS APPLICATION IS SUBJECT TO WITHDRAWAL FROM ISSUE AT THE INITIATIVE OF THE OFFICE OR UPON PETITION BY THE APPLICANT. SEE 37 CFR 1.313 AND MPEP 1308.

THE ISSUE FEE AND PUBLICATION FEE (IF REQUIRED) MUST BE PAID WITHIN THREE MONTHS FROM THE MAILING DATE OF THIS NOTICE OR THIS APPLICATION SHALL BE REGARDED AS ABANDONED. THIS STATUTORY PERIOD CANNOT BE EXTENDED. SEE 35 U.S.C. 151. THE ISSUE FEE DUE INDICATED ABOVE DOES NOT REFLECT A CREDIT FOR ANY PREVIOUSLY PAID ISSUE FEE IN THIS APPLICATION. IF AN ISSUE FEE HAS PREVIOUSLY BEEN PAID IN THIS APPLICATION (AS SHOWN ABOVE), THE RETURN OF PART B OF THIS FORM WILL BE CONSIDERED A REQUEST TO REAPPLY THE PREVIOUSLY PAID ISSUE FEE TOWARD THE ISSUE FEE NOW DUE.

HOW TO REPLY TO THIS NOTICE:

I. Review the ENTITY STATUS shown above. If the ENTITY STATUS is shown as SMALL or MICRO, verify whether entitlement to that entity status still applies.

If the ENTITY STATUS is the same as shown above, pay the TOTAL FEE(S) DUE shown above.

If the ENTITY STATUS is changed from that shown above, on PART B - FEE(S) TRANSMITTAL, complete section number 5 titled "Change in Entity Status (from status indicated above)".

For purposes of this notice, small entity fees are 40% the amount of undiscounted fees, and micro entity fees are 20% the amount of undiscounted fees.

II. PART B - FEE(S) TRANSMITTAL, or its equivalent, must be completed and returned to the United States Patent and Trademark Office (USPTO) with your ISSUE FEE and PUBLICATION FEE (if required). If you are charging the fee(s) to your deposit account, section "4b" of Part B - Fee(s) Transmittal should be completed. If an equivalent of Part B is filed, a request to reapply a previously paid issue fee must be clearly made, and delays in processing may occur due to the difficulty in recognizing the paper as an equivalent of Part B.

III. All communications regarding this application must give the application number. Please direct all communications prior to issuance to Mail Stop ISSUE FEE unless advised to the contrary.

IMPORTANT REMINDER: Maintenance fees are due in utility patents issuing on applications filed on or after Dec. 12, 1980. It is patentee's responsibility to ensure timely payment of maintenance fees when due. More information is available at www.uspto.gov/PatentMaintenanceFees.

PART B - FEE(S) TRANSMITTAL

Complete and send this form, together with applicable fee(s), by mail or fax, or via the USPTO patent electronic filing system.

By mail, send to: Mail Stop ISSUE FEE
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

By fax, send to: (571)-273-2885

INSTRUCTIONS: This form should be used for transmitting the ISSUE FEE and PUBLICATION FEE (if required). Blocks 1 through 5 should be completed where appropriate. All further correspondence will be mailed to the current correspondence address as indicated unless corrected below or directed otherwise in Block 1, by (a) specifying a new correspondence address; and/or (b) indicating a separate "FEE ADDRESS" for maintenance fee notifications. **Because electronic patent issuance may occur shortly after issue fee payment, any desired continuing application should preferably be filed prior to payment of this issue fee in order not to jeopardize copendency.**

CURRENT CORRESPONDENCE ADDRESS (Note: Use Block 1 for any change of address)

21971 7590 01/06/2026
WILSON SONSINI GOODRICH & ROSATI
650 PAGE MILL ROAD
PALO ALTO, CA 94304-1050

Note: A certificate of mailing can only be used for domestic mailings of the Fee(s) Transmittal. This certificate cannot be used for any other accompanying papers. Each additional paper, such as an assignment or formal drawing, must have its own certificate of mailing or transmission.

Certificate of Mailing or Transmission

I hereby certify that this Fee(s) Transmittal is being deposited with the United States Postal Service with sufficient postage for first class mail in an envelope addressed to the Mail Stop ISSUE FEE address above, or being transmitted to the USPTO via the USPTO patent electronic filing system or by facsimile to (571) 273-2885, on the date below.

(Typed or printed name)
(Signature)
(Date)

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961

TITLE OF INVENTION: METHODS FOR TREATING TESTICULAR AND OVARIAN ADRENAL REST TUMORS

APPLN. TYPE	ENTITY STATUS	ISSUE FEE DUE	PUBLICATION FEE DUE	PREV. PAID ISSUE FEE	TOTAL FEE(S) DUE	DATE DUE
nonprovisional	UNDISCOUNTED	\$1290	\$0.00	\$0.00	\$1290	04/06/2026

EXAMINER	ART UNIT	CLASS-SUBCLASS
PIHONAK, SARAH	1627	514-233200

1. Change of correspondence address or indication of "Fee Address" (37 CFR 1.363).

☐ Change of correspondence address (or Change of Correspondence Address form PTO/AIA/122 or PTO/SB/122) attached.

☐ "Fee Address" indication (or "Fee Address" Indication form PTO/AIA/47 or PTO/SB/47; Rev 03-02 or more recent) attached. **Use of a Customer Number is required.**

2. For printing on the patent front page, list

(1) The names of up to 3 registered patent attorneys or agents OR, alternatively,

(2) The name of a single firm (having as a member a registered attorney or agent) and the names of up to 2 registered patent attorneys or agents. If no name is listed, no name will be printed.

1 _____

2 _____

3 _____

3. ASSIGNEE NAME AND RESIDENCE DATA TO BE PRINTED ON THE PATENT (print or type)

PLEASE NOTE: Unless an assignee is identified below, no assignee data will appear on the patent. If an assignee is identified below, the document must have been previously recorded, or filed for recordation, as set forth in 37 CFR 3.11 and 37 CFR 3.81(a). Completion of this form is NOT a substitute for filing an assignment.

(A) NAME OF ASSIGNEE

(B) RESIDENCE: (CITY and STATE OR COUNTRY)

Please check the appropriate assignee category or categories (will not be printed on the patent): ☐ Individual ☐ Corporation or other private group entity ☐ Government

4a. Fees submitted: ☐ Issue Fee ☐ Publication Fee (if required)

4b. Method of Payment: (Please first reapply any previously paid fee shown above)

☐ Electronic Payment via the USPTO patent electronic filing system ☐ Enclosed check ☐ Non-electronic payment by credit card (Attach form PTO-2038)

☐ The Director is hereby authorized to charge the required fee(s), any deficiency, or credit any overpayment to Deposit Account No. _____

5. **Change in Entity Status** (from status indicated above)

☐ Applicant certifying micro entity status. See 37 CFR 1.29

☐ Applicant asserting small entity status. See 37 CFR 1.27

☐ Applicant changing to regular undiscounted fee status.

NOTE: Absent a valid certification of Micro Entity Status (see forms PTO/SB/15A and 15B), issue fee payment in the micro entity amount will not be accepted at the risk of application abandonment.

NOTE: If the application was previously under micro entity status, checking this box will be taken to be a notification of loss of entitlement to micro entity status.

NOTE: Checking this box will be taken to be a notification of loss of entitlement to small or micro entity status, as applicable.

NOTE: This form must be signed in accordance with 37 CFR 1.31 and 1.33. See 37 CFR 1.4 for signature requirements and certifications.

Authorized Signature _____

Date _____

Typed or printed name _____

Registration No. _____



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961
21971 7590 01/06/2026 WILSON SONSINI GOODRICH & ROSATI 650 PAGE MILL ROAD PALO ALTO, CA 94304-1050			EXAMINER PIHONAK, SARAH	
			ART UNIT 1627	PAPER NUMBER
DATE MAILED: 01/06/2026				

Determination of Patent Term Adjustment under 35 U.S.C. 154 (b) (Applications filed on or after May 29, 2000)

The Office has discontinued providing a Patent Term Adjustment (PTA) calculation with the Notice of Allowance.

Section 1(h)(2) of the AIA Technical Corrections Act amended 35 U.S.C. 154(b)(3)(B)(i) to eliminate the requirement that the Office provide a patent term adjustment determination with the notice of allowance. See Revisions to Patent Term Adjustment, 78 Fed. Reg. 19416, 19417 (Apr. 1, 2013). Therefore, the Office is no longer providing an initial patent term adjustment determination with the notice of allowance. The Office will continue to provide a patent term adjustment determination with the Issue Notification Letter that is mailed to applicant approximately three weeks prior to the issue date of the patent, and will include the patent term adjustment on the patent. Any request for reconsideration of the patent term adjustment determination (or reinstatement of patent term adjustment) should follow the process outlined in 37 CFR 1.705.

Any questions regarding the Patent Term Extension or Adjustment determination should be directed to the Office of Patent Legal Administration at (571)-272-7702. Questions relating to issue and publication fee payments should be directed to the Customer Service Center of the Office of Patent Publication at 1-(888)-786-0101 or (571)-272-4200.

OMB Clearance and PRA Burden Statement for PTOL-85 Part B

The Paperwork Reduction Act (PRA) of 1995 requires Federal agencies to obtain Office of Management and Budget approval before requesting most types of information from the public. When OMB approves an agency request to collect information from the public, OMB (i) provides a valid OMB Control Number and expiration date for the agency to display on the instrument that will be used to collect the information and (ii) requires the agency to inform the public about the OMB Control Number's legal significance in accordance with 5 CFR 1320.5(b).

The information collected by PTOL-85 Part B is required by 37 CFR 1.311. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 30 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, Virginia 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450. Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Privacy Act Statement

The Privacy Act of 1974 (P.L. 93-579) requires that you be given certain information in connection with your submission of the attached form related to a patent application or patent. The United States Patent and Trademark Office (USPTO) collects the information in this record under authority of 35 U.S.C. 2. The USPTO's system of records is used to manage all applicant and owner information including name, citizenship, residence, post office address, and other information with respect to inventors and their legal representatives pertaining to the applicant's/owner's activities in connection with the invention for which a patent is sought or has been granted. The applicable Privacy Act System of Records Notice for the information collected in this form is COMMERCE/PAT-TM-7 Patent Application Files, available in the Federal Register at 78 FR 19243 (March 29, 2013).

<https://www.govinfo.gov/content/pkg/FR-2013-03-29/pdf/2013-07341.pdf>

Routine uses of the information in this record may include disclosure to:

- 1) law enforcement, in the event that the system of records indicates a violation or potential violation of law;
- 2) a federal, state, local, or international agency, in response to its request;
- 3) a contractor of the USPTO having need for the information in order to perform a contract;
- 4) the Department of Justice for determination of whether the Freedom of Information Act (FOIA) requires disclosure of the record;
- 5) a Member of Congress submitting a request involving an individual to whom the record pertains, when the individual has requested the Member's assistance with respect to the subject matter of the record;
- 6) a court, magistrate, or administrative tribunal, in the course of presenting evidence, including disclosures to opposing counsel in the course of settlement negotiations;
- 7) the Administrator, General Services Administration (GSA), or their designee, during an inspection of records conducted by GSA under authority of 44 U.S.C. 2904 and 2906, in accordance with the GSA regulations and any other relevant (i.e., GSA or Commerce) directive, where such disclosure shall not be used to make determinations about individuals;
- 8) another federal agency for purposes of National Security review (35 U.S.C. 181) and for review pursuant to the Atomic Energy Act (42 U.S.C. 218(c));
- 9) the Office of Personnel Management (OPM) for personnel research purposes; and
- 10) the Office of Management and Budget (OMB) for legislative coordination and clearance.

If you do not furnish the information requested on this form, the USPTO may not be able to process and/or examine your submission, which may result in termination of proceedings, abandonment of the application, and/or expiration of the patent.

Notice of Allowability	Application No. 19/043,264	Applicant(s) HOWERTON et al.	
	Examiner SARAH PIHONAK	Art Unit 1627	AIA (First Inventor to File) Status Yes

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address--

All claims being allowable, PROSECUTION ON THE MERITS IS (OR REMAINS) CLOSED in this application. If not included herewith (or previously mailed), a Notice of Allowance (PTOL-85) or other appropriate communication will be mailed in due course. **THIS NOTICE OF ALLOWABILITY IS NOT A GRANT OF PATENT RIGHTS.** This application is subject to withdrawal from issue at the initiative of the Office or upon petition by the applicant. See 37 CFR 1.313 and MPEP 1308.

1. ☒ This communication is responsive to 12/10/25.
☐ A declaration(s)/affidavit(s) under **37 CFR 1.130(b)** was/were filed on _____.

2. ☐ An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.

3. ☒ The allowed claim(s) is/are 1-21. As a result of the allowed claim(s), you may be eligible to benefit from the **Patent Prosecution Highway** program at a participating intellectual property office for the corresponding application. For more information, please see http://www.uspto.gov/patents/init_events/pph/index.jsp or send an inquiry to PPHfeedback@uspto.gov.

4. ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
Certified copies:
a) ☐ All b) ☐ Some* c) ☐ None of the:
1. ☐ Certified copies of the priority documents have been received.
2. ☐ Certified copies of the priority documents have been received in Application No. _____.
3. ☐ Copies of the certified copies of the priority documents have been received in this national stage application from the International Bureau (PCT Rule 17.2(a)).
* Certified copies not received: _____.

Applicant has THREE MONTHS FROM THE "MAILING DATE" of this communication to file a reply complying with the requirements noted below. Failure to timely comply will result in ABANDONMENT of this application.
THIS THREE-MONTH PERIOD IS NOT EXTENDABLE.

5. ☐ CORRECTED DRAWINGS (as "replacement sheets") must be submitted.
☐ including changes required by the attached Examiner's Amendment / Comment or in the Office action of Paper No./Mail Date _____.

Identifying indicia such as the application number (see 37 CFR 1.84(c)) should be written on the drawings in the front (not the back) of each sheet. Replacement sheet(s) should be labeled as such in the header according to 37 CFR 1.121(d).

6. ☐ DEPOSIT OF and/or INFORMATION about the deposit of BIOLOGICAL MATERIAL must be submitted. Note the attached Examiner's comment regarding REQUIREMENT FOR THE DEPOSIT OF BIOLOGICAL MATERIAL.

Attachment(s)

1. ☐ Notice of References Cited (PTO-892)	5. ☐ Examiner's Amendment/Comment
2. ☒ Information Disclosure Statements (PTO/SB/08), Paper No./Mail Date _____.	6. ☒ Examiner's Statement of Reasons for Allowance
3. ☐ Examiner's Comment Regarding Requirement for Deposit of Biological Material _____.	7. ☐ Other _____.
4. ☐ Interview Summary (PTO-413), Paper No./Mail Date. _____.	

/SARAH PIHONAK/ Primary Examiner, Art Unit 1627	
--	--

Notice of Pre-AIA or AIA Status

The present application, filed on or after March 16, 2013, is being examined under the first inventor to file provisions of the AIA.

A request for continued examination under 37 CFR 1.114, including the fee set forth in 37 CFR 1.17(e), was filed in this application after allowance or after an Office action under *Ex Parte Quayle*, 25 USPQ 74, 453 O.G. 213 (Comm'r Pat. 1935). Since this application is eligible for continued examination under 37 CFR 1.114, and the fee set forth in 37 CFR 1.17(e) has been timely paid, prosecution in this application has been reopened pursuant to 37 CFR 1.114. Applicant's submission filed on 12/10/25 has been entered.

Status of Claims

1. Claims 1-21 were allowed for reasons noted in the notice of allowance mailed on 11/17/25.

Information Disclosure Statement

2. The IDS filed on 12/10/25 has been considered.

Correspondence

Any inquiry concerning this communication or earlier communications from the examiner should be directed to SARAH PIHONAK whose telephone number is (571)270-7710. The examiner can normally be reached Monday-Friday 9:00-5:30 EST.

Examiner interviews are available via telephone, in-person, and video conferencing using a USPTO supplied web-based collaboration tool. To schedule an interview, applicant is encouraged to use the USPTO Automated Interview Request (AIR) at <http://www.uspto.gov/interviewpractice>.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Kortney Klinkel can be reached at 571-270-5239. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Information regarding the status of published or unpublished applications may be obtained from Patent Center. Unpublished application information in Patent Center is available to registered users. To file and manage patent submissions in Patent Center, visit: <https://patentcenter.uspto.gov>. Visit <https://www.uspto.gov/patents/apply/patent-center> for more information about Patent Center and <https://www.uspto.gov/patents/docx> for information about filing in DOCX format. For additional questions, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

SARAH . PIHONAK
Primary Examiner
Art Unit 1627

/SARAH PIHONAK/
Primary Examiner, Art Unit 1627



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER FOR PATENTS
P.O. Box 1450
Alexandria, Virginia 22313-1450
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
19/043,264	01/31/2025	Alexis HOWERTON	50535-710.306	6961
21971 7590 01/16/2026 WILSON SONSINI GOODRICH & ROSATI 650 PAGE MILL ROAD PALO ALTO, CA 94304-1050			EXAMINER PIHONAK, SARAH	
			ART UNIT 1627	PAPER NUMBER
			NOTIFICATION DATE 01/16/2026	DELIVERY MODE ELECTRONIC

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

Notice of the Office communication was sent electronically on above-indicated "Notification Date" to the following e-mail address(es):

patentdocket@wsgr.com

CORRECTED Notice of Allowability	Application No. 19/043,264	Applicant(s) HOWERTON et al.	
	Examiner SARAH PIHONAK	Art Unit 1627	AIA (First Inventor to File) Status Yes

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address--

All claims being allowable, PROSECUTION ON THE MERITS IS (OR REMAINS) CLOSED in this application. If not included herewith (or previously mailed), a Notice of Allowance (PTOL-85) or other appropriate communication will be mailed in due course. **THIS NOTICE OF ALLOWABILITY IS NOT A GRANT OF PATENT RIGHTS.** This application is subject to withdrawal from issue at the initiative of the Office or upon petition by the applicant. See 37 CFR 1.313 and MPEP 1308.

1. ☒ This communication is responsive to 1/12/26.
☐ A declaration(s)/affidavit(s) under **37 CFR 1.130(b)** was/were filed on _____.

2. ☐ An election was made by the applicant in response to a restriction requirement set forth during the interview on _____; the restriction requirement and election have been incorporated into this action.

3. ☒ The allowed claim(s) is/are 1-21. As a result of the allowed claim(s), you may be eligible to benefit from the **Patent Prosecution Highway** program at a participating intellectual property office for the corresponding application. For more information, please see http://www.uspto.gov/patents/init_events/pph/index.jsp or send an inquiry to PPHfeedback@uspto.gov.

4. ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
Certified copies:
 a) ☐ All b) ☐ Some* c) ☐ None of the:
 1. ☐ Certified copies of the priority documents have been received.
 2. ☐ Certified copies of the priority documents have been received in Application No. _____.
 3. ☐ Copies of the certified copies of the priority documents have been received in this national stage application from the International Bureau (PCT Rule 17.2(a)).
 * Certified copies not received: _____.

Applicant has THREE MONTHS FROM THE "MAILING DATE" of this communication to file a reply complying with the requirements noted below. Failure to timely comply will result in ABANDONMENT of this application.
THIS THREE-MONTH PERIOD IS NOT EXTENDABLE.

5. ☐ CORRECTED DRAWINGS (as "replacement sheets") must be submitted.
☐ including changes required by the attached Examiner's Amendment / Comment or in the Office action of Paper No./Mail Date _____.

Identifying indicia such as the application number (see 37 CFR 1.84(c)) should be written on the drawings in the front (not the back) of each sheet. Replacement sheet(s) should be labeled as such in the header according to 37 CFR 1.121(d).

6. ☐ DEPOSIT OF and/or INFORMATION about the deposit of BIOLOGICAL MATERIAL must be submitted. Note the attached Examiner's comment regarding REQUIREMENT FOR THE DEPOSIT OF BIOLOGICAL MATERIAL.

Attachment(s)

1. ☐ Notice of References Cited (PTO-892) 2. ☒ Information Disclosure Statements (PTO/SB/08), Paper No./Mail Date _____ 3. ☐ Examiner's Comment Regarding Requirement for Deposit of Biological Material _____ 4. ☐ Interview Summary (PTO-413), Paper No./Mail Date. _____	5. ☐ Examiner's Amendment/Comment 6. ☐ Examiner's Statement of Reasons for Allowance 7. ☐ Other _____
---	--

/SARAH PIHONAK/ Primary Examiner, Art Unit 1627	
--	--