



The next 150 years of congenital adrenal hyperplasia

Adina F. Turcu^a, Richard J. Auchus^{a,b,*}

^a Division of Metabolism, Diabetes, and Endocrinology, Department of Internal Medicine, University of Michigan, Ann Arbor, MI 48019, United States

^b Department of Pharmacology, University of Michigan, Ann Arbor, MI 48019, United States



ARTICLE INFO

Article history:

Received 26 March 2015

Received in revised form 26 May 2015

Accepted 31 May 2015

Available online 3 June 2015

ABSTRACT

Congenital adrenal hyperplasias (CAH) are a group of autosomal recessive defects in cortisol biosynthesis. Substantial progress has been made since the description of the first report, 150 years ago. This article reviews some of the recent advances in the genetics, diagnosis and treatment of CAH. In addition, we underline the aspects where further progress is required, including, among others, better diagnostic modalities for the mild phenotype and for some of the rare forms of disease, elucidation of epigenetic factors that lead to different phenotypes in patients with identical genotype and expanding on treatment options for controlling the adrenal androgen excess.

©2015 Elsevier Ltd. All rights reserved.

1. Introduction

In a recent issue of *Endocrinology*, Luisa Delle Piane and colleagues put the spotlight on the case regarded as the first report of non-salt wasting congenital adrenal hyperplasia (CAH), initially described in 1865 by Luigi de Crecchio [1]. The article details the autopsy of a prematurely deceased virilized female with enlarged adrenal glands, and the conundrum it presented to the team of pathologists involved [2]. Much has been learned over the past 150 years about CAH, now recognized as one of the most common inherited diseases. However, contemporary practitioners and researchers still have questions to answer and hypotheses to test even in the modern medical era. In this article, we focus on the recent advances in CAH and on those aspects where progress over the next century is imperative.

2. Congenital adrenal hyperplasia-brief overview

CAH is an umbrella term for inherited enzymatic deficiencies in cortisol synthesis (Fig. 1). The defective cortisol production alleviates the negative feedback to the hypothalamus and pituitary gland, resulting in excessive secretion of corticotropin-releasing hormone (CRH) and adrenocorticotropin (ACTH), respectively. The raised ACTH, in turn, cannot overcome the block in cortisol

synthesis, but its trophic action leads to enlargement of the adrenal glands. The enzymatic defects in all forms of CAH can be complete or partial, resulting into a broad spectrum of clinical presentations. The most severe forms are conventionally called “classic” CAH and are the easiest to recognize. Conversely, the milder forms or “nonclassic” CAH are often difficult to diagnose or may be overlooked, as their features overlap between each other, as well as with other clinical entities.

The most common form of CAH is 21-hydroxylase (P450c21, CYP21A2) deficiency (21OHD, Table 1), accounting for over 90% of all cases [3]. Classic 21OHD affects approximately 1 in 16,000 newborns worldwide [4]. Nonclassic 21OHD occurs in roughly 1 of 1000 Caucasians and even more frequently in populations of specific ethnicities, such as Ashkenazi Jews (1:27), Hispanics (1:53), Yugoslavs (1:62) and Italians (1:300) [5]. Complete absence of CYP21A2 activity results in both glucocorticoid and mineralocorticoid deficiencies, as well as in severe adrenal-derived androgen excess. The androgen excess is clinically evident in newborn girls, whose external genitalia are virilized. Patients with nonclassic 21OHD do not have adrenal insufficiency, and they typically present with evidence of androgen excess, such as premature pubarche, hirsutism, acne, and irregular menses, at various ages.

A second form of CAH is 11 β -hydroxylase (CYP11B1) deficiency (11OHD), which represents up to 5–8% of all CAH cases in some series of high-risk populations [6,7], although the incidence of this disorder in the general population has never been ascertained. Its incidence is estimated at 1 in 100,000 live births in the general population but is 20 times higher in Moroccan Jews, due to a founder mutation [8]. CYP11B1 catalyzes the conversion of 11-deoxycorticosterone (DOC) and 11-deoxycortisol into

* Corresponding author at: Division of Metabolism, Diabetes, and Endocrinology, Department of Internal Medicine, University of Michigan School of Medicine, Room 5560A, MSRBII, 1150 W. Medical Center Drive, Ann Arbor, MI 48019, United States. Tel.: +1 7347647764; fax: +1 7349366684.

E-mail address: rauchus@med.umich.edu (R.J. Auchus).




Fig. 1. Pathways of adrenal steroid hormone synthesis. STAR, steroidogenic acute regulatory protein; CYP11A1, cytochrome P450 cholesterol side-chain cleavage; HSD3B2, 3 β -hydroxysteroid dehydrogenase type 2; CYP17A1, 17 α -hydroxylase/17,20-lyase; CYB5A, cytochrome *b*₅; CYP21A2, 21-hydroxylase; CYP11B1, 11 β -hydroxylase; CYP11B2, aldosterone synthase; AKR1C3, 17 β -hydroxysteroid dehydrogenase type 5; DOC, 11-deoxycorticosterone; AD, androstenedione; T, testosterone.

corticosterone and cortisol, respectively, reactions whose substrates are products of CYP21A2. As in 21OHD, patients exhibit decreased cortisol synthesis and adrenal androgen overproduction. In contrast to 21OHD, the distinctive features of classic 11OHD are hypertension and less commonly hypokalemia, owing to the mineralocorticoid action of DOC, which manifests in two-thirds of patients often in mid-childhood and also makes patients with 11OHD less prone to adrenal crisis than those with 21OHD. Nonclassic 11OHD is difficult to distinguish clinically from nonclassic 21OHD but is far less common.

A third form of CAH is 3 β -hydroxysteroid dehydrogenase/isomerase type 2 (3 β HSD2) deficiency and is characterized by both mineralo- and glucocorticoid deficiency. This enzyme catalyzes the

reactions preceding those of CYP21A2, being responsible for the C3 dehydrogenation and simultaneous Δ^5 to Δ^4 transfer of the double bond into the A ring of the core steroid structure. Humans have 2 homologous 3 β HSD enzymes: type 1, expressed in placenta and peripheral tissues (skin, prostate and breast), and type 2, expressed in the adrenal glands and gonads [9–11]. Severe impairment of 3 β HSD2 results in both mineralocorticoid and glucocorticoid deficiencies and limits steroid flux to dehydroepiandrosterone (DHEA). In peripheral tissues containing 3 β HSD1 and downstream enzymes, DHEA is metabolized to androgens, which causes mild virilization in newborn girls, such as slight clitoral enlargement without the labioscrotal fusion found in 21OHD and 11OHD. In contrast with the latter two, boys are undervirilized, due to the impaired androgen synthesis in the testes, which also requires 3 β HSD2. In women, nonclassic 3 β HSD deficiency was thought to be common but is actually extremely rare, and most children with premature adrenarche do not have a mutation in the *HSD3B2* gene (see Section 3.1 and 3.2.2). Nonclassic 3 β HSD deficiency is difficult to distinguish from nonclassic 21OHD and 11OHD or PCOS clinically, all presenting with hirsutism and oligomenorrhea.

Deficiency of 17 α -hydroxylase/17,20-lyase (CYP17A1) is rare, with most cases described coming from Brazil and Asia. CYP17A1 is required for both cortisol and androgen synthesis. The only unaffected pathway is that to progesterone, aldosterone and other mineralocorticoids, and the weak glucocorticoid corticosterone, similar to the rodent adrenal that lacks this enzyme. In order to produce enough corticosterone to substitute for cortisol, DOC rises markedly, and patients present with hypertension, hypokalemia and hypogonadism. Both 46,XY and 46,XX individuals have feminine external genitalia and primarily present during pubertal age, with amenorrhea and absence of secondary sexual characteristics. Occasionally, 46,XY cases of 17OHD are diagnosed in infancy due to inguinal hernias. Partial defects of CYP17A1 activities have been found in women with poor breast development and/or menstrual abnormalities [12,13] and in males with ambiguous external genitalia, absence of male secondary sexual

Table 1
Forms of congenital adrenal hyperplasia.

Defective enzyme	Gene/Chromosome	Incidence and Populations	Clinical features	Biomarkers
21-hydroxylase (P450c21)	<i>CYP21A2</i> /6p21.3	Classic 1:16,000 Nonclassic <1:1,000 More frequent in Ashkenazi Jews, Mediterraneans, Hispanics, and Yupik Eskimos	Adrenal insufficiency in classic forms Variable degrees of virilization	17OH-progesterone, AD, T
11 β -hydroxylase (P450c11 β)	<i>CYP11B1</i> /8q24.3	1:100,000 in Caucasians 1:7,000 in Moroccan Jews	Hypertension in most patients; hypokalemia; virilization	DOC, 11-deoxycortisol, AD, T
3 β -hydroxysteroid dehydrogenase type 2	<i>HSD3B2</i> /1p13.1	Rare	Volume depletion, hyponatremia, and hyperkalemia 46,XX: virilization 46,XY: undervirilization	Pregnenolone, 17OH-pregnenolone, DHEA, DHEAS
17-hydroxylase/17,20-lyase (P450c17)	<i>CYP17A1</i> /10q21-q22	1:50,000 worldwide; More common in Brazil and Asia	Hypertension, hypokalemia, and hypogonadism 46,XX: primary amenorrhea and absence of secondary sexual characteristics 46,XY: undervirilization, abdominal testes	Progesterone, DOC, corticosterone; LH and FSH
Steroidogenic acute regulatory protein (STAR)	<i>STAR</i> /8p11.2	Rare More frequent in Japanese, Palestinians, Koreans	Adrenal insufficiency; enlarged, lipid-laden adrenal glands. Female phenotype of external genitalia in both sexes	All steroids decreased
Cholesterol side-chain cleavage enzyme (P450scc)	<i>CYP11A1</i> /15q23-q24	Rare	Adrenal insufficiency; adrenal glands may appear absent	All steroids decreased
P450-oxidoreductase deficiency (POR)	<i>POR</i> /7q11.2	Rare More common in Japan and Korea	Volume depletion, skeletal malformations (Antley-Bixler); maternal virilization 46,XX: mild-to-moderate virilization 46,XY: undervirilization	Highly variable profiles, multiple partial defects

DOC, 11-deoxycorticosterone; AD, androstenedione; T, testosterone; DHEA, dehydroepiandrosterone; DHEAS, DHEA sulfate; LH, luteinizing hormone; FSH, follicle-stimulating hormone.

characteristics, and gynecomastia at puberty [14], but a “non-classic” form has not been described. Conceivably, patients with very mild or nonclassic 17OHD might be dismissed as having low-renin hypertension and primary hypogonadism and never evaluated for 17OHD. Isolated 17,20-lyase deficiency has been described in patients with mutations in the *CYP17A1* [15,16], *POR* [17] or *CYB5A* [18,19] genes, but these patients do not have CAH because cortisol synthesis is normal. Isolated 17,20-lyase deficiency can be difficult to distinguish from partial androgen insensitivity or from rare steroidogenic defects such as *AKR1C2* + *AKR1C4* [20].

Lipoid CAH (LCAH), the most severe defect in steroidogenesis, is named for the massively enlarged, lipid-laden adrenals characteristic of this disease. LCAH is caused by a defect in the steroidogenic acute regulatory protein (StAR), which prevents the mobilization of cholesterol into steroidogenic pathways and results in negligible production of all steroids. Cholesterol side-chain cleavage enzyme (*P450scc*) deficiency yields a similar severe global defect in steroidogenesis as LCAH but without the enlarged adrenals found with StAR deficiency [21,22]. Affected children typically present with life-threatening adrenal insufficiency in early infancy, and males appear phenotypically female due to impaired testicular androgen synthesis in utero. Nonclassic forms of both lipoid CAH and *P450scc* deficiency have been described, presenting with late-onset isolated glucocorticoid insufficiency and normal external genitalia [23–26].

Finally, *P450-oxidoreductase* (*POR*) deficiency combines features of 21OHD, 17OHD, and aromatase (*CYP19A1*) deficiencies, in various combinations and severities [27]. In addition, these patients may present skeletal malformations, such as craniosynostosis, radiohumeral or radio-ulnar synostosis, and femoral bowing, as part of the Antley–Bixler syndrome [28]. In contrast, one of the initial cases described was a phenotypically normal woman with infertility, illustrating the wide spectrum of this disease and suggesting that “nonclassic” disease could be defined by the absence of skeletal and/or genital anomalies.

3. Diagnosis of CAH

3.1. Hormonal testing

For each enzymatic defect, the precursor to product ratio is the mainstay of diagnosis. Cosyntropin stimulation maximizes these ratios and is particularly important for all cases with indeterminate baseline results [29].

Newborn screening for 21OHD was first implemented in 1978 [30,31], and is currently available throughout the United States and many other countries. Screening reduces the time to diagnosis in infants, particularly for boys who are often not diagnosed at birth and suffer crises several days later [32,33]. The most important role of early diagnosis is to reduce morbidity and mortality for severely affected babies, although this aspect remains controversial and might also depend on the economic and healthcare status of each country [34–36]. First-tier testing measures 17OHP in dried blood spots by an immunofluorometric assay. A random 17OHP >20,000 ng/dL is suggestive of 21OHD; however, false-positive results are commonly seen in premature and severely ill infants [37,38]. Factors contributing to false-positive screening include activation of the hypothalamic–pituitary–adrenal axis in response to perinatal stress, the relative immaturity of adrenal *CYP11B1* activity in preterm infants [39], which also elevates 17OHP [39,40], and interfering steroids in the immunoassays. False-negative rates of up to 22% have been reported in infant screening [41,42], particularly when mothers had been exposed to glucocorticoids prenatally. Although higher false-negative rates have been reported in girls [42], it is possible that boys not identified with newborn screening remained undiagnosed for

several years. Weight- and gestational age-adjusted cutoffs for 17OHP have been implemented to improve the positive predictive value of screening [43–45], and more specific second-tier screening procedures to adjudicate abnormal first-tier screens have been successfully implemented in some states [46].

Beyond neonatal screening, children with clinical evidence of androgen excess—such as pubic and axillary hair growth, oily skin, rapid somatic growth, and advanced skeletal maturation—initially undergo testing of a morning baseline 17OHP [47]. Intermediate screening values (200–1000 ng/dL) are followed by retesting after cosyntropin stimulation. The diagnosis of classic 21OHD is based on stimulated 17OHP levels above 10,000 ng/dL, while nonclassic 21OHD requires a 17OHP >1000 ng/dL⁴⁷.


Classic 11OHD and 3βHSD2 deficiency overlap clinically with classic 21OHD, and in all 3 conditions, 17OHP can be elevated. Due to their low prevalence, the diagnosis of 11OHD is considered when hypertension and/or hypokalemia are present, or when the 17OHP is lower than expected for the degree of androgen excess in a 21OHD patient. 3βHSD2 deficiency is considered if virilization is mild relative to the degree of salt wasting. Nonclassic 21OHD is vastly more common than other forms of nonclassic CAH, but these conditions are even more difficult to distinguish clinically than their classic forms. A diagnosis of nonclassic 11OHD or 3βHSD2 deficiencies should be pursued in patients with androgen excess during childhood only after more common causes (such as exposure to exogenous androgens, PCOS, 21OHD, adrenal or gonadal tumors, chorionic gonadotropin-secreting tumor) have been excluded. Nonclassic 11OHD is diagnosed when 11-deoxycortisol rises above 1800 ng/dL and cortisol is >18 μg/dL after cosyntropin [7]. The diagnosis of nonclassic 3βHSD deficiency requires a 17-hydroxypregnenolone >3000 ng/dL and a cortisol >18 μg/dL after cosyntropin, with a 17-hydroxypregnenolone/cortisol ratio >10 standard deviations above normal [48].

In most centers, steroid measurements are performed by immunoassays, which are limited by cross-reactivity and can yield unreliable results, particularly when ordinarily minor steroids are markedly elevated. More accurate testing can be attained by using liquid chromatography/tandem mass spectrometry (LC–MS/MS) [49–51]. In addition to increased specificity and sensitivity, LC–MS/MS also affords quantitation of multiple steroids in a single measurement. Elevated 21-deoxycortisol by LC–MS/MS has been shown to increase the sensitivity of newborn screening [50] and to discriminate heterozygote carriers of *CYP21A2* mutations from nonclassic 21OHD better than 17OHP [52]. Simultaneous measurement of 11-deoxycortisol can distinguish 11OHD from 21OHD in second-tier screening. Other novel biomarkers for diagnosis, such as 16α-hydroxyprogesterone and 11β-hydroxyprogesterone, have been recently proposed (Fig. 2) [53]. Multi-steroid panels simultaneously testing all forms of CAH clinically in question from a small volume serum sample will likely dominate in the future. For now, however, the widespread use of LC–MS/MS is constrained by its limited availability, technical demands and high cost. Alternatively, comprehensive analysis of urinary steroid metabolites using gas chromatography–mass spectrometry (GC/MS) affords patterns characteristic for each form of CAH [54]. While GC/MS has been used for decades, the availability of this methodology is limited, and sample throughput is slow and tedious.

3.2. Genetics of CAH and role of genetic testing

3.2.1. Genetics of CAH

All CAH forms are monogenic, autosomal-recessive disorders. The gene encoding human *CYP21A2* is located on chromosome 6p21.3, within the human leucocyte antigen (HLA) major histocompatibility complex and adjacent to the genes for the fourth component of complement [55–57]. In addition to this active gene, humans also



The *HSD3B2* gene, located on chromosome 1p13.1, is expressed almost exclusively in the adrenal and gonads [10]. The highly homologous type 13βHSD gene (*HSD3B1*) is located in vicinity on the

Although human CYP17A1 catalyzes two separate reactions, it is encoded by a single gene located on chromosome 10q24.3, which consists of 8 exons and encodes a 508-amino-acid protein [85–87]. Over 90 CYP17A1 mutations have been described throughout the entire gene. Populations in which 17OHD is more prevalent, such as Brazilians [88], Canadian Mennonites, Dutch Frieslanders [89], Japanese [90], and patients from East Asia [91], however, have specific reoccurring mutations, due to founder effects. Rare mutations located in the redox-partner interaction site or active site can lead to isolated 17,20-lyase deficiency [15,16,92,93].

The role of genetic testing is best established in cases that remain equivocal after cosyntropin stimulation or for prenatal genetic

counseling. Up to 70% of patients with nonclassic 21OHD carry a classic 21OHD mutation on one of their alleles, rendering them a carrier for classic 21OHD [94,95]. Children of women with nonclassic 21OHD have an approximately 2.5% risk of being born with classic 21OHD and a 15% risk for having nonclassic 21OHD [96]. The same increased risk of 21OHD theoretically applies equally to fathers with nonclassic 21OHD, but similar data for men are lacking. The risk increases significantly in ethnic groups known to have a higher prevalence of the disease [96]. Thus, particularly in cases of a known affected prospective parent, preconception genetic testing of the second parent might help families to understand the risk of an affected offspring and to receive appropriate genetic counseling.

Another role for genetic testing evolved from efforts to prevent virilization of female fetuses affected with classic 21OHD. Prenatal treatment of the mother carrying an affected child with dexamethasone can reduce genital virilization compared to sisters who were not treated in utero. Prenatal diagnosis of 21OHD has been accomplished by chorionic villus sampling at approximately 14 weeks of gestation or by amniocentesis at approximately 20 weeks [97]. Both approaches are invasive and increase the risk of miscarriages [98–100]. To be successful, however, dexamethasone treatment must be started by 8 weeks' gestation, when genital anatomy is sensitive to dihydrotestosterone action [101,102]. Thus, presumptive treatment must be started before the prenatal diagnosis can be established. Statistically, only 1 in 8 fetuses is an affected girl, so up to 88% of pregnancies are treated unnecessarily. Prenatal dexamethasone was associated with impaired verbal working memory and related cognitive metrics in one study [103], and more recent data have suggested that exposure to prenatal synthetic glucocorticoids interferes with the normal brain development [104,105]. Thus, prenatal therapy is currently regarded as experimental by academic societies [47,106].

In recent years, early noninvasive genetic testing has been proposed. Lo and colleagues first documented the presence of cell-free fetal DNA (cff-DNA) in the maternal circulation in 1997 [107]. Quantitative polymerase chain reaction analysis of the SRY gene in maternal plasma was introduced in 2001, as a sensitive method for fetal gender determination in women with CAH [108]. Technological advances in genetic testing have recently made the diagnosis of 21OHD possible as early as 6 weeks of gestation, by targeted massively parallel sequencing performed on cff-DNA circulating in maternal plasma [109]. Rarely, prenatal DNA sequencing can lead to erroneous diagnosis of CAH, as in duplication of the *CYP21A2* gene [110]. Moreover, massively parallel sequencing remains technically challenging, costly and only available in selected centers.

As genetic testing becomes cheaper and more widely available, one can anticipate that targeted sequencing of several genes of interest could clarify the diagnosis in patients with atypical clinical presentations and borderline biochemical workup. Diagnosis of all forms of CAH currently relies almost exclusively on hormonal testing, which is fraught with pitfalls. For example, earlier studies of hirsute women over-diagnosed nonclassic 3β HSD2 deficiency, when relying on measurement of Δ^5 steroids [111–114]. Genetic testing uncovered mutations in the *HSD3B2* gene in only a vanishingly small number of such women [48,115]. As genetic databases continue to grow, more robust links between genotype and phenotype will develop. In the interim however, the clinical and biochemical context remain critical for the diagnosis of this heterogeneous group of diseases.

4. Treatment of CAH

4.1. Conventional and experimental glucocorticoid and mineralocorticoid therapy

The two general goals of treatment in all forms of CAH are to replace the deficient hormones on one hand, and to offset the

undesirable effects of excessive hormonal production on the other. With a variety of available formulations, corticosteroid replacement has become relatively facile, with some limitations. For example, the aldosterone substitute fludrocortisone acetate is not available in mainland China; hydrocortisone tablets are not sold in many countries including Brazil; and liquid forms of hydrocortisone must be obtained from compounding pharmacies without quality control in the United States. Conversely, management of adrenal androgen excess in both classic and nonclassic CAH remains challenging. Several strategies to counteract the excessive adrenal androgen synthesis have been explored, as outlined below.

The most commonly used approach has been that of suppressing ACTH by strategic dosing of glucocorticoids. The typical short- or intermediate-acting glucocorticoid regimens, which are sufficient for replacement in non-CAH related adrenal insufficiency, often fail to blunt the early morning rise of ACTH, which is the principal drive for adrenal androgen overproduction. Physicians have resorted to supraphysiologic or non-physiologic (nocturnal) doses of glucocorticoids in an attempt to counteract the excessive androgen synthesis, thus promoting bone loss, obesity and features of metabolic syndrome in these patients [116,117]. Sustained-released hydrocortisone preparations have been recently developed, attempting to mimic the cortisol circadian rhythm and to suppress the early morning ACTH elevation [118,119]. In a phase two clinical trial, a modified-release formulation of hydrocortisone administered once daily mimicked the normal diurnal cortisol rhythm more closely than conventional hydrocortisone dosing; however, androstenedione rose higher than with conventional hydrocortisone in the afternoon. After reformulation, a second phase two trial with twice-daily dosing achieved lower hydrocortisone dose equivalent, as well as lower 17OHP and androstenedione levels, compared to conventional therapies [118]. Continuous subcutaneous hydrocortisone infusion via a pump, mimicking a circadian secretory profile, has been used experimentally in young patients with increased cortisol clearance, and this approach was shown to decrease 17OHP and adrenal androgen production with a lower total daily dose [120,121].

4.2. Experimental therapies beyond glucocorticoids

Another alternative for decreasing ACTH was undertaken in a recent proof-of-concept single-blind, placebo-controlled, single center, fixed-sequence, single-dose trial study, using the corticotropin-releasing factor receptor type 1 antagonist NBI77860 [122]. At 300 and 600 mg, a nocturnal dose lowered morning ACTH and 17OHP by >50% from baseline in 4 of 8 participants, and the action correlated with drug exposure. Further studies are needed to determine the effects of NBI77860 on key parameters of disease control, such as serum androgens, when given in repeated doses at steady state.

A second strategy is that of directly inhibiting androgen synthesis or antagonizing androgen action. The combination of flutamide, an antiandrogen, and testolactone, an aromatase inhibitor, permitted the use of lower doses of hydrocortisone and fludrocortisone acetate and normalized linear growth and bone maturation in children followed for 2 years [123,124]. Long-term results from this trial are anticipated soon; however, latter-generation anti-androgens and aromatase inhibitors have supplanted both flutamide and testolactone for the treatment of prostate and breast cancers, respectively. A GnRH antagonist has been successful in improving height in children with 21OHD and precocious puberty [125]. Abiraterone acetate, a potent CYP17A1 inhibitor approved for use in castration-resistant prostate cancer, normalized androstenedione in adult women with classic 21OHD and elevated androgens when added to physiologic

hydrocortisone and fludrocortisone acetate replacement in one short-term study [126].

Monitoring and titrating treatment remains a major clinical challenge, and no consensus exists among practitioners. Although 17OHP has long been used to guide the management of 21OHD, its serum concentration correlates poorly with DHEA and androstenedione [127]. DHEAS, the major C₁₉ adrenal product, can be paradoxically low in 21OHD patients with inadequate control [128–130]. Moreover, there is no good correlation between the routinely measured androgens (androstenedione and, in women, testosterone) and clinical evidence of androgen excess [131,132]. Research conducted in the recent years has brought into consideration non-conventional androgens and precursors, such as steroids derived from the so-called “back-door pathway” [133] (Fig. 2). This pathway starts with two consecutive 5 α -, then 3 α -reductions of 17OHP and leads via androsterone and 5 α -androstan-3 α ,17 β -diol to dihydrotestosterone, the most potent endogenous androgen, while bypassing androstenedione and testosterone. Kamrath and colleagues found increased excretion of 5 α -reduced products and intermediates of the backdoor pathway in 142 children and young adults with CAH, compared with 138 similarly aged controls, particularly during the first year of life [127]. Consequently, this pathway has been proposed to contribute to the virilization of female fetuses with 21OHD [134], as has been shown in POR deficiency [135]. In addition, the adrenal glands also produce other active androgens, such as 11 β -hydroxytestosterone. Substantial amounts of 11-oxygenated C₁₉ steroids were documented in adrenal vein samples obtained from normal adrenals, with 11 β -hydroxyandrostenedione being particularly abundant [136]. In a cell-based androgen transactivation assay, 11 β -hydroxytestosterone and 11-ketotestosterone activated the androgen receptors at concentration approximately ten times higher than testosterone, but 30 times lower than androstenedione and 11 β -hydroxyandrostenedione [136]. The contribution of these steroids to the androgen excess of 21OHD remains to be further investigated. As these 11-oxygenated C₁₉ steroids are products of CYP11B1, an adrenal-specific enzyme, these compounds might serve as specific biomarkers of adrenal-derived androgen excess and facilitate titration of therapy.

5. Summary

We have reviewed the state of knowledge of CAH 150 years after the initial description and highlighted some recent advances in the field. Based on this review, we propose a research agenda and

related questions for improved understanding and management of CAH (Box 1).

1. The mild phenotype: What would be the phenotype of a patient with nonclassic 17OHD? How many cases of nonclassic 11OHD are we missing? What is the minimum enzyme activity that distinguishes unaffected and nonclassic CAH patients?
2. Modifier genes: How can siblings with the same CAH genotype show very different phenotypes and disease control? What genes control drug metabolism, response to androgens, and peripheral metabolism of C₁₉-adrenal products? Why does salt wasting lessen in many but not all adults with classic 21OHD? Why do not all women with nonclassic 21OHD suffer from oligomenorrhea and subfertility?
3. Biomarkers for diagnosis and treatment: Can a panel of adrenal-derived 21-deoxysteroids be used to diagnose nonclassic 21OHD without cosyntropin stimulation and to eliminate false-positive newborn screens for classic 21OHD? Can similar panels be developed for 11OHD, 17OHD, POR, and 3 β HSD2 deficiencies? What are the best steroids to use to monitor and titrate treatment?
4. Better treatments: What are the optimal glucocorticoid regimens? How do we position hydrocortisone as continuous subcutaneous infusion or modified-release oral preparations in treatment algorithms? Can we eliminate Cushingoid side effects and maintain control of androgen excess with the addition of abiraterone acetate, NBI77860, or other novel therapeutics to physiologic replacement doses of hydrocortisone?
5. Genetic diagnosis: Can cf-DNA in the maternal circulation be used to diagnose CAH and to determine sex chromosome complement for fetuses at risk for having CAH before 8 weeks' gestation? What is the value of whole-exome sequencing in children with suspected CAH?

This incomplete list represents a substantial amount of work. To answer these questions and accomplish these goals will take research funding, collaboration among and creativity from investigators, participation of patients and their families, and partnerships with the pharmaceutical industry. With advances in technology, biochemistry, and genetics, the time is right to make major advances in CAH, which will have important implications for other related and often more common diseases. These patients need better options today, and we cannot take another 150 years to bring these advances to fruition.

Acknowledgement

This work was supported by F32DK 103461 to A.F.T.

References

- [1] L. Delle Piane, P.F. Rinaudo, W.L. Miller, 150 years of congenital adrenal hyperplasia: translation and commentary of de Creccchio's classic paper from 1865, *Endocrinology* 156 (2015) 1210–1217.
- [2] L. De Creccchio, Sopra un caso di apparenze virili in una donna, *Il Morgagni* (1865) 151–89.
- [3] P.W. Speiser, P.C. White, Congenital adrenal hyperplasia, *New Engl. J. Med.* 349 (2003) 776–788.
- [4] B.L. Therrell Jr., S.A. Berenbaum, V. Manter-Kapanke, et al., Results of screening 1.9 million Texas newborns for 21-hydroxylase-deficient congenital adrenal hyperplasia, *Pediatrics* 101 (1998) 583–590.
- [5] P.W. Speiser, B. Dupont, P. Rubinstein, A. Piazza, A. Kastelan, M.I. New, High frequency of nonclassical steroid 21-hydroxylase deficiency, *Am. J. Hum. Genet.* 37 (1985) 650–667.
- [6] P.C. White, M.I. New, B. Dupont, Congenital adrenal hyperplasia. (1), *New Engl. J. Med.* 316 (1987) 1519–1524.
- [7] P.C. White, K.M. Curnow, L. Pascoe, Disorders of steroid 11 β -hydroxylase isozymes, *Endocr. Rev.* 15 (1994) 421–438.

Box 1. Areas of future research in congenital adrenal hyperplasia (CAH).

Research agenda

- **Diagnosis**
 - Biomarker panels for improved diagnosis of all nonclassic CAH forms
 - Early prenatal genetic diagnosis
- **Treatment**
 - Characterization of biomarkers for treatment monitoring and titration
 - Improved treatment modalities, to allow suppression of androgen excess while avoiding supraphysiologic doses of glucocorticoids.
- **Elucidation of modifier genes/epigenetic contributors to phenotypes**

- [8] A. Rosler, E. Leiberman, T. Cohen, High frequency of congenital adrenal hyperplasia (classic 11 β -hydroxylase deficiency) among Jews from Morocco, *Am. J. Med. Genet.* 42 (1992) 827–834.
- [9] J. Simard, F. Durocher, F. Mebarki, et al., Molecular biology and genetics of the 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 isomerase gene family, *J. Endocrinol.* 150 Suppl (1996) S189–S207.
- [10] E. Rheume, Y. Lachance, H.F. Zhao, et al., Structure and expression of a new complementary DNA encoding the almost exclusive 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 -isomerase in human adrenals and gonads, *Mol. Endocrinol.* 5 (1991) 1147–1157.
- [11] S. Gingras, R. Moriggi, B. Groner, J. Simard, Induction of 3 β -hydroxysteroid dehydrogenase/ Δ^5 - Δ^4 isomerase type 1 gene transcription in human breast cancer cell lines and in normal mammary epithelial cells by interleukin-4 and interleukin-13, *Mol. Endocrinol.* 13 (1999) 66–81.
- [12] P.N. Singhellakis, D. Panidis, J. Papadimas, et al., Spontaneous sexual development and menarche in a female with 17 α -hydroxylase deficiency, *J. Endocrinol. Invest.* 9 (1986) 177–183.
- [13] Y. Katayama, S. Kado, S. Wada, et al., A case of 17 α -hydroxylase deficiency with retained menstruation, *Endocr. J.* 41 (1994) 213–218.
- [14] M.I. New, Male pseudohermaphroditism due to 17 α -hydroxylase deficiency, *J. Clin. Invest.* 49 (1970) 1930–1941.
- [15] D.H. Geller, R.J. Auchus, B.B. Mendonca, W.L. Miller, The genetic and functional basis of isolated 17,20-lyase deficiency, *Nat. Genet.* 17 (1997) 201–205.
- [16] D.P. Sherbet, D. Tiosano, K.M. Kwist, Z. Hochberg, R.J. Auchus, CYP17 mutation E305G causes isolated 17,20-lyase deficiency by selectively altering substrate binding, *J. Biol. Chem.* 278 (2003) 48563–48569.
- [17] E. Hershkovitz, R. Parvari, S.A. Wudy, et al., Homozygous mutation G539R in the gene for P450 oxidoreductase in a family previously diagnosed as having 17,20-lyase deficiency, *J. Clin. Endocrinol. Metab.* 93 (2008) 3584–3588.
- [18] R.C. Kok, M.A. Timmerman, K.P. Wolfenbuttel, S.L. Drop, F.H. de Jong, Isolated 17,20-lyase deficiency due to the cytochrome b₅ mutation W27X, *J. Clin. Endocrinol. Metab.* 95 (2010) 994–999.
- [19] J. Idkowiak, T. Randell, V. Dhir, et al., A missense mutation in the human cytochrome b₅ gene causes 46,XY disorder of sex development due to true isolated 17,20 lyase deficiency, *J. Clin. Endocrinol. Metab.* 97 (2012) E465–E475.
- [20] C.E. Fluck, M. Meyer-Boni, A.V. Pandey, et al., Why boys will be boys: two pathways of fetal testicular androgen biosynthesis are needed for male sexual differentiation, *Am. J. Hum. Genet.* 89 (2011) 201–218.
- [21] H.S. Bose, T. Sugawara, J.F.3rd Strauss, W.L. Miller, International Congenital Lipoid Adrenal Hyperplasia Consortium, The pathophysiology and genetics of congenital lipoid adrenal hyperplasia, *New Engl. J. Med.* 335 (1996) 1870–1878.
- [22] M.K. Tee, M. Abramsohn, N. Loewenthal, et al., Varied clinical presentations of seven patients with mutations in CYP11A1 encoding the cholesterol side-chain cleavage enzyme, P450_{sc}, *J. Clin. Endocrinol. Metab.* 98 (2013) 713–720.
- [23] B.Y. Baker, L. Lin, C.J. Kim, et al., Nonclassic congenital lipoid adrenal hyperplasia: a new disorder of the steroidogenic acute regulatory protein with very late presentation and normal male genitalia, *J. Clin. Endocrinol. Metab.* 91 (2006) 4781–4785.
- [24] P. Rubtsov, M. Karmanov, P. Sverdlova, P. Spirin, A. Tiupakov, A novel homozygous mutation in CYP11A1 gene is associated with late-onset adrenal insufficiency and hypospadias in a 46,XY patient, *J. Clin. Endocrinol. Metab.* 94 (2009) 936–939.
- [25] T. Sahakitrungruang, M.K. Tee, P.R. Blackett, W.L. Miller, Partial defect in the cholesterol side-chain cleavage enzyme P450_{sc} (CYP11A1) resembling nonclassic congenital lipoid adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 96 (2011) 792–798.
- [26] S. Parajes, C. Kamrath, I.T. Rose, et al., A novel entity of clinically isolated adrenal insufficiency caused by a partially inactivating mutation of the gene encoding for P450 side chain cleavage enzyme (CYP11A1), *J. Clin. Endocrinol. Metab.* 96 (2011) E1798–E1806.
- [27] N. Huang, A.V. Pandey, V. Agrawal, et al., Diversity and function of mutations in P450 oxidoreductase in patients with Antley-Bixler syndrome and disordered steroidogenesis, *Am. J. Hum. Genet.* 76 (2005) 729–749.
- [28] C.E. Fluck, T. Tajima, A.V. Pandey, et al., Mutant P450 oxidoreductase causes disordered steroidogenesis with and without Antley-Bixler syndrome, *Nat. Genet.* 36 (2004) 228–230.
- [29] M.I. New, F. Lorenzen, A.J. Lerner, et al., Genotyping steroid 21-hydroxylase deficiency: hormonal reference data, *J. Clin. Endocrinol. Metab.* 57 (1983) 320–326.
- [30] S. Pang, J. Hotchkiss, A.L. Drash, L.S. Levine, M.I. New, Microfilter paper method for 17 α -hydroxyprogesterone radioimmunoassay: its application for rapid screening for congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 45 (1977) 1003–1008.
- [31] S.Y. Pang, M.A. Wallace, L. Hofman, et al., Worldwide experience in newborn screening for classical congenital adrenal hyperplasia due to 21-hydroxylase deficiency, *Pediatrics* 81 (1988) 866–874.
- [32] B.L. Therrell, Newborn screening for congenital adrenal hyperplasia, *Endocrinol. Metab. Clin. North Am.* 30 (2001) 15–30.
- [33] A. Popov, I. Beloev, Early diagnosis and prevention of internal diseases, *Vutr. Boles.* 16 (1977) 1–10.
- [34] K.A. Strnadova, F. Votava, J. Lebl, et al., Prevalence of congenital adrenal hyperplasia among sudden infant death in the Czech Republic and Austria, *Eur. J. Pediatr.* 166 (2007) 1–4.
- [35] S.D. Grosse, G. Van Vliet, How many deaths can be prevented by newborn screening for congenital adrenal hyperplasia? *Horm. Res.* 67 (2007) 284–291.
- [36] B.E. Hird, L. Tetlow, S. Tobin, L. Patel, P.E. Clayton, No evidence of an increase in early infant mortality from congenital adrenal hyperplasia in the absence of screening, *Arch. Dis. Child.* 99 (2014) 158–164.
- [37] P. Cavarzere, D. Samara-Boustani, I. Flechtner, et al., Transient hyper-17-hydroxyprogesteronemia: a clinical subgroup of patients diagnosed at neonatal screening for congenital adrenal hyperplasia, *Eur. J. Endocrinol.* 161 (2009) 285–292.
- [38] B. Coulm, J. Coste, V. Tardy, et al., Efficiency of neonatal screening for congenital adrenal hyperplasia due to 21-hydroxylase deficiency in children born in mainland France between 1996 and 2003, *Arch. Pediatr. Adolesc. Med.* 166 (2012) 113–120.
- [39] M.M. Lee, L. Rajagopalan, G.J. Berg, T. Moshang Jr., Serum adrenal steroid concentrations in premature infants, *J. Clin. Endocrinol. Metab.* 69 (1989) 1133–1136.
- [40] R.V. Hingre, S.J. Gross, K.S. Hingre, D.M. Mayes, R.A. Richman, Adrenal steroidogenesis in very low birth weight preterm infants, *J. Clin. Endocrinol. Metab.* 78 (1994) 266–270.
- [41] K. Sarafoglou, K. Banks, J. Kylo, S. Pittock, W. Thomas, Cases of congenital adrenal hyperplasia missed by newborn screening in Minnesota, *JAMA* 307 (2012) 2371–2374.
- [42] T.S. Varness, D.B. Allen, G.L. Hoffman, Newborn screening for congenital adrenal hyperplasia has reduced sensitivity in girls, *J. Pediatr.* 147 (2005) 493–498.
- [43] M. Steigert, E.J. Schoenle, A. Biason-Lauber, T. Torresani, High reliability of neonatal screening for congenital adrenal hyperplasia in Switzerland, *J. Clin. Endocrinol. Metab.* 87 (2002) 4106–4110.
- [44] B. Olgemoller, A.A. Roscher, B. Liebl, R. Fingerhut, Screening for congenital adrenal hyperplasia: adjustment of 17-hydroxyprogesterone cut-off values to both age and birth weight markedly improves the predictive value, *J. Clin. Endocrinol. Metab.* 88 (2003) 5790–5794.
- [45] H.J. van der Kamp, C.G. Oudshoorn, B.H. Elvers, et al., Cutoff levels of 17 α -hydroxyprogesterone in neonatal screening for congenital adrenal hyperplasia should be based on gestational age rather than on birth weight, *J. Clin. Endocrinol. Metab.* 90 (2005) 3904–3907.
- [46] C.L. Chan, K. McFann, L. Taylor, D. Wright, P.S. Zeitler, J.M. Barker, Congenital adrenal hyperplasia and the second newborn screen, *J. Pediatr.* 163 (2013) 109–113 e1.
- [47] P.W. Speiser, R. Azziz, L.S. Baskin, et al., Congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency: an Endocrine Society clinical practice guideline, *J. Clin. Endocrinol. Metab.* 95 (2010) 4133–4160.
- [48] L.M. Mermejo, L.L. Elias, S. Marui, A.C. Moreira, B.B. Mendonca, M. de Castro, Refining hormonal diagnosis of type II 3 β -hydroxysteroid dehydrogenase deficiency in patients with premature pubarche and hirsutism based on HSD3B2 genotyping, *J. Clin. Endocrinol. Metab.* 90 (2005) 1287–1293.
- [49] E. Schwarz, A. Liu, H. Randall, et al., Use of steroid profiling by UPLC-MS/MS as a second tier test in newborn screening for congenital adrenal hyperplasia: the Utah experience, *Pediatr. Res.* 66 (2009) 230–235.
- [50] N. Janzen, M. Peter, S. Sander, et al., Newborn screening for congenital adrenal hyperplasia: additional steroid profile using liquid chromatography-tandem mass spectrometry, *J. Clin. Endocrinol. Metab.* 92 (2007) 2581–2589.
- [51] N. Janzen, S. Sander, M. Terhardt, M. Peter, J. Sander, Fast and direct quantification of adrenal steroids by tandem mass spectrometry in serum and dried blood spots, *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 861 (2008) 117–122.
- [52] F.A. Costa-Barbosa, V.F. Tonetto-Fernandes, V.M. Carvalho, et al., Superior discriminating value of ACTH-stimulated serum 21-deoxycortisol in identifying heterozygote carriers for 21-hydroxylase deficiency, *Clin. Endocrinol. (Oxford)* 73 (2010) 700–706.
- [53] A.F. Turcu, J. Rege, R. Chomic, J. Liu, H.K. Nishimoto, T. Else, A.G. Moraitis, G.S. Palapattu, W.E. Rainey, R.J. Auchus, *J. Clin. Endocrinol. Metab.* 100 (2015) 2283–2290.
- [54] N. Krone, B.A. Hughes, G.G. Lavery, P.M. Stewart, W. Arlt, C.H. Shackleton, Gas chromatography/mass spectrometry (GC/MS) remains a pre-eminent discovery tool in clinical steroid investigations even in the era of fast liquid chromatography tandem mass spectrometry (LC/MS/MS), *J. Steroid. Biochem. Mol. Biol.* 121 (2010) 496–504.
- [55] M.C. Carroll, R.D. Campbell, R.R. Porter, Mapping of steroid 21-hydroxylase genes adjacent to complement component C4 genes in HLA, the major histocompatibility complex in man, *Proc. Natl. Acad. Sci. U.S.A.* 82 (1985) 521–525.
- [56] P.C. White, D. Grossberger, B.J. Onufer, et al., Two genes encoding steroid 21-hydroxylase are located near the genes encoding the fourth component of complement in man, *Proc. Natl. Acad. Sci. U.S.A.* 82 (1985) 1089–1093.
- [57] P.C. White, M.I. New, B. Dupont, Structure of human steroid 21-hydroxylase genes, *Proc. Natl. Acad. Sci. U.S.A.* 83 (1986) 5111–5115.
- [58] P.C. White, P.W. Speiser, Congenital adrenal hyperplasia due to 21-hydroxylase deficiency, *Endocr. Rev.* 21 (2000) 245–291.
- [59] E. Mornet, J. Dupont, A. Vitek, P.C. White, Characterization of two genes encoding human steroid 11 β -hydroxylase (P-450_{11 β}), *J. Biol. Chem.* 264 (1989) 20961–20967.

- [60] P.C. White, J. Dupont, M.I. New, E. Leiberger, Z. Hochberg, A. Rosler, A mutation in CYP11B1 (Arg-448→His) associated with steroid 11 β -hydroxylase deficiency in Jews of Moroccan origin, *J. Clin. Invest.* 87 (1991) 1664–1667.
- [61] K.M. Curnow, L. Slutsker, J. Vitek, et al., Mutations in the CYP11B1 gene causing congenital adrenal hyperplasia and hypertension cluster in exons 6, 7, and 8, *Proc. Natl. Acad. Sci. U.S.A.* 90 (1993) 4552–4556.
- [62] A. Helmberg, B. Ausserer, R. Kofler, Frame shift by insertion of 2 basepairs in codon 394 of CYP11B1 causes congenital adrenal hyperplasia due to steroid 11 β -hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 75 (1992) 1278–1281.
- [63] Y. Naiki, T. Kawamoto, Y. Mitsuuchi, et al., A nonsense mutation (TGG [Trp116]→TAG [Stop]) in CYP11B1 causes steroid 11 β -hydroxylase deficiency, *Eur. J. Clin. Endocrinol. Metab.* 77 (1993) 1677–1682.
- [64] C.A. Skinner, G. Rumsby, Steroid 11 β -hydroxylase deficiency caused by a five base pair duplication in the CYP11B1 gene, *Hum. Mol. Genet.* 3 (1994) 377–378.
- [65] L.X. Yang, K. Toda, K. Miyahara, et al., Classic steroid 11 β -hydroxylase deficiency caused by a C→G transversion in exon 7 of CYP11B1, *Biochem. Biophys. Res. Commun.* 216 (1995) 723–728.
- [66] Y. Nakagawa, M. Yamada, H. Ogawa, Y. Igarashi, Missense mutation in CYP11B1 (CGA[Arg-384]→GGA[Gly]) causes steroid 11 β -hydroxylase deficiency, *Eur. J. Endocrinol.* 132 (1995) 286–289.
- [67] S. Geley, K. Kapelari, K. Johrer, et al., CYP11B1 mutations causing congenital adrenal hyperplasia due to 11 β -hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 81 (1996) 2896–2901.
- [68] D.P. Merke, T. Tajima, A. Chhabra, et al., Novel CYP11B1 mutations in congenital adrenal hyperplasia due to steroid 11 β -hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 83 (1998) 270–273.
- [69] Y.S. Zhu, J.J. Cordero, S. Can, et al., Mutations in CYP11B1 gene: phenotype–genotype correlations, *Am. J. Med. Genet. A* 122A (2003) 193–200.
- [70] N. Krone, F.G. Riepe, D. Gotze, et al., Congenital adrenal hyperplasia due to 11-hydroxylase deficiency: functional characterization of two novel point mutations and a three-base pair deletion in the CYP11B1 gene, *J. Clin. Endocrinol. Metab.* 90 (2005) 3724–3730.
- [71] N. Krone, Y. Grischuk, M. Muller, et al., Analyzing the functional and structural consequences of two point mutations (P94L and A368D) in the CYP11B1 gene causing congenital adrenal hyperplasia resulting from 11-hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 91 (2006) 2682–2688.
- [72] S. Polat, A. Kulle, Z. Karaca, et al., Characterisation of three novel CYP11B1 mutations in classic and non-classic 11 β -hydroxylase deficiency, *Eur. J. Endocrinol.* 170 (2014) 697–706.
- [73] K. Joehrer, S. Geley, E.M. Strasser-Wozak, et al., CYP11B1 mutations causing non-classic adrenal hyperplasia due to 11 β -hydroxylase deficiency, *Hum. Mol. Genet.* 6 (1997) 1829–1834.
- [74] N. Reisch, W. Hogler, S. Parajes, et al., A diagnosis not to be missed: nonclassic steroid 11 β -hydroxylase deficiency presenting with premature adrenarche and hirsutism, *J. Clin. Endocrinol. Metab.* 98 (2013) E1620–E1625.
- [75] S. Menabo, S. Polat, L. Baldazzi, et al., Congenital adrenal hyperplasia due to 11 β -hydroxylase deficiency: functional consequences of four CYP11B1 mutations, *Eur. J. Hum. Genet.* 22 (2014) 610–616.
- [76] S. Parajes, L. Loidi, N. Reisch, et al., Functional consequences of seven novel mutations in the CYP11B1 gene: four mutations associated with nonclassic and three mutations causing classic 11 β -hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 95 (2010) 779–788.
- [77] J. Simard, M.L. Ricketts, S. Gingras, P. Soucy, F.A. Feltus, M.H. Melner, Molecular biology of the 3 β -hydroxysteroid dehydrogenase/ Δ^5 – Δ^4 isomerase gene family, *Endocr. Rev.* 26 (2005) 525–582.
- [78] M.W. McBride, A.J. McVie, S.M. Burrige, et al., Cloning, expression, and physical mapping of the 3 β -hydroxysteroid dehydrogenase gene cluster (HSD3BP1–HSD3BP5) in human, *Genomics* 61 (1999) 277–284.
- [79] J.F. Cara, T. Moshang Jr., A.M. Bongiovanni, B.S. Marx, Elevated 17-hydroxyprogesterone and testosterone in a newborn with 3 β -hydroxysteroid dehydrogenase deficiency, *New Engl. J. Med.* 313 (1985) 618–621.
- [80] L. Zhang, J.I. Mason, Y. Naiki, et al., Characterization of two novel homozygous missense mutations involving codon 6 and 259 of type II 3 β -hydroxysteroid dehydrogenase (3 β HSD) gene causing, respectively, nonsalt-wasting and salt-wasting 3 β HSD deficiency disorder, *J. Clin. Endocrinol. Metab.* 85 (2000) 1678–1685.
- [81] E. Rheume, R. Sanchez, J. Simard, et al., Molecular basis of congenital adrenal hyperplasia in two siblings with classical nonsalt-losing 3 β -hydroxysteroid dehydrogenase deficiency, *J. Clin. Endocrinol. Metab.* 79 (1994) 1012–1018.
- [82] A.J. Russell, A.M. Wallace, M.G. Forest, M.D. Donaldson, C.R. Edwards, R.G. Sutcliffe, Mutation in the human gene for 3 β -hydroxysteroid dehydrogenase type II leading to male pseudohermaphroditism without salt loss, *J. Mol. Endocrinol.* 12 (1994) 225–237.
- [83] R. Sanchez, E. Rheume, N. Laflamme, R.L. Rosenfield, F. Labrie, J. Simard, Detection and functional characterization of the novel missense mutation Y254D in type II 3 β -hydroxysteroid dehydrogenase (3 β HSD) gene of a female patient with nonsalt-losing 3 β HSD deficiency, *J. Clin. Endocrinol. Metab.* 78 (1994) 561–567.
- [84] B.B. Mendonca, A.J. Russell, M. Vasconcelos-Leite, et al., Mutation in 3 β -hydroxysteroid dehydrogenase type II associated with pseudohermaphroditism in males and premature pubarche or cryptic expression in females, *J. Mol. Endocrinol.* 12 (1994) 119–122.
- [85] K.J. Matteson, J. Picado-Leonard, B.C. Chung, T.K. Mohandas, W.L. Miller, Assignment of the gene for adrenal P450c17 (steroid 17 α -hydroxylase/17,20 lyase) to human chromosome 10, *J. Clin. Endocrinol. Metab.* 63 (1986) 789–791.
- [86] J. Picado-Leonard, W.L. Miller, Cloning and sequence of the human gene for P450c17 (steroid 17 α -hydroxylase/17,20 lyase): similarity with the gene for P450c21, *DNA* 6 (1987) 439–448.
- [87] B.C. Chung, J. Picado-Leonard, M. Haniu, et al., Cytochrome P450c17 (steroid 17 α -hydroxylase/17,20 lyase): cloning of human adrenal and testis cDNAs indicates the same gene is expressed in both tissues, *Proc. Natl. Acad. Sci. U.S.A.* 84 (1987) 407–411.
- [88] M. Costa-Santos, C.E. Kater, R.J. Auchus, Brazilian Congenital Adrenal Hyperplasia Multicenter Study Group, Two prevalent CYP17 mutations and genotype–phenotype correlations in 24 Brazilian patients with 17-hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 89 (2004) 49–60.
- [89] T. Imai, T. Yanase, M.R. Waterman, E.R. Simpson, J.J. Pratt, Canadian Mennonites and individuals residing in the Friesland region of The Netherlands share the same molecular basis of 17 α -hydroxylase deficiency, *Hum. Genet.* 89 (1992) 95–96.
- [90] K. Miura, K. Yasuda, T. Yanase, et al., Mutation of cytochrome P-45017 α gene (CYP17) in a Japanese patient previously reported as having glucocorticoid-responsive hyperaldosteronism: with a review of Japanese patients with mutations of CYP17, *J. Clin. Endocrinol. Metab.* 81 (1996) 3797–3801.
- [91] C.E. Fardella, L.H. Zhang, P. Mahachoklertwattana, D. Lin, W.L. Miller, Deletion of amino acids Asp487–Ser488–Phe489 in human cytochrome P450c17 causes severe 17 α -hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 77 (1993) 489–493.
- [92] R.J. Auchus, W.L. Miller, Molecular modeling of human P450c17 (17 α -hydroxylase/17,20-lyase): insights into reaction mechanisms and effects of mutations, *Mol. Endocrinol.* 13 (1999) 1169–1182.
- [93] E.L. Van Den Akker, J.W. Koper, A.L. Boehmer, et al., Differential inhibition of 17 α -hydroxylase and 17,20-lyase activities by three novel missense CYP17 mutations identified in patients with P450c17 deficiency, *J. Clin. Endocrinol. Metab.* 87 (2002) 5714–5721.
- [94] M. Bidet, C. Bellanne-Chantelot, M.B. Galand-Portier, et al., Clinical and molecular characterization of a cohort of 161 unrelated women with nonclassical congenital adrenal hyperplasia due to 21-hydroxylase deficiency and 330 family members, *J. Clin. Endocrinol. Metab.* 94 (2009) 1570–1578.
- [95] G.P. Finkelstein, W. Chen, S.P. Mehta, et al., Comprehensive genetic analysis of 182 unrelated families with congenital adrenal hyperplasia due to 21-hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 96 (2011) E161–E172.
- [96] C. Moran, R. Azziz, N. Weintrob, et al., Reproductive outcome of women with 21-hydroxylase-deficient nonclassic adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 91 (2006) 3451–3456.
- [97] S. Nimkarn, M.I. New, Prenatal diagnosis and treatment of congenital adrenal hyperplasia owing to 21-hydroxylase deficiency, *Nat. Clin. Pract. Endocrinol. Metab.* 3 (2007) 405–413.
- [98] K. Sundberg, J. Bang, S. Smidt-Jensen, et al., Randomised study of risk of fetal loss related to early amniocentesis versus chorionic villus sampling, *Lancet* 350 (1997) 697–703.
- [99] F. Mujezinovic, Z. Alfirevic, Procedure-related complications of amniocentesis and chorionic villous sampling: a systematic review, *Obstet. Gynecol.* 110 (2007) 687–694.
- [100] M. Kollmann, M. Haeusler, J. Haas, B. Csapo, U. Lang, P. Klaritsch, Procedure-related complications after genetic amniocentesis and chorionic villus sampling, *Ultraschall Med.* 34 (2013) 345–348.
- [101] S.F. Witchel, W.L. Miller, Prenatal treatment of congenital adrenal hyperplasia—not standard of care, *J. Genet. Couns.* 21 (2012) 615–624.
- [102] M.I. New, A. Carlson, J. Obeid, et al., Prenatal diagnosis for congenital adrenal hyperplasia in 532 pregnancies, *J. Clin. Endocrinol. Metab.* 86 (2001) 5651–5657.
- [103] T. Hirvikoski, A. Nordenstrom, T. Lindholm, et al., Cognitive functions in children at risk for congenital adrenal hyperplasia treated prenatally with dexamethasone, *J. Clin. Endocrinol. Metab.* 92 (2007) 542–548.
- [104] E.P. Davis, C.A. Sandman, C. Buss, D.A. Wing, K. Head, Fetal glucocorticoid exposure is associated with preadolescent brain development, *Biol. Psychiatry* 74 (2013) 647–655.
- [105] W.L. Miller, S.F. Witchel, Prenatal treatment of congenital adrenal hyperplasia: risks outweigh benefits, *Am. J. Obstet. Gynecol.* 208 (2013) 354–359.
- [106] T. Hirvikoski, A. Nordenstrom, A. Wedell, M. Ritzen, S. Lajic, Prenatal dexamethasone treatment of children at risk for congenital adrenal hyperplasia: the Swedish experience and standpoint, *J. Clin. Endocrinol. Metab.* 97 (2012) 1881–1883.
- [107] Y.M. Lo, N. Corbetta, P.F. Chamberlain, et al., Presence of fetal DNA in maternal plasma and serum, *Lancet* 350 (1997) 485–487.
- [108] R.J. Rijnders, C.E. van der Schoot, B. Bossers, M.A. de Vroede, G.C. Christiaens, Fetal sex determination from maternal plasma in pregnancies at risk for congenital adrenal hyperplasia, *Obstet. Gynecol.* 98 (2001) 374–378.
- [109] M.I. New, Y.K. Tong, T. Yuen, et al., Noninvasive prenatal diagnosis of congenital adrenal hyperplasia using cell-free fetal DNA in maternal plasma, *J. Clin. Endocrinol. Metab.* 99 (2014) E1022–E1030.
- [110] O. Lekarev, K. Tafuri, A.H. Lane, et al., Erroneous prenatal diagnosis of congenital adrenal hyperplasia owing to a duplication of the CYP21A2 gene, *J. Perinatol.* 33 (2013) 76–78.

- [111] R.A. Lobo, U. Goebelsmann, Evidence for reduced 3β -ol-hydroxysteroid dehydrogenase activity in some hirsute women thought to have polycystic ovary syndrome, *J. Clin. Endocrinol. Metab.* 53 (1981) 394–400.
- [112] S.Y. Pang, A.J. Lerner, E. Stoner, et al., Late-onset adrenal steroid 3β -hydroxysteroid dehydrogenase deficiency. I. A cause of hirsutism in pubertal and postpubertal women, *J. Clin. Endocrinol. Metab.* 60 (1985) 428–439.
- [113] S.F. Siegel, D.N. Finegold, R. Lanes, P.A. Lee, ACTH stimulation tests and plasma dehydroepiandrosterone sulfate levels in women with hirsutism, *New Engl. J. Med.* 323 (1990) 849–854.
- [114] T. Eldar-Geva, A. Hurwitz, P. Vecsei, Z. Palti, A. Milwidsky, A. Rosler, Secondary biosynthetic defects in women with late-onset congenital adrenal hyperplasia, *New Engl. J. Med.* 323 (1990) 855–863.
- [115] C. Lutfallah, W. Wang, J.I. Mason, et al., Newly proposed hormonal criteria via genotypic proof for type II 3β -hydroxysteroid dehydrogenase deficiency, *J. Clin. Endocrinol. Metab.* 87 (2002) 2611–2622.
- [116] W. Arlt, D.S. Willis, S.H. Wild, et al., Health status of adults with congenital adrenal hyperplasia: a cohort study of 203 patients, *J. Clin. Endocrinol. Metab.* 95 (2010) 5110–5121.
- [117] G.P. Finkelstein, M.S. Kim, N. Sinaii, et al., Clinical characteristics of a cohort of 244 patients with congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 97 (2012) 4429–4438.
- [118] A. Mallappa, N. Sinaii, P. Kumar, et al., A phase 2 study of chronocort, a modified-release formulation of hydrocortisone, in the treatment of adults with classic congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 100 (2015) 1137–1145.
- [119] G. Johannsson, R. Bergthorsdottir, A.G. Nilsson, H. Lennernas, T. Hedner, S. Skrtic, Improving glucocorticoid replacement therapy using a novel modified-release hydrocortisone tablet: a pharmacokinetic study, *Eur. J. Endocrinol.* 161 (2009) 119–130.
- [120] S.M. Bryan, J.W. Honour, P.C. Hindmarsh, Management of altered hydrocortisone pharmacokinetics in a boy with congenital adrenal hyperplasia using a continuous subcutaneous hydrocortisone infusion, *J. Clin. Endocrinol. Metab.* 94 (2009) 3477–3480.
- [121] P.C. Hindmarsh, The child with difficult to control congenital adrenal hyperplasia: is there a place for continuous subcutaneous hydrocortisone therapy, *Clin. Endocrinol. (Oxford)* 81 (2014) 15–18.
- [122] R.J. Auchus, A.F. Turcu, J.L. Spencer-Segal, et al., A Pharmacokinetic and Biomarker Study of the Corticotropin-Releasing Factor Receptor Antagonist NBI-77860 in Adult Females with Classic 21-Hydroxylase Deficiency, *Endocr. Rev.* 36 (2015) OR06-6.
- [123] L. Laue, D.P. Merke, J.V. Jones, K.M. Barnes, S. Hill, G.B. Cutler Jr., A preliminary study of flutamide, testolactone, and reduced hydrocortisone dose in the treatment of congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 81 (1996) 3535–3539.
- [124] D.P. Merke, M.F. Keil, J.V. Jones, J. Fields, S. Hill, G.B. Cutler Jr., Flutamide, testolactone, and reduced hydrocortisone dose maintain normal growth velocity and bone maturation despite elevated androgen levels in children with congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 85 (2000) 1114–1120.
- [125] K. Lin-Su, M.G. Vogiatzi, I. Marshall, et al., Treatment with growth hormone and luteinizing hormone releasing hormone analog improves final adult height in children with congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 90 (2005) 3318–3325.
- [126] R.J. Auchus, E.O. Buschur, A.Y. Chang, et al., Abiraterone acetate to lower androgens in women with classic 21-hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 99 (2014) 2763–2770.
- [127] C. Kamrath, Z. Hochberg, M.F. Hartmann, T. Remer, S.A. Wudy, Increased activation of the alternative backdoor pathway in patients with 21-hydroxylase deficiency: evidence from urinary steroid hormone analysis, *J. Clin. Endocrinol. Metab.* 97 (2012) E367–E375.
- [128] F. Ferreira, J.M. Martins, S. do Vale, R. Esteves, G. Nunes, I. Carmo, Rare and severe complications of congenital adrenal hyperplasia due to 21-hydroxylase deficiency: a case report, *J. Med. Case Rep.* 7 (2013) 39.
- [129] I. Nermoen, I. Folling, K. Vegge, et al., Two adults with adrenal myelolipoma and 21-hydroxylase deficiency, *Case Rep. Med.* 2009 (2009) 916891.
- [130] L.M. Mermejo, J. Elias Junior, F.P. Saggioro, et al., Giant adrenal myelolipoma associated with 21-hydroxylase deficiency: unusual association mimicking an androgen-secreting adrenocortical carcinoma, *Arq. Bras. Endocrinol. Metabol.* 54 (2010) 419–424.
- [131] P.W. Speiser, J. Dupont, D. Zhu, et al., Disease expression and molecular genotype in congenital adrenal hyperplasia due to 21-hydroxylase deficiency, *J. Clin. Invest.* 90 (1992) 584–595.
- [132] N. Krone, A. Braun, A.A. Roscher, D. Knorr, H.P. Schwarz, Predicting phenotype in steroid 21-hydroxylase deficiency? Comprehensive genotyping in 155 unrelated, well defined patients from southern Germany, *J. Clin. Endocrinol. Metab.* 85 (2000) 1059–1065.
- [133] J.D. Wilson, R.J. Auchus, M.W. Leihy, et al., 5α -androstane- $3\alpha,17\beta$ -diol is formed in tammar wallaby pouch young testes by a pathway involving 5α -pregnane- $3\alpha,17\alpha$ -diol-20-one as a key intermediate, *Endocrinology* 144 (2003) 575–580.
- [134] R.J. Auchus, The backdoor pathway to dihydrotestosterone, *Trends Endocrinol. Metab.* 15 (2004) 432–438.
- [135] K. Homma, T. Hasegawa, T. Nagai, et al., Urine steroid hormone profile analysis in cytochrome P450 oxidoreductase deficiency: implication for the backdoor pathway to dihydrotestosterone, *J. Clin. Endocrinol. Metab.* 91 (2006) 2643–2649.
- [136] J. Rege, Y. Nakamura, F. Satoh, et al., Liquid chromatography-tandem mass spectrometry analysis of human adrenal vein 19-carbon steroids before and after ACTH stimulation, *J. Clin. Endocrinol. Metab.* 98 (2013) 1182–1188.