

LETTER TO THE EDITOR

Sublingual minoxidil for the treatment of male and female pattern hair loss: a randomized, double-blind, placebo-controlled, phase 1B clinical trial

Dear Editor

Minoxidil is an approved medication for severe hypertension androgenetic alopecia.¹ Oral minoxidil (OM) is a potent vasodilator used to treat hypertension. It is a pro-drug, activated by hepatic dehydroepiandrosterone sulfotransferase (SULT2A1) to minoxidil sulfate.² OM doses of 2.5 and 5 mg produced peak plasma concentrations of 16.8 and 37.2 ng/mL within 30 min post-dose.³ Side-effects include hypertrichosis, lower limb oedema, postural hypotension and tachycardia. Minoxidil lotion (ML) is approved for the treatment of hair loss. It is also a pro-drug converted in hair bulbs by thermostable phenol sulfotransferase (SULT1A1) to minoxidil sulphate. There is a considerable inter-subject variability in levels of both hepatic SULT2A1 and follicular SULT1A1. Low SULT1A1 predicts weak hair regrowth with both ML and OM.^{4,5} Weak hair growth due to low follicular SULT1A1 can be overcome with OM through dose escalation but cannot be overcome with ML due to low solubility and saturation absorption kinetics.⁶

As sublingual dosing bypasses hepatic first-pass metabolism, sublingual minoxidil (SM) would be anticipated to increase follicular minoxidil sulphate bioavailability and consequently hair growth.⁷ As hepatic sulphation of minoxidil enhances the haemodynamic effect, SM would also be anticipated to reduce haemodynamic side effects and consequently improve safety.

To investigate SM as an alternative to OM, we conducted a prospective, randomized placebo-controlled, double-blinded dose-escalation phase 1B clinical trial (HREC approval number 2017-09-669; ANZCTR number: ACTRN12618000606280). Forty men with Hamilton Norwood stage III vertex to V or women with Sinclair stages 2–5 hair loss, aged 30–65 years, were enrolled and completed the core study receiving either SM 0.45 mg daily or placebo for 24 weeks. Twelve participants rolled into a 24-week open-label extension study and received SM 1.35 or 4.05 mg daily. Co-primary end-points were macrophotography and terminal hair count on phototrichogram.

Macro photographs were scored using a 7-point scale. At 24 weeks, nine patients (45%) receiving 0.45-mg SM had improved frontal hair density, and 11 patients (55%) showed

vertex improvement. One out of six patients (17%) treated with 1.35-mg SM had improved frontal, and 3 out of 6 (50%) had an improved vertex hair density. Four of six patients (67%) treated with 4.05-mg dose showed an improvement in both frontal and vertex hair density (Fig. 1).

Phototrichograms demonstrated a mean increase in terminal hair count/cm² of 4 for the frontal and 9 for the vertex scalp with the 0.45-mg dose. The 1.35-mg dose produced a mean increased terminal hair count/cm² of 10 and 26, and 4.05-mg SM produced a mean increase terminal hair count/cm² of 38 and 88 for the frontal and vertex scalp, respectively (Fig. 2). Compared to placebo, the difference in mean hair count 0 achieved statistical significance for both the frontal and vertex scalp at all doses except the 0.45-mg dose over the vertex.

Failure to achieve statistical significance at the 0.45-mg dose on the vertex scalp is attributed to a solitary patient in the placebo group, whose vertex scalp hair density unexpectedly increased by 76 hair per cm².

Mean peak minoxidil plasma concentration following the initial sublingual dose was at 1.62 ng/mL (range 0.3–5.3 ng/mL), 30 min post-dose. Minoxidil was undetectable in plasma after 24 h. This is more than an order of magnitude below the plasma concentration threshold (20 ng/mL) for the development of any haemodynamic effects.⁸ No significant change in BP was detected in the placebo or treatment arms.

In conclusion, SM produced a dose-dependent increase in mean terminal hair count on the frontal and vertex scalp and an improvement in hair density. Further studies with larger patient cohorts are warranted to determine the optimal dose of SM and comparing the relative efficacy of OM and SM, as well as the pharmacokinetics of SM.

Acknowledgement

We acknowledge the contribution of Dr Nekma Meah, Dr Dmitri Wall, Dr Bevin Bhoyrul, Dr Lara Carvalho, Dr Janina Poa as blinded assessors for the global scalp photographs. The patients in this manuscript have given written informed consent to the publication of their case details.

Conflict of interest

No conflict of interests has been identified for the remaining authors.

Author contribution

RS: Director and Founder Samson Medical Pty Ltd and holds the patent for the use of sublingual minoxidil and low dose oral




Figure 1 Representative global photographs of the frontal scalp of patients in each arm at Week 0 and Week 24.



minoxidil for promoting hair growth and treatment of hair loss or excessive hair shedding. Pharmaceutical advisory board Eli Lilly, Pfizer, Leo Pharmaceutical. Speaker bureau Abbvie,

Novartis. Principal investigator in clinical trials for Amgen, Novartis, Arcutis, Aerotech, Merck and Co, Celgene, Coherus Bioscience, Janssen, Regeneron, MedImmune, Glaxo Smith


Figure 2 Representative phototrichograms of frontal and vertex macro photographs of patients on sublingual minoxidil (a) 0.45 mg arm (b) 1.35 mg arm and (c) 4.05 mg arm at Week 0 and Week 24.

(a) 0.45mg

Week 0




Week 24




(b) 1.35mg

Week 0



Week 24




(c) 4.05mg

Week 0



Week 24



Kline, Samson Clinical, Boehringer Ingelheim, Oncobiologics, Roche, Ascend, Demira, Astra Zeneca, Akesobio, Reistone, UCB, Sanofi, Connect, Arena, Sun Pharma, Bristol Myer Squibb, Galderma.

Funding

Nil.

L. Bokhari,^{1,*}  L.N. Jones,¹  R.D. Sinclair^{1,2} 

¹Sinclair Dermatology, 2/2 Wellington Parade, East Melbourne, Vic., 3002, Australia, ²Department of Medicine, University of Melbourne

*Correspondence: Department of Medicine, St Vincent's Hospital, Clinical Sciences Building, Level 4 / 29 Regent Street, Fitzroy, VIC 3065, Australia.

E-mail: laita.bokhari@sinclairdermatology.com.au

References

- Lowenthal DT, Affrime MB. Pharmacology and pharmacokinetics of minoxidil. *J Cardiovasc Pharmacol* 1980; 2(Suppl 2): S93–S106.
- Anderson RJ, Kudlacek PE, Clemens DL. Sulfation of minoxidil by multiple human cytosolic sulfotransferases. *Chem Biol Interact* 1998; 109: 53–67.
- Fleishaker JC, Andreadis NA, Welshman IR, Wright CE. The pharmacokinetics of 2.5- to 10-mg oral doses of minoxidil in healthy volunteers. *J Clin Pharmacol* 1989; 29: 162–167.
- Ramos PM, Sinclair R, Miot HA, Goren A. Sulfotransferase activity in plucked hair follicles predicts response to topical minoxidil treatment in Brazilian female pattern hair loss patients. *Dermatol Ther* 2020; 33: e13195.
- Ramos PM, Gohad P, McCoy J, Wambier C, Goren A. Minoxidil Sulfotransferase Enzyme (SULT1A1) genetic variants predicts response to oral minoxidil treatment for female pattern hair loss. *J Eur Acad Dermatol Venereol* 2020; 35: e24–e26.
- Jimenez-Cauhe J, Saceda-Corralo D, Rodrigues-Barata R *et al.* Effectiveness and safety of low-dose oral minoxidil in male androgenetic alopecia. *J Am Acad Dermatol* 2019; 81: 648–649.
- Sinclair R, Trindade de Carvalho L, Ferial Ismail F, Meah N. Treatment of male and female pattern hair loss with sublingual minoxidil: a retrospective case-series of 64 patients. *J Eur Acad Dermatol Venereol* 2020; 34: e795–e796.
- Ferry JJ, Turner SW, Albert DG, Dietz AJ, Luderer JR. Hemodynamic effects of minoxidil following intravenous infusions in untreated hypertensive patients. *Clin Pharmacol Ther* 1996; 59: 166.

DOI: 10.1111/jdv.17623