

CANCER

Colocalization of Inflammatory Response with B7-H1 Expression in Human Melanocytic Lesions Supports an Adaptive Resistance Mechanism of Immune Escape

Janis M. Taube,^{1,2*} Robert A. Anders,² Geoffrey D. Young,^{3,4} Haiying Xu,¹ Rajni Sharma,² Tracee L. McMiller,⁴ Shuming Chen,⁴ Alison P. Klein,^{2,5} Drew M. Pardoll,⁵ Suzanne L. Topalian,^{4*} Lieping Chen^{1,5,6*}

Although many human cancers such as melanoma express tumor antigens recognized by T cells, host immune responses often fail to control tumor growth for as yet unexplained reasons. Here, we found a strong association between melanocyte expression of B7-H1 (PD-L1), an immune-inhibitory molecule, and the presence of tumor-infiltrating lymphocytes (TILs) in human melanocytic lesions: 98% of B7-H1⁺ tumors were associated with TILs compared with only 28% of B7-H1⁻ tumors. Indeed, B7-H1⁺ melanocytes were almost always localized immediately adjacent to TILs. B7-H1/TIL colocalization was identified not only in melanomas but also in inflamed benign nevi, indicating that B7-H1 expression may represent a host response to tissue inflammation. Interferon- γ , a primary inducer of B7-H1 expression, was detected at the interface of B7-H1⁺ tumors and TILs, whereas none was found in B7-H1⁻ tumors. Therefore, TILs may actually trigger their own inhibition by secreting cytokines that drive tumor B7-H1 expression. Consistent with this hypothesis, overall survival of patients with B7-H1⁺ metastatic melanoma was significantly prolonged compared with that of patients with B7-H1⁻ metastatic melanoma. Therefore, induction of the B7-H1/PD-1 pathway may represent an adaptive immune resistance mechanism exerted by tumor cells in response to endogenous antitumor activity and may explain how melanomas escape immune destruction despite endogenous antitumor immune responses. These observations suggest that therapies that block this pathway may benefit patients with B7-H1⁺ tumors.

INTRODUCTION

Melanoma is considered an immunogenic neoplasm, and tumor-specific T cells can often be isolated from both peripheral blood and melanoma lesions. These T cells recognize tumor-associated or tumor-specific antigens (1). However, most melanomas will progress if left untreated, and spontaneous regression is rare (2), which suggests that endogenous immunity fails to control tumor growth. Local factors within the tumor microenvironment, such as secreted inhibitory cytokines (for example, transforming growth factor- β), regulatory T cells and myeloid-derived suppressor cells, and cell membrane-bound immunomodulatory proteins, have been proposed to contribute to creating a suppressive milieu that protects tumor cells from immune destruction. Indeed, the co-inhibitory molecule B7-H1 (also called PD-L1), which is often up-regulated on tumor cells, impairs T cell responses—leading to anergy, exhaustion, or apoptosis upon engagement with its cognate co-inhibitory receptor PD-1, which is often highly expressed on tumor-infiltrating lymphocytes (TILs) (3–8).

Current clinical efforts in immunotherapy are directed toward overcoming systemic and local immunoregulation, which is thought to have thwarted cancer vaccines, adoptive cellular therapies, and other anti-tumor immunotherapies in the past (9). Ipilimumab, a blocking mono-

clonal antibody (mAb) directed against the co-inhibitory T cell receptor CTLA-4 (cytotoxic T lymphocyte-associated antigen 4), systemically activates T cells, which led to enhancement of antitumor T cell responses and demonstrated a survival benefit in patients with advanced metastatic melanoma in randomized trials (10, 11). However, serious immune-related adverse events have been experienced by 15 to 30% of patients receiving this therapy (12–14).

The modest therapeutic window for anti-CTLA-4 therapy is due in part to the complete lack of tumor-specific expression of its cognate ligands, B7.1 and B7.2. In contrast, the ligand-receptor interactions of the B7-H1/PD-1 axis occur more selectively within the tumor microenvironment. The B7 Homolog One (B7-H1, also called PD-L1) protein is not detectable in most normal tissues (5, 15–17); however, its expression can be induced to a high level in various cell types by pro-inflammatory cytokines, particularly interferon- γ (IFN- γ) (5, 18), and B7-H1 is highly expressed on many human tumors, including melanoma (3). In addition to the immunosuppressive effects of B7-H1 binding to PD-1 on activated T and B cells, it also mediates suppressive functions via interactions with CD80 on activated T cells (19). Whereas CTLA-4^{-/-} mice die at a young age from uncontrolled lymphoproliferation, B7-H1^{-/-} mice demonstrate only a mild lymphocyte accumulation in vital organs with no signs of autoimmune disease (20, 21). PD-1^{-/-} mice develop late-onset organ-specific autoimmunity, which is strain-dependent (22). Therefore, blockade of the B7-H1/PD-1 axis may be a more tumor microenvironment-specific and less toxic approach to reversing tumor-induced immune tolerance in the clinic than CTLA-4 blockade.

B7-H1 expression has been observed in various solid malignancies, including melanoma, squamous cell carcinoma of the head and neck, and

¹Department of Dermatology, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA. ²Department of Pathology, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA. ³Department of Otolaryngology, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA. ⁴Department of Surgery, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA. ⁵Department of Oncology, Johns Hopkins Medical Institutions, Baltimore, MD 21287, USA. ⁶Department of Immunobiology, Yale University, New Haven, CT 06519, USA.

*To whom correspondence should be addressed. E-mail: jtaube1@jhmi.edu (J.M.T.); stopali1@jhmi.edu (S.L.T.); lieping.chen@yale.edu (L.C.)

carcinomas of the esophagus, ovary, gastrointestinal tract, breast, lung, and kidney (3, 23). In some tumor types, such as renal cell, esophageal, gastric, and pancreatic carcinomas, cell surface expression (also referred to as “membranous” expression) by more than 5 to 10% of primary tumor cells has been reported to be an independent predictor of adverse patient outcomes (24–26). Additionally, expression of B7-H1 by TILs has been reported to correlate with decreased patient survival (27, 28), suggesting that B7-H1 expression in the tumor microenvironment may exert an immunosuppressive function. Moreover, early clinical trials of a fully human mAb against PD-1 have demonstrated clinical activity in patients with advanced melanoma and other cancers, associated with generally manageable side effects (29).

Here, we used immunohistochemistry (IHC) with a specific mAb to human B7-H1 to examine patterns of membranous B7-H1 expression by both melanocytes/melanoma cells and TILs in 150 melanocytic lesions of various malignant potentials, including nevi, melanomas in situ, primary melanomas of various histologic subtypes, and regional and distant metastases. A significant correlation was observed between the presence of TILs and B7-H1 expression in the tumor microenvironment, which was associated with local production of the inflammatory cytokine IFN- γ . These data support an adaptive mechanism rather than constitutive oncogene-driven expression of B7-H1.

RESULTS

Patient demographics and follow-up

Patient demographic information is provided in Table 1. Tissue samples were obtained from 150 patients (74 women and 76 men) treated at Johns Hopkins Hospital over 20 years. Patients had a median age of 53 years (range, 7 to 94 years). Neither patient gender nor the pathologic subtype of the melanocytic lesion was significantly associated with the degree of melanocyte B7-H1 expression. At the time of last follow-up, 54 of 110 patients with melanoma in situ, invasive primary melanoma, or metastatic melanoma were alive. Median follow-up for the patients who were alive was 100 months (range, 24 to 319 months).

Table 1. Correlation of B7-H1 expression by melanocytes with clinicopathologic features in 150 patients.

	Number of patients			<i>p</i> *
	Total	B7-H1 ⁺ (≥5%)	B7-H1 [−] (<5%)	
All patients	150	57	93	0.867
Gender				
Male	76	28	48	
Female	74	29	45	0.378
Lesion type				
Nevus	40	14	26	
Melanoma in situ	11	6	5	
Invasive primary melanoma	43	13	30	
Metastasis	56	24	32	

* χ^2 Analysis.

Of the 56 melanoma patients who expired, the median time to death was 18 months (range, 1 to 220 months).

Patterns of B7-H1 expression in melanocytic lesions

Membranous B7-H1 expression by melanocytes within the tumors (including benign melanocytes and malignant melanoma cells) demonstrated three major patterns on IHC, as shown in Fig. 1: the absence of B7-H1 (Fig. 1A); regional expression of B7-H1 on melanocytes, highly colocalized with TILs (Fig. 1, B and C, and fig. S1, B and C); and B7-H1 expression in the presence of rare or no TILs (Fig. 1D and fig. S1D). B7-H1 expression at the interface with TILs was by far the predominant pattern and often occurred in discrete geographic foci (Fig. 2, A and B). Among 150 melanocytic lesions spanning the spectrum of benign nevi to metastatic melanoma, 57 (38%) were B7-H1⁺. Of these, only a single case of metastatic melanoma demonstrated broad B7-H1 expression unassociated with TILs. Additionally, only one case of primary melanoma and two metastases demonstrated mixed patterns, including distinct regions of B7-H1⁺ tumor cells either associated or not with TILs.

Membranous B7-H1 expression was also observed in the TILs and associated macrophages/histiocytes in the host response to tumor




Fig. 1. Patterns of B7-H1 expression observed in melanocytic lesions stained with the anti-B7-H1 mAb 5H1. (A) No B7-H1 expression (brown chromogen) by melanocytes in a benign nevus. Original magnification, $\times 200$ (scale bar, 50 μ m). Note the paucity of TILs in this case. (B) B7-H1 expression by both melanocytes and TILs at the advancing edge of an invasive primary melanoma, nodular histologic subtype. Original magnification, $\times 40$ (scale bar, 200 μ m). (C) Original magnification of the boxed area shown in (B), $\times 400$ (scale bar, 20 μ m). Overall, 10% of total melanoma cells in this specimen expressed B7-H1 (scoring was performed as described in Materials and Methods). The inflammatory host response to tumor was graded as “moderate” in this case and included B7-H1⁺ TILs. (D) Diffuse B7-H1 expression by melanocytes in a subcutaneous melanoma metastasis to the scalp, associated with singular TILs. Original magnification, $\times 200$ (scale bar, 50 μ m). Representative photomicrographs of CD3 staining for TILs corresponding to each of these examples are shown in fig. S1. The interface pattern with B7-H1 expression by both melanocytes and TILs at the advancing edge predominated in this study.

Downloaded from https://www.science.org on October 15, 2026

(Fig. 1C and fig. S2). In some cases, lymphohistiocytic aggregates unassociated with melanocytes also expressed B7-H1. For example, 3 of the 10 common melanocytic nevi examined showed lymphohistiocytic aggregates within the biopsy specimen (most often located perivascularly and not infiltrating among melanocytes), with some degree of B7-H1 positivity in the macrophages and in occasional lymphocytes.

Quantification of B7-H1 expression by melanocytes and the associated host inflammatory response

Formalin-fixed, paraffin-embedded (FFPE) tissue sections from each case studied were evaluated for B7-H1 expression. Among 150 melanocytic lesions, 93 were negative for membranous B7-H1 expression

(defined as $<5\%$ B7-H1⁺ cells) (Fig. 3A). A 5% expression threshold was chosen to define B7-H1 positivity, in keeping with previous studies (26, 30). Thirty-eight percent (57 of 150) of the study specimens were above this threshold, including 35% (14 of 40) of nevi and 39% (43 of 110) of malignant lesions (Table 1). Thus, B7-H1 expression on melanocytic cells did not correlate with benign compared with malignant lesions.

A similar analysis was conducted for expression of B7-H1 in infiltrating host inflammatory cells (TILs and associated macrophages/histiocytes), as expressed by the adjusted inflammation score (AIS; density of inflammation multiplied by the percent B7-H1 expression by infiltrating host inflammatory cells) (Fig. 3B). Only true intratumoral

immune infiltrates, and not perivascular inflammation, were recorded in the AIS. A threshold AIS of 5 was chosen to parallel the threshold designated for melanocytes. Seventy-seven of 150 specimens (51%) exhibited an AIS of <5 , reflecting either absent/scant infiltrating immune cells or absent/scant B7-H1 expression on infiltrating cells. Seventy-three of 150 (49%) samples, including nevi, demonstrated an AIS of ≥ 5 , and 51% (56 of 110) of malignant lesions were above this threshold. Therefore, similar to findings with B7-H1 expression by melanocytes, the AIS did not correlate with the malignant potential of the lesion.

Association of B7-H1 expression by melanocytes with immune infiltrates

To explore a potential association between B7-H1 expression by melanocytes and the geographic proximity of infiltrating immune cells (Figs. 1, B and C, and 2 and figs. S1 and S2), we compared the degree of B7-H1 cell surface (membranous) expression based on IHC (negative, defined as $<5\%$ positive tumor cells; positive, $\geq 5\%$) to the intensity of associated infiltrates (Table 2). TILs and associated macrophages/histiocytes were scored as mild (rare), moderate (focal), or severe (diffuse), based on hematoxylin and eosin (H&E) staining. We found that 35% (14 of 40) of benign nevi (2 inflamed Spitz nevi, 2 severely dysplastic nevi, and 10 halo nevi) expressed B7-H1 (mean melanocyte expression in positive samples, 30%; range, 10 to 70%). Notably, 100% (14 of 14) of the B7-H1⁺ cases were associated with at least mild TILs compared to only 15% (4 of 26) of B7-H1⁻ cases ($P < 0.0001$).

Similarly, 35% (19 of 54) of malignant primary lesions (in situ and invasive melanomas) were B7-H1⁺ (mean melanocyte expression, 14%; range, 5 to 50%),




Fig. 2. Geographic association of B7-H1 expression and TILs. **(A)** Primary nodular melanoma with associated “severe” grade of lymphocytic infiltration, highlighted by CD8 immunostaining. The dark line demarcates the area of colocalizing T cells and B7-H1⁺ tumor cells. Overall, 30% of the tumor cells demonstrated cell surface B7-H1 expression. Note that the melanoma cell cytoplasm is heavily pigmented due to abundant melanin (see isotype control). Original magnifications, $\times 200$ for left and middle column panels (scale bars, 50 μm) and $\times 400$ for right column panels (scale bars, 25 μm). **(B)** Metastatic deposit of melanoma in a lymph node, associated with “mild” grade of lymphocyte infiltration shown by CD3 immunostaining. Original magnifications, $\times 100$ for left column panels (scale bars, 100 μm), $\times 200$ for middle column panels (scale bars, 50 μm), and $\times 400$ for right column panels (scale bars, 25 μm). Arrows highlight the geographic area of concordance of CD3⁺ TILs with B7-H1 expression by melanoma, and also point to the area shown at higher magnification in the right-hand panels. Five percent of tumor cells demonstrated B7-H1 expression in this case.

and 100% (19 of 19) of the positive cases were associated with at least mild TILs compared to 43% (15 of 35) of B7-H1⁻ cases ($P < 0.0001$). Among metastatic melanoma lesions, 43% (24 of 56) were B7-H1⁺ (mean melanoma cell expression in positive samples, 15%; range, 5 to 60%), and 96% of those samples (23 of 24) were associated with at least mild TILs compared to 22% (7 of 32) of B7-H1⁻ cases ($P < 0.0001$). In total, we found that 98% (56 of 57) of B7-H1⁺ cases were

associated with at least mild TILs compared to 28% (26 of 93) of B7-H1⁻ cases ($P < 0.0001$).

When all 150 specimens were analyzed, B7-H1 expression of $\geq 5\%$ was significantly associated with the presence of inflammatory infiltrates ($P < 0.0001$, Fisher's exact test) (Table 2). This association was confirmed in subset analyses for nevi, primary in situ and invasive melanoma, and metastatic disease. The degree of the inflammatory infiltrate (mild, moderate, severe) did not vary by melanoma stage (in situ, invasive, or metastatic) ($P = 0.60$, Fisher's exact test) but was significantly correlated with the amount of B7-H1 expression by tumor (Spearman's $\rho = 0.71$, $P < 0.001$). These data demonstrate a strong association between B7-H1 expression and immune cell infiltration in both nevi and malignant melanoma.

We also assessed whether immune infiltrates associated with melanocytes expressed B7-H1, as indicated by the AIS. By univariate analysis, the AIS was highly associated with B7-H1 expression by all types of melanocytic lesion studied ($P < 0.001$). When analyzed by subtype of malignant lesion, this association was statistically significant for invasive primary melanomas ($P < 0.001$) and metastatic melanomas ($P < 0.001$) and had borderline significance for in situ disease ($P = 0.08$). Therefore, when a melanocytic lesion is associated with a host immune response, both components are likely to express B7-H1.




Fig. 3. Distribution of intensity of B7-H1 expression by melanocytic cells and immune infiltrates in 150 lesions. **(A)** Percent B7-H1 expression by melanocytes $\geq 5\%$ was considered positive. **(B)** AIS, defined as the intensity of intratumoral inflammation including TILs and histiocytes (graded 0 to 3, see text) multiplied by the percent B7-H1⁺ inflammatory cells.

IFN- γ detected at the interface of TILs and B7-H1⁺ tumor cells by laser capture microdissection and quantitative reverse transcription-polymerase chain reaction

The high correlation between TIL infiltration and B7-H1 expression by melanocytes among 150 lesions of various malignant potentials, and geographically within individual tumors, suggested that, as with inflamed tissues, B7-H1 expression was induced by inflammatory cytokines such as IFN- γ . To investigate the presence of IFN- γ -producing immune cells that could potentially drive B7-H1 expression in melanomas, we performed quantitative reverse transcription-polymerase chain reaction (qRT-PCR) for the leukocyte common antigen CD45 (protein tyrosine phosphatase receptor type C) and IFN- γ in laser capture microdissection (LCM)-dissected specimens from both B7-H1⁻ and B7-H1⁺ tumors at the interface with TILs (Table 3). Fresh peripheral blood mononuclear cells (PBMCs) and activated T cells, and cultured melanoma cells, were used as positive and negative controls for IFN- γ and CD45 expression, respectively. Although CD45 mRNA was detected by qRT-PCR in all five B7-H1⁻ specimens examined, indicating the

Table 2. Correlation of B7-H1 expression by melanocytes with the presence of immune cell infiltration.

Histology	Total	Number of cases/total cases (%)				P*
		B7-H1 ⁺⁺		B7-H1 ⁻		
		TIL ⁺⁺	TIL ⁻	TIL ⁺	TIL ⁻	
Benign nevi	40	14/14 (100)	0/14 (0)	4/26 (15)	22/26 (85)	<0.0001
Primary melanomas (in situ or invasive)	54	19/19 (100)	0/19 (0)	15/35 (43)	20/35 (57)	<0.0001
Metastases	56	23/24 (96)	1/24 (4)	7/32 (22)	25/32 (78)	<0.0001
All	150	56/57 (98)	1/57 (2)	26/93 (28)	67/93 (72)	<0.0001

*Fisher's exact test, two-sided, was conducted on the 2×2 matrix defined by B7-H1 ($\pm$) expression and TIL ($\pm$) for each lesion type.
 †Including mild, moderate, and severe lymphocyte infiltrates and their associated histiocytes/macrophages.

‡More than 5% melanocytes with membranous expression on IHC.

Table 3. Specific expression of IFN- γ in B7-H1⁺ melanomas. IFN- γ was detected by qRT-PCR as described in Materials and Methods. These results are representative of two separate experiments. UD, undetectable (C_T greater than 45 cycles); NA, not applicable.

Patient number	Tissue source	C_T (mean)*	SEM†	C_T (mean)	SEM
B7-H1 ⁻ tumors					
1	Lymph node metastasis	UD‡	NA	33.19	0.45
2	Small bowel metastasis	UD§	NA	34.46	0.38
3	Lymph node metastasis	UD	NA	32.93	0.11
4	Lymph node metastasis	UD	NA	35.53	0.28
5	Cutaneous primary lesion	UD	NA	36.11	0.13
B7-H1 ⁺ tumors					
6	Lymph node metastasis	34.67	0.49	32.41	0.16
7	Lymph node metastasis	35.00	0.32	33.55	0.36
8	Lung metastasis	33.15	0.30	31.91	0.78
9	Lung metastasis	31.80	0.07	28.43	0.02
10	Lymph node metastasis	35.17	0.22	32.51	0.22
Controls					
NA	PBMCs	33.12	0.35	28.55	0.66
NA	Activated CD3 ⁺ T cells	23.60	0.08	28.33	0.11
NA	1359-mel	UD	NA	UD	NA

*Lower values reflect greater expression. †SEM of triplicate samples. ‡One of three triplicate reactions for patient 1 tumor had $C_T = 41.7$, and two of three were undetectable and average undetectable. §One of three triplicate reactions for patient 2 tumor had $C_T = 43.1$, and two of three were undetectable and average undetectable.

presence of hematopoietically derived cells, IFN- γ was not detected in these specimens. In contrast, both CD45 and IFN- γ were detected in all five B7-H1⁺ tumors. These results strongly support the notion that functional production by TIL of IFN- γ , a cytokine shown to rapidly induce B7-H1 expression by melanoma cells (fig. S3), is an important inducer of B7-H1 in melanomas in vivo.

Association of B7-H1 expression by melanocytes and host infiltrating cells, with melanoma subtypes, Breslow thickness, pT stage, and TNM stage

To gain further insights into the clinical biology of B7-H1 expression in melanoma, we examined the relationship between B7-H1 expression patterns and tumor subtype and stage. Forty-five percent (13 of 29) of superficial spreading and nodular melanomas were B7-H1⁺ (mean melanocyte expression, 16%), whereas other subtypes, including lentigo maligna melanoma ($n = 3$), acral lentiginous melanoma ($n = 7$), and desmoplastic melanoma ($n = 4$), were B7-H1⁻, suggesting an association of B7-H1 expression with histologic subtype ($P = 0.033$, Fisher's exact test).

When the Breslow thickness of primary invasive melanomas was examined, there was no correlation with B7-H1 expression by univariate analysis ($P = 0.23$, Wilcoxon rank-sum test). In addition, for 54 in situ or invasive primary melanoma lesions examined, B7-H1 expression did not vary by pathologic primary tumor stage (T1 to T4) or the highest clinicopathologic TNM stage of the patient during a median 8-year follow-up ($P = 0.43$ and 0.33 , respectively, Fisher's exact

test, one-sided) (table S1), indicating that B7-H1 expression is not linked to melanoma stage. Furthermore, the AIS for primary tumors did not associate with Breslow thickness ($P = 0.30$), pathologic tumor stage, or clinicopathologic TNM stage ($P = 0.64$ and 0.66 , respectively). When B7-H1 expression in metastases was analyzed by anatomic location, there was no association between expression and location in lymph nodes, subcutaneous tissue, lung, or other visceral sites ($P = 0.516$, Fisher's exact test, one-sided).

Association of B7-H1 expression in melanomas with overall patient survival

If indeed B7-H1 expression marked the existence of an active anti-tumor response, it might be predicted that, even though this resistance mechanism ultimately allowed tumors to persist, tumor growth might be partially restrained, leading to improved survival relative to patients with little antitumor response (and hence no B7-H1 expression). We therefore performed a survival analysis for the 99 patients with primary invasive or metastatic melanoma included in our study. As described above, the clinicopathologic factors such as Breslow depth, pathologic T stage, and TNM stage were not associated with B7-H1 expression by univariate analysis; thus, a multivariate analysis was not indicated. Among 43 patients who had primary invasive melanoma, those whose melanoma cells expressed B7-H1 showed no significant difference in overall survival when compared to those with B7-H1⁻ lesions ($P = 0.33$, Fig. 4A). In contrast, there was a significant difference in survival for 56 patients with metastatic melanoma, where B7-H1 expression by tumor cells ($\geq 5\%$) correlated with improved survival ($P = 0.032$, Fig. 4B).

Because of the association between B7-H1 expression and inflammatory infiltrates, we examined the relationship of the AIS to melanoma patient survival. By univariate analysis, patients with primary melanomas with an AIS of ≥ 5 demonstrated no significant difference in survival when compared to those with a score of < 5 ($P = 0.25$, Fig. 4C). However, significantly improved overall survival was noted for patients whose metastases were associated with an AIS of ≥ 5 ($P = 0.014$, Fig. 4D). When the impact of TILs alone on overall survival was determined, a similar result was obtained. Patients with primary melanomas demonstrated no difference in survival as related to the presence or absence of immune infiltrates ($P = 0.32$, Fig. 4E), whereas improved survival was seen in patients with metastatic disease who had TILs present ($P = 0.017$, Fig. 4F). The close parallels among the influence of TILs, the AIS, and tumor cell B7-H1 expression on survival are consistent with the hypothesis that these factors are inter-related and may influence patient outcomes.

The correlation of B7-H1⁺ tumor cells and AIS scores of ≥ 5 with improved survival for patients with metastatic, but not localized, melanoma may reflect the subsequent therapies received by these patients. Among the 56 patients with metastatic melanoma (stage III or IV) in this study, treatment records were available for 53. Forty-three percent of these patients (23 of 53) received some form of systemic immunotherapy [IFN- α , high-dose interleukin-2 (IL-2), vaccine, anti-PD-1 mAb, or combinations thereof] after their diagnostic biopsy demonstrating metastatic disease. Among those patients whose tumors were B7-H1⁻, 8 of 22 (36%) received immunotherapy compared to 15 of 31 (48%) with B7-H1⁺ tumors. These differences were not significant ($P = 0.415$), suggesting that patients with "inflamed" tumors expressing B7-H1 might be more likely to respond to immunotherapy.




Fig. 4. Correlation of overall patient survival with B7-H1 expression in melanomas. (A) There was no significant difference in survival related to melanoma cell B7-H1 expression in 43 patients with primary invasive melanomas ($P = 0.33$). (B) However, among 56 patients with metastatic disease, those whose metastatic tumors expressed B7-H1 had significantly improved survival compared to the B7-H1⁻ cohort ($P = 0.032$). Similar results were seen in analyzing the AIS, representing B7-H1 expression in lymphohistiocytic tumor infiltrates, as well as in analyzing the presence of TILs (mild, moderate, and severe lymphocyte infiltrates with associated histiocytes/macrophages). (C and E) No correlation of AIS or TILs was observed with survival in patients with invasive primary melanomas [$P = 0.25$ (C) and 0.32 (E)]. (D and F) However, there was a significant correlation in patients with metastatic disease [$P = 0.014$ (D) and 0.017 (F)]. B7-H1⁺, $\geq 5\%$ of tumor cells with membranous expression; B7-H1⁻, $< 5\%$ membranous tumor cell expression.

The potential impact of high-aggregate B7-H1 expression on overall survival was examined in 99 patients with invasive or metastatic melanoma lesions. There was no difference in survival for patients whose tumors demonstrated $>20\%$ B7-H1 expression on melanoma cells and/or an AIS of >50 compared to those with lower-aggregate B7-H1 expression ($P = 0.58$). Together, these findings validate our selection of expression thresholds by tumor cells and infiltrating immune cells, and indicate that even focal B7-H1 expression in the immediate environment of metastatic melanoma is associated with improved survival.

DISCUSSION

Our study reveals a highly significant concordance of membranous expression of B7-H1 by nevus and melanoma cells with the presence of immune infiltrates and provides direct evidence of geographic colocalization. Furthermore, expression of the inflammatory cytokine

IFN- γ , known to be a primary cytokine driving B7-H1 expression, was detected specifically at the interface of B7-H1⁺ tumor cells and infiltrating immune cells, but not in B7-H1⁻ tumors. These findings suggest that infiltrating immune cells can produce factors driving B7-H1 expression as a negative feedback mechanism, resulting in what may be considered adaptive immune resistance exerted by the tumor. This mechanism is distinct from the conventional view of immune resistance by either immune escape via mutation of tumor-specific epitopes (31, 32) or constitutive oncogene-driven up-regulation of immune-inhibitory ligands (33). IFN- γ is secreted by human melanoma-specific T lymphocytes, and this cytokine rapidly induces B7-H1 in cultured melanomas. Although our studies associate IFN- γ production with B7-H1⁺ melanomas in vivo, other secreted factors in the tumor microenvironment such as IL-10, IL-6, and common γ -chain cytokines may also be involved (34, 35). These findings help to explain why immunogenic melanoma fails to be controlled by the immune system, even when boosted by antigen-based cancer vaccines, and implicate a new immune escape mechanism in the cancer microenvironment.

The up-regulation of B7-H1 in the cancer microenvironment should not be interpreted simply as the dominance of immune suppression because this up-regulation is also associated with endogenous inflammatory immune responses. The balance of the host's immune response and negative feedback inhibition will therefore determine outcome. We did not

detect a significant correlation between overall survival and B7-H1 expression in patients with primary invasive melanomas. In contrast, we documented a significantly improved overall survival in patients with B7-H1⁺ metastatic lesions. With the exception of one published report in cervical cancer (36), this finding is different from a number of reports from other tumor types that have claimed B7-H1 expression as a negative overall prognostic factor (24–26).

This seemingly paradoxical observation, in which expression of an immunosuppressive molecule correlates with improved outcomes, may be resolved if B7-H1 expression is viewed as reflecting the presence of endogenous antitumor immunity (infiltrating lymphocytes producing cytokines in response to tumor-specific or tumor-selective antigens) among melanoma patients who are candidates for immunotherapy. Indeed, 43% of the patients with metastatic disease in our study received some form of immunotherapy after diagnostic tumor biopsy, including IFN- α , high-dose IL-2, vaccine, PD-1 blockade, or combinations thereof. A growing literature suggests the positive correlation of an inflamed tumor microenvironment with response to

immunotherapies such as cancer vaccines or ipilimumab (37, 38), and our findings further suggest that these immunotherapies should be interpreted as protection or potentiation of ongoing immunity rather than the generation of de novo immunity against cancer. Note that medical center-specific treatment practices may be critical to these observations. Indeed, about one-third of patients with metastatic melanoma show evidence of objective tumor regressions in response to anti-PD-1 therapy (39). Ongoing studies in our laboratories are investigating whether tumor regressions in this series of patients correlate with tumor cell B7-H1 expression and the presence and type of TILs.

Although intratumoral lymphocytes are a common feature in many different types of cancer, the prognostic significance of this finding varies by tumor type. For example, TILs are associated with a better prognosis in patients with colorectal, ovarian, pancreatic, esophageal, and small-cell lung carcinoma (40–44), but are an adverse prognostic feature in patients with renal cell carcinoma (45). Notably, in colon cancer, the type, density, and location of immune infiltrates have been shown to be a better predictor of survival than TNM classification (46). In melanoma, TILs may be seen in all stages of disease, but are not prognostically significant in early stages, that is, in situ or radial growth phase melanomas; they do reach significance in vertical growth phase melanomas and those that have metastasized (47, 48). Our findings provide an interpretation for these seemingly contradictory observations. Although the presence of TILs may indicate immunogenicity of tumors and ongoing immune responses, clinical outcomes may depend on B7-H1 expression in addition to other factors such as composition of infiltration, persistence of inflammation, and the treatment received.

We observed only rare cases showing diffuse tumor cell B7-H1 expression, unrelated or out of proportion to TILs. These may represent melanomas that have developed intrinsic pathways driving B7-H1 gene expression. PTEN (phosphatase and tensin homolog deleted from chromosome 10) deficiency has been shown to support posttranscriptional B7-H1 expression in gliomas (34), but factors driving constitutive expression in melanoma have not been defined. Distinct histologic subtypes of melanoma are associated with characteristic tumorigenic signaling pathways: for example, superficial spreading and nodular melanomas are more likely to have activating BRAF mutations than acral lentiginous melanomas, which are often associated with c-Kit mutations. In our study, 45% of superficial spreading and nodular melanomas expressed B7-H1, whereas other subtypes including acral lentiginous melanoma, lentigo maligna melanoma, and desmoplastic melanoma did not, suggesting the presence of distinct signaling pathways. Studies focused on the transcriptional regulation of B7-H1 expression in nodular and superficial spreading melanomas and metastatic lesions, as well as studies of cytokine expression profiles in microdissected boundary areas between tumor cells and TILs, are currently under way in our laboratories to explore these issues. Notably, the correlation between IFN- γ and B7-H1 expression was 100% among the microdissected samples tested. However, roughly one-third of lesions with observable TIL (on IHC and/or H&E staining) did not express B7-H1, suggesting that a T helper 1 (T_H1) pattern of cytokine expression characterized by IFN- γ may be needed to induce B7-H1 expression.

The clinical significance of B7-H1 expression in melanoma has not yet been firmly established. Two reports have been published to date, with conflicting results (49, 50), perhaps because of different histologic subtypes studied or methodologies used. Gadiot *et al.* used the polyclonal antibody 4059 for IHC detection, scored a 1% threshold of ex-

pression by melanocytes as a positive result, and did not distinguish membranous and cytoplasmic patterns of cellular expression. In that study, most patients had superficial spreading or nodular-type melanomas, and B7-H1 expression by melanocytes was associated with a trend toward better survival. In contrast, Hino *et al.* used mAb clone 27A2 for IHC and compared “high” to “low” cytoplasmic expression based on automated color density measurements. Most patients had the acral lentiginous subtype of melanoma, which is associated with c-Kit mutations, rather than with BRAF mutations, which predominate in cohorts of European descent (51, 52). Hino *et al.* reported high-intensity B7-H1 expression as an independent risk factor for decreased survival. Both studies had about a 4- to 5-year median follow-up time. Neither reported the geographic association of B7-H1 expression with TILs or B7-H1 expression by other cells in the tumor microenvironment.

Both cytoplasmic and membranous B7-H1 staining have been reported in multiple tumor types; however, B7-H1 is a type I transmembrane molecule (6). Cytoplasmic staining may represent intracellular stores of B7-H1, which may be deployed to the cell surface depending on appropriate stimulation. Our previous study describing B7-H1 expression in various human cancers included a small sample size of melanomas ($n = 22$), among which all were deemed at least focally positive in frozen sections (5); however, the previous study assessed both cytoplasmic and membranous positivity, whereas the current study assesses only membranous expression. We hypothesize that cell surface expression of B7-H1 is the most immediately biologically relevant as a potential biomarker predicting clinical response to PD-1 blockade (29).

These findings highlight a therapeutic opportunity in administering agents blocking the PD-1/B7-H1 pathway to patients with metastatic melanoma. B7-H1 is a critical immunomodulating component within the melanoma microenvironment. Expression of B7-H1 on tumor cells at the interface with immune infiltrates, in comparison to constitutive tumor expression in areas devoid of infiltrates, may represent distinct underlying immune resistance mechanisms and warrant further study on the molecular level. A deeper understanding of mechanisms driving the observed expression patterns is essential for developing rational treatment combinations with mAbs blocking PD-1 or B7-H1, which are already in clinical trials, and other potentially synergistic therapeutic agents.

MATERIALS AND METHODS

Case selection

After Johns Hopkins Institutional Review Board approval, 150 melanocytic lesions of various histologic subtypes and stages were identified from 150 individual patients through a search of the Johns Hopkins Hospital surgical pathology archives. Nevi of various histologic subtypes ($n = 40$, including 10 common, 2 blue, 13 halo, 8 Spitz, and 7 dysplastic nevi), melanomas in situ ($n = 11$), invasive melanomas of different histologic subtypes ($n = 43$, including 14 superficial spreading, 15 nodular, 3 lentigo maligna, 4 desmoplastic, and 7 acral lentiginous melanomas), and metastases to various sites ($n = 56$, including 34 lymph node, 8 cutaneous/soft tissue, 6 lung, and 8 other visceral sites) were evaluated.

Clinical and pathologic features

The original histologic diagnosis was confirmed on archival H&E-stained slides, and pathologic features such as histologic subtype,

Breslow thickness, and pathologic T stage were collected. The pathologic T stage is based on microstaging of the primary tumor and includes the parameters of Breslow thickness, ulceration, and mitotic activity. A single representative paraffin block for each case was chosen. Where necessary, cases were restaged to current American Joint Committee on Cancer criteria (53). The highest TNM stage and overall survival data were obtained for each patient. For patients with metastatic disease, information on treatment regimens was gathered. The duration of follow-up was calculated from the date of the diagnostic procedure to the date of cancer progression, last follow-up, or death.

Immunohistochemistry

For the purpose of this retrospective study, archival FFPE specimens were cut into 5- μ m sections and mounted on glass slides. IHC for B7-H1 was performed with a murine anti-human B7-H1 mAb [clone 5H1, isotype mouse IgG1 (immunoglobulin G1)] (5) at a concentration of 2 μ g/ml according to a standard protocol. See the Supplementary Materials and fig. S4 for additional details. IHC for B7-H1 was also performed on select slides with a rabbit polyclonal antibody to B7-H1 designated 4059 (ProSci) as previously described (50). Additional details about 4059 are provided in figs. S5 and S6. IHC for select cell lineage markers including CD3, S100, and CD68 for T cells, melanocytes, and macrophages, respectively, was performed on adjacent 5- μ m sections according to standard automated protocols.

Quantification of B7-H1 expression by melanocytes/melanoma cells and the associated host inflammatory response

Percentages of tumor cells exhibiting a membranous staining pattern for B7-H1 were independently quantified as <5%, 5 to 9%, and then in 10% increments up to 100% by two pathologists (J.M.T. and R.A.A.) who were blinded to patient outcomes. Differences in scoring were adjudicated. The host response to tumor, composed predominantly of lymphocytes with scattered histiocytes (tissue macrophages, dendritic cells, and Langerhans cells), referred to collectively as “immune cell infiltrates,” was also scored for percent B7-H1 membranous expression. The intensity of the infiltrate was graded as none (0), mild (score of 1, rare lymphocytes), moderate (2, focal infiltration of tumor by lymphohistiocytic aggregates), or severe (3, diffuse infiltration). An AIS was defined as the intensity of intratumoral inflammation multiplied by the percent of inflammatory cells expressing B7-H1 (27). B7-H1 expression by inflammatory cells that were not immediately associated with melanocytic cells in a given tumor section was not included in calculating the AIS. Percent B7-H1 expression by tumor cells and the AIS, reflecting B7-H1 in the inflammatory host response, as well as the presence of TILs were correlated with overall patient survival.

LCM and qRT-PCR for IFN- γ

Specimens were cut into 7- μ m sections on Arcturus PEN membrane glass slides (Applied Biosystems). LCM was performed with a Leica LMD6000 Laser Capture Microdissection microscope. In melanoma biopsies that were B7-H1⁺, central areas containing nonnecrotic tumor cells were captured. In tumors that were B7-H1⁺, tumor cells were excised along with associated infiltrating immune cells (fig. S7). RT-PCR was performed with primer/probe preparations for either human IFN- γ or CD45, according to standard protocols. Triplicate C_T averages and SEs of the mean are reported. See the Supplementary Materials for additional details.

Western blotting

537-mel was cultured in the presence or absence of IFN- γ (500 IU/ml) for 3 days. Lysates of whole cells as well as fractions of cytosolic/nuclear proteins and membrane proteins were prepared with M-PER and MEM-PER Reagent, respectively (Pierce). Cell lysates (10 μ g of total protein per lane) were separated by 10% bis-tris SDS-PAGE (polyacrylamide gel electrophoresis) under reducing conditions (Invitrogen). Human IgG1 (Sigma) and recombinant B7 Fc chimeric proteins (B7.1, B7.2, B7-H1, and B7-DC from R&D Systems Inc.) were loaded at 300 ng per lane and resolved by 4 to 12% bis-tris SDS-PAGE under reducing conditions. Proteins were transferred to a polyvinylidene difluoride membrane, which was blocked for 1 hour with 5% milk in tris-buffered saline–Tween 20. Membranes were incubated with either rabbit anti-human B7-H1 polyclonal antibody (ProSci 4059) or mouse anti-human B7-H1 mAb (clone 5H1) (0.5 μ g/ml) at 4°C overnight. The secondary antibody was either anti-rabbit IgG-HRP (horseradish peroxidase) (1:10,000 dilution) or anti-mouse IgG-HRP (1:5000), respectively. Proteins were detected by the ECL Plus Chemiluminescent Detection kit (GE Healthcare).

Statistical analysis

Overall survival was calculated from the date of the diagnostic biopsy to the date of last follow-up or death with the Kaplan-Meier method and analyzed with the log-rank test. Associations of B7-H1 expression and clinicopathologic features were evaluated with Fisher's exact test, Student's *t* tests, and χ^2 tests. Statistical analyses were performed with the STATA V11 software package. All tests were two-sided except as indicated, and *P* values of <0.05 were considered significant.

SUPPLEMENTARY MATERIALS

www.sciencetranslationalmedicine.org/cgi/content/full/4/127/127ra37/DC1

Materials and Methods

Fig. S1. Geographic patterns of CD3⁺ TILs corresponding to the patterns of B7-H1 expression in cases shown in Fig. 1.

Fig. S2. Immunohistochemical characterization of cell types and architecture at the interface of B7-H1 expression and immune infiltrates in a melanoma lesion.

Fig. S3. Kinetics of B7-H1 induction by IFN- γ in cultured human melanoma cells.

Fig. S4. Comparison of B7-H1 detection in fresh and FFPE tissues using mAb 5H1.

Fig. S5. Comparison of B7-H1 detection by the mAb 5H1 versus the polyclonal antibody 4059.

Fig. S6. Comparative specificities of anti-B7-H1 mAb 5H1 and polyclonal antibody 4059 by Western blotting.

Fig. S7. B7-H1⁺ tumor and associated TILs sampled by laser capture microdissection.

Table S1. B7-H1 expression by melanocytes and infiltrating immune cells in 54 in situ and invasive primary melanomas does not correlate with pT or TNM stage.

REFERENCES AND NOTES

1. H. Benlalam, N. Labarrière, B. Linard, L. Derré, E. Diez, M. C. Pandolfino, M. Bonneville, F. Jotereau, Comprehensive analysis of the frequency of recognition of melanoma-associated antigen (MAA) by CD8 melanoma infiltrating lymphocytes (TIL): Implications for immunotherapy. *Eur. J. Immunol.* **31**, 2007–2015 (2001).
2. K. Kamposioras, G. Pentheroudakis, D. Pectasides, N. Pavlidis, Malignant melanoma of unknown primary site. To make the long story short. A systematic review of the literature. *Crit. Rev. Oncol. Hematol.* **78**, 112–126 (2011).
3. W. Zou, L. Chen, Inhibitory B7-family molecules in the tumour microenvironment. *Nat. Rev. Immunol.* **8**, 467–477 (2008).
4. Y. Ishida, Y. Agata, K. Shibahara, T. Honjo, Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J.* **11**, 3887–3895 (1992).

5. H. Dong, S. E. Strome, D. R. Salomao, H. Tamura, F. Hirano, D. B. Flies, P. C. Roche, J. Lu, G. Zhu, K. Tamada, V. A. Lennon, E. Celis, L. Chen, Tumor-associated B7-H1 promotes T-cell apoptosis: A potential mechanism of immune evasion. *Nat. Med.* **8**, 793–800 (2002).
6. H. Dong, G. Zhu, K. Tamada, L. Chen, B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion. *Nat. Med.* **5**, 1365–1369 (1999).
7. G. J. Freeman, A. J. Long, Y. Iwai, K. Bourque, T. Chernova, H. Nishimura, L. J. Fitz, N. Malenkovich, T. Okazaki, M. C. Byrne, H. F. Horton, L. Fouser, L. Carter, V. Ling, M. R. Bowman, B. M. Carreno, M. Collins, C. R. Wood, T. Honjo, Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J. Exp. Med.* **192**, 1027–1034 (2000).
8. Y. Iwai, M. Ishida, Y. Tanaka, T. Okazaki, T. Honjo, N. Minato, Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 12293–12297 (2002).
9. S. T. Topalian, G. J. Weiner, D. M. Pardoll, Cancer immunotherapy comes of age. *J. Clin. Oncol.* **29**, 4828–4836 (2011).
10. F. S. Hodi, S. J. O'Day, D. F. McDermott, R. W. Weber, J. A. Sosman, J. B. Haanen, R. Gonzales, C. Robert, D. Schadendorf, J. C. Hassel, W. Akerly, A. J. van den Eertwegh, J. Lutzky, P. Lorigan, J. M. Vaubel, G. P. Linette, D. Hogg, C. H. Ottensmeier, C. Lebbé, C. Peschel, I. Quirt, J. I. Clark, J. D. Wolchok, J. S. Weber, J. Tian, M. J. Yellin, G. M. Nichol, A. Hoos, W. J. Urba, Improved survival with ipilimumab in patients with metastatic melanoma. *N. Engl. J. Med.* **363**, 711–723 (2010).
11. C. Robert, L. Thomas, I. Bondarenko, S. O'Day, J. W. M. D. C. Garbe, C. Lebbe, J. F. Baurain, A. Testori, J. J. Grob, N. Davidson, J. Richards, M. Maio, A. Hauschild, W. H. Miller Jr., P. Gascon, M. Lotem, K. Harmankaya, R. Ibrahim, S. Francis, T. T. Chen, R. Humphrey, A. Hoos, J. D. Wolchok, Ipilimumab plus dacarbazine for previously untreated metastatic melanoma. *N. Engl. J. Med.* **364**, 2517–2526 (2011).
12. S. G. Downey, J. A. Klapper, F. O. Smith, J. C. Yang, R. M. Sherry, R. E. Royal, U. S. Kammula, M. S. Hughes, T. E. Allen, C. L. Levy, M. Yellin, G. Nichol, D. E. White, S. M. Steinberg, S. A. Rosenberg, Prognostic factors related to clinical response in patients with metastatic melanoma treated by CTL-associated antigen-4 blockade. *Clin. Cancer Res.* **13**, 6681–6688 (2007).
13. K. E. Beck, J. A. Blansfield, K. Q. Tran, A. L. Feldman, M. S. Hughes, R. E. Royal, U. S. Kammula, S. L. Topalian, R. M. Sherry, D. Kleiner, M. Quezada, I. Lowy, M. Yellin, S. A. Rosenberg, J. C. Yang, Enterocolitis in patients with cancer after antibody blockade of cytotoxic T-lymphocyte-associated antigen 4. *J. Clin. Oncol.* **24**, 2283–2289 (2006).
14. P. Waterhouse, J. M. Penninger, E. Timms, A. Wakeham, A. Shahinian, K. P. Lee, C. B. Thompson, H. Griesser, T. W. Mak, Lymphoproliferative disorders with early lethality in mice deficient in *Ctla-4*. *Science* **270**, 985–988 (1995).
15. J. A. Brown, D. M. Dorfman, F. R. Ma, E. L. Sullivan, O. Munoz, C. R. Wood, E. A. Greenfield, G. J. Freeman, Blockade of programmed death-1 ligands on dendritic cells enhances T cell activation and cytokine production. *J. Immunol.* **170**, 1257–1266 (2003).
16. S. C. Liang, Y. E. Latchman, J. E. Buhlmann, M. F. Tomczak, B. H. Horwitz, G. J. Freeman, A. H. Sharpe, Regulation of PD-1, PD-L1, and PD-L2 expression during normal and auto-immune responses. *Eur. J. Immunol.* **33**, 2706–2716 (2003).
17. T. Yamazaki, H. Akiba, H. Iwai, Expression of programmed death 1 ligands by murine T cells and APC. *J. Immunol.* **169**, 5538–5545 (2002).
18. T. J. Curiel, S. Wei, H. Dong, X. Alvarez, P. Cheng, P. Mottram, R. Krzysek, K. L. Knutson, B. Daniel, M. C. Zimmermann, O. David, M. Burrow, A. Gordon, N. Dhurandhar, L. Myers, R. Berggren, A. Hemminki, R. D. Alvarez, D. Emille, D. T. Curiel, L. Chen, W. Zou, Blockade of B7-H1 improves myeloid dendritic cell-mediated antitumor immunity. *Nat. Med.* **9**, 562–567 (2003).
19. M. J. Butte, M. E. Keir, T. B. Phamduy, A. H. Sharpe, G. J. Freeman, Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses. *Immunity* **27**, 111–122 (2007).
20. E. A. Tivol, F. Borriello, A. N. Schweitzer, W. P. Lynch, J. A. Bluestone, A. H. Sharpe, Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity* **3**, 541–547 (1995).
21. H. Dong, G. Zhu, K. Tamada, D. B. Flies, J. M. van Deursen, L. Chen, B7-H1 determines accumulation and deletion of intrahepatic CD8⁺ T lymphocytes. *Immunity* **20**, 327–336 (2004).
22. H. Nishimura, T. Okazaki, Y. Tanaka, K. Nakatani, M. Hara, A. Matsumori, S. Sasayama, A. Mizoguchi, H. Hiari, N. Minato, T. Honjo, Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice. *Science* **291**, 319–322 (2001).
23. D. B. Flies, L. Chen, The new B7s: Playing a pivotal role in tumor immunity. *J. Immunother.* **30**, 251–260 (2007).
24. C. Wu, Y. Zhu, J. Jiang, J. Zhao, X. G. Zhang, N. Xu, Immunohistochemical localization of programmed death-1 ligand-1 (PD-L1) in gastric carcinoma and its clinical significance. *Acta Histochem.* **108**, 19–24 (2006).
25. Y. Ohigashi, M. Sho, Y. Yamada, Y. Tsurui, K. Hamada, N. Ikeda, T. Mizuno, R. Yoriki, H. Kashizuka, K. Yane, F. Tsumima, N. Otsuki, H. Yagita, M. Azuma, Y. Nakajima, Clinical significance of programmed death-1 ligand-1 and programmed death-1 ligand-2 expression in human esophageal cancer. *Clin. Cancer Res.* **11**, 2947–2953 (2005).
26. R. H. Thompson, S. M. Kuntz, B. C. Leibovich, H. Dong, C. M. Lohse, W. S. Webster, Tumor B7-H1 is associated with poor prognosis in renal cell carcinoma patients with long-term follow-up. *Cancer Res.* **66**, 3381–3385 (2006).
27. R. H. Thompson, M. D. Gillett, J. C. Cheville, C. M. Lohse, H. Dong, W. S. Webster, K. G. Krejci, J. R. Lobo, S. Sengupta, L. Chen, H. Zinke, M. L. Blute, S. E. Strome, B. C. Leibovich, E. D. Kwon, Costimulatory B7-H1 in renal cell carcinoma patients: Indicator of tumor aggressiveness and potential therapeutic target. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 17174–17179 (2004).
28. H. Ghebeh, S. Mohammed, A. Al-Omar, A. Qattan, C. Lehe, G. Al-Qudahi, N. Elkum, M. Alshabanah, S. Bin Amer, A. Tulbah, D. Ajarim, T. Al-Tweigeri, S. Dermime, The B7-H1 (PD-L1) T lymphocyte-inhibitory molecule is expressed in breast cancer patients with infiltrating ductal carcinoma: Correlation with important high-risk prognostic factors. *Neoplasia* **8**, 190–198 (2006).
29. J. R. Brahmer, C. G. Drake, I. Wollner, J. D. Powderly, J. Picus, W. H. Sharfman, E. Stankevich, A. Pons, T. M. Salay, T. L. McMiller, M. M. Gilson, C. Wang, M. Selby, J. M. Taube, R. Anders, L. Chen, A. J. Korman, D. M. Pardoll, I. Lowy, S. L. Topalian, Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: Safety, clinical activity, pharmacodynamics, and immunologic correlates. *J. Clin. Oncol.* **28**, 3167–3175 (2010).
30. R. H. Thompson, H. Dong, E. D. Kwon, Implications of B7-H1 expression in clear cell carcinoma of the kidney for prognostication and therapy. *Clin. Cancer Res.* **13**, 709s–715s (2007).
31. R. D. Schreiber, L. J. Old, M. J. Smyth, Cancer immunoediting: Integrating immunity's roles in cancer suppression and promotion. *Science* **331**, 1565–1570 (2011).
32. G. P. Dunn, A. T. Bruce, H. Ikeda, L. J. Old, R. D. Schreiber, Cancer immunoediting: From immunosurveillance to tumor escape. *Nat. Immunol.* **3**, 991–998 (2002).
33. A. T. Parsa, J. S. Waldron, A. Panner, C. A. Crane, I. F. Parney, J. J. Barry, Loss of tumor suppressor PTEN function increases B7-H1 expression and immunoresistance in glioma. *Nat. Med.* **13**, 84–88 (2007).
34. S. J. Wölflle, J. Strebovsky, H. Bartz, A. Sähr, C. Arnold, C. Kaiser, A. H. Dalpke, K. Heeg, PD-L1 expression on tolerogenic APCs is controlled by STAT-3. *Eur. J. Immunol.* **41**, 413–424 (2011).
35. A. L. Kinter, E. J. Godbout, J. P. McNally, I. Sereti, G. A. Roby, M. A. O'Shea, A. S. Fauci, The common γ -chain cytokines IL-2, IL-7, IL-15, and IL-21 induce the expression of programmed death-1 and its ligands. *J. Immunol.* **181**, 6738–6746 (2008).
36. R. Karim, E. S. Jordanova, S. J. Piersma, G. G. Kenter, L. Chen, J. M. Boer, C. J. Melief, S. H. van der Burg, Tumor-expressed B7-H1 and B7-DC in relation to PD-1+ T-cell infiltration and survival of patients with cervical carcinoma. *Clin. Cancer Res.* **15**, 6341–6347 (2009).
37. T. F. Gajewski, J. Louahed, V. G. Brichard, Gene signature in melanoma associated with clinical activity: A potential clue to unlock cancer immunotherapy. *Cancer J.* **16**, 399–403 (2010).
38. R. R. Ji, S. D. Chasalow, L. Wang, O. Hamid, H. Schmidt, J. Cogswell, S. Alaparthi, D. Berman, M. Jure-Kunkel, N. O. Siemers, J. R. Jackson, V. Shahabi, An immune-active tumor micro-environment favors clinical response to ipilimumab. *Cancer Immunol. Immunother.* **10.1007/s00262-011-1172-6** (2011).
39. M. Sznol, J. D. Powderly, D. C. Smith, J. R. Brahmer, C. G. Drake, D. F. McDermott, D. P. Lawrence, J. D. Wolchok, S. L. Topalian, I. Lowy, Safety and antitumor activity of biweekly MDX-1106 (anti-PD-1, BMS-936558/ONO-4538) in patients with advanced refractory malignancies [abstract 2506]. *J. Clin. Oncol. (Meet. Abstr.)* **28**, 15s (2010).
40. L. Zhang, J. R. Conejo-Garcia, D. Katsaros, P. A. Gimotty, M. Massobrio, G. Regnani, A. Makrigiannakis, H. Gray, K. Schlienger, M. N. Liebman, S. C. Rubin, G. Coukos, Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N. Engl. J. Med.* **348**, 203–213 (2003).
41. A. Fukunaga, M. Miyamoto, Y. Cho, S. Murakami, Y. Kawarada, T. Oshikiri, K. Kato, T. Kurokawa, M. Suzuki, Y. Nakakubo, K. Hiraoka, T. Itoh, T. Morikawa, S. Okushiba, S. Kondo, H. Katoh, CD8⁺ tumor-infiltrating lymphocytes together with CD4⁺ tumor-infiltrating lymphocytes and dendritic cells improve the prognosis of patients with pancreatic adenocarcinoma. *Pancreas* **28**, e26–e31 (2004).
42. A. K. Eerola, Y. Soini, P. Pääkkö, A high number of tumor-infiltrating lymphocytes are associated with a small tumor size, low tumor stage, and a favorable prognosis in operated small cell lung carcinoma. *Clin. Cancer Res.* **6**, 1875–1881 (2000).
43. K. Schumacher, W. Haensch, C. Röelfsaad, P. M. Schlag, Prognostic significance of activated CD8⁺ T cell infiltrations within esophageal carcinomas. *Cancer Res.* **61**, 3932–3936 (2001).
44. F. Pagès, A. Berger, M. Camus, F. Sanchez-Cabo, A. Costes, R. Molitor, B. Mlecnik, A. Kirilovskiy, M. Nilsson, D. Damotte, T. Meatchi, P. Bruneval, P. H. Cugnenc, Z. Trajanoski, W. H. Fridman, J. Galon, Effector memory T cells, early metastasis, and survival in colorectal cancer. *N. Engl. J. Med.* **353**, 2654–2666 (2005).
45. W. S. Webster, C. M. Lohse, R. H. Thompson, H. Dong, X. Frigola, D. L. Dicks, S. Sengupta, I. Frank, B. C. Leibovich, M. L. Blute, J. C. Cheville, E. D. Kwon, Mononuclear cell infiltration in clear-cell renal cell carcinoma independently predicts patient survival. *Cancer* **107**, 46–53 (2006).
46. J. Galon, A. Costes, F. Sanchez-Cabo, A. Kirilovskiy, B. Mlecnik, C. Lagorce-Pagès, M. Tosolini, M. Camus, A. Berger, P. Wind, F. Zinzindohoué, P. Bruneval, P. H. Cugnenc, Z. Trajanoski, W. H. Fridman, F. Pagès, Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* **313**, 1960–1964 (2006).
47. C. G. Clemente, M. C. Mihm Jr., R. Bufalino, S. Zurrida, P. Collini, N. Cascinelli, Prognostic value of tumor infiltrating lymphocytes in the vertical growth phase of primary cutaneous melanoma. *Cancer* **77**, 1303–1310 (1996).
48. M. C. Mihm Jr., C. G. Clemente, N. Cascinelli, Tumor infiltrating lymphocytes in lymph node melanoma metastases: A histopathologic prognostic indicator and an expression of local immune response. *Lab. Invest.* **74**, 43–47 (1996).

49. R. Hino, K. Kabashima, Y. Kato, H. Yagi, M. Nakamura, T. Honjo, T. Okazaki, Y. Tokura, Tumor cell expression of programmed cell death-1 ligand 1 is a prognostic factor for malignant melanoma. *Cancer* **116**, 1757–1766 (2010).
50. J. Gadiot, A. I. Hooijkaas, A. D. M. Kaiser, H. van Tinteren, H. van Boven, C. Blank, Overall survival and PD-L1 expression in metastasized malignant melanoma. *Cancer* **117**, 2192–2201 (2011).
51. J. A. Curtin, J. Fridlyand, T. Kageshita, H. N. Patel, K. J. Busam, H. Kutzner, K. H. Cho, S. Aiba, E. B. Bröcker, P. E. LeBoit, D. Pinkel, B. C. Bastian, Distinct sets of genetic alterations in melanoma. *N. Engl. J. Med.* **353**, 2135–2147 (2005).
52. J. A. Curtin, K. Busam, D. Pinkel, B. C. Bastian, Somatic activation of KIT in distinct subtypes of melanoma. *J. Clin. Oncol.* **24**, 4340–4346 (2006).
53. C. M. Balch, J. E. Gershenwald, S. J. Soong, J. F. Thompson, M. B. Atkins, D. R. Byrd, A. C. Buzaid, A. J. Cochran, D. G. Coit, S. Ding, A. M. Eggermont, K. T. Flaherty, P. A. Gimotty, J. M. Kirkwood, K. M. McMasters, M. C. Mihm Jr., D. L. Morton, M. I. Ross, A. J. Sober, V. K. Sondak, Final version of 2009 AJCC melanoma staging and classification. *J. Clin. Oncol.* **27**, 6199–6206 (2009).

Acknowledgments: We thank R. Li, C. Umbricht, Y. Liu, and F. Housseau for helpful discussions, as well as J. Califano and J. Handa for their support (all Johns Hopkins University School of Medicine). **Funding:** This study was supported by grants from the National Cancer Institute (CA016359, CA97085, and CA85721 to L.C.), the NIH (R01DK080736 and R01DK081417 to R.A.A.), the Melanoma

Research Alliance (to D.M.P., S.L.T., and L.C.), the Barney Family Foundation (to S.L.T.), the Michael Rolfe Foundation for Pancreatic Cancer Research (to R.A.A.), and the Dermatology Foundation (to J.M.T.). **Author contributions:** J.M.T., R.A.A., G.D.Y., S.C., D.M.P., S.L.T., and L.C. conceived and designed the experiments. J.M.T., R.A.A., G.D.Y., H.X., R.S., T.L.M., and S.C. performed the experiments. J.M.T., R.A.A., G.D.Y., T.L.M., S.C., A.P.K., D.M.P., S.L.T., and L.C. analyzed the data. J.M.T., D.M.P., S.L.T., and L.C. wrote the manuscript. **Competing interests:** S.L.T. is a consultant to (un-compensated) and S.L.T., J.M.T., R.A.A., and H.X. receive research support from Bristol-Myers Squibb. The 5H1 antibody (made in L.C.'s laboratory, U.S. 7,797,710 and U.S. 7,892,540) will be distributed via a standard University Material Transfer Agreements for research, which is consistent with *Science Translational Medicine's* material sharing policy. The other authors declare that they have no competing interests.

Submitted 9 January 2012

Accepted 23 February 2012

Published 28 March 2012

10.1126/scitranslmed.3003689

Citation: J. M. Taube, R. A. Anders, G. D. Young, H. Xu, R. Sharma, T. L. McMiller, S. Chen, A. P. Klein, D. M. Pardoll, S. L. Topalian, L. Chen, Colocalization of inflammatory response with B7-H1 expression in human melanocytic lesions supports an adaptive resistance mechanism of immune escape. *Sci. Transl. Med.* **4**, 127ra37 (2012).

Colocalization of Inflammatory Response with B7-H1 Expression in Human Melanocytic Lesions Supports an Adaptive Resistance Mechanism of Immune Escape

Janis M. Taube, Robert A. Anders, Geoffrey D. Young, Haiying Xu, Rajni Sharma, Tracee L. McMiller, Shuming Chen, Alison P. Klein, Drew M. Pardoll, Suzanne L. Topalian, and Lieping Chen

Sci. Transl. Med. 4 (127), . DOI: 10.1126/scitranslmed.3003689

The Great Escape

In the movie *The Great Escape*, “problem” prisoners with multiple escape attempts are put in an “escape-proof” POW camp, where they use their cleverness and specialized skills to outwit their captors. However, when it comes to escaping, even Steve McQueen doesn’t have anything on cancer cells. Although human cancers express tumor antigens recognized by the immune system, host immune responses often fail to control tumor growth. Taube *et al.* now explain one way in which tumor cells may adapt to escape from immune surveillance.

The researchers found a strong association between expression of the immune-inhibitory molecule B7-H1 (PD-L1) on melanocytes and immune cell infiltration into tumors in patients with different stages of melanoma. The B7-H1+ melanocytes were found directly adjacent to the immune cells, with interferon- γ detected at the melanocyte–immune cell interface. Interferon- γ , which is secreted by the immune cells, induces B7-H1 expression; thus, the tumor may adapt by causing immune cells to trigger their own inhibition. Indeed, patients with B7-H1+ metastatic melanoma had prolonged overall survival when compared with B7-H1 γ metastatic melanoma patients, perhaps suggesting that B7-H1 expression by the tumors is stimulated by a more successful immune response. It remains to be seen whether blocking B7-H1 in these patients will further improve survival. But it is clear that for both prisoners and tumors, adaptation is the key to escape.

View the article online

<https://www.science.org/doi/10.1126/scitranslmed.3003689>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

Science Translational Medicine (ISSN 1946-6242) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science Translational Medicine* is a registered trademark of AAAS.

Copyright © 2012, American Association for the Advancement of Science

Supplementary Materials for

Colocalization of Inflammatory Response with B7-H1 Expression in Human Melanocytic Lesions Supports an Adaptive Resistance Mechanism of Immune Escape

Janis M. Taube,* Robert A. Anders, Geoffrey D. Young, Haiying Xu, Rajni Sharma, Tracee L. McMiller, Shuming Chen, Alison P. Klein, Drew M. Pardoll, Suzanne L. Topalian,* Lieping Chen*

*To whom correspondence should be addressed. E-mail: jtaube1@jhmi.edu (J.M.T.); stopali1@jhmi.edu (S.L.T.); lieping.chen@yale.edu (L.C.)

Published 28 March 2012, *Sci. Transl. Med.* **4**, 127ra37 (2012)
DOI: 10.1126/scitranslmed.3003689

The PDF file includes:

Materials and Methods

Fig. S1. Geographic patterns of CD3⁺ TILs corresponding to the patterns of B7-H1 expression in cases shown in Fig. 1.

Fig. S2. Immunohistochemical characterization of cell types and architecture at the interface of B7-H1 expression and immune infiltrates in a melanoma lesion.

Fig. S3. Kinetics of B7-H1 induction by IFN- γ in cultured human melanoma cells.

Fig. S4. Comparison of B7-H1 detection in fresh and FFPE tissues using mAb 5H1.

Fig. S5. Comparison of B7-H1 detection by the mAb 5H1 versus the polyclonal antibody 4059.

Fig. S6. Comparative specificities of anti-B7-H1 mAb 5H1 and polyclonal antibody 4059 by Western blotting.

Fig. S7. B7-H1⁺ tumor and associated TILs sampled by laser capture microdissection.

Table S1. B7-H1 expression by melanocytes and infiltrating immune cells in 54 in situ and invasive primary melanomas does not correlate with pT or TNM stage.

Supplementary Materials and Methods

Immunohistochemistry (IHC)

We showed previously that B7-H1 protein could be detected on the cell surface of various cancers by IHC on frozen tissue sections with the mAb 5H1 [5]. For archived formalin-fixed paraffin embedded (FFPE) slides, however, an additional re-naturation step is required for this mAb [27]. We compared the staining pattern and sensitivity of these two approaches. To do so, immunodeficient NOD-SCID IL2Rgamma^{null} mice were inoculated with B7-H1-transfected human 624-mel cultured melanoma cells to induce progressively growing tumors. Tumor nodules were harvested and bisected: half of the material was embedded in OCT compound and frozen, and the other half was fixed in formalin and embedded in paraffin. As shown in Figure S4, similar membrane staining patterns were observed in both frozen and FFPE sections. Although the sensitivity of the methods is generally comparable, we observed slightly less intensity for staining in FFPE than frozen sections.

For the purpose of this retrospective study, archival FFPE specimens were cut into 5µm sections and mounted on glass slides. IHC for B7-H1 was performed using a murine anti-human B7-H1 mAb (clone 5H1, isotype mouse IgG1) [5] at a concentration of 2 µg/ml according to a standard protocol (See Supplementary Material). After deparaffinization, antigen retrieval was performed using a Tris-EDTA buffer, pH 9.0 at 120°C for 10 minutes in a Decloaking Chamber (Biocare Medical). Endogenous peroxidase, biotin and proteins were blocked (CAS system K1500, DAKO; Avidin/biotin Blocking Kit, SP-2001, Vector Laboratories; Serotec Block ACE). The primary antibody was applied and allowed to incubate at 4°C for 20 hours. Secondary antibody (biotinylated anti-mouse IgG1, 553441 BD) at a concentration of 1 µg/ml was applied for 30 minutes at room temperature. The signal was then developed with amplification according to the manufacturer's protocol (CAS system K1500, DAKO). Sections were counterstained with hematoxylin, dehydrated in graded ethanol and cleared in xylene, and a coverslip was applied.

Positive control specimens for B7-H1 IHC were created by transfecting cultured human melanoma 624-mel cells [5], which do not constitutively express B7-H1, with a recombinant plasmid encoding full length human B7-H1. Cell surface B7-H1 expression on cultured cells

was confirmed with flow cytometry. NOD-SCID IL2Rgamma^{null} mice were inoculated with the B7-H1-transfected 624-mel cells. Xenografted tumors were harvested and preserved as both fresh-frozen and paraffin-embedded specimens. A B7-H1 fusion protein was used to competitively block binding of the 5H1 mAb to tissue sections, confirming the specificity of 5H1 (Figure S4C). Animal studies were approved by the Institutional Review Board at Johns Hopkins University and adhered to the NIH Guidelines for the Care and Use of Laboratory Animals.


IHC for B7-H1 was also performed on select slides using a rabbit polyclonal Ab to B7-H1 designated 4059 (ProSci) as previously described [50]. When we compared the staining pattern of mAb 5H1 and polyclonal Ab 5049 on FFPE specimens, 5H1 staining showed a clear membranous staining pattern, whereas Ab 4059 demonstrated diffuse cytoplasmic staining (Figure S5). Non-specific isotype controls using purified mouse IgG1 or rabbit IgG1 were performed for each case stained with either the 5H1 mAb or 4059 polyclonal Ab, respectively. Western blot analysis indicated that both antibodies specifically bound a purified human recombinant B7-H1/IgG fusion protein (Figure S6A), but in lysates of melanoma cells the Ab 4059 detected multiple proteins in addition to a 50 kDa protein corresponding to the expected mass of glycosylated B7-H1 (Figure S6B). Therefore, mAb 5H1 was used for all IHC experiments in the current study.

Laser capture microdissection (LCM) and quantitative RT-PCR for IFN- γ

Formalin-fixed human melanoma biopsy specimens and cultured cell pellets were embedded in paraffin blocks. Cultured cells included the human melanoma cell line 1359-mel, and isolated normal donor CD3⁺ T cells (Pan T Cell Isolation Kit II, Miltenyi Biotech) that were activated for 48 hr by anti-CD3 and -CD28 mAbs (1 ug/ml each). Fresh normal donor peripheral blood mononuclear cells (PBMCs), including lymphocytes and monocytes, were isolated from leukapheresis specimens by density centrifugation. Cells in suspension were collected by centrifugation, washed twice with phosphate-buffered saline (PBS), resuspended in 50-100 ul of PBS, dispensed atop solidified 2% agarose plugs in microcentrifuge tubes, and spun for 5-7 min at 1500 rpm. Formalin was slowly added to each pellet, and samples were allowed to sit at room temperature for at least 24 hr before embedding in paraffin blocks.


Specimens were cut into 7µm sections on Arcturus PEN membrane glass slides (Applied Biosystems). Slides were de-paraffinized and stained with a modified hematoxylin and eosin (H&E) protocol. LCM was performed using a Leica LMD6000 Laser Capture Micro-dissection microscope. In melanoma biopsies that were B7-H1(-), central areas containing non-necrotic tumor cells were dissected with intent to avoid surrounding stromal elements. In tumors that were B7-H1+, areas of expression were identified by IHC on neighboring sections and tumor cells were excised along with associated infiltrating immune cells (Figure S7). Total RNA was isolated from LCM specimens using the High Pure RNA Paraffin Kit (Roche) with the following modifications: residual PEN membrane pieces were removed by centrifugation after adding binding buffer, and a DNase incubation step was added to remove contaminating genomic DNA. Total RNA concentration was determined using a NanoDrop spectrophotometer. Fifty ng of total RNA from each specimen was reverse-transcribed in a 20 ul reaction volume using qScript cDNA SuperMix (Quanta Biosciences) per protocol. One ul from each RT reaction was combined with 5ul TaqMan Universal Master Mix II (Applied Biosystems), 3.5ul Molecular Grade Water, and 0.5 ul of primer/probe preparations specific for either human IFN-γ (Applied Biosystems), or CD45 (PTPRC) to standardize to myeloid-derived cells (Applied Biosystems). PCR reactions were run in triplicate on 384-well plates using a 7900 HT Fast Real Time PCR system (Applied Biosystems). Cycle thresholds were determined using a manual cutoff of 0.04. Triplicate C_T averages and standard errors of the mean are reported.

Supplementary Figure 1




Supplementary Figure 1. Geographic patterns of CD3⁺ TILs corresponding to the patterns of B7-H1 expression in cases shown in Fig. 1. (A) Rare CD3⁺ TILs identified by IHC in a benign nevus that was B7-H1(-), indicated by arrows. 200x original magnification (size bar 50 um). (B) A 'moderate' infiltrate of CD3⁺ TILs at the advancing edge of an invasive primary melanoma, nodular histologic subtype, associated with B7-H1+ tumor cells (see Figure 1B). 40x original magnification (size bar 200 um). (C) 400x original magnification (20 um size bar) of the boxed area shown in panel B. (D) Only singular CD3⁺ TILs were observed, predominantly in perivascular areas and not infiltrating among tumor cells, in a diffusely B7-H1+ melanoma metastasis. 200x original magnification (size bar 50 um).

Supplementary Figure 2




Supplementary Figure 2. Immunohistochemical characterization of cell types and architecture at the interface of B7-H1 expression and immune infiltrates in a melanoma lesion. All panels 400x original magnification (size bar 20 μ m). B7-H1 expression is present on the surface of tumor cells (long arrow) and infiltrating lymphocytes (short arrow). S100 immunostaining highlights melanoma cells aggregating in nests. CD68 highlights macrophages/histiocytes accompanying TILs. CD3+ TILs comprise the majority of cells in the immune infiltrate. Lymphocytes are distinguished from melanoma cells by much smaller size and nuclear homogeneity. The nesting architecture of S100+ melanoma cells is distinguished from the singular arrangement of CD3+ lymphocytes and CD68+ histiocytes.

Supplementary Figure 3




Supplementary Figure 3. Kinetics of B7-H1 induction by IFN- γ in cultured human melanoma cells. The cultured cell line 1102-mel was incubated in the presence of 0, 100, or 500 international units (IU)/ml of IFN- γ (Biogen) for the indicated time periods. Then, adherent cells were harvested with trypsin at timed intervals, and B7-H1 expression was determined by flow cytometry with the anti-human B7-H1 mAb MIH1 (eBioscience Inc.). These results show induction of B7-H1 expression as early as 6 hr after IFN- γ exposure in vitro.

Supplementary Figure 4




Supplementary Figure 4. Comparison of B7-H1 detection in fresh and FFPE tissues using mAb 5H1. NOD-SCID IL2Rgamma^{null} mice were inoculated with B7-H1 transfected 624-mel cells. Tumor was harvested and bisected: half was embedded in OCT compound and frozen, and half was formalin-fixed and paraffin-embedded. IHC for B7-H1 using the 5H1 mAb shows a similar expression pattern in both (A) frozen and (B) paraffin-embedded tissues. Prevalence of cell surface staining and staining intensity in the paraffin-embedded section is slightly less than in the frozen specimen. (C) B7-H1 fusion protein was used to competitively block binding of the 5H1 mAb, confirming specificity for B7-H1. 400x original magnification (size bar 20 um), all panels.

Supplementary Figure 5




Supplementary Figure 5. Comparison of B7-H1 detection by the mAb 5H1 versus the polyclonal antibody 4059. IHC for B7-H1 was performed using mAb 5H1 or polyclonal Ab 4059, as described in Methods, on FFPE specimens of xenografted B7-H1 transfected 624-mel. (A) Predominantly membranous staining was detected with the 5H1 mAb. (B) Ab 4059 showed diffuse cytoplasmic staining with rare, focal membranous staining. Isotype staining controls using (C) purified mouse IgG1 or (D) rabbit IgG1 were performed for 5H1 or 4059, respectively. 400x original magnification (size bar 20 μ m), all panels. Evaluation of cell surface expression by IHC mandates employing a mAb such as 5H1, which is capable of detecting cell surface B7-H1 in FFPE specimens. The 5H1 mAb cannot stain FFPE sections without a protein re-naturing process, indicating that 5H1 binds a conformation-dependent epitope on B7-H1.

Supplementary Figure 6A




Supplementary Figure 6B



Supplementary Figure 6. Comparative specificities of anti-B7-H1 mAb 5H1 and polyclonal antibody 4059 by Western blotting. (A) Both antibodies are specific for a 70 kDa recombinant human B7-H1/IgG fusion protein. Lane 1, human IgG1; lane 2, hB7.1/IgG; lane 3, hB7.2/IgG; lane 4, hB7-H1/IgG; lane 5, hB7-DC/IgG. (B) In melanoma cell lysates, mAb 5H1 specifically detects a 50 kDa band consistent with glycosylated B7-H1, whereas polyclonal Ab 4059 detects additional proteins of various molecular weights, which were not further characterized. Lane 1, cultured human melanoma 537-mel, whole cell lysate; lane 2, membrane fraction of lysate; lane 3, cytosolic/nuclear fraction of lysate; lanes 4-6, same as lanes 1-3, performed on 537-mel cells cultured with IFN- γ 500 IU/ml x 72 hr.

Supplementary Figure 7



Supplementary Figure 7. B7-H1⁺ tumor and associated TILs sampled by laser capture microdissection. In tumors that were B7-H1⁺, (A) areas of expression were identified by IHC, and on neighboring sections, areas of tumor cells along with associated infiltrating immune cells were demarcated (B) and excised (C). Size bar 500 um for all panels.

Supplementary Table 1. B7-H1 expression by melanocytes and infiltrating immune cells in 54 in situ and invasive primary melanomas does not correlate with pT or TNM stage

	B7-H1(-)	B7-H1(+)	Adjusted inflam. Score <5	Adjusted inflam. Score ≥5
pT Stage				
pis (n=11)	5	6	5	6
1a (n=11)	6	5	3	8
2a (n=3)	2	1	1	2
2b (n=1)	1	0	0	1
3a (n=8)	6	2	6	2
3b (n=4)	3	1	3	1
4a (n=10)	6	4	3	7
4b (n=6)	6	0	4	2
pT1 and pT2 (n=15)	9	6	4	11
pT3 and pT4 (n=28)	21	7	16	12
TNM stage				
0 (n=11)	5	6	5	6
IA (n=11)	6	5	3	8
IB (n=3)	2	1	1	2
IIA (n=7)	6	1	5	2
IIB (n=9)	7	2	6	3
IIIA (n=2)	1	1	1	1
IIIB (n=5)	4	1	2	3
IIIC (n=4)	2	2	1	3
IV (n=2)	2	0	1	1